

Soft pattern of gravitational Rutherford scattering from heavy target mass expansion*

Yu Jia^{1,2†} Jichen Pan (潘济陈)^{1,2‡} Jia-Yue Zhang^{3,1§}

¹Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China

²School of Physics, University of Chinese Academy of Sciences, Beijing 100049, China

³Theory Center, Jefferson Lab, Newport News, Virginia 23606, USA

Abstract: We investigate the soft behavior of the tree-level Rutherford scattering processes mediated via t -channel one-graviton exchange. We consider two types of Rutherford scattering processes, *e.g.*, a low-energy massless structureless projectile (up to spin-1) hits a static massive composite particle carrying various spins (up to spin-2), and a slowly-moving light projectile hits a heavy static composite target. The unpolarized cross sections in the first type are found to exhibit universal forms at the first two orders in $1/M$ expansion, yet differ at the next-to-next-to-leading order, though some terms at this order still remain universal or depend on the target spin in a definite manner. The unpolarized cross sections in the second type are universal at the lowest order in projectile velocity expansion and through all orders in $1/M$, independent of the spins of both projectile and target. The universality partially breaks down at relative order- v^2/M^2 , albeit some terms at this order still depend on the target spin in a specific manner.

Keywords: Rutherford scattering, Form factor, Soft limit, High spin

DOI:

I. INTRODUCTION

As is taught in virtually every quantum field theory textbook, a generic tree-level QED process with emission of a low-energy photon exhibits simplifying feature [1]. In the soft limit, the full QED amplitude can be factorized into the product of the simpler one with the external photon removed times a universal eikonal factor. The universal pattern governing the emission of soft photon can be readily carried over to the case of emission of soft graviton [2], and can also be extended through the next-to-leading-order (NLO) in the small k expansion (where k^μ denotes the four-momentum of the emitted photon or graviton), which is generically referred to as the LBK theorem [3–5]. Recently there has been attempt to reproduce the LBK theorem entailing soft graviton emission from the perspective of the soft-collinear effective theory [6].

The LBK theorem only applies to the case of on-shell photon/graviton emitted from the external legs, in which the matter fields correspond to structureless point-like

particles. Nevertheless, it is also of theoretical curiosity about the soft pattern of the processes involving composite particles and the soft photon/graviton emerging in the internal line, in which the LBK theorem is no longer applicable. On the physical ground, one anticipates that in a process entailing a heavy composite particle, the soft limit implicates that the very long wavelength of the (real or virtual) photon/graviton is unable to resolve the detailed internal structure of the composite particle, so its properties can be simply summarized in terms of a few low order multipoles, correspondingly the expanded cross sections might exhibit some simple textures. A classical example is the soft limit of the Compton scattering, with the leading contribution represented by the Thomson cross section, depending only on the total electric charge of the composite target, while the NLO contribution in $1/M$ expansion becomes sensitive to its magnetic dipole [7, 8]. Another illuminating example is the Rutherford scattering process, where a low-energy projectile bombards a static, heavy, composite target particle bearing arbitrary spin, mediated by a t -channel photon exchange. Recently

Received 30 April 2024; Accepted 13 July 2024

* This work is supported in part by the National Natural Science Foundation of China under Grants No. 11925506. The work of J.-Y. Z. is also supported in part by the US Department of Energy (DOE) Contract No. DE-AC05-06OR23177, under which Jefferson Science Associates, LLC operates Jefferson Lab.

[†] E-mail: jjay@ihep.ac.cn

[‡] E-mail: panjichen@ihep.ac.cn

[§] E-mail: jzhang@jlab.org



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Article funded by SCOAP³ and published under licence by Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd

the soft limit of the electromagnetic Rutherford scattering process has been comprehensively investigated by two of the authors, and some simple patterns about the target spin dependence have been identified upon performing the heavy target mass expansion [10].

The central theme of this work is to extend the preceding analysis in electromagnetic case [10] to the gravitational Rutherford scattering process, *i.e.*, a structureless point-like projectile bombs on a static, heavy, composite target particle bearing arbitrary spins, now with the graviton being the force carrier. We consider two types of benchmark gravitational Rutherford scattering processes, *e.g.*, a low-energy massless structureless projectile hits a static massive composite particle with spin up to 2, and a non-relativistic light projectile bombs on a heavy static composite target. The major observation of this work is similar to what is found in its electromagnetic counterpart [10]: the unpolarized cross sections in the first type are universal (independent of the target spin) at the first two orders in $1/M$ expansion, yet differ at the next-to-next-to-leading order (NNLO), though some terms at this order still remain universal or depend on the target spin in a definite manner. The unpolarized cross sections in the second type are universal at the lowest order in projectile velocity expansion and through all orders in $1/M$, insensitive to both projectile and target's spin. The universality partially breaks down at relative order- v^2/M^2 , though some terms at this order are still universal or depend on the target spin in a recognizable manner.

The rest of the paper is distributed as follows. In Sec. II, we present the expression of the tree-level amplitude for gravitational Rutherford scattering process involving a heavy composite spinning target particle, and specify the parametrization of the gravitational form factor of massive target particle carrying various spin. In Sec. III, we consider the low-energy massless point-like projectile with spin-0, $\frac{1}{2}$ and 1, striking on a heavy composite target particle with spin ranging from 0 to 2. We organize the unpolarized cross section in the heavy target mass expansion up to NNLO, and identify some universal pattern about the target spin dependence. In Sec. IV, we consider another type of gravitational Rutherford scattering process, where the projectile is replaced by a slowly-moving light structureless particle with spin ranging from 0 to 1. We identify some universal pattern of the cross section in the double expansion of the projectile velocity and $1/M$. We summarize in Sec. V. In Appendix A, we present the spin sum formula for the target particles with various spin. In Appendix B, we tackle the gravitational Rutherford scattering based on the heavy black hole effective theory (HBET), taking the massless spinless projectile and heavy spinless target particle as example. The reason why the NLO amplitude vanishes becomes transparent from the perspective of effective field theory.

II. AMPLITUDE OF GRAVITATIONAL RUTHERFORD SCATTERING INVOLVING A HEAVY COMPOSITE TARGET PARTICLE

In general relativity, the energy-momentum tensor of the matter field plays the role of the gravitational sources. The symmetric Belinfante-Rosenfeld energy-momentum tensor is defined as

$$T_{\mu\nu} = \frac{2}{\sqrt{-g}} \frac{\delta S}{\delta g^{\mu\nu}}, \quad (1)$$

where S denotes the gravitational action of the matter field, $g^{\mu\nu}(x)$ denotes the spacetime metric field. In the weak gravity case, one approximates $g_{\mu\nu}(x) = \eta_{\mu\nu} + \kappa h_{\mu\nu}(x)$ with $\eta_{\mu\nu} = \text{diag}(1, -1, -1, -1)$ being the Minkowski metric. Here $\kappa = \sqrt{32\pi G_N}$, with Newton's constant $G_N = 6.709 \times 10^{-39} \text{ GeV}^{-2}$. In the linearized approximation, the matter field couples with the graviton simply through the interaction $\mathcal{L}_{int} = \frac{\kappa}{2} h_{\mu\nu} T^{\mu\nu}$.

In this work, we focus on the gravitational Rutherford scattering process $I(k)N(p) \rightarrow I(k')N(p')$, where I and N represents a point-like projectile and a heavy composite target particle, respectively. We are working in the laboratory frame where the target particle is at rest, so that the four-momentum of the target particle in the initial state becomes $p^\mu = (M, \mathbf{0})$. As depicted in Fig. 1, the tree-level gravitational Rutherford scattering process is induced by the t -channel graviton exchange, and the corresponding amplitude reads

$$\mathcal{M} = \frac{\kappa^2 \mathcal{P}_{\mu\nu\rho\sigma}}{4q^2} \langle I(k') | T^{\mu\nu} | I(k) \rangle \langle N(p', \lambda') | T^{\rho\sigma} | N(p, \lambda) \rangle, \quad (2)$$

with $\mathcal{P}_{\mu\nu\rho\sigma} \equiv \eta_{\mu\rho}\eta_{\nu\sigma} + \eta_{\mu\sigma}\eta_{\nu\rho} - \eta_{\mu\nu}\eta_{\rho\sigma}$ corresponding to the harmonic gauge. $q = k - k'$ represents the momentum carried by the virtual graviton, λ, λ' denote the polarization




Fig. 1. Tree-level Feynman diagram for gravitational Rutherford scattering process $IN \rightarrow IN$. The double wavy line represents the graviton propagator. The thick double line represents the heavy target particle, and the heavy dot denotes the gravitational vertex given in (5).

indices for the massive spinning target particle. For simplicity, we have suppressed the spin indices of the projectile particle.

We will consider three different types of projectile particles, the spin-0, 1/2, 1 point-like particles. With the mass denoted by m , the corresponding energy-momentum tensors read

$$T^{\mu\nu} = \partial^\mu \phi \partial^\nu \phi - \frac{\eta^{\mu\nu}}{2} (\partial^\rho \phi \partial_\rho \phi - m^2 \phi^2), \quad \text{spin-0} \quad (3a)$$

$$T^{\mu\nu} = \frac{i}{4} \bar{\psi} \left[\gamma^\mu (\partial^\nu - \overleftarrow{\partial}^\nu) + \gamma^\nu (\partial^\mu - \overleftarrow{\partial}^\mu) \right] \psi, \quad \text{spin-}\frac{1}{2} \quad (3b)$$

$$T^{\mu\nu} = -F^{\mu\lambda} F^\nu{}_\lambda + \frac{1}{4} \eta^{\mu\nu} F^{\rho\sigma} F_{\rho\sigma} - \frac{1}{2} m^2 \eta^{\mu\nu} A^\lambda A_\lambda + m^2 A^\mu A^\nu. \quad \text{spin-1} \quad (3c)$$

The corresponding gravitational matrix elements involving the projectile particles can be readily worked out:

$$\langle I(k') | T^{\mu\nu} | I(k) \rangle = k^\mu k^\nu + k^\nu k^\mu - \eta^{\mu\nu} (k \cdot k' - m^2), \quad \text{spin-0} \quad (4a)$$

$$\langle I(k') | T^{\mu\nu} | I(k) \rangle = \frac{1}{4} \bar{u}(k') (\gamma^\mu (k'^\nu + k^\nu))$$

$$+ \gamma^\nu (k'^\mu + k^\mu) u(k), \quad \text{spin-}\frac{1}{2} \quad (4b)$$

$$\begin{aligned} \langle I(k') | T^{\mu\nu} | I(k) \rangle = & \varepsilon_\sigma^*(k') [P^{\mu\nu\rho\sigma} (k \cdot k' - m^2) + \eta^{\mu\nu} k^\sigma k'^\rho \\ & + \eta^{\rho\sigma} (k^\mu k'^\nu + k^\nu k'^\mu) - \eta^{\nu\rho} k^\sigma k'^\mu - \eta^{\mu\rho} k^\sigma k'^\nu \\ & - \eta^{\nu\sigma} k^\mu k'^\rho - \eta^{\mu\sigma} k^\nu k'^\rho] \varepsilon_\rho(k). \quad \text{spin-1} \end{aligned} \quad (4c)$$

The gravitational matrix elements involving the composite target particles in (2) are in general nonperturbative objects, which vary with target species. In literature they are usually referred to as the *gravitational form factors* (GFFs) [11, 12]. Since the GFFs encode some essential mechanical properties of a hadron such as mass, spin and shear force distributions [14–16], people's interest toward hadron's GFFs has revived in recent years. Although it is unfeasible to detect the gravitational Rutherford scattering in the foreseeable future, it is of high priority of the current and forthcoming ep facilities such as JLab, EIC and EicC to extract nucleon' GFFs in an indirect way [17]¹⁾. Recently the GFFs of the proton and pion at small momentum transfer have been investigated from the lattice QCD simulation [18, 19].

In this work we consider five types a composite target particles of mass M , with spin varying from 0 to 2. In line with the Lorentz group representation, the corresponding gravitational matrix elements involving various target particles can be decomposed into the linear combination of different GFFs [20]²⁾:

$$\langle N(p', \lambda') | T^{\mu\nu} | N(p, \lambda) \rangle_{s=0} = 2P^\mu P^\nu F_{10} \left(\frac{q^2}{M^2} \right) + 2(q^\mu q^\nu - \eta^{\mu\nu} q^2) F_{20} \left(\frac{q^2}{M^2} \right), \quad (5a)$$

$$\langle N(p', \lambda') | T^{\mu\nu} | N(p, \lambda) \rangle_{s=\frac{1}{2}} = \bar{u}(p', \lambda') \left[2P^\mu P^\nu F_{10} \left(\frac{q^2}{M^2} \right) + 2(q^\mu q^\nu - \eta^{\mu\nu} q^2) F_{20} \left(\frac{q^2}{M^2} \right) + P^{[\mu} \frac{i}{2} \sigma^{\nu]\rho} q_\rho F_{40} \left(\frac{q^2}{M^2} \right) \right] u(p, \lambda), \quad (5b)$$

$$\begin{aligned} \langle N(p', \lambda') | T^{\mu\nu} | N(p, \lambda) \rangle_{s=1} = & -\varepsilon_{\alpha'}^*(p', \lambda') \left[2P^\mu P^\nu \left(\eta^{\alpha'\alpha} F_{10} \left(\frac{q^2}{M^2} \right) - \frac{q^{\alpha'} q^\alpha}{2M^2} F_{11} \left(\frac{q^2}{M^2} \right) \right) + 2(q^\mu q^\nu - \eta^{\mu\nu} q^2) \left(\eta^{\alpha'\alpha} F_{20} \left(\frac{q^2}{M^2} \right) \right. \right. \\ & - \frac{q^{\alpha'} q^\alpha}{2M^2} F_{21} \left(\frac{q^2}{M^2} \right) \left. \right) - 2M^2 \eta^{\mu\nu} \left(\frac{q^{\alpha'} q^\alpha}{2M^2} F_{31} \left(\frac{q^2}{M^2} \right) \right) - P^{[\mu} \eta^{\nu]\alpha'} q^{\alpha'} F_{40} \left(\frac{q^2}{M^2} \right) \\ & \left. - (q^{[\mu} \eta^{\nu]\alpha'} q^{\alpha'} - \eta^{\mu\nu} q^{\alpha'} q^{\alpha'} - \eta^{\alpha'[\mu} \eta^{\nu]\alpha} q^2) F_{50} \left(\frac{q^2}{M^2} \right) \right] \varepsilon_\alpha(p, \lambda), \end{aligned} \quad (5c)$$

1) Note that it has been recently proposed that the nucleon GFF may be accessed in the future electron-ion collider via the interference effect between the photon-induced and the massive-graviton-induced amplitude in some beyond Standard Model scenarios [9].

2) Current conservation enables one to drop the F_{30} and $F_{6,i}$ terms, as well as implies $F_{31} + F_{50} = 0$ [20].

$$\begin{aligned}
\langle N(p', \lambda') | T^{\mu\nu} | N(p, \lambda) \rangle_{s=\frac{3}{2}} = & -\bar{u}_{\alpha'}(p', \lambda') \left[2P^\mu P^\nu \left(\eta^{\alpha'\alpha} F_{10} \left(\frac{q^2}{M^2} \right) - \frac{q^{\alpha'} q^\alpha}{2M^2} F_{11} \left(\frac{q^2}{M^2} \right) \right) + 2(q^\mu q^\nu - \eta^{\mu\nu} q^2) \left(\eta^{\alpha'\alpha} F_{20} \left(\frac{q^2}{M^2} \right) \right. \right. \\
& - \frac{q^{\alpha'} q^\alpha}{2M^2} F_{21} \left(\frac{q^2}{M^2} \right) - 2M^2 \eta^{\mu\nu} \left(\frac{q^{\alpha'} q^\alpha}{2M^2} F_{31} \left(\frac{q^2}{M^2} \right) \right) + P^{\mu i} \frac{i}{2} \sigma^{\nu\rho} q_\rho \left(\eta^{\alpha'\alpha} F_{40} \left(\frac{q^2}{M^2} \right) \right. \\
& \left. \left. - \frac{q^{\alpha'} q^\alpha}{2M^2} F_{41} \left(\frac{q^2}{M^2} \right) \right) - (q^{\mu\eta^{\nu|\alpha'}} q^{\alpha'} - \eta^{\mu\nu} q^{\alpha'} q^\alpha - \eta^{\alpha'\mu} \eta^{\nu\alpha} q^2) F_{50} \left(\frac{q^2}{M^2} \right) \right] u_\alpha(p, \lambda),
\end{aligned} \tag{5d}$$

$$\begin{aligned}
\langle N(p', \lambda') | T^{\mu\nu} | N(p, \lambda) \rangle_{s=2} = & \varepsilon_{\alpha'_1 \alpha'_2}^*(p', \lambda') \left[2P^\mu P^\nu \left(\eta^{\alpha'_1 \alpha'_2} F_{10} \left(\frac{q^2}{M^2} \right) - \frac{q^{\alpha'_1} q^{\alpha'_2}}{2M^2} F_{11} \left(\frac{q^2}{M^2} \right) + \frac{q^{\alpha'_1} q^{\alpha'_2}}{2M^2} \frac{q^{\alpha'_2} q^{\alpha'_2}}{2M^2} F_{12} \left(\frac{q^2}{M^2} \right) \right) \right. \\
& + 2(q^\mu q^\nu - \eta^{\mu\nu} q^2) \left(\eta^{\alpha'_1 \alpha'_2} F_{20} \left(\frac{q^2}{M^2} \right) - \frac{q^{\alpha'_1} q^{\alpha'_2}}{2M^2} F_{21} \left(\frac{q^2}{M^2} \right) + \frac{q^{\alpha'_1} q^{\alpha'_2}}{2M^2} \frac{q^{\alpha'_2} q^{\alpha'_2}}{2M^2} F_{22} \left(\frac{q^2}{M^2} \right) \right) \\
& - 2M^2 \eta^{\mu\nu} \left(\frac{q^{\alpha'_1} q^{\alpha'_2}}{2M^2} F_{31} \left(\frac{q^2}{M^2} \right) - \frac{q^{\alpha'_1} q^{\alpha'_2}}{2M^2} \frac{q^{\alpha'_2} q^{\alpha'_2}}{2M^2} F_{32} \left(\frac{q^2}{M^2} \right) \right) \\
& - P^{\mu i} \eta^{\nu|\alpha'_2} q^{\alpha'_2} \left(\eta^{\alpha'_1 \alpha'_2} F_{40} \left(\frac{q^2}{M^2} \right) - \frac{q^{\alpha'_1} q^{\alpha'_2}}{2M^2} F_{41} \left(\frac{q^2}{M^2} \right) \right) - (q^{\mu\eta^{\nu|\alpha'_2}} q^{\alpha'_2} - \eta^{\mu\nu} q^{\alpha'_2} q^{\alpha'_2} - \eta^{\alpha'_2 \mu} \eta^{\nu\alpha'_2} q^2) \\
& \left. \times \left(\eta^{\alpha'_1 \alpha'_1} F_{50} \left(\frac{q^2}{M^2} \right) - \frac{q^{\alpha'_1} q^{\alpha'_1}}{2M^2} F_{51} \left(\frac{q^2}{M^2} \right) + q^{[\alpha'_2 \eta^{\nu|\mu} \eta^{\nu\alpha'_1}] \alpha'_1} F_{70} \left(\frac{q^2}{M^2} \right) \right] \varepsilon_{\alpha_1 \alpha_2}(p, \lambda),
\end{aligned} \tag{5e}$$

where $P \equiv (p + p')/2$ is the average momentum of the target particle between the initial and final states, $q \equiv p' - p$ denotes the transfer momentum. $a^{\mu b^\nu} \equiv a^\mu b^\nu + a^\nu b^\mu$, $a^{[\mu} b^{\nu]} \equiv a^\mu b^\nu - a^\nu b^\mu$, and $\sigma^{\mu\nu} = \frac{i}{2} [\gamma^\mu, \gamma^\nu]$. u , ε^μ , u^μ , $\varepsilon^{\alpha\beta}$ signify the wave functions of the spin-1/2, 1, 3/2, and 2 particles, respectively. Note various GFFs are normalized to be dimensionless Lorentz scalars that solely depend on the ratio q^2/M^2 . Note that we have suppressed terms that are forbidden by the current conservation.

From (5) one observes that for target particle with spin s , the number of independent GFFs is $2(s+1) + 3[s] - \Theta(s-1)$ [20]¹⁾. Analogous to electromagnetic form factors, various GFFs with zero momentum transfer encapsulate the properties of the gravitational multipoles of the composite target particles. For the target particle carrying an arbitrary spin s , the F_{10} and F_{40} have an absolute normalization at zero momentum transfer, $F_{10}(0) = 1$ and $F_{40}(0) = s$, as dictated by the energy-momentum and angular momentum conservation, respectively. The $F'_{10}(0)$ ²⁾, $F_{11}(0)$, $F_{20}(0)$ and $F_{50}(0)$ terms are related to angular momentum [13, 14], pressure and shear force [14–16] of the target particle. The mass radius of the target particle can also be obtained from the linear combination of these form factors with zero momentum transfer.

III. LOW-ENERGY GRAVITATIONAL RUTHERFORD SCATTERING WITH MASSLESS PROJECTILE

We first consider the case of the massless projectile of spin j . The corresponding differential unpolarized cross section in the laboratory frame is given by

$$\frac{d\sigma}{d\cos\theta} = \frac{1}{2|\mathbf{k}'|} \cdot \frac{1}{2M} \cdot \frac{\mathbf{k}'^2}{8\pi|\mathbf{k}|M} \cdot \frac{1}{2j+1} \frac{1}{2s+1} \sum_{\text{spins}} |\mathcal{M}|^2, \tag{6}$$

where θ denotes the polar angle between the reflected and incident projectile. The magnitude of the three-momentum of the outgoing projectile, $|\mathbf{k}'|$, can be expressed in terms of $|\mathbf{k}|$, M , and $\cos\theta$:

$$|\mathbf{k}'| = \frac{|\mathbf{k}|}{1 + \frac{|\mathbf{k}|}{M} (1 - \cos\theta)}. \tag{7}$$

Squaring the amplitude in (2), summing over the polarization in the final state and averaging upon the polarizations in the initial state utilizing the spin sum formula in Appendix for target particles, one encounters rather

1) The Heaviside step function is defined as $\Theta(x) = 1$ for $x \geq 0$, otherwise vanishes. The symbol $[s]$ signifies the floor function of s . Later the symbol $\lceil s \rceil$ will be used to represent the ceiling function of s .

2) The Taylor expansion of the GFF around the origin is understood to be $F_n(q^2/M^2) = F_n(0) + F'_n(0)q^2/M^2 + O(1/M^4)$.

lengthy and cumbersome-looking expressions. It is difficult to identify any clear pattern about the dependence on the heavy target particle spin. Nevertheless, since we are solely concerned with the low-energy limit $|\mathbf{k}| \ll M$, it becomes elucidating to carry out the heavy target mass expansion for the unpolarized cross sections. As we will see, the soft behavior of the gravitational Rutherford scattering becomes transparent and one is able to recognize some universal patterns.

A. Massless spin-0 projectile

Expanding (6) in powers of $1/M$, one immediately observes that the first two terms of the unpolarized cross sections are universal, *e.g.*, independent of the target particle spin:

$$\left(\frac{d\sigma}{d\cos\theta}\right)_{\text{LO}}^s = \frac{\kappa^4 M^2 F_{10}^2}{512\pi \sin^4 \frac{\theta}{2}}, \quad (8a)$$

$$\left(\frac{d\sigma}{d\cos\theta}\right)_{\text{NLO}}^s = -\frac{\kappa^4 M |\mathbf{k}| F_{10}^2}{128\pi \sin^2 \frac{\theta}{2}}, \quad (8b)$$

with the occurring GFFs evaluated at the zero momentum transfer. For simplicity, we have adopted F_n as the shorthand for $F_n(0)$ from now on. Note that $F_{10} = 1$ for any type of composite target particles. The leading order (LO) term is identical to the cross section obtained from the light-bending angle in classical general relativity[21, 22]. This is intuitively as expected, because in the soft limit, the long-wavelength graviton can only feel the total mass of the composite target particle, insensitive to any further details about its internal structure. Interestingly, the next-to-leading order (NLO) term still remains universal, which originates from expanding the phase factor $e^{i\mathbf{k}\cdot\mathbf{r}}/k^2$ to NLO in $1/M$. It is instructive to understand why only a single GFF F_{10} contributes at NLO from the angle of effective field theory. We will devote Appendix B to such an analysis.

In contrast, at the next-to-next-to-leading-order (NNLO) in heavy target mass expansion, the differential cross sections starts to depend on the target particle spin:

$$\begin{aligned} \left(\frac{d\sigma}{d\cos\theta}\right)_{\text{NNLO}}^s &= -\frac{\kappa^4 \mathbf{k}^2}{64\pi \sin^2 \frac{\theta}{2}} \left\{ F_{10} F'_{10} - \frac{1}{2} F_{10} F_{20} (1 - \cos\theta) + \frac{1}{8} F_{10}^2 \left[7 \cos\theta - \frac{2}{3} \left(\frac{21}{2} + s + [s] \right) \right] \right. \\ &\quad \left. + \Theta\left(s - \frac{1}{2}\right) \left[\frac{(-1)^{2s} + 7}{24} F_{10} F_{40} - f_s^{(0)} F_{40}^2 (\cos\theta + 1) \right] \right. \\ &\quad \left. - \frac{1}{6} \Theta(s-1) [2 \cos\theta F_{10} F_{50} + F_{10} F_{11}] - \Theta(s-2) \frac{1}{6} F_{10} F_{70} (1 + \cos\theta) \right\}, \quad s = 0, \frac{1}{2}, 1, \frac{3}{2}, 2 \end{aligned} \quad (9)$$

with

$$f_{\frac{1}{2}}^{(0)} = \frac{1}{16}, \quad (10a)$$

$$f_1^{(0)} = \frac{1}{6}, \quad (10b)$$

$$f_{\frac{3}{2}}^{(0)} = \frac{5}{144}, \quad (10c)$$

$$f_2^{(0)} = \frac{1}{8}. \quad (10d)$$

We observe that $F'_{10} F_{10}$, $F_{10} F_{20}$, $F_{10}^2 \cos\theta$ terms still remain universal, *i.e.*, independent of the target spin. In fact, the $F'_{10} F_{10}$ terms actually have the same origin of the LO and NLO cross sections, which correspond to higher-order term in Taylor expansion of $F_{10}^2(q^2/M^2)$ in the squared LO amplitude and the phase space measure. The GFFs F'_{10} , F_{11} , F_{20} and F_{50} reflect the mechanical proper-

ties of the composite target particles such as angular momentum [13, 14], pressure and shear force [14–16]. Our results indicate that at NNLO in heavy target mass expansion, the cross section starts to depend on the detailed three-dimensional internal structure of target particle other than its mass. Curiously, the coefficient of the $F_{10} F_{40}$ term seems to reflect the spin-statistic characteristic of the target particle, which alternates from 1 (fermions) to $4/3$ (bosons). Although we only enumerate five different kinds of target spin, it is conceivable that the aforementioned patterns should hold true for arbitrary target spin.

B. Massless spin-1/2 projectile

We can repeat our investigation in Sec. III A by replacing the projectile with a point-like massless Dirac fermion. After conducting the heavy target mass expansion, we again observe that the unpolarized cross sections exhibit universal forms at LO and NLO in $1/M$ expansion:

$$\left(\frac{d\sigma}{d\cos\theta}\right)_{\text{LO}}^s = \frac{\kappa^4 M^2 F_{10}^2 \cos^2 \frac{\theta}{2}}{512\pi \sin^4 \frac{\theta}{2}}, \quad (11a)$$

$$\left(\frac{d\sigma}{d\cos\theta}\right)_{\text{NLO}}^s = -\frac{\kappa^4 M |\mathbf{k}| F_{10}^2 \cos^2 \frac{\theta}{2}}{128\pi \sin^2 \frac{\theta}{2}}. \quad (11b)$$

It is worth emphasizing that, although these expressions are independent of the target spin, they are sensitive to the projectile spin. To be definite, there arises an extra factor $\cos^2 \frac{\theta}{2}$ in the numerators of (11) with respect to (8) in the case of spinless projectile. In the forward limit $\theta \approx 0$, the occurrence of this extra factor does not make a difference. Nevertheless, this extra factor brings in a strong suppression of the LO and NLO cross sections around the backward scattering limit ($\theta \approx \pi$) for a

spin- $\frac{1}{2}$ projectile with respect to the spin-0 projectile, as is evident by comparing (8) and (11). This feature can be readily understood by the fact that, analogous to QED, gravitational interaction also conserves the helicity of massless matter particle. Therefore, the conservation of angular momentum along the beam direction forbids the backward scattering of projectile to occur, since the spin of the heavy static target particle cannot be changed by the soft gravitational scattering.

As in Sec. III A, the universality becomes partially violated at NNLO. For various composite target particles with spin ranging from 0 to 2, the NNLO contributions to the cross sections become

$$\begin{aligned} \left(\frac{d\sigma}{d\cos\theta}\right)_{\text{NNLO}}^s &= \frac{\kappa^4 \mathbf{k}^2}{64\pi \sin^2 \frac{\theta}{2}} \left\{ -F_{10} F'_{10} \cos^2 \frac{\theta}{2} + \frac{1}{8} F_{10}^2 \cos^2 \frac{\theta}{2} \left[13 \sin^2 \frac{\theta}{2} + \frac{2}{3} (s + \lceil s \rceil) \right] \right. \\ &\quad - \Theta\left(s - \frac{1}{2}\right) \left[\frac{(-1)^{2s} + 7}{24} F_{10} F_{40} \cos^2 \frac{\theta}{2} - f_s^{(\frac{1}{2})} F_{40}^2 (\cos 2\theta + 8 \cos \theta + 23) \right] \\ &\quad \left. + \frac{1}{6} \Theta(s-1) F_{10} (2F_{50} + F_{11}) \cos^2 \frac{\theta}{2} + \Theta(s-2) \frac{1}{3} F_{10} F_{70} \cos^2 \frac{\theta}{2} \right\}, \quad s = 0, \frac{1}{2}, 1, \frac{3}{2}, 2 \end{aligned} \quad (12)$$

with

$$f_{\frac{1}{2}}^{(\frac{1}{2})} = \frac{1}{256}, \quad (13a)$$

$$f_1^{(\frac{1}{2})} = \frac{1}{96}, \quad (13b)$$

$$f_{\frac{3}{2}}^{(\frac{1}{2})} = \frac{5}{2304}, \quad (13c)$$

$$f_2^{(\frac{1}{2})} = \frac{1}{128}. \quad (13d)$$

Similar to the pattern revealed in the case of massless spinless projectile, we observe that the coefficients of the $F'_{10} F_{10}$, $F_{10} F_{20}$, $F_{10}^2 \sin^2 \frac{\theta}{2} \cos^2 \frac{\theta}{2}$ remain independent of the target spin. The $F'_{10} F_{10}$ term actually has the same origin of the LO and NLO cross sections. The coefficient of the $F_{10} F_{40}$ term again reflects the spin-statistic characteristic of the target particle, which alternates from 1 (fermions) to $4/3$ (bosons).

C. Massless spin-1 projectile

We can repeat the the preceding analysis by turning to

a massless spin-1 projectile. Upon heavy target mass expansion, analogous to the cases of the spin-0 and spin- $\frac{1}{2}$ projectile, we again observe the similar universal pattern at LO and NLO:

$$\left(\frac{d\sigma}{d\cos\theta}\right)_{\text{LO}}^s = \frac{\kappa^4 M^2 F_{10}^2 \cos^4 \frac{\theta}{2}}{512\pi \sin^4 \frac{\theta}{2}}, \quad (14a)$$

$$\left(\frac{d\sigma}{d\cos\theta}\right)_{\text{NLO}}^s = -\frac{\kappa^4 M |\mathbf{k}| F_{10}^2 \cos^4 \frac{\theta}{2}}{128\pi \sin^2 \frac{\theta}{2}}, \quad (14b)$$

except there appears an extra factor $\cos^4 \frac{\theta}{2}$ in the numerators with respect to the case of spinless projectile. In the forward limit $\theta \approx 0$, the emergence of this extra factor does not make a difference. Similar to the case of the spin- $\frac{1}{2}$ projectile, this extra factor brings in a stronger suppression of the LO and NLO cross sections near the backward scattering limit ($\theta \approx \pi$) for a spin-1 projectile. The reason is the same as explained before, since the gravitational interaction preserves the helicity of massless spin-1 matter particle (photon), the conservation of angular momentum along the beam direction forbids the backward scattering of the massless projectile to occur.

The universality becomes partially violated at NNLO. For target particle with spin ranging from 0 to 2, the NNLO contributions to the unpolarized cross sections become

$$\begin{aligned} \left(\frac{d\sigma}{d\cos\theta}\right)_{\text{NNLO}}^s &= -\frac{\kappa^4 \mathbf{k}^2 \cos^2 \frac{\theta}{2}}{256\pi \sin^2 \frac{\theta}{2}} \left\{ F_{10}^2 \cos^2 \frac{\theta}{2} \left[3\cos\theta - \frac{1}{3}(9-s-[s]) \right] + 4F_{10}F'_{10} \cos^2 \frac{\theta}{2} \right. \\ &\quad \left. + \Theta\left(s - \frac{1}{2}\right) \left[\frac{(-1)^{2s} + 7}{6} F_{10}F_{40} \cos^2 \frac{\theta}{2} + f_s^{(1)} F_{40}^2 (\cos\theta - 3) \right] \right. \\ &\quad \left. - \frac{2}{3} F_{10} [\Theta(s-1)(F_{11} + 2F_{50}) + \Theta(s-2)2F_{70}] \cos^2 \frac{\theta}{2} \right\}, \quad s = 0, \frac{1}{2}, 1, \frac{3}{2}, 2 \end{aligned} \quad (15)$$

with

$$f_{\frac{1}{2}}^{(1)} = \frac{1}{4}, \quad (16a)$$

$$f_1^{(1)} = \frac{2}{3}, \quad (16b)$$

$$f_{\frac{3}{2}}^{(1)} = \frac{5}{36}, \quad (16c)$$

$$f_2^{(1)} = \frac{1}{2}. \quad (16d)$$

Similar to the projectile of spin-0 and $\frac{1}{2}$, we observe that the coefficients of $F'_{10}F_{10}$, $F_{10}F_{20}$, $F_{10}^2 \cos\theta \cos^2 \frac{\theta}{2}$ terms are independent of the target spin. Actually the $F'_{10}F_{10}$ term has the same origin of the LO and NLO cross sections. Again, the coefficient of the $F_{10}F_{40}$ term again reflects the spin-statistic characteristic of the target particle, which alternates from 1 (fermions) to 4/3 (bosons).

IV. GRAVITATIONAL RUTHERFORD SCATTERING WITH NON-RELATIVISTIC LIGHT PROJECTILE

In this section, we turn to the original prototype of

$$\left(\frac{d\sigma}{d\cos\theta}\right)_{(v^0)}^s = \frac{F_{10}^2 \kappa^4 m^4 M(M+m)^2 \left(\sqrt{M^2 - m^2 \sin^2 \theta} + m \cos\theta\right)^2}{512\pi \mathbf{k}^4 \sqrt{M^2 - m^2 \sin^2 \theta} \left(M - \cos\theta \sqrt{M^2 - m^2 \sin^2 \theta} + m \sin^2 \theta\right)^2} = \frac{\kappa^4 M^2 m^4 F_{10}^2}{2048\pi \mathbf{k}^4 \sin^4 \frac{\theta}{2}} + \mathcal{O}\left(\frac{m^6}{M^4 \mathbf{k}^4}\right). \quad (18)$$

which scales as $1/\mathbf{k}^4$, exactly identical to the familiar Rutherford formula obtained from the classical mechanics with Newtonian's gravitational law [28]. Note this expression is insensitive to both target and projectile's spins, since the spin degree of freedom decouples in the nonrelativistic limit. This is in contrast to the LO expressions in the case of massless projectile in heavy target mass expansion, which is insensitive to the target spin yet depends on the projectile spin.

Rutherford scattering process, *i.e.*, a slowly moving light projectile hits a heavy static target, albeit with the interaction mediated by exchanging a graviton exchange rather than exchanging a photon. We assume the projectile is point-like particle with mass $m \ll M$.

The differential cross section for the nonrelativistic Rutherford scattering in the laboratory frame is given by

$$\begin{aligned} \frac{d\sigma}{d\cos\theta} &= \frac{1}{32\pi M} \left[p'^0 + k^0 \left(1 - \frac{|\mathbf{k}|}{|\mathbf{k}'|} \cos\theta \right) \right]^{-1} \\ &\quad \times \frac{|\mathbf{k}'|}{|\mathbf{k}|} \frac{1}{2j+1} \frac{1}{2s+1} \sum_{\text{spins}} |\mathcal{M}|^2, \end{aligned} \quad (17)$$

where the projectile bears spin j , θ denotes the polar angle between the reflected and incident projectile.

Similar to Sec. III, let us consider again three different types of projectiles and five different types of targets. The resulting expressions of unpolarized cross sections are generally rather lengthy and unilluminating. Since there are three widely separated scales inherent in this process, $\mathbf{k} \ll m \ll M$, the appropriate way of extracting the soft behavior is to simultaneously expand the differential cross sections in powers of $v = |\mathbf{k}|/m$ (velocity of the projectile) and m/M . The necessity of conducting double expansion renders this case more complicated than the case of low-energy massless projectile as discussed in Sec. III.

Interestingly, at the lowest order in velocity yet to all orders in $1/M$, the differential cross sections scales assumes a uniform form:

At NLO in velocity expansion, the differential cross sections scale as $1/\mathbf{k}^2$, whose explicit expressions are still rather complicated yet vary with different projectile and target species. Nevertheless, once the heavy target mass expansion is performed, some clear pattern starts to emerges. In the following, we will consider three different types of light projectiles with spin ranging from 0 to 1.

A. Slowly-moving spin-0 projectile

At relative order- v^2 , after conducting the heavy target mass expansion, the differential unpolarized cross section becomes particularly simple:

$$\left(\frac{d\sigma}{d\cos\theta}\right)_{(v^2)}^s = \frac{\kappa^4 M^2 m^2 F_{10}}{256\pi k^2 \sin^2 \frac{\theta}{2}} \times \left[\frac{F_{10}}{2 \sin^2 \frac{\theta}{2}} - \frac{m}{M} F_{10} + \frac{m^2}{M^2} g_s^{(0)} + \dots \right], \quad (19)$$

with

$$\begin{aligned} g_s^{(0)} = & -F_{20} - F'_{10} + \frac{F_{10}}{4} \left[\frac{1}{3} (3 + s + [s]) - 2 \cos\theta \right] \\ & - \Theta \left(s - \frac{1}{2} \right) \frac{1}{24} [(-1)^{2s} + 7] F_{40} \\ & + \Theta(s-1) \left(\frac{1}{6} F_{11} + F_{50} \right) + \Theta(s-2) \frac{2}{3} F_{70}, \\ s = & 0, \frac{1}{2}, 1, \frac{3}{2}, 2. \end{aligned} \quad (20)$$

Clearly the $O(v^2/M^n)$ ($n = 0, 1$) terms remain independent of the target spin. At $O(v^2/M^2)$, the universality becomes partially violated. Nevertheless, the F'_{10} , F_{20} , and $F_{10} \cos\theta$ terms still do not depend on the target particle spin. The coefficient of the F_{40} term again reflects the spin-statistic characteristic of the target particle, which alternates from 1 (fermions) to 4/3 (bosons).

B. Slowly-moving spin- $\frac{1}{2}$ projectile

We can repeat our investigation by replacing the projectile with a slowly-moving Dirac fermion. The $O(v^0)$ cross section is given by (18), since the spin degree of freedom decouples in the nonrelativistic limit. At relative order- v^2 , after carrying out the heavy target mass expansion, the differential cross section again possesses a simple form:

$$\left(\frac{d\sigma}{d\cos\theta}\right)_{(v^2)}^s = \frac{\kappa^4 M^2 m^2 F_{10}}{256\pi k^2 \sin^2 \frac{\theta}{2}} \left[\frac{F_{10}}{16} \frac{3 \cos\theta + 5}{\sin^2 \frac{\theta}{2}} - \frac{m F_{10}}{8M} (3 \cos\theta + 5) - \frac{m^2}{M^2} g_s^{(\frac{1}{2})} + \dots \right], \quad (21)$$

with

$$\begin{aligned} g_s^{(\frac{1}{2})} = & F'_{10} + F_{20} + \frac{1}{48} F_{10} \left[15 \cos\theta - 4 \left(\frac{3}{4} + s + [s] \right) \right] \\ & + \Theta \left(s - \frac{1}{2} \right) \frac{1}{24} [(-1)^{2s} + 7] F_{40} \\ & - \Theta(s-1) \left(F_{50} + \frac{1}{6} F_{11} \right) - \Theta(s-2) \frac{2}{3} F_{70}, \\ s = & 0, \frac{1}{2}, 1, \frac{3}{2}, 2. \end{aligned} \quad (22)$$

The $O(v^2/M^n)$ ($n = 0, 1$) terms remain universal. The universality has been partially violated in the $O(v^2/M^2)$ term. However, even at this order, the F'_{10} , F_{20} , and $F_{10} \cos\theta$ terms still appears to be independent of the target particle spin. The coefficient of the F_{40} term again reflects the spin-statistic characteristic of the target particle, which alternates from 1 (fermions) to 4/3 (bosons).

C. Slowly-moving spin-1 projectile

As a final example, we consider a slowly-moving light spin-1 projectile. Needless to say, the $O(v^0)$ differential cross section is again described by (18). At relative order- v^2 , after carrying out the heavy target mass expansion, the expanded differential cross sections bear the following structure:

$$\left(\frac{d\sigma}{d\cos\theta}\right)_{(v^2)}^s = \frac{\kappa^4 M^2 m^2 F_{10}}{1536\pi k^2 \sin^2 \frac{\theta}{2}} \left[\frac{F_{10} (2 \cos\theta + 1)}{\sin^2 \frac{\theta}{2}} - \frac{2m}{M} (2 \cos\theta + 1) F_{10} - \frac{m^2}{M^2} g_s^{(1)} + \dots \right], \quad (23)$$

where

$$\begin{aligned} g_s^{(1)} = & F_{10} \left[\cos\theta + \frac{1}{2} (1 - s - [s]) \right] + 6F'_{10} + 6F_{20} \\ & + \Theta \left(s - \frac{1}{2} \right) \frac{1}{4} [(-1)^{2s} + 7] F_{40} \\ & - \Theta(s-1) (F_{11} - 6F_{50}) - \Theta(s-2) 4F_{70}, \\ s = & 0, \frac{1}{2}, 1, \frac{3}{2}, 2. \end{aligned} \quad (24)$$

Clearly the $O(v^2/M^n)$ ($n = 0, 1$) terms remain universal. At $O(v^2/M^2)$, the universality becomes partially violated, notwithstanding that the F'_{10} , F_{20} , and $F_{10} \cos\theta$ terms still do not depend on the target spin. The coefficient of the F_{40} term again reflects the spin-statistic characteristic of the target particle, which alternates from 1 (fermions) to 4/3 (bosons).

V. SUMMARY

In this work, we have conducted a comprehensive study of the soft pattern of the tree-level gravitational Rutherford scattering processes. Two classes of Rutherford scattering processes have been considered, *i.e.*, a low-energy massless projectile strikes on a static, heavy, composite target carrying spin up to 2, and a slowly-moving light structureless projectile bombs on a static, heavy, spinning composite target particle.

The soft limits of both classes of gravitational Rutherford scattering processes have exhibited some universal and simple patterns. For the former type, given a massless projectile with a certain spin, the first two terms in

the heavy target mass expansion remain universal, and the NNLO term starts to develop target spin dependence. Nevertheless, several terms at NNLO still remain universal or have some definite pattern of dependence on the target spin. For the latter, one has to carry out the double expansion in projectile velocity and $1/M$ in order to identify a simple soft limit. At the lowest order in v yet to all orders in $1/M$, the differential cross section has a universal form, insensitive to both of the projectile and target spin. At the relative order- v^2 , the first two terms in $1/M$ expansion are still independent of the target spin. The universality starts to be partially violated in the $O(v^2/M^2)$ piece, though some terms at this order still remain independent of the target spin, or bears a definite pattern of target spin dependence.

It is curious that at NNLO in heavy target mass expansion, the prefactors of the $F_{40} = s$ term in both types of gravitational Rutherford scattering processes encapsulate some peculiar spin-statistics characteristic, which alternate from a constant for fermionic target to another constant for bosonic target. It is interesting to examine whether this pattern holds true for the composite target with arbitrarily high spin.

ACKNOWLEDGMENTS

We thank Kangyu Chai for participating in the early stage of this work. We are grateful to Zhewen Mo for discussions.

APPENDIX A: POLARIZATION SUM FORMULA

In deriving the unpolarized cross sections, the following spin sum formulas concerning massive target particles are useful:

$$\sum_{\lambda} u(p, \lambda) \bar{u}(p, \lambda) = \frac{\not{p} + M}{2M}, \quad (\text{A1a})$$

$$\sum_{\lambda} \varepsilon_{\alpha}(p, \lambda) \varepsilon_{\alpha'}^*(p, \lambda) = \hat{\eta}_{\alpha\alpha'}, \quad (\text{A1b})$$

$$\begin{aligned} \sum_{\lambda} u_{\alpha}(p, \lambda) \bar{u}_{\alpha'}(p, \lambda) &= -\frac{\not{p} + M}{2M} \left(\eta_{\alpha\alpha'} - \frac{1}{3} \gamma_{\alpha} \gamma_{\alpha'} \right. \\ &\quad \left. - \frac{2p_{\alpha} p_{\alpha'}}{3M^2} + \frac{\gamma_{\alpha'} p_{\alpha} - \gamma_{\alpha} p_{\alpha'}}{3M} \right), \end{aligned} \quad (\text{A1c})$$

$$\begin{aligned} \sum_{\lambda} \varepsilon_{\alpha_1 \alpha_2}(p, \lambda) \varepsilon_{\alpha'_1 \alpha'_2}^*(p, \lambda) &= \hat{\eta}_{\alpha_1 \alpha'_1} \hat{\eta}_{\alpha_2 \alpha'_2} \\ &\quad + \hat{\eta}_{\alpha_1 \alpha'_2} \hat{\eta}_{\alpha_2 \alpha'_1} - \frac{2}{3} \hat{\eta}_{\alpha_1 \alpha_2} \hat{\eta}_{\alpha'_1 \alpha'_2}, \end{aligned} \quad (\text{A1d})$$

with $\hat{\eta}_{\alpha\beta} \equiv -\eta_{\alpha\beta} + \frac{p_{\alpha} p_{\beta}}{M^2}$. Note that the Dirac spinor is normalized as $\bar{u}(p, r) u(p, s) = \delta^{rs}$.

APPENDIX B: THE VANISHING OF THE NLO AMPLITUDE IN THE HEAVY BLACK HOLE EFFECTIVE THEORY

Since the first detection of the gravitational wave (GW) by LIGO and VIRGO in 2015, precise predictions of the GW templates become an imperative task. As an efficient theoretical framework to organize the post-Newtonian and post-Minkowski expansion, the heavy black hole effective theory (HBET) has recently been developed [24, 25], which is analogous to the heavy quark effective theory (HQET) tailored for heavy quark physics. Recently it has been applied to GW emission in the scattering of binary spinless black holes or neutron stars with arbitrary masses at next-to-leading order in the post-Minkowski expansion [26].

The original HBET Lagrangian is designed to describe a heavy structureless particle (black hole) interacting with soft gravitons, with the expansion parameter being $1/M$ [24, 25]. We will use this machinery to explain why the NLO amplitude in $1/M$ expansion vanishes for gravitational Rutherford scattering.

For simplicity, let us consider heavy spinless target particle, which is represented by a complex scalar field. The underlying theory describing a heavy structureless particle interacting with gravity is simply assumed to be

$$S = \int d^4x \sqrt{-g} (g^{\mu\nu} \partial_{\mu} \phi^* \partial_{\nu} \phi - M^2 |\phi|^2). \quad (\text{B1})$$

Mimicking the derivation of HQET from QCD, one integrates out the heavy anti-particle degree of freedom by substituting the equation of motion into (B1) and expands the Lagrangian in powers of $1/M$. The effective action in general spacetime background then reads [24, 25]

$$\begin{aligned} S_{\text{HBET}} &= \int d^4x \frac{\sqrt{-g}}{2} \left[M (v_{\mu} v_{\nu} g^{\mu\nu} - 1) \varphi_v^* \varphi_v \right. \\ &\quad + \frac{i}{2} g^{\mu\nu} (v_{\mu} (\varphi_v^* \partial_{\nu} \varphi_v - (\partial_{\nu} \varphi_v^*) \varphi_v) \\ &\quad + v_{\nu} (\varphi_v^* \partial_{\mu} \varphi_v - (\partial_{\mu} \varphi_v^*) \varphi_v)) \\ &\quad \left. + \frac{i}{2} (v_{\mu} v_{\nu} g^{\mu\nu} - 1) ((v \cdot \partial \varphi_v^*) \varphi_v - \varphi_v^* v \cdot \partial \varphi_v) \right]. \end{aligned} \quad (\text{B2})$$

Since we are interested in the Rutherford scattering in Minkowski spacetime, the weak field approximation $\sqrt{-g} = 1 - \frac{\kappa}{2} \eta_{\mu\nu} h^{\mu\nu} + O(h^2)$ is invoked. The subscript of φ is the velocity label of the heavy target particle, with $v^{\mu} = (1, \mathbf{0})$ in the laboratory frame. It should be cautioned

that (B2) was invented for a point-like target particle [24, 25]. To describe a heavy composite target particle, one should assign general Wilson coefficients c_i to those higher-dimensional operators, which reflect its nontrivial internal structure. Keeping the free kinetic term of the heavy scalar field, as well as organizing the $\varphi_v^* \varphi_v$ -graviton interactions in powers of $1/M$, we then obtain

$$\begin{aligned} \mathcal{L}_{\text{HBET}'} = & \frac{i}{2} (\varphi_v^* v \cdot \partial \varphi_v - (v \cdot \partial \varphi_v^*) \varphi_v) + c_1 \frac{M\kappa}{2} h^{\mu\nu} v_\mu v_\nu \\ & + \frac{i\kappa}{4} h^{\mu\nu} (c_{2,2} v_\mu v_\nu + c_{2,1} \eta_{\mu\nu}) ((v \cdot \partial \varphi_v^*) \varphi_v - \varphi_v^* v \cdot \partial \varphi_v) \\ & + c_{2,1} \frac{i\kappa}{4} h^{\mu\nu} (v_\mu (\varphi_v^* \partial_\nu \varphi_v - (\partial_\nu \varphi_v^*) \varphi_v) \\ & + v_\nu (\varphi_v^* \partial_\mu \varphi_v - (\partial_\mu \varphi_v^*) \varphi_v)), \end{aligned} \quad (\text{B3})$$

where the subscript HBET' implies that the original HBET is generalized to account for the composite heavy target particle.

From (B3) one readily reads off the Feynman rules for the $\varphi_v^* \varphi_v$ -graviton vertices through NLO in $1/M$:

$$V_{\text{LO}}^{\varphi_v^* \varphi_v h} = c_1 \frac{iM\kappa}{2} v^\mu v^\nu, \quad (\text{B4a})$$

$$\begin{aligned} V_{\text{NLO}}^{\varphi_v^* \varphi_v h} = & \frac{i\kappa}{4} [c_{2,1} v^\mu (\tilde{p}^\nu + \tilde{p}'^\nu) + c_{2,1} v^\nu (\tilde{p}^\mu + \tilde{p}'^\mu) \\ & - (c_{2,2} v^\mu v^\nu + c_{2,1} \eta^{\mu\nu}) (v \cdot \tilde{p} + v \cdot \tilde{p}')], \end{aligned} \quad (\text{B4b})$$

For a point-like target particle, one simply has $c_1 = c_{2,1} = c_{2,2} = 1$ ¹⁾. Note that the \tilde{p} and \tilde{p}' signify the residual momenta of target particle, *e.g.*, $p = Mv + \tilde{p}$. Assuming the projectile to be a massless spinless point particle, combining (4a) and the $\varphi_v^* \varphi_v h$ vertices enumer-

ated in (B4), we then obtain the HBET prediction to the gravitational Rutherford scattering amplitudes through NLO in $1/M$ expansion:

$$\mathcal{M}_{\text{EFT}} = \frac{c_1 \kappa^2 M}{4(\cos\theta - 1)} + \frac{c_{2,2} \kappa^2 (|\mathbf{k}| - |\mathbf{k}'|)}{16 \sin^2(\frac{\theta}{2})} + \mathcal{O}\left(\frac{1}{M}\right), \quad (\text{B5})$$

However, in light of the relation between $|\mathbf{k}|$ and $|\mathbf{k}'|$ as given in (7), one readily observes that the second (nominally NLO) term in (B5) is suppressed with respect to the LO term actually by a factor of $1/M^2$, rather than $1/M$. Therefore, we conclude

$$\mathcal{M}_{\text{EFT}} = \frac{c_1 \kappa^2 M}{4(\cos\theta - 1)} + \mathcal{O}\left(\frac{1}{M}\right). \quad (\text{B6})$$

This EFT analysis provides a clear perspective to understand why the NLO amplitude vanishes.

The EFT prediction (B6) should be identical to the LO result in (8a), which has been derived earlier in terms of the GFFs of composite target particle,

$$\mathcal{M} = \frac{\kappa^2 M^2 F_{10}}{2(\cos\theta - 1)} + \mathcal{O}\left(\frac{1}{M^0}\right). \quad (\text{B7})$$

This criterion enforces $c_1 = F_{10} = 1$, once the $2M$ factor is compensated for the non-relativistic state normalization. The requirement $c_1 = 1$, irrespective to whether the heavy target particle is fundamental or composite, may be attributed to the reparametrization invariance in HBET.

Since the $\mathcal{O}(1/M)$ HBET vertex yields a vanishing contribution to the Rutherford scattering amplitude, therefore the NLO contribution to the unpolarized cross section in (8b) solely arises from the expansion of the phase space factor.

References

- [1] M. D. Schwartz, *Quantum Field Theory and the Standard Model*, Cambridge University Press, 2014, ISBN 978-1-107-03473-0,
- [2] S. Weinberg, *Phys. Rev.* **140**, B516 (1965)
- [3] F. Bloch and A. Nordsieck, *Phys. Rev.* **52**, 54 (1937)
- [4] T. Kinoshita, *J. Math. Phys.* **3**, 650 (1962)
- [5] T. D. Lee and M. Nauenberg, *Phys. Rev.* **133**, B1549 (1964)
- [6] M. Beneke, P. Hager and R. Szafron, *JHEP* **03**, 199 (2022), arXiv:2110.02969[hep-th]
- [7] F. E. Low, *Phys. Rev.* **96**, 1428 (1954)
- [8] M. Gell-Mann and M. L. Goldberger, *Phys. Rev.* **96**, 1433 (1954)
- [9] Y. Hatta, *Phys. Rev. D* **109**(5), L051502 (2024), arXiv:2311.14470[hep-ph]
- [10] Y. Jia and J. Y. Zhang, [arXiv: 2303.18243 [hep-ph]].
- [11] I. Y. Kobzarev and L. B. Okun, *Zh. Eksp. Teor. Fiz.* **43**, 1904 (1962)
- [12] H. Pagels, *Phys. Rev.* **144**, 1250 (1966)
- [13] E. Leader and C. Lorcé, *Phys. Rept.* **541**(3), 163 (2014), arXiv:1309.4235[hep-ph]
- [14] M. V. Polyakov, *Phys. Lett. B* **555**, 57 (2003), arXiv:hep-ph/0210165[hep-ph]
- [15] M. V. Polyakov and H. D. Son, *JHEP* **09**, 156 (2018), arXiv:1808.00155[hep-ph]
- [16] M. V. Polyakov and P. Schweitzer, *Int. J. Mod. Phys. A* **33**(26), 1830025 (2018), arXiv:1805.06596[hep-ph]

¹⁾ Note that the authors of [24, 25] consider the real scalar field. They simply discard the rapidly oscillating terms proportional to $e^{\pm 2im \cdot v}$ and obtain $c_1 = c_{2,1} = 1$ and $c_{2,2} = 0$.

- [17] V. D. Burkert, L. Elouadrhiri, F. X. Girod, C. Lorcé, P. Schweitzer and P. E. Shanahan, *Rev. Mod. Phys.* **95**(4), 041002 (2023), arXiv:2303.08347[hep-ph]
- [18] D. C. Hackett, P. R. Oare, D. A. Pefkou and P. E. Shanahan, *Phys. Rev. D* **108**(11), 114504 (2023), arXiv:2307.11707[hep-lat]
- [19] D. C. Hackett, D. A. Pefkou and P. E. Shanahan, [arXiv:2310.08484 [hep-lat]].
- [20] S. Cotogno, C. Lorcé, P. Lowdon and M. Morales, *Phys. Rev. D* **101**(5), 056016 (2020), arXiv:1912.08749[hep-ph]
- [21] N. E. J. Bjerrum-Bohr, J. F. Donoghue, B. R. Holstein, L. Planté and P. Vanhove, *Phys. Rev. Lett.* **114**(6), 061301 (2015), arXiv:1410.7590[hep-th]
- [22] J. F. Donoghue and B. R. Holstein, *Am. J. Phys.* **54**, 827 (1986)
- [23] B. P. Abbott *et al.* [LIGO Scientific and Virgo], *Phys. Rev. Lett.* **116**(6), 061102 (2016), arXiv:1602.03837[gr-qc]
- [24] P. H. Damgaard, K. Haddad and A. Helset, *JHEP* **11**, 070 (2019), arXiv:1908.10308[hep-ph]
- [25] R. Aoude, K. Haddad and A. Helset, *JHEP* **05**, 051 (2020), arXiv:2001.09164[hep-th]
- [26] A. Brandhuber, G. R. Brown, G. Chen, S. De Angelis, J. Gowdy and G. Travaglini, *JHEP* **06**, 048 (2023), arXiv:2303.06111[hep-th]
- [27] B. Maybee, D. O'Connell and J. Vines, *JHEP* **12**, 156 (2019), arXiv:1906.09260[hep-th]
- [28] C. W. Misner, K. S. Thorne and J. A. Wheeler, *Gravitation*, Princeton University Press, 2017, ISBN 978-0-7167-0344-0.