

Correlation between Zero-Sound Modes and Nuclear Equation of State Stiffness Detected by Light Vector Boson*

Jing Ye (叶婧)[†] Wei-Zhou Jiang[‡]

School of Physics, Southeast University, Nanjing 211189, China

Abstract: We investigate zero-sound modes in nuclear matter and neutron star matter using the relativistic random phase approximation (RRPA), focusing on the role of a light vector U boson in stiffening the nuclear equation of state (EOS). The occurrence of zero sound is closely linked to the stiffness of the nuclear EOS. The additional vector repulsive potential induced by the U boson results in the occurrence of zero sound at high densities in models with a soft high-density EOS, and the density range for zero sound extends to higher densities with larger coupling strength of the U boson. This confirms the unambiguous role of stiffening the high-density EOS in resulting in the appearance of zero sound.

Keywords: equation of state, zero sound, random phase approximation, U boson, nuclear matter

DOI: 10.1088/1674-1137/adfa81

CSTR:

I. INTRODUCTION

By combining observational data from radio pulsar observatories [1–4], gravitational wave (GW) signals [4, 5], short gamma-ray bursts [6], and kilonovae [7], meaningful celestial constraints on the stiffness of the equation of state (EOS) for compact stars can be derived. While extremely stiff (high-density) EOSs have been ruled out by experimental nucleon flow data [8] and gravitational wave observations due to their incompatibility with neutron star radii constraints [9], exceedingly soft high-density EOSs are also disfavored due to their inability to reproduce massive neutron stars [10, 11] and sufficiently massive accretion disks required for short gamma-ray bursts and kilonova emissions [12]. Observations of massive neutron stars, with masses around $2M_{\odot}$ [13–15], suggest that their EOS must be relatively stiff, thereby constraining possible non-nucleonic degrees of freedom in neutron star matter [10, 16, 17]. On the other hand, the celestial constraints on the nuclear EOS cannot be completely exempt from the degeneracy arising from non-nucleonic degrees of freedom and some celestial factors. For instance, studies on bosonic dark matter-admixed neutron stars show consistency with maximum mass and tidal deformability constraints [18, 19]. Furthermore, extensive research on the non-Newtonian gravitational effect in neutron stars and strange stars suggests the possibility of soft high-density EOSs supporting massive compact stars [20–25]. Such an effect is often simulated by

the exchange of weakly interacting light bosons beyond the standard model, potentially altering gravitational interactions and affecting EOS constraints.

In particular, a neutral, weakly coupled spin-1 gauge boson, proposed in supersymmetric extensions of the Standard Model [26, 27], has been suggested as a candidate boson that modifies the inverse-square law of gravity [25, 28, 29]. Many terrestrial experiments have been conducted to search for it [30]. It has also been used as the interaction propagator of the MeV dark matter to explain the 511 keV gamma-ray emission from the galactic bulge [31–36]. More interestingly, the introduction of a vector gauge boson of the U(1) symmetry, referred to as the U boson, can significantly stiffen the soft EOS of asymmetric nuclear matter, influencing the structure of neutron stars [23, 28, 37–39]. Such stiffening is advantageous, as the softening of the EOS due to spontaneous phase transitions in the interior core of neutron stars must be compensated to sustain large-mass stars.

Vector bosons, including the vector meson and U boson, are involved in density fluctuation and collective modes as a response to the perturbation of nuclear systems. The collective modes can be used to constrain the properties of nuclear EOS. For instance, the nuclear matter compression modulus [40–42], the symmetry energy [43, 44], and the effective mass of the nucleon [45, 46] are correlated tightly with the isoscalar giant multipole resonances, the isovector giant dipole resonance, and the isoscalar giant quadrupole resonance, respectively. It is

Received 12 May 2025; Accepted 12 August 2025

* This work was supported in part by the National Natural Science Foundation of China under Grant No. 12375112

[†] E-mail: yejing_@seu.edu.cn

[‡] E-mail: Wei-Zhou Jiang, wzjiang@seu.edu.cn

©2026 Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd. All rights, including for text and data mining, AI training, and similar technologies, are reserved.

interesting to investigate how changes in these bosonic mediators, associated with nuclear matter stiffness, influence the collective excitation modes. As a collective mode, zero-sound mode in nuclear matter has been investigated in some early works [47–52]. Recently, we have demonstrated that the relativistic mean-field (RMF) model with a stiff EOS facilitates the emergence of zero-sound modes at high densities [53, 54]. The zero sound in nuclear matter is found to be intimately linked to the stiffness of the EOS. Namely, the occurrence of zero sound prefers a stiff EOS, regardless of whether the matter is symmetric or asymmetric. Though the soft EOS models, on the other hand, do not support the occurrence of zero sounds, stiffening the EOS by adjusting the c_3 parameter of the vector meson self-interaction term, which increases the repulsive interaction, facilitates the revival of zero-sound modes at high densities. In light of the role of the vector meson in reviving the zero sound, it is interesting to examine whether the EOS stiffening induced by the U boson can have a similar effect on the occurrence of zero-sound modes. By doing so, we can examine whether the relationship is consistent between various EOS stiffenings and the occurrence of zero sound.

The following facts may further highlight the significance of this work. Zero sounds, as potential candidates for Goldstone bosons, may annihilate into neutrino-anti-neutrino pairs via Z^0 boson exchange [55]. This mechanism may yield an additional neutrino emission channel, potentially affecting the cooling behavior and long-term evolution of neutron stars [56, 57]. Moreover, the dual properties of the vector U-boson in modifying the EOS and gravity may bridge the relation between the zero sound and modified gravity in a dense astrophysical medium. Such investigations can shed new light on the fundamental aspects and correlated relationships of new physics beyond the Standard Model.

In this work, we investigate the effects of the U boson on zero-sound modes in nuclear matter and neutron star matter using the relativistic random phase approximation (RRPA) method based on the RMF models. The RRPA formulations have been widely and successfully applied to study various collective modes in finite nuclei [40, 58–62]. It is necessary to mention that the zero sound obtained with the RRPA is consistent with those from Landau's kinetic theory [49, 63]. The structure of the remaining paper is as follows. In Sec. II, we briefly introduce the RMF models and the RRPA formalism. Sec. III presents numerical results and discussions. Lastly, a summary is given in Sec. IV.

II. FORMALISM

A. RMF models of effective interactions

In the RMF approach, the Lagrangian can generally

be given as [64–66]:

$$\begin{aligned} \mathcal{L} = & \bar{\psi} [i\gamma_\mu \partial^\mu - M + g_\sigma \sigma - g_\omega \gamma_\mu \omega^\mu - g_\rho \gamma_\mu \tau_3 b_0^\mu] \psi \\ & + \frac{1}{2} (\partial_\mu \sigma \partial^\mu \sigma - m_\sigma^2 \sigma^2) - \frac{1}{3} g_2 \sigma^3 - \frac{1}{4} g_3 \sigma^4 \\ & - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{1}{2} m_\omega^2 \omega_\mu \omega^\mu + \frac{1}{4} c_3 (\omega_\mu \omega^\mu)^2 \\ & - \frac{1}{4} B_{\mu\nu} B^{\mu\nu} + \frac{1}{2} m_\rho^2 b_{0\mu} b_0^\mu \\ & + 4\Lambda_V g_\rho^2 g_\omega^2 \omega_\mu \omega^\mu b_{0\mu} b_0^\mu + \mathcal{L}_u, \end{aligned} \quad (1)$$

where $\psi, \sigma, \omega, b_0$ are the fields of the nucleon, the scalar, vector, and isovector-vector mesons, respectively, with their masses M, m_σ, m_ω , and m_ρ . $F_{\mu\nu}$ and $B_{\mu\nu}$ are the respective strength tensors of the vector mesons ω and ρ , respectively:

$$F_{\mu\nu} = \partial_\mu \omega_\nu - \partial_\nu \omega_\mu, \quad B_{\mu\nu} = \partial_\mu b_{0\nu} - \partial_\nu b_{0\mu}. \quad (2)$$

In Eq. (1), the additional Lagrangian $\mathcal{L}_u$ of the light vector U boson, which is postulated to account for the possible fifth force or modification to Newtonian gravity [20, 67–69], is included to comparatively check the repulsive role in the zero-sounds in nuclear matter. Similar to the form of the vector meson, $\mathcal{L}_u$ is written as [23]

$$\mathcal{L}_u = -\bar{\psi} g_u \gamma_\mu u^\mu \psi - \frac{1}{4} U_{\mu\nu} U^{\mu\nu} + \frac{1}{2} m_u^2 u_\mu u^\mu, \quad (3)$$

where g_u, m_u , and $U_{\mu\nu}$ are the coupling constant, mass, and strength tensor of the U boson, which is in a form similar to that of the ω meson.

The nonlinear equations of motion for nucleons and mesons are deduced from the standard Euler-Lagrange formula. In the mean-field approximation, meson fields in uniform nuclear matter are replaced by their expectation values, obtained through iteration. With these mean fields, the effective meson masses m_i^* , which are derived from the second derivative of the Lagrangian, are given as follows:

$$m_\sigma^{*2} = m_\sigma^2 + 2g_2 \sigma_0 + 3g_3 \sigma_0^2, \quad (4a)$$

$$m_\omega^{*2} = m_\omega^2 + 3c_3 \omega_0^2 + 8\Lambda_V g_\omega^2 g_\rho^2 b_0^2, \quad (4b)$$

$$m_\rho^{*2} = m_\rho^2 + 8\Lambda_V g_\omega^2 g_\rho^2 \omega_0^2. \quad (4c)$$

The resulting energy density ε and pressure P of nuclear matter are written as follows:

$$\begin{aligned} \varepsilon = & \sum_{i=p,n} \frac{2}{(2\pi)^3} \int^{k_{F_i}} d^3k E_i^* + \frac{1}{2} m_\omega^2 \omega_0^2 + \frac{1}{2} m_\sigma^2 \sigma_0^2 \\ & + \frac{1}{2} m_\rho^2 b_0^2 + \frac{1}{3} g_2 \sigma_0^3 + \frac{1}{4} g_3 \sigma_0^4 + \frac{3}{4} c_3 \omega_0^4 + \frac{1}{2} \frac{g_u^2}{m_u^2} \rho_B^2 \\ & + 12 \Lambda_V g_\rho^2 g_\omega^2 \omega_0^2 b_0^2, \end{aligned} \quad (5)$$

$$\begin{aligned} P = & \frac{1}{3} \sum_{i=p,n} \frac{2}{(2\pi)^3} \int^{k_{F_i}} d^3k \frac{\mathbf{k}^2}{E_i^*} + \frac{1}{2} m_\omega^2 \omega_0^2 - \frac{1}{2} m_\sigma^2 \sigma_0^2 \\ & + \frac{1}{2} m_\rho^2 b_0^2 - \frac{1}{3} g_2 \sigma_0^3 - \frac{1}{4} g_3 \sigma_0^4 + \frac{1}{4} c_3 \omega_0^4 + \frac{1}{2} \frac{g_u^2}{m_u^2} \rho_B^2 \\ & + 4 \Lambda_V g_\rho^2 g_\omega^2 \omega_0^2 b_0^2, \end{aligned} \quad (6)$$

$$\text{with } E_i^* = \sqrt{\mathbf{k}^2 + (M_i^*)^2}.$$

B. Polarization tensors in RPA

In the RRPA method, one can, in principle, involve the particle-hole and particle-antiparticle excitations of the fermions through various polarization tensors, while within the framework of the RMF models, the particle-antiparticle excitations are usually ignored in a no-sea approximation [40]. The interacting polarization is determined through the Dyson equation. For instance, the interacting longitudinal component of the vector polarization tensor $\Pi_{\mu\nu}$, which is usually used to search for the collective modes, is given as

$$\tilde{\Pi}_L = \Pi_L + \Pi_L D_L \tilde{\Pi}_L, \quad (7)$$

where the polarization Π_L and meson propagator D_L are in matrix form. The lowest-order longitudinal polarization Π_L in nuclear matter is written as

$$\Pi_L = \begin{pmatrix} \Pi_s^n + \Pi_s^p & \Pi_m^p & \Pi_m^n \\ \Pi_m^p & \Pi_{00}^p & 0 \\ \Pi_m^n & 0 & \Pi_{00}^n \end{pmatrix}, \quad (8)$$

where the individual polarization entries are given by:

$$i\Pi_s^a(q, q_0) = \int \frac{d^4p}{(2\pi)^4} \text{Tr}[G^a(p)G^a(p+q)], \quad (9a)$$

$$i\Pi_m^a(q, q_0) = \int \frac{d^4p}{(2\pi)^4} \text{Tr}[G^a(p)\gamma_0 G^a(p+q)], \quad (9b)$$

$$i\Pi_{00}^a(q, q_0) = \int \frac{d^4p}{(2\pi)^4} \text{Tr}[G^a(p)\gamma_0 G^a(p+q)\gamma_0], \quad (9c)$$

With Tr indicating the trace over Dirac indices and

$a = p, n$, the fermion Green function reads

$$\begin{aligned} G^a(k) = & (\gamma_\mu k^\mu + M_a^*) \left[\frac{1}{k_\mu^2 - M_a^{*2} + i\varepsilon} \right. \\ & \left. + \frac{i\pi}{E_{ka}^*} \delta(k_0 - E_{ka}^*) \theta(k_F^a - |\mathbf{k}|) \right], \end{aligned} \quad (10)$$

where $M_a^* = M_a - g_\sigma \sigma_0$ is the nucleon effective mass, and k_F is the Fermi momentum.

With the crossing coupling term of the ρ and ω mesons in Eq. (1), the meson propagator D_L is given as [53, 70]

$$D_L = \begin{pmatrix} \chi_\sigma & 0 & 0 \\ 0 & \tilde{\chi}_\omega + \tilde{\chi}_\rho + 2\tilde{\chi}_{\omega\rho} & \tilde{\chi}_\omega - \tilde{\chi}_\rho \\ 0 & \tilde{\chi}_\omega - \tilde{\chi}_\rho & \tilde{\chi}_\omega + \tilde{\chi}_\rho - 2\tilde{\chi}_{\omega\rho} \end{pmatrix}. \quad (11)$$

Expressions for the various meson propagators with nonlinear meson couplings are provided as follows:

$$\chi_\sigma = \frac{g_\sigma^2}{q_\mu^2 - m_\sigma^{*2}}, \quad (12a)$$

$$\chi_\omega = \frac{g_\omega^2 (q_\mu^2 - m_\omega^{*2})}{(q_\mu^2 - m_\omega^{*2})(q_\mu^2 - m_\omega^{*2}) - (16\Lambda_V g_\omega^2 g_\rho^2 \omega_0 b_0)^2}, \quad (12b)$$

$$\chi_\rho = \frac{g_\rho^2 (q_\mu^2 - m_\rho^{*2})}{(q_\mu^2 - m_\rho^{*2})(q_\mu^2 - m_\rho^{*2}) - (16\Lambda_V g_\omega^2 g_\rho^2 \omega_0 b_0)^2}, \quad (12c)$$

$$\chi_{\omega\rho} = \frac{-16\Lambda_V g_\omega^3 g_\rho^3 \omega_0 b_0}{(q_\mu^2 - m_\rho^{*2})(q_\mu^2 - m_\omega^{*2}) - (16\Lambda_V g_\omega^2 g_\rho^2 \omega_0 b_0)^2}, \quad (12d)$$

$$\tilde{\chi}_i = \frac{q_\mu^2}{q^2} \chi_i, \quad i = \omega, \rho. \quad (12e)$$

When the vector repulsion of the U boson is included, χ_ω is replaced by χ'_ω to modify the Dyson equation.

$$\chi'_\omega = \chi_\omega + \chi_u, \quad \chi_u = \frac{g_u^2}{q_\mu^2 - m_u^{*2}}. \quad (13)$$

The nuclear systems undergo transitions at small-amplitude density fluctuations by encountering the zeros of the following dielectric function.

$$\varepsilon_L = \det(1 - D_L \Pi_L) = 0. \quad (14)$$

III. RESULTS AND DISCUSSIONS

In our work, we adopt two well-fitting but relatively soft parameter sets, TM1 and FSUGarnet, to investigate the impact of the U boson on zero-sound modes. To improve the consistency of neutron star radius predictions in the TM1 model with observational data, we soften the symmetry energy by introducing an isoscalar-isovector (ω - ρ) coupling term (Λ_V) [53, 66, 71]. The modified TM1 model is referred to as TM1w02. The parameters and saturation properties of these two models are listed in Table 1.

In Fig. 1, the pressure as a function of the energy density of symmetric nuclear matter is shown with the inclusion of the U boson in parameter sets TM1w02 and FSUGarnet. The EOS trajectories are depicted for a set of ratios g_u^2/m_u^2 . In the RMF approximation, the contribution of the U boson is determined linearly by the ratio of the coupling constant to its mass, i.e., g_u/m_u , as seen in Eqs. (5) and (6). The actual values of g_u and m_u lack sufficient constraints and are largely uncertain. As argued by Krivoruchenko et al. [20], $m_u \leq 30$ MeV and the g_u^2/m_u^2 value of the U boson should be less than about 300 GeV^{-2} , which roughly corresponds to the value of the ordinary vector ω meson. To escape detection in terrestrial measurements on nuclei, it was pointed out that the mass of the weakly coupled U boson is on the order of 1 MeV [23]. In this work, the ratio is chosen to be beneath 300 GeV^{-2} . It is observed that the inclusion of the U boson can significantly stiffen the nuclear EOS with the TM1w02 and FSUGarnet. This is physically evident since the vector U boson provides additional repulsion in addition to the vector mesons.

Zero sounds are determined by the zeros of the dielectric function and are the collective modes following the dispersion relation with the zero-energy limit. At small momentum transfer q , zero sounds have similar properties, and for numerical conciseness, the momentum $q = 10 \text{ MeV}$ is typically chosen in our calculations. In previous studies [53], it has been found that zero-sound modes at lower densities (typically $\rho_B \lesssim 2\rho_0$) are dominated by the ρ meson and are referred to as isovector-dominated or isospin zero-sound modes. These modes are highly insensitive to the stiffness of the EOS. In contrast,

zero-sound modes that emerge at higher densities are primarily governed by the σ and ω mesons and are thus termed isoscalar-dominated zero-sound modes. The occurrence of these high-density modes has been shown to be strongly correlated with the EOS stiffness in both symmetric and asymmetric nuclear matter. In soft EOS models, these modes emerge only when the nonlinear self-interaction term of the ω meson ($c_3\omega_0^4$, see Eq. (1)) is reduced to yield a stiffer EOS. In this work, we examine whether the EOS stiffening induced by the U boson can have a similar effect on the occurrence of isoscalar-dominated zero-sound modes and further revisit the relationship between high-density zero-sound modes and the stiffness of the EOS. In contrast to the RMF EOS, the mass and coupling constant of the U boson are both contributive in RPA (see Eq. (13)). As mentioned above, the mass of the weakly coupling U boson is roughly of the order of 1 MeV [23]. Without loss of generality, the U boson mass is assumed to be 2 MeV or 4 MeV in the present RPA calculation, given the ratio g_u^2/m_u^2 beneath 300 GeV^{-2} .

It is interesting to examine how the U boson-induced modification to the EOS affects the behavior of zero sound. As shown in Figs. 2 and 3, the isoscalar-dominated zero-sound modes arise in the models TM1w02 and FSUGarnet at high density due to the stiffening role of the U boson in the EOS. Without the inclusion of the U boson, such isoscalar-dominated modes do not appear in these models at high density; only isovector-dominated zero-sound modes, mainly driven by the ρ meson, emerge at lower densities, which are generally insensitive to the stiffness of the EOS. This results from the fact that the U boson provides an excess repulsion to overwhelm the attractive interaction at high density. The reproduction of the zero-sound modes by the U boson repulsion is similar to that caused by stiffening the repulsion of the ω meson [53], providing additional evidence that the occurrence of zero-sound modes can signal the stiffness of the nuclear EOS. As the EOS becomes stiffer by increasing the coupling term of the U boson, we can see from the inset of Figs. 2 and 3 that the high-density zero-sound modes start to appear at $g_u^2/m_u^2 = 95$ (200) GeV^{-2} in TM1w02 (FSUGarnet), with the region expansion of zero-sound modes for larger coupling strength. It is worth noting that the coupling constant of the U boson for the onset of zero sounds in FSUGarnet is larger than that in TM1w02. This difference is mainly attributed to the lar-

Table 1. Parameters and saturation properties for the parameter sets TM1w02 and FSUGarnet. The meson masses m_i ($i = \sigma, \omega, \rho$), the incompressibility K_0 , and the symmetry energy E_{sym} are in units of MeV. The saturation density ρ_0 is in units of fm^{-3} .

	g_σ	g_ω	g_ρ	m_σ	m_ω	m_ρ	g_2	g_3	c_3	Λ_V	ρ_0	K_0	M^*/M	E_{sym}
TM1w02	10.029	12.614	5.277	511.198	783.000	770	7.233	0.618	71.31	0.02	0.145	281.20	0.63	30.7
FSUGarnet	10.505	13.700	6.945	496.939	782.500	763	9.576	-7.207	137.981	0.04338	0.153	229.628	0.58	30.9




Fig. 1. (Color online) The relationship between pressure and energy density for various equations of state (EOS) with different g_u^2/m_u^2 in TM1w02 and FSUGarnet




Fig. 2. (Color online) Isospin zero-sound modes (at low density) and isoscalar dominated zero-sound modes (at high density) in symmetric nuclear matter with the TM1w02 model. The U boson is included with varying coupling ratio parameters g_u^2/m_u^2 , and its mass is fixed at 2 MeV. The inset separately shows isoscalar dominated zero-sound modes for different U boson coupling parameters.

ger coupling coefficient Λ_V between the ω and ρ mesons, which increases the effective mass of the ρ meson but suppresses the magnitude of the ρ meson propagator. Consequently, this leads to a suppression of the contribution of the $\chi_u \chi_\rho$ term to the vector part of the dielectric




Fig. 3. (Color online) The same as in Fig. 2, but for the RMF model FSUGarnet with different U boson coupling strengths.

function, and a larger coupling constant of the U boson is needed for compensation. The onset of isoscalar-dominated zero-sound modes occurs at a slightly higher baryon density in FSUGarnet than in TM1w02, due to differences in their scalar meson self-interactions and effective meson masses, which affect the delicate balance between vector repulsion and scalar attraction. This balance plays a crucial role in determining the zeros of the dielectric function and thus the emergence of collective modes.

For isovector-dominated zero-sound modes, their behavior is independent of the stiffness of the EOS, and therefore the introduction of the U boson has no impact on them.

Figure 4 shows that when the mass of the U boson is fixed at 4 MeV, the values of g_u^2/m_u^2 , at which the isoscalar-dominated zero-sound modes begin to appear, are 50 (100) GeV^{-2} for TM1w02 (FSUGarnet). The reduction in the threshold value of g_u^2/m_u^2 is evident. As shown in Eq. (13), the magnitude of χ_u is proportional to both the mass m_u and the ratio g_u^2/m_u^2 . As m_u increases, the g_u^2/m_u^2 decreases for the same χ_u required by the emergence of isoscalar-dominated zero-sound modes.

In contrast to the case in symmetric matter, the distinction between isoscalar and isovector dominated zero-sound modes becomes blurred in asymmetric matter. For the soft EOS models, such as TM1w02, the emergence of zero-sound modes in asymmetric nuclear matter similarly requires stiffening the EOS. As shown in Fig. 5, the inclusion of the U boson enables the appearance of zero-sound modes in asymmetric matter. Notably, a lower isospin asymmetry parameter α ($\alpha = (\rho_n - \rho_p)/\rho_B$) leads to a lower onset density of the zero-sound modes, as isospin asymmetry influences the delicate balance of the dielectric function. Similar to symmetric matter, in asymmetric




Fig. 4. (Color online) Isoscalar-dominated zero-sound modes in symmetric nuclear matter with the TM1w02 and FSUGarnet models. The mass of the U boson is fixed at 4 MeV.




Fig. 5. (Color online) Zero sound in asymmetric nuclear matter ($\alpha = 0.3, 0.5$) with TM1w02. The mass of the U boson is set to 2 MeV, and the interaction ratio g_u^2/m_u^2 is set to 150 GeV^{-2} .

matter, the larger the U boson coupling, the wider the density range over which zero-sound modes appear, as shown in Fig. 6 for the TM1w02 model. Note that Fig. 6 adopts $m_u = 4$ MeV, while Fig. 5 uses $m_u = 2$ MeV, for which zero-sound modes appear only when $g_u^2/m_u^2 \gtrsim 150 \text{ GeV}^{-2}$. Unlike the case in symmetric matter, where zero sound consists of the undamped (the one with larger q_0) and damped branches, multiple zero-sound branches ap-




Fig. 6. (Color online) Zero sounds in asymmetric nuclear matter with TM1w02 for various g_u^2/m_u^2 . The mass of the U boson is fixed at 4 MeV. The left panels represent the results with $\alpha = 0.3$, and the right panels with $\alpha = 0.5$. The orange lines represent the undamped branches of zero sound, and the green lines represent the damped branches.

pear in asymmetric matter. Among them, the branch with the largest q_0 is undamped, while the others are damped. Therefore, it can be concluded that the stiffening of the EOS by the U boson allows zero-sound modes to emerge in both symmetric and asymmetric matter.

Furthermore, we investigate the effect of the U boson on zero-sound modes in the β -equilibrated neutron star matter. Zero-sound modes in neutron star matter can possibly be associated with neutron star cooling and pulsar radiation [55]. Investigating the correlation between zero sound and the stiffness of the EOS is essential for understanding astrophysical phenomena in the multimessenger era. Figure 7 shows the zero-sound modes in neutron star matter with various g_u^2/m_u^2 . We can see that the damped zero-sound modes are similar across different values of g_u^2/m_u^2 , showing little sensitivity to the degree of EOS stiffening, whereas the undamped modes exhibit significant dependence on the coupling strength of the U boson. As g_u^2/m_u^2 increases, the range for the undamped zero-sound modes gradually extends to higher densities.

It is worth noting that the repulsion of the U boson results in an increase in the predicted maximum mass of neutron stars. This is advantageous in satisfying the lower limit of the maximum mass of neutron stars around $2M_\odot$ [2, 3]. Meanwhile, the additional repulsion of the U bo-




Fig. 7. (Color online) The same as in Fig. 6, but for neutron star matter.

son also tends to increase the predicted neutron star radii [21, 23], potentially exceeding observational bounds. This tension can be alleviated by adopting a density-dependent U boson coupling to merely reduce the star radius [72]. Moreover, the zero-sound modes have been proposed as potential Goldstone bosons that possibly couple to neutrino emission channels via Z^0 -boson exchange [55]. Thus, it suggests that facilitating the onset of zero sound by the U boson may affect neutron star cooling and long-term evolution through additional neutrino emission, which warrants further investigation.

Lastly, it is necessary to point out that in the FSUGarnet model, the dependence of the zero sound on the U boson repulsion in asymmetric nuclear matter and neutron star matter is similar to that in the TM1w02 model, although a larger g_u^2/m_u^2 is needed for the FSUGarnet model as in symmetric matter. To save space, the results with the FSUGarnet model are not presented here.

IV. SUMMARY

In this work, we have studied the effects of a light vector U boson on zero-sound modes in nuclear matter

and neutron star matter using the RRPA method. By employing two typical RMF models with a soft high-density EOS, our results show that the stiffening of the EOS by the U boson promotes the emergence of zero-sound modes at high densities.

In symmetric nuclear matter, low-density isovector zero-sound modes are not affected by the U boson, whereas the isoscalar-dominated zero-sound modes at high density are recreated in the RMF models with a soft EOS through the inclusion of the repulsion provided by the U boson. The larger coupling strength of the U boson further broadens their density range for zero sound to the high-density end.

In asymmetric nuclear matter, the role of the U boson in recreating zero-sound modes in RMF models with a soft EOS is similar to that in symmetric matter. The repulsion provided by the U boson supports the emergence of multiple zero-sound branches, either damped or undamped. Moreover, the inherent isospin asymmetry disrupts the balance of the different terms in the dielectric function, resulting in a blurred distinction between isovector- and isoscalar-dominated zero-sound modes.

In neutron star matter, multiple zero-sound modes in RMF models with a soft EOS are recovered by the U boson, similar to that in asymmetric nuclear matter. Interestingly, the presence of the U boson extends the density region for undamped zero-sound modes. If zero sounds couple to neutrino emission, the U boson that facilitates the creation of zero sounds may affect neutron star dynamics.

In comparison to zero sound creation through EOS stiffening by adjusting the ω self-interaction strength, the inclusion of the U boson achieves similar consequences. This confirms the unambiguous role of stiffening the high-density EOS in resulting in the appearance of zero sounds. As the deviation from the inverse square law of gravity due to the inclusion of the U boson can be alternatively expressed by its modification to the EOS [21], our study implies that the zero sound facilitated by the U boson in dense matter may be correlated with modified gravity, which would make the study of zero sound in dense matter particularly informative.

ACKNOWLEDGEMENTS

Additionally, we acknowledge the Big Data Computing Center of Southeast University for providing facility support for the partial numerical calculations in this study.

References

- [1] H. T. Cromartie, E. Fonseca, S. M. Ransom, P. B. Demorest, Z. Arzoumanian, H. Blumer, P. R. Brook, M. E. DeCesar, T. Dolch, J. A. Ellis, *et al.*, *Nature Astronomy* **4**, 72 (2020)
- [2] P. B. Demorest, T. Pennucci, S. Ransom, M. Roberts, and J. Hessels, *nature* **467**, 1081 (2010)

- [3] J. Antoniadis, P. C. Freire, N. Wex, T. M. Tauris, R. S. Lynch, M. H. Van Kerkwijk, M. Kramer, C. Bassa, V. S. Dhillon, T. Driebe, *et al.*, *Science* **340**, 1233232 (2013)
- [4] E. Fonseca, T. T. Pennucci, J. A. Ellis, I. H. Stairs, D. J. Nice, S. M. Ransom, P. B. Demorest, Z. Arzoumanian, K. Crowter, T. Dolch, *et al.*, *The Astrophysical Journal* **832**, 167 (2016)
- [5] B. P. Abbott, R. Abbott, T. Abbott, F. Acernese, K. Ackley, C. Adams, T. Adams, P. Addesso, R. X. Adhikari, V. B. Adya, *et al.*, *Physical Review Letters* **119**, 161101 (2017)
- [6] A. Goldstein, P. Veres, E. Burns, M. Briggs, R. Hamburg, D. Kocevski, C. Wilson Hodge, R. Preece, S. Poolakkil, O. Roberts, *et al.*, *The Astrophysical Journal Letters* **848**, L14 (2017)
- [7] B. D. Metzger, Kilonovae, *Living reviews in relativity* **20**, 3 (2017).
- [8] P. Danielewicz, R. Lacey, and W. G. Lynch, *Science* **298**, 1592 (2002)
- [9] B. P. Abbott, R. Abbott, T. Abbott, F. Acernese, K. Ackley, C. Adams, T. Adams, P. Addesso, R. X. Adhikari, V. B. Adya, *et al.*, *Physical Review Letters* **121**, 161101 (2018)
- [10] J. M. Lattimer and M. Prakash, *Physics Reports* **442**, 109 (2007)
- [11] W. Z. Jiang, B. A. Li, and L. W. Chen, *The Astrophysical Journal* **756**, 56 (2012)
- [12] D. Radice, A. Perego, K. Hotokezaka, S. A. Fromm, S. Bernuzzi, and L. F. Roberts, *The Astrophysical Journal* **869**, 130 (2018)
- [13] P. B. Demorest, T. Pennucci, S. Ransom, M. Roberts, and J. Hessels, *Nature* **467**, 1081 (2010)
- [14] J. Antoniadis, P. C. Freire, N. Wex, T. M. Tauris, R. S. Lynch, M. H. Van Kerkwijk, M. Kramer, C. Bassa, V. S. Dhillon, T. Driebe, *et al.*, *Science* **340**, 1233232 (2013)
- [15] H. T. Cromartie, E. Fonseca, S. M. Ransom, P. B. Demorest, Z. Arzoumanian, H. Blumer, P. R. Brook, M. E. DeCesar, T. Dolch, J. A. Ellis, *et al.*, *Nature Astronomy* **4**, 72 (2020)
- [16] F. ?zel, *Nature* **441**, 1115 (2006)
- [17] T. Klähn, D. Blaschke, F. Sandin, C. Fuchs, A. Faessler, H. Grigorian, G. Röpke, and J. Trümper, *Physics Letters B* **654**, 170 (2007)
- [18] C. Jockel and L. Sagunski, *Particles* **7**, 52 (2024)
- [19] D. Rafiei Karkevandi, M. Shahrbafe, S. Shakeri, and S. Typel, *Particles* **7**, 201 (2024)
- [20] M. Krivoruchenko, F. Šimkovic, and A. Faessler, *Physical Review D* **79**, 125023 (2009)
- [21] D. H. Wen, B. A. Li, and L. W. Chen, *Physical Review Letters* **103**, 211102 (2009)
- [22] A. Sulaksono, MARLIANA, and KASMUDIN, *Modern Physics Letters A* **26**, 367 (2011)
- [23] D. R. Zhang, P. L. Yin, W. Wang, Q. C. Wang, and W. Z. Jiang, *Physical Review C* **83**, 035801 (2011)
- [24] W. Lin, B. A. Li, L. W. Chen, D. H. Wen, and J. Xu, *Journal of Physics G* **41**, 075203 (2014)
- [25] S. H. Yang, C. M. Pi, X. P. Zheng, and F. Weber, *The Astrophysical Journal* **902**, 32 (2020)
- [26] P. Fayet, *Physics Letters B* **95**, 285 (1980)
- [27] P. Fayet, *Nuclear Physics B* **187**, 184 (1981)
- [28] S. H. Yang, C. M. Pi, X. P. Zheng, and F. Weber, *Physical Review D* **103**, 043012 (2021)
- [29] J. Xu, B. A. Li, L. W. Chen, and H. Zheng, Nonnewtonian gravity in finite nuclei, in *Journal of Physics: Conference Series*, Vol. 420 (IOP Publishing, 2013) p. 012155.
- [30] G. C. Yong and B. A. Li, *Physics Letters B* **723**, 388 (2013)
- [31] P. Jean, J. Knödlseider, V. Lonjou, M. Allain, J.-P. Roques, G. Skinner, B. Teegarden, G. Vedrenne, P. von Ballmoos, B. Cordier, *et al.*, *Astronomy & Astrophysics* **407**, L55 (2003)
- [32] C. Boehm, P. Fayet, and J. Silk, *Physical Review D* **69**, 101302 (2004)
- [33] C. Boehm, D. Hooper, J. Silk, M. Casse, and J. Paul, *Physical Review Letters* **92**, 101301 (2004)
- [34] N. Borodatchenkova, D. Choudhury, and M. Drees, *Physical review letters* **96**, 141802 (2006)
- [35] S. h. Zhu, *Physical Review D* **75**, 115004 (2007)
- [36] P. Fayet, *Physical Review D* **75**, 115017 (2007)
- [37] H. Zheng and L. W. Chen, *Physical Review D* **85**, 043013 (2012)
- [38] J. Xu, B. A. Li, L. W. Chen, and H. Zheng, *Journal of Physics G* **40**, 035107 (2013)
- [39] W. Z. Jiang, B. A. Li, and F. Fattoyev, *The European Physical Journal A* **51**, 1 (2015)
- [40] Z. y. Ma, N. Van Giai, A. Wandelt, D. Vretenar, and P. Ring, *Nuclear Physics A* **686**, 173 (2001)
- [41] J. Stone, N. Stone, and S. Moszkowski, *Physical Review C* **89**, 044316 (2014)
- [42] U. Garg and G. Colo, *Progress in Particle and Nuclear Physics* **101**, 55 (2018)
- [43] L. Trippa, G. Colo, and E. Vigezzi, *Physical Review C* **77**, 061304 (2008)
- [44] J. M. Lattimer and Y. Lim, *The Astrophysical Journal* **771**, 51 (2013)
- [45] M. N. Harakeh and A. Woude, *Giant Resonances: fundamental high-frequency modes of nuclear excitation*, Vol. 24 (Oxford University Press on Demand, 2001).
- [46] X. Roca Maza, M. Brenna, B. Agrawal, P. Bortignon, G. Colo, L. G. Cao, N. Paar, and D. Vretenar, *Physical Review C* **87**, 034301 (2013)
- [47] S. A. Chin, *Annals of Physics* **108**, 301 (1977)
- [48] K. Lim and C. Horowitz, *Nuclear Physics A* **501**, 729 (1989)
- [49] J. Caillon, P. Gabinski, and J. Labarsouque, *Nuclear Physics A* **696**, 623 (2001)
- [50] F. Braghin and D. Vautherin, *Physics Letters B* **333**, 289 (1994)
- [51] F. L. Braghin, D. Vautherin, and A. Abada, *Physical Review C* **52**, 2504 (1995)
- [52] F. L. Braghin, *Nuclear Physics A* **665**, 13 (2000)
- [53] J. Ye, J. Margueron, N. Li, and W. Jiang, *Phys. Rev. C* **108**, 044312 (2023)
- [54] J. Ye and W. Jiang, Thermal properties of zero sound in asymmetrical nuclear matter, accepted by *Phys. Rev. C* (2025).
- [55] P. F. Bedaque, G. Rupak, and M. J. Savage, *Physical Review C* **68**, 065802 (2003)
- [56] J. M. Lattimer and M. Prakash, *Science* **304**, 536 (2004)
- [57] J. Pons, S. Reddy, M. Prakash, J. Lattimer, and J. Miralles, *The Astrophysical Journal* **513**, 780 (1999)
- [58] Z. Ma, H. Toki, and N. Van Giai, *Nuclear Physics A* **627**, 1 (1997)
- [59] C. De Conti, A. Galeao, and F. Krmpotić, *Physics Letters B* **444**, 14 (1998)
- [60] D. Vretenar, N. Paar, T. Nikšić, and P. Ring, *Physical Review Letters* **91**, 262502 (2003)
- [61] H. Liang, N. Van Giai, and J. Meng, *Physical review letters* **101**, 122502 (2008)

- [62] D. Yang, L. G. Cao, and Z. Y. Ma, *Chinese Physics C* **37**, 124102 (2013)
- [63] T. Matsui, *Nuclear Physics A* **370**, 365 (1981)
- [64] C. Horowitz and B. D. Serot, *Nuclear Physics A* **399**, 529 (1983)
- [65] B. D. Serot and J. D. Walecka, Relativistic nuclear manybody theory, in *Recent Progress in Many-Body Theories* (Springer, 1992) pp. 49–92.
- [66] W. Z. Jiang and Y. L. Zhao, *Physics Letters B* **617**, 33 (2005)
- [67] P. Fayet, *Physics Letters B* **95**, 285 (1980)
- [68] P. Fayet, *Nuclear Physics B* **347**, 743 (1990)
- [69] E. Adelberger, B. Heckel, and A. Nelson, *Annual Review of Nuclear and Particle Science* **53**, 77 (2003)
- [70] J. Carriere, C. Horowitz, and J. Piekarewicz, *Astrophys. J.* **593**, 463 (2003)
- [71] C. Horowitz and J. Piekarewicz, *Physical Review Letters* **86**, 5647 (2001)
- [72] D. R. Zhang, W. Z. Jiang, S. N. Wei, R. Y. Yang, and Q. F. Xiang, *The European Physical Journal A* **52**, 142 (2016)