

Tensor formalism for the partial wave analysis of reactions with resonances decaying into four pseudoscalar mesons*

M.A. Matveev A.T. Sitnikov A.V. Sarantsev[†]

NRC "Kurchatov Institute", Petersburg Nuclear Physica Institute, 188300 Gatchina, Russia

Abstract: We construct a formalism which describes the resonances decaying into four pseudoscalar meson final states. This method is fully covariant and can be directly applied for the partial-wave analysis of high statistical data. Two topologies of the process are considered: two intermediate resonances each decaying into two final mesons and cascade decay via three meson intermediate states. In particular, we consider the production of such states in the central collision reactions and in radiative J/Ψ decay.

Keywords: partial-wave analysis, radiative J/ψ decay, four-meson final state, Glueballs, Resonance, strong QCD

DOI: **CSTR:**

The main information about meson states comes from the πN scattering with high energy pion beams, from the antiproton-nucleon annihilation, from the decay of relatively stable, heavy hadrons and from the production of mesons in the central collision. In most of the reactions analyzed, resonances decay into final states with two stable particles (see, for example, [1–5]), or the reactions can be considered as quasi-two-particle scattering (for example, antiproton-proton annihilation at rest into three pseudoscalar mesons [6–8]). We note that some of the developments for multichannel decays were made for the analysis of radiative decay J/Ψ [9], the decay of heavy mesons [10–12] for the annihilation of the antiproton-proton at rest into five pions [13]. However, no systematic formalism for the analysis of the data in which meson resonances decay to a four-meson final state had been developed. Nevertheless, such a decay mode is a dominant one for the resonances in many partial waves already at masses around 1400 MeV.

In the scalar isoscalar sector, the decay mode 4π is already a dominant one for the $f_0(1370)$ state. Moreover, in the elastic scattering data $\pi\pi$ (extracted from the reaction πN) this state can only appear due to rescattering with other scalar states in the 4π channel and cannot be clearly observed. This property simulated a number of discussions about the existence of this state. However, this state was observed in the antiproton-proton in the analysis of the data with the final states $3\pi^0$ [6] and 5π

[13]. We mention that the scalar isoscalar sector is rather difficult to analyze. In this sector, one expects a strong mixing between non-strange and strange quark-antiquark components and a production of exotic states. For example, many authors treat lowest-scalar states as molecular-like states or as four-quark bound systems. The lowest bound states of two gluons: glueballs are also expected in this sector [14]. Such a picture makes the analysis of this sector a rather complicated procedure, and information about four-meson final states can be vital for understanding the spectrum and properties of these mesons.

In the present paper, we construct the covariant approach for the analysis of the resonances decaying into four pseudoscalar mesons and consider the production of these resonances in J/Ψ radiative decay and in the NN central collision reactions (pomeron-pomeron scattering). The J/Ψ radiative decay is one of the main sources of the search for glueball states. In fact, the partial wave decomposition of the BES III data on the J/ψ radiative decay into two pseudo-scalar mesons [4, 5] demonstrated a very complicated resonant structures in the isoscalar-scalar partial wave in the mass region 1500-2100 MeV. The combined analysis of these data with the scattering data from $\pi\pi$ and the data on the annihilation of the proton-antiproton at rest in three pseudoscalar mesons revealed the contribution of ten scalar states [15]. The distribution of resonance production intensities demonstrated a peak in the mass region of 1850 MeV, which was explained by

Received 2 June 2025; Accepted 17 October 2025

* The paper is supported by the RSF grant 24-22-00322

[†] E-mail: sarantsev_AV@pnpi.nrcki.ru (Corresponding author)



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Article funded by SCOAP³ and published under licence by Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd

the presence of two-gluon components in the observed scalar states. This idea was confirmed by the calculation of the mixing angles between $n\bar{n}$, $s\bar{s}$ and the glueball components [16]. In the alternative analysis of these data [17] it was declared that the resonance with mass around 1750 MeV is produced dominantly and is the main candidate to the scalar glueball. The analysis of the data with the four-pion channel should resolve this issue.

Another issue is the search for the tensor glueball. If the J/Ψ radiative decay into two pseudoscalar mesons reveal the production of the scalar glueball in the mass region 1700-1900 MeV then one should expect the production of the tensor glueball (for example from lattice calculations [18, 19]) in the mass region 2200-2500 MeV. The tensor partial waves extracted from the BES III data showed a strong production of $f_2(1275)$ (decaying into 2π final state) and $f_2'(1525)$ (decaying into $K\bar{K}$ final state) and practically no structure at higher energies. In fact, no clear signal was found in the analysis [20]. The only solution to this problem is related to the fact that tensor states in this mass region decay dominantly into four pseudoscalar mesons, and signals from the tensor glueball should be searched in the data on the J/Ψ radiative decay into these final states.

Another prominent source for the production of the glueball states is in the meson production at nucleon-nucleon central collision reactions. In these processes, the states are predominantly produced from the pomeron-pomeron collision. Taking into account that the pomeron is an effective way to describe the gluon lattice, it is relevant to expect that the states with a large gluon component will be strongly produced in such a reaction. Therefore, our method should be useful for planning new experiments for the new constructed colliders like the NICA.

The covariant approach based on the tensor formalism was put forward by Zemach [21] and developed further in [22], [23]. The mathematical framework allowing the construction of angular momentum operators using recursive expressions was provided in [24]. This method was developed for the case of fermions in [25]. We should mention that in a number of the analyses the authors used the approach based on the helicity formalism. In that formalism, particle propagators are represented as a product of polarization tensors (summed over possible polarizations). In this case, the total amplitude is equal to the sum of products of amplitudes describing two particle scattering, which can be calculated in the helicity basis. However, such two-particle amplitudes should be carefully rotated, which is not a trivial task. For the case of the J/Ψ -radiative decay into two pseudoscalar mesons, the exact correspondence between the helicity and the tensor approach was given in [26]. However, in the case of the resonance decay into a four-particle state, the helicity approach is much more complicated than a tensor

formalism.

I. THE COVARIANT SPIN-ORBITAL FORMALISM

A. Orbital angular momentum tensor

The most detailed description of the tensor formalism is given in [25]. We briefly recall it here. Consider the decay of a composite system with spin J and momentum P ($P^2 = s$) into two spinless particles with momenta k_1 and k_2 . The only quantities measured in such a reaction are the particle momenta. The angular dependent part of the wave function of the composite state is described by tensors constructed out of these momenta and the metric tensor. Such tensors (we denote them as $X_{\mu_1 \dots \mu_L}^{(L)}$, where L is the orbital momentum) are called orbital angular momentum tensors and correspond to irreducible representations of the Lorentz group. They satisfy the following properties:

- Symmetry under permutation of any two indices:

$$X_{\mu_1 \dots \mu_i \dots \mu_j \dots \mu_L}^{(L)} = X_{\mu_1 \dots \mu_j \dots \mu_i \dots \mu_L}^{(L)}. \quad (1)$$

- Orthogonality to the total momentum of the system, $P = k_1 + k_2$.

$$P_{\mu_i} X_{\mu_1 \dots \mu_i \dots \mu_L}^{(L)} = 0 \quad (2)$$

- The traceless property for convolution of any two indices with metric tensor:

$$g_{\mu_i \mu_j} X_{\mu_1 \dots \mu_i \dots \mu_j \dots \mu_L}^{(L)} = 0. \quad (3)$$

The orthogonality condition (2) is automatically fulfilled if the tensors are constructed from the relative momenta k_μ^\perp and the tensor $g_{\mu\nu}^\perp$ orthogonal to the total momentum of the system:

$$k_\mu^\perp = \frac{1}{2}(k_1 - k_2)_\nu g_{\mu\nu}^\perp \quad g_{\mu\nu}^\perp = g_{\mu\nu} - \frac{P_\mu P_\nu}{s} \quad (4)$$

In the center-of-mass system (cms), where $P = (P_0, \vec{P}) = (\sqrt{s}, 0)$, the vector k^\perp is space-like: $k^\perp = (0, \vec{k})$.

The orbital tensor for $L = 0$ is a scalar value (for example, a unit), and the tensor for the orbital momentum $L = 1$ is a vector that can only be constructed from k_μ^\perp . The orbital angular momentum tensors for L up to three have the form:

$$\begin{aligned}
X^{(0)} &= 1, & X_{\mu}^{(1)} &= k_{\mu}^{\perp}, \\
X_{\mu_1\mu_2}^{(2)} &= \frac{3}{2} \left(k_{\mu_1}^{\perp} k_{\mu_2}^{\perp} - \frac{1}{3} k_{\perp}^2 g_{\mu_1\mu_2}^{\perp} \right), \\
X_{\mu_1\mu_2\mu_3}^{(3)} &= \frac{5}{2} \left[k_{\mu_1}^{\perp} k_{\mu_2}^{\perp} k_{\mu_3}^{\perp} - \frac{k_{\perp}^2}{5} (g_{\mu_1\mu_2}^{\perp} k_{\mu_3}^{\perp} + g_{\mu_1\mu_3}^{\perp} k_{\mu_2}^{\perp} + g_{\mu_2\mu_3}^{\perp} k_{\mu_1}^{\perp}) \right].
\end{aligned} \tag{5}$$

The tensors $X_{\mu_1 \dots \mu_L}^{(L)}$ for $L \geq 1$ can be constructed from the tensors with lower orbital momenta in the form of a recurrent expression:

$$\begin{aligned}
X_{\mu_1 \dots \mu_L}^{(L)} &= k_{\perp}^{\perp} Z_{\mu_1 \dots \mu_L}^{\alpha}, \\
Z_{\mu_1 \dots \mu_L}^{\alpha} &= \frac{2L-1}{L^2} \left(\sum_{i=1}^L X_{\mu_1 \dots \mu_{i-1} \mu_{i+1} \dots \mu_L}^{(L-1)} g_{\mu_i \alpha}^{\perp} \right. \\
&\quad \left. - \frac{2}{2L-1} \sum_{\substack{i,j=1 \\ i < j}}^L g_{\mu_i \mu_j}^{\perp} X_{\mu_1 \dots \mu_{i-1} \mu_{i+1} \dots \mu_{j-1} \mu_{j+1} \dots \mu_L}^{(L-1)} \right).
\end{aligned} \tag{6}$$

The normalization of the tensors is fixed by the convolution equality:

$$X_{\mu_1 \dots \mu_L}^{(L)} k_{\mu_L}^{\perp} = k_{\perp}^2 X_{\mu_1 \dots \mu_{L-1}}^{(L-1)}. \tag{7}$$

Iterating eq. (6) one obtains the following expression for the tensor $X_{\mu_1 \dots \mu_L}^{(L)}$:

$$\begin{aligned}
X_{\mu_1 \dots \mu_L}^{(L)}(k^{\perp}) &= \alpha(L) \left[k_{\mu_1}^{\perp} k_{\mu_2}^{\perp} k_{\mu_3}^{\perp} k_{\mu_4}^{\perp} \dots k_{\mu_L}^{\perp} \right. \\
&\quad \left. - \frac{k_{\perp}^2}{2L-1} \left(g_{\mu_1\mu_2}^{\perp} k_{\mu_3}^{\perp} k_{\mu_4}^{\perp} \dots k_{\mu_L}^{\perp} \right. \right. \\
&\quad \left. \left. + g_{\mu_1\mu_3}^{\perp} k_{\mu_2}^{\perp} k_{\mu_4}^{\perp} \dots k_{\mu_L}^{\perp} + \dots \right) \right. \\
&\quad \left. + \frac{k_{\perp}^4}{(2L-1)(2L-3)} \left(g_{\mu_1\mu_2}^{\perp} g_{\mu_3\mu_4}^{\perp} k_{\mu_5}^{\perp} k_{\mu_6}^{\perp} \dots k_{\mu_L}^{\perp} \right. \right. \\
&\quad \left. \left. + g_{\mu_1\mu_2}^{\perp} g_{\mu_3\mu_5}^{\perp} k_{\mu_4}^{\perp} k_{\mu_6}^{\perp} \dots k_{\mu_L}^{\perp} + \dots \right) + \dots \right].
\end{aligned} \tag{8}$$

where

$$\alpha(L) = \prod_{i=1}^L \frac{2i-1}{i} = \frac{(2L-1)!!}{L!}. \tag{9}$$

Using normalization condition we obtain:

$$X_{\mu_1 \dots \mu_L}^{(L)} X_{\mu_1 \dots \mu_L}^{(L)} = \alpha(L) (k_{\perp}^2)^L \tag{10}$$

B. The boson projection operator

The projection operator $O_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L}(P)$ for the partial wave

with angular momentum L is defined as:

$$\int \frac{d\Omega}{4\pi} X_{\mu_1 \dots \mu_L}^{(L)}(k^{\perp}) X_{\nu_1 \dots \nu_L}^{(L)}(k^{\perp}) = \frac{\alpha(L)}{2L+1} (k_{\perp}^2)^L O_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L}(P). \tag{11}$$

This tensor satisfies the following relations:

$$\begin{aligned}
X_{\mu_1 \dots \mu_L}^{(L)}(k^{\perp}) O_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L}(P) &= X_{\nu_1 \dots \nu_L}^{(L)}(k^{\perp}), \\
O_{\alpha_1 \dots \alpha_L}^{\mu_1 \dots \mu_L}(P) O_{\nu_1 \dots \nu_L}^{\alpha_1 \dots \alpha_L}(P) &= O_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L}(P).
\end{aligned} \tag{12}$$

This tensor has the same symmetry, orthogonality and traceless properties as X -tensors (for the same set of up and down indices) but the O -operator does not depend on the relative momentum of the constituents and does not describe decay processes. It represents the structure of the propagator of the composite system. Taking into account the definition of the projection operators (12) and the properties of the X -tensors (8) we obtain:

$$k_{\mu_1} \dots k_{\mu_L} O_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L} = \frac{1}{\alpha(L)} X_{\nu_1 \dots \nu_L}^{(L)}(k^{\perp}). \tag{13}$$

This equation presents the basic property of the projection operator: it projects any tensor with L indices onto the tensor which satisfied all properties of the partial wave considered.

The projection operator can also be calculated from tensors with lower rank using the recurrent expression:

$$\begin{aligned}
O_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L} &= \frac{1}{L^2} \left(\sum_{i,j=1}^L g_{\mu_i \nu_j}^{\perp} O_{\nu_1 \dots \nu_{j-1} \nu_{j+1} \dots \nu_L}^{\mu_1 \dots \mu_{i-1} \mu_{i+1} \dots \mu_L} - \frac{4}{(2L-1)(2L-3)} \right. \\
&\quad \left. \sum_{\substack{i < j \\ k < m}}^L g_{\mu_i \mu_j}^{\perp} g_{\nu_k \nu_m}^{\perp} O_{\nu_1 \dots \nu_{k-1} \nu_{k+1} \dots \nu_{m-1} \nu_{m+1} \dots \nu_L}^{\mu_1 \dots \mu_{i-1} \mu_{i+1} \dots \mu_{j-1} \mu_{j+1} \dots \mu_L} \right)
\end{aligned} \tag{14}$$

The low order projection operators are:

$$\begin{aligned}
O &= 1 & O_{\nu}^{\mu} &= g_{\mu\nu}^{\perp} \\
O_{\alpha\beta}^{\mu\nu} &= \frac{1}{2} \left(g_{\mu\alpha}^{\perp} g_{\nu\beta}^{\perp} + g_{\mu\beta}^{\perp} g_{\nu\alpha}^{\perp} - \frac{2}{3} g_{\mu\nu}^{\perp} g_{\alpha\beta}^{\perp} \right)
\end{aligned} \tag{15}$$

The scattering of the two spinless particles in the partial wave with the total spin $J=L$ (for example a $\pi\pi \rightarrow \pi\pi$ transition) is described as a convolution of the operators $X^{(L)}(k)$ and $X^{(L)}(q)$ where k and q are relative momenta before and after the interaction.

$$X_{\mu_1 \dots \mu_L}^{(L)}(k^{\perp}) O_{\nu_1 \dots \nu_L}^{\mu_1 \dots \mu_L} X_{\mu_1 \dots \mu_L}^{(L)}(q^{\perp}) = \alpha(L) \left(\sqrt{k_{\perp}^2} \sqrt{q_{\perp}^2} \right)^L P_L(z). \tag{16}$$

Here $P_L(z)$ are Legendre polynomials and $z = (k^\perp q^\perp) / (\sqrt{k_\perp^2} \sqrt{q_\perp^2})$ which are, in c.m.s., functions of the cosine of the angle between initial and final particles.

C. The decay of bound system into two pseudoscalar meson states

The system of the two pseudoscalar particles can form a partial wave with spin and parity J^P where $J = L$ and $P = (-1)^L$. The decay vertex of such system is described by the orbital momentum tensor only:

$$V_{\mu_1 \dots \mu_L}^J = X_{\mu_1 \dots \mu_L}^{(L)}(k^\perp), \quad (17)$$

In the two-meson decay the G -parity corresponds to the product of the G -parities of the final particles and the isospin can have values from $|I_1 - I_2|$ to $I_1 + I_2$. If the final particles are neutral one the C -parity of the two meson system is equal to the product of the C -parities of the final particles. In the case where the final particles are the particle and its own antiparticle $C = (-1)^L$. The identical particles cannot form the bound states with the odd partial waves: such amplitude will be anti-symmetrical and disappear when particles are permuted. Then for the scattering of two pseudoscalar particles, the combined system can have the quantum numbers listed in Table 1. For the two-pion system, the isospin and G parity can be $I^G = 0^+, 1^+, 2^+$, for the $\pi\eta$ system $I^G = 1^-$ and for the $\eta\eta(\eta\eta')$ system $I^G = 0^+$. In the quark model, where mesons are considered to be bound states of the quark and antiquark, the isospin can be $I = 0, 1$ G -parity is connected with C -parity as $G = C(-1)^I$ and states with $CP = -1$ and with $P = (-1)^J$ are forbidden.

D. The decay of the resonance into three spinless particles.

Let us consider the composite system decaying into the final three pseudo-scalar mesons via an intermediate two-body system with spin $I_{12}^{G_{12}} J_{12}^{P_{12} C_{12}}$. The tensor which describes the intrinsic spin is constructed by the convolution of the of the orbital momentum tensor which describes the two-particle intermediate state and the projection operator of three particle system:

$$S_{\mu_1 \dots \mu_n}^{(3)}(J_{12}) = X_{\nu_1 \dots \nu_n}^{(L_{12})}(k_{12}^\perp) O_{\mu_1 \dots \mu_n}^{\nu_1 \dots \nu_n}(P_3). \quad (18)$$

Here k_1 and k_2 are the momenta of the particles from decay of the intermediate system and

$$k_{12\mu}^\perp = \frac{(k_1 + k_2)_\nu}{2} g_{\mu\nu}^{\perp P_{12}} \quad g_{\mu\nu}^{\perp P_{12}} = g_{\mu\nu} - \frac{P_{12\mu} P_{12\nu}}{P_{12}^2} \quad (19)$$

where $P_{12} = k_1 + k_2$, $P_3 = k_1 + k_2 + k_3$ and $n = J_{12} = L_{12}$. If

Table 1. The partial waves in the channel of two pseudo-scalar particles.

L	0	1	2	3	4	
$\pi^0\pi^0$	0^{++}		2^{++}		4^{++}	...
$\pi^+\pi^-$	0^{++}	1^-	2^{++}	3^-	4^{++}	...
$\pi^\pm\pi^0$	0^+	1^- ,	2^+	3^-	4^+	...
$\eta\eta$	0^{++}		2^{++}		4^{++}	...
$\pi^0\eta$	0^{++}	1^{++}	2^{++}	3^{+-}	4^{++}	...
$\pi^\pm\eta$	0^+	1^-	2^+	3^-	4^+	...
$\eta\eta'$	0^{++}	1^{++}	2^{++}	3^{+-}	4^{++}	...

the total spin of the three-particle system is equal to J_3 the orbital momentum between the intermediate state and the spectator particle can be $L_3 = |J_3 - J_{12}|, \dots, J_3 + J_{12}$. If the combination $(L_3 + J_{12} - J_3)$ is an even number then the decay vertex can be constructed as:

$$V_{\mu_1 \dots \mu_{J_3}}^{(+1)}(Q_3) = S_{\mu_1 \dots \mu_i \nu_1 \dots \nu_m}^{(3)}(J_{12}) X_{\nu_1 \dots \nu_m \mu_{i+1} \dots \mu_{J_3}}^{(L_3)}(k_3^\perp) \quad (20)$$

where k_3^\perp is the momentum of the spectator particle orthogonal to the total momentum of the three particle system. The number of convoluted indices $m = (J_{12} + L_3 - J_3)/2$ and $i = J_{12} - m$. The multiindex Q_3 matches the principal quantum numbers $Q_3 \equiv J_3, L_3, J_{12}$. If the combination $(J_{12} + L_3 - J_3)$ is the odd number then the amplitudes is formed by means of the antisymmetric tensor:

$$V_{\mu_1 \dots \mu_{J_3}}^{(-1)}(Q_3) = \varepsilon_{\mu_1 \alpha \beta P_3} \times S_{\alpha \mu_2 \dots \mu_i \nu_1 \dots \nu_m}^{(3)}(J_{12}) X_{\beta \nu_1 \dots \nu_m \mu_{i+1} \dots \mu_{J_3}}^{(L_3)}(k_3^\perp), \quad (21)$$

$$m = (J_{12} + L_3 - J_3 - 1)/2, \quad i = J_{12} - m \text{ and}$$

$$\varepsilon_{\mu\alpha\beta P_3} \equiv \varepsilon_{\mu\alpha\beta\nu} P_{3\nu}. \quad (22)$$

The final tensor should be symmetrical, traceless and orthogonal to the total momentum of the three particle system which can be done by convolution with the projection operator:

$$A_{\mu_1 \dots \mu_n}^{(3,\alpha)}(Q_3) = V_{\nu_1 \dots \nu_n}^{(\alpha)}(Q_3) O_{\mu_1 \dots \mu_n}^{\nu_1 \dots \nu_n}(P_3). \quad (23)$$

Therefore there are two classes of the vertices with $\alpha = +1, -1$ which we will refer below as natural (tensor) and unnatural (pseudo-tensor) structures. The parity of these states are defined as:

$$P = (-1)^{L_3 + J_{12}} \prod_{i=1}^3 P_i, \quad (24)$$

where P_i denote the parities of the final states. Table 2 enumerates the potential states for the final state $3\pi^0$, considering the intermediate states of the scalar and tensor.

Table 2. List of the partial wave states produced in $3\pi^0$ channel. The partial waves are denoted as $J_{m\alpha}^{PC}$ (see eqs.(20,21,23)). The standard $q\bar{q}$ states in $3\pi^0$ channel have isospin $I = 1$.

	$J_{12} = 0$	$J_{12} = 2$
$L_3 = 0$	0_{0+}^-	2_{0+}^-
$L_3 = 1$	1_{0+}^+	$1_{1+}^{++} 2_{0+}^{++} 3_{0+}^{++}$
$L_3 = 2$	2_{0+}^-	$0_{2+}^- 1_{1+}^- 2_{1+}^- 3_{0+}^- 4_{0+}^-$
$L_3 = 3$	3_{0+}^+	$1_{2+}^{++} 2_{1+}^{++} 3_{1+}^{++} 4_{0+}^{++} 5_{0+}^{++}$

E. The decay of the resonance into four spinless particles through three body intermediate state.

This case is very similar to the one described above. If the 3-body bound state has quantum numbers $I_3^G J_3^{P_3 C_3}$ then the intrinsic spin can be described as projection of the three body amplitude eq.23 into 4-particle system:

$$S_{\mu_1 \dots \mu_{J_3}}^{(4,\alpha)}(Q_3) = A_{\nu_1 \dots \nu_{J_3}}^{(3,\alpha)}(Q_3) O_{\mu_1 \dots \mu_{J_3}}^{\nu_1 \dots \nu_{J_3}}(P), \quad (25)$$

where $P = k_1 + k_2 + k_3 + k_4$. The orbital momentum between the 3-body state and a spectator particle can be $L_4 = |J_4 - J_3|, \dots, J_4 + J_3$. If the combination $(L_4 + J_3 - J_4)$ is an even number, then the decay vertex can be written as convolution of the intrinsic spin tensor with orbital momentum tensor :

$$V_{\mu_1 \dots \mu_{J_4}}^{(+1,\alpha)}(Q_4) = S_{\mu_1 \dots \mu_{J_3} \nu_1 \dots \nu_m}^{(4,\alpha)}(Q_3) X_{\nu_1 \dots \nu_m \mu_{i+1} \dots \mu_{J_4}}^{(L_4)}(k_4^\perp) \quad (26)$$

where k_4^\perp is the momentum of the spectator particle orthogonal to the momentum of the four particle system. The number of convoluted indices is equal to $m = (J_3 + L_4 - J_4)/2$ and $i = J_3 - m$. The multiindex Q_4 lists all relevant quantum numbers $Q_4 = J_4, L_4, J_3, L_3, J_{12}$. If the combination $(J_3 + L_4 - J_4)$ is the odd number:

$$V_{\mu_1 \dots \mu_{J_4}}^{(-1,\alpha)}(Q_4) = \varepsilon_{\mu_1 \eta \beta P} \times S_{\eta \mu_2 \dots \mu_i \nu_1 \dots \nu_m}^{(4,\alpha)}(Q_3) X_{\beta \nu_1 \dots \nu_m \mu_{i+1} \dots \mu_{J_4}}^{(L_4)}(k_4^\perp), \quad (27)$$

$m = (J_3 + L_4 - J_4 - 1)/2$ and $i = J_3 - m$. The final tensor should be symmetrical, traceless and orthogonal to the total momentum of the four particle system which can be done by means the projection operator:

$$A_{\mu_1 \dots \mu_{J_4}}^{(4,\beta,\alpha)}(Q_4) = V_{\nu_1 \dots \nu_{J_4}}^{(\beta,\alpha)}(Q_4) O_{\mu_1 \dots \mu_{J_4}}^{\nu_1 \dots \nu_{J_4}}(P), \quad (28)$$

where $\beta, \alpha = -1, +1$. Thus we have 4 classes of amp-

litudes:

$$+1, +1 : J_3 + L_4 - J_4 = 2n \quad J_{12} + L_3 - J_3 = 2m \quad (29)$$

$$+1, -1 : J_3 + L_4 - J_4 = 2n \quad J_{12} + L_3 - J_3 = 2m + 1 \quad (30)$$

$$-1, +1 : J_3 + L_4 - J_4 = 2n + 1 \quad J_{12} + L_3 - J_3 = 2m \quad (31)$$

$$-1, -1 : J_3 + L_4 - J_4 = 2n + 1 \quad J_{12} + L_3 - J_3 = 2m + 1 \quad (32)$$

The parity of the 4-particle state are defined as:

$$P = (-1)^{L_4 + L_3 + J_{12}} \prod_{i=1}^4 P_i. \quad (33)$$

The partial wave amplitudes in the $4\pi^0$ channel in the case of cascade decay are listed in Table 3 for the intermediate three body states up to $J_3 = 2$. The exotic states which can be produced in the $4\pi^0$ and $3\pi^0$ channels are also included in this Table.

Table 3. List of the partial wave amplitudes $J_{n\beta}^{PC}$ state for cascade decay into $4\pi^0$.

J_3^{PC}	0^-	1^{++}	1^-
$L_4 = 0$	0_{0+}^{++}	1_{0+}^-	1_{0+}^{++}
$L_4 = 1$	1_{0+}^-	$0_{1+}^{++} 1_{0+}^{++} 2_{0+}^{++}$	$0_{0+}^{++} 1_{0+}^{++} 2_{0+}^{++}$
$L_4 = 2$	2_{0+}^{++}	$1_{1+}^- 2_{0+}^- 3_{0+}^-$	$1_{1+}^- 2_{0+}^- 3_{0+}^-$
J_3^{PC}	2^{++}	2^-	
$L_4 = 0$	2_{0+}^-	2_{0+}^{++}	
$L_4 = 1$	$1_{0+}^{++} 2_{0+}^{++} 3_{0+}^{++}$	$1_{1+}^- 2_{0+}^- 3_{0+}^-$	
$L_4 = 2$	$0_{2+}^- 1_{1+}^- 2_{1+}^- 3_{0+}^- 4_{0+}^-$	$0_{2+}^{++} 1_{1+}^{++} 2_{1+}^{++} 3_{0+}^{++} 4_{0+}^{++}$	

F. The decay of the resonance into two resonances decaying into two spinless particles

Consider the decay of the resonance into two resonances with spin J_{12} and J_{34} that decay into two spinless particles with momenta k_1, k_2 and k_3, k_4 , respectively. The decay of these two states is described with $X_{\mu_1 \dots \mu_{J_{12}}}^{(J_{12})}(k_{12}^\perp)$ and $X_{\mu_1 \dots \mu_{J_{34}}}^{(J_{34})}(k_{34}^\perp)$ tensors, where

$$k_{ij\mu}^\perp = \frac{k_{iv} - k_{jv}}{2} g_{\mu\nu}^{\perp P_{ij}} \quad g_{\mu\nu}^{\perp P_{ij}} = g_{\mu\nu} - \frac{P_{ij\mu} P_{ij\nu}}{P_{ij}^2}, \quad (34)$$

where $P_{ij} = k_i + k_j$. The intrinsic spin can have values

$S = |J_{12} - J_{34}|, \dots, J_{12} + J_{34}$. If $J_{12} + J_{34} - S = 2m$, the tensor that describes this spin state can be formed by convolution of the m indices.

$$V_{\mu_1 \dots \mu_S}^{(+1)}(Q_S) = X_{\mu_1 \dots \mu_{J_{12}-m} \nu_1 \dots \nu_m}^{(J_{12})}(k_{12}^\perp) X_{\nu_1 \dots \nu_m \mu_{J_{12}-m+1} \dots \mu_S}^{(J_{34})}(k_{34}^\perp) \quad (35)$$

where $Q_S = S, J_{12}, J_{34}$. If the difference between intrinsic spin and the sum of spin resonances is an odd number $J_{12} + J_{34} - S = 2m + 1$ the vertex is formed by means of the antisymmetric tensor.

$$V_{\mu_1 \dots \mu_S}^{(-1)}(Q_S) = \varepsilon_{\mu_1 \eta \beta P} X_{\eta \mu_2 \dots \mu_{J_{12}-m} \nu_1 \dots \nu_m}^{(J_{12})}(k_{12}^\perp) \times X_{\beta \nu_1 \dots \nu_m \mu_{J_{12}-m+1} \dots \mu_S}^{(J_{34})}(k_{34}^\perp), \quad (36)$$

where $P = k_1 + k_2 + k_3 + k_4$. The symmetrization of indices and traceless property can be satisfied in the standard way by convolution with the projection operator:

$$S_{\mu_1 \dots \mu_S}^{(22,\alpha)}(Q_S) = V_{\nu_1 \dots \nu_S}^{(\alpha)}(Q_S) O_{\mu_1 \dots \mu_S}^{\nu_1 \dots \nu_S}(P), \quad (37)$$

If the spin of the 4-particle partial wave is equal J_4 , the orbital momentum between the two resonances can be $L_4 = |J_4 - S|, \dots, J_4 + S$. If the combination $(L_4 + S - J_4) = 2n$ is the even number then the decay vertex can be written as:

$$V_{\mu_1 \dots \mu_{J_4}}^{(+1,\alpha)}(Q_{22}) = S_{\mu_1 \dots \mu_{S-n} \nu_1 \dots \nu_n}^{(22,\alpha)}(Q_S) X_{\nu_1 \dots \nu_n \mu_{S-n+1} \dots \mu_{J_4}}^{(L_4)}(k^\perp) \quad (38)$$

where k^\perp is the relative momentum between two intermediate resonances:

$$k_\mu^\perp = \frac{1}{2}(k_1 + k_2 - k_3 - k_4)_\nu g_{\mu\nu}^\perp \quad (39)$$

and multi-index $Q_{22} = J_4, L_4, S, J_{12}, J_{34}$. If the combination $(J_3 + L_4 - J_4)$ is the odd number:

$$V_{\mu_1 \dots \mu_{J_4}}^{(-1,\alpha)}(Q_{22}) = \varepsilon_{\mu_1 \eta \beta P} \times S_{\eta \mu_2 \dots \mu_{S-n} \nu_1 \dots \nu_n}^{(22,\alpha)}(Q_S) X_{\beta \nu_1 \dots \nu_n \mu_{S-n+1} \dots \mu_{J_4}}^{(L_4)}(k^\perp), \quad (40)$$

$n = (S + L_4 - J_4 - 1)/2$. The final tensor should be symmetrical, traceless and orthogonal to the total momentum of the four particle system which can be done with projection operator:

$$A_{\mu_1 \dots \mu_{J_4}}^{(22,\beta,\alpha)}(Q_{22}) = V_{\nu_1 \dots \nu_{J_4}}^{(\beta,\alpha)}(Q_{22}) O_{\mu_1 \dots \mu_{J_4}}^{\nu_1 \dots \nu_{J_4}}(P), \quad (41)$$

Thus we have 4 classes of amplitudes:

$$+1, +1 : S + L_4 - J_4 = 2n \quad J_{12} + J_{34} - S = 2m \quad (42)$$

$$+1, -1 : S + L_4 - J_4 = 2n \quad J_{12} + J_{34} - S = 2m + 1 \quad (43)$$

$$-1, +1 : S + L_4 - J_4 = 2n + 1 \quad J_{12} + J_{34} - S = 2m \quad (44)$$

$$-1, -1 : S + L_4 - J_4 = 2n + 1 \quad J_{12} + J_{34} - S = 2m + 1 \quad (45)$$

The parity of the 4-particle state are defined as:

$$P = (-1)^{L_4 + J_{12} + J_{34}} \prod_{i=1}^4 P_i. \quad (46)$$

G. Construction of tensors for the decay of the resonance into four pseudoscalar mesons

The formulas given in previous sections allowed one to construct directly the code for the decay of a resonance into any four pseudoscalar meson state. Let us consider as an example the decay of the tensor resonance into two tensor final states.

It is seen that two final tensor states can form intrinsic spin $S^P = 0^+, 1^+, 2^+, 3^+, 4^+$ (see Table 4). If the intrinsic spin is equal to 2^+ the tensor for this state is constructed with $S_{m\alpha}^P = 2_{1+}^+$ which means that one index is convoluted and no antisymmetric tensors are involved. Then:

$$S_{\mu\nu}^{(22,+)}(Q_S) = O_{\eta\chi}^{\mu\nu}(P) X_{\eta\xi}^{(2)}(k_{12}^\perp) X_{\xi\chi}^{(2)}(k_{34}^\perp). \quad (47)$$

If the initial state is the tensor one, it can be constructed with $L_4 = 0$ or $L_4 = 2$ (see Table 5). In the first case, the $n\beta = 0+$ and the tensor for the decay into four mesons has a simple form:

$$V_{\mu\nu}^{(2+,1)} = S_{\mu\nu}^{(22,+)}(Q_S) \quad (48)$$

In the second case $n\beta = 1+$ and one convolution with orbital momentum tensor:

$$V_{\mu\nu}^{(2+,2)} = X_{\mu\chi}^{(2)}(k^\perp) S_{\chi\nu}^{(22,+)}(Q_S) \quad (49)$$

Here, for convenience, we have changed the upper indices for the tensor $V_{\mu_1 \dots \mu_J}^{J^P, m}$ given spin and parity of the 4-particle resonance and just the tensor number.

Appendix presents the tensor list associated with the

Table 4. List of possible spin combinations $S_{m\alpha}^P$ states ($C=+1$) in four particle decay via two resonance decays.

J_{12}^{PC}	0^{++}	0^{++}	2^{++}	2^{++}
J_{34}^{PC}	0^{++}	2^{++}	2^{++}	4^{++}
$S_{m\alpha}^P$	0_{0+}^+	2_{0+}^+	$0_{2+}^+ 1_{1-}^+ 2_{1+}^+ 3_{0-}^+ 4_{0+}^+$	$2_{2+}^+ 3_{1-}^+ 4_{1+}^+ 5_{0-}^+ 6_{0+}^+$

Table 5. List of states $J_{n\beta}^{PC}$ in $4\pi^0$ channel decaying into two resonances forming intrinsic spin S^{PC} .

S^{PC}	0^{++}	1^{++}	2^{++}
$L_4 = 0$	0_{0+}^{++}	1_{0+}^{++}	2_{0+}^{++}
$L_4 = 1$	1_{0+}^{++}	$0_{1+}^{++} 1_{0-}^{++} 2_{0+}^{++}$	$1_{1+}^{++} 2_{0-}^{++} 3_{0+}^{++}$
$L_4 = 2$	2_{0+}^{++}	$1_{1+}^{++} 2_{0-}^{++} 3_{0+}^{++}$	$0_{2+}^{++} 1_{1-}^{++} 2_{1+}^{++} 3_{0-}^{++} 4_{0+}^{++}$
S^{PC}		3^{++}	4^{++}
$L_4 = 0$		3_{0+}^{++}	4_{0+}^{++}
$L_4 = 1$		$2_{1+}^{++} 3_{0-}^{++} 4_{0+}^{++}$	$3_{1+}^{++} 4_{0-}^{++} 5_{0+}^{++}$
$L_4 = 2$		$1_{2+}^{++} 2_{1-}^{++} 3_{1+}^{++} 4_{0-}^{++} 5_{0+}^{++}$	$2_{2+}^{++} 3_{1-}^{++} 4_{1+}^{++} 5_{0-}^{++} 6_{0+}^{++}$

decay of a resonance into a final state of $4\pi^0$. An extra symmetry arises from the permutations of pions. When a resonance decays into two other resonances, amplitudes in which the sum of intrinsic spin and orbital momentum is odd are nullified. The scenario is more intricate in cascade decays, and all possible configurations are enumerated.

H. The production of the states decaying into 4 pseudo-scalar mesons.

A partial wave amplitude is a scalar value. The convolution of the tensors with an antisymmetric tensor creates a pseudo-tensor structure (unnatural). Thus, the scalar value should be either convolution of any number of tensors (natural structures) or a convolution of the even number of unnatural structures. Therefore, if the produced state is a natural state, the decay amplitude can have only $(++)$ or $(--)$ structures. If the produced state is an unnatural state, the decay should be described by the $(+-)$ or $(-+)$ combination.

1. The production of the state from pomeron-pomeron, $f_0 f_0$ or $\pi^0 \pi^0$ collision

Consider the production of resonances in central collision reactions. In that case, the states can be produced, for example, from the pomeron-pomeron collision. The similar mechanism is responsible for the production of resonances in two-scalar or two neutral pseudo-scalar mesons. In such processes, only states with $G=+1$ and $C=+1$ are produced. In the standard quark model only resonances with isospin 0 and even spin can be produced in such reaction: $I^G J^{PC} = 0^+ J^{++}$ and $J = 2n$. The production vertex

is described as the orbital momentum tensor $X_{\mu_1 \dots \mu_{J_4}}^{(J_4)}(q^\perp)$ where q is the relative momenta of the collided particles. Then the partial wave amplitude for the partial wave with spin J_4 has the form:

$$A_{J_4}^{(t\alpha\beta)} = X_{\mu_1 \dots \mu_{J_4}}^{(J_4)}(q^\perp) A_{\mu_1 \dots \mu_{J_4}}^{(t\alpha\beta)}(Q_t), \quad (50)$$

where index t describes topologies $t=4$ (cascade) and $t=22$ (decay into two resonances). Consider the case with the final state $4\pi^0$ (or 4η). Then only resonances with $I^G J^{PC} = 0^+ J_{ij}^{++}$ where $J_{ij} = 2n$ can be produced in the two-particle channel. If we have two scalar resonances $J_{12} = J_{34} = 0$ in the final states only $++$ amplitudes are possible:

$$\begin{aligned} A_{J_4}^{(22^{++})} &= X_{\mu_1 \dots \mu_{J_4}}^{(J_4)}(q^\perp) O_{\nu_1 \dots \nu_{J_4}}^{\mu_1 \dots \mu_{J_4}}(P) X_{\nu_1 \dots \nu_{J_4}}^{(J_4)}(k^\perp) \\ &= X_{\mu_1 \dots \mu_{J_4}}^{(J_4)}(q^\perp) X_{\mu_1 \dots \mu_{J_4}}^{(J_4)}(k^\perp) \end{aligned} \quad (51)$$

we obtain the standard Legendre polynomial dependence in the rest system of the 4-particle state (see eq.16). In the cascade topology the amplitudes have $++$ or $--$ signatures. The examples of other amplitudes are given in the Appendix.

2. The production of the states decaying into 4 pseudo-scalar particles in the J/Ψ radiative decays.

The states produced in radiative decay J/Ψ can be natural or unnatural states. All states have isospin 0, $G=+1$ and $C=+1$. This process can be considered as the production of resonances in collisions $J/\Psi\gamma$. In this case the intrinsic spin can have values $S = 0, 1, 2$, and the corresponding spin tensors are:

$$\begin{aligned} S^0 &= g_{\mu\nu} \epsilon_\mu^\Psi \epsilon_\nu^{\gamma*} & S=0 \\ S^1_\eta &= \epsilon_{\eta\mu\nu\rho} \epsilon_\mu^\Psi \epsilon_\nu^{\gamma*} & S=1 \\ S^2_{\eta\xi} &= \epsilon_\mu^\Psi \epsilon_\nu^{\gamma*} O_{\eta\xi}^{\mu\nu}(P) & S=2 \end{aligned} \quad (52)$$

In the spin-orbital basis (SL) the partial waves can be described as $^{2S+1}L_J$. In this basis, the partial waves with $S=0$ have $L=J$ and parity $P=(-1)^J$. These production vertices are described as:

$$V_{\mu_1 \dots \mu_J}^0 = S^0 X_{\mu_1 \dots \mu_J}^{(J)}(q^\perp) \quad {}^1J_J \quad (53)$$

and all amplitudes are the natural ones. The natural amplitudes can be constructed with intrinsic spin 1 only with $L=J$ (3L_J):

$$V_{\mu_1 \dots \mu_J}^{1,0} = \varepsilon_{\mu_1 \eta \xi P} S_\eta^1 X_{\xi \mu_2 \dots \mu_J}^{(J)}(q^\perp) \quad {}^3 J_J. \quad (54)$$

This vertex can be symmetrized and traceless by convolution with the projection operator. However, we already applied this operator to the decay vertices, and it is not necessary to apply it to the production vertices. For the amplitudes with $S=2$ the natural amplitudes are produced with $L=J-2, J, J+2$:

$$\begin{aligned} V_{\mu_1 \dots \mu_J}^{5,-2} &= S_{\mu_1 \mu_2}^2 X_{\mu_3 \dots \mu_J}^{(J-2)}(q^\perp) \quad {}^5 (J-2)_J \\ V_{\mu_1 \dots \mu_J}^{5,0} &= S_{\mu_1 \eta}^2 X_{\eta \mu_2 \dots \mu_J}^{(J)}(q^\perp) \quad {}^5 J_J \\ V_{\mu_1 \dots \mu_J}^{5,+2} &= S_{\xi \eta}^2 X_{\xi \eta \mu_1 \dots \mu_J}^{(J+2)}(q^\perp) \quad {}^5 (J+2)_J \end{aligned} \quad (55)$$

Recall that the decay of the natural states can be described with $-$ or $++$ amplitudes in the decay into 4 pseudoscalar mesons. In the gauge-invariant limit for the partial waves with $J \geq 2$ only three amplitudes are linearly independent. For $J=0$ only one amplitude is linearly independent and for $J=1$ we have two linearly independent amplitudes (see the appendix).

The unnatural amplitudes are formed with intrinsic spin $S=1,2$ and orbital momentum $L=J-1$ and $L=J+1$:

$$\begin{aligned} V_{\mu_1 \dots \mu_J}^{3,-1} &= S_{\mu_1}^1 X_{\mu_2 \dots \mu_J}^{(J-1)}(q^\perp) \quad {}^3 (J-1)_J \\ V_{\mu_1 \dots \mu_J}^{3,+1} &= S_\eta^1 X_{\eta \mu_1 \dots \mu_J}^{(J+1)}(q^\perp) \quad {}^3 (J+1)_J \\ V_{\mu_1 \dots \mu_J}^{5,-1} &= \varepsilon_{\mu_1 \eta \xi P} S_{\eta \mu_2}^2 X_{\xi \mu_3 \dots \mu_J}^{(J-1)}(q^\perp) \quad {}^5 (J-1)_J \\ V_{\mu_1 \dots \mu_J}^{5,+1} &= \varepsilon_{\mu_1 \eta \xi P} S_{\eta \chi}^2 X_{\xi \chi \mu_2 \dots \mu_J}^{(J+1)}(q^\perp) \quad {}^5 (J+1)_J \end{aligned} \quad (56)$$

In the case of the decay a vector meson into virtual photon (or a massive vector particle like ω -meson) and four pseudoscalar mesons, all these vertices are linear independent. However, in the case of the real photon and $J \geq 2$ only three amplitudes are linearly independent. For $J=0$ only one amplitude is linearly independent and for $J=1$ we have two linearly independent amplitudes. The amplitudes' basis set might vary; when dealing with an energy-independent fit, the specific choice of amplitudes does not matter for the analysis. Conversely, in the context of an energy-dependent fit, it is essential to manage the amplitudes asymptotic behavior. From this point of view, we suggested using the amplitudes with the lowest orbital momentum and different intrinsic spins. The details of the amplitude linear dependence for the case of the real photon are given in the Appendix.

II. CONCLUSION

We have developed the formalism for the partial-wave analysis of the data with four pseudoscalar meson states. The method is covariant and can be directly ap-

plied to the event-by-event analysis of the data, for example, in the framework of the maximum likelihood method. In particular, we consider the $4\pi^0$ production in the radiative decay J/Ψ and the production of the four pseudo-scalar mesons in the central collision.

III. APPENDIX

A. Linearly dependent vertices for the J/Ψ radiative decay

Consider the general structure of J/Ψ radiative decay. The production of the state with spin J can be described as a convolution projection operator with vertex:

$$O_{\mu_1 \dots \mu_J}^{\gamma_1 \dots \gamma_J}(P) V_{\mu_1 \dots \mu_J}(q^\perp) \quad (57)$$

where q^\perp is momentum of J/Ψ orthogonal to the momentum of the resonance. Taking into account the orthogonality and traceless properties of the projection operator the general structure of the vertex for the natural states is:

$$\begin{aligned} V_{\mu_1 \dots \mu_J} &= (\epsilon^\Psi \epsilon^{\gamma^*}) q_{\mu_1}^\perp \dots q_{\mu_J}^\perp F_1 + \epsilon_{\mu_1}^\Psi \epsilon_{\mu_2}^{\gamma^*} q_{\mu_3}^\perp \dots q_{\mu_J}^\perp F_2 \\ &+ \epsilon_{\mu_1}^\Psi (\epsilon^{\gamma^*} q^\perp) q_{\mu_2}^\perp \dots q_{\mu_J}^\perp F_3 + (\epsilon^\Psi q^\perp) \epsilon_{\mu_1}^{\gamma^*} q_{\mu_2}^\perp \dots q_{\mu_J}^\perp F_4 \\ &+ (\epsilon^\Psi q^\perp) (\epsilon^{\gamma^*} q^\perp) q_{\mu_1}^\perp \dots q_{\mu_J}^\perp F_5 \end{aligned} \quad (58)$$

where F_i are scalar functions. Recall that the polarization vector of the particle is orthogonal to its momentum, and we can use q^\perp as universal momentum in convolution with both polarization vectors. In the case of the real photon, it's polarization vector is orthogonal to all momenta in the vertex and therefore $(\epsilon^{\gamma^*} q^\perp) = 0$. As a consequence, only structures with F_1, F_2 and F_4 will contribute to the decay. In the case of the scalar state ($J=0$) the vertex has only structures F_1, F_5 and only F_1 is not zero in this limit. For the vector state ($J=1$) only structures F_1, F_3, F_4, F_5 can contribute to the vertex and only F_1 and F_4 survive.

For the production of unnatural states, the vertex has the structure:

$$\begin{aligned} V_{\mu_1 \dots \mu_J} &= \varepsilon_{q^\perp \alpha \beta P} \epsilon_\alpha^\Psi \epsilon_\beta^{\gamma^*} q_{\mu_1}^\perp \dots q_{\mu_J}^\perp F_1 \\ &+ \varepsilon_{\mu_1 \alpha \beta P} \epsilon_\alpha^\Psi \epsilon_\beta^{\gamma^*} q_{\mu_2}^\perp \dots q_{\mu_J}^\perp F_2 \\ &+ \varepsilon_{\mu_1 \alpha q^\perp P} (\epsilon_\alpha^\Psi \epsilon_{\mu_2}^{\gamma^*} + \epsilon_{\mu_2}^\Psi \epsilon_\alpha^{\gamma^*}) q_{\mu_3}^\perp \dots q_{\mu_J}^\perp F_3 \\ &+ \varepsilon_{\mu_1 \alpha q^\perp P} (\epsilon_\alpha^\Psi (\epsilon^{\gamma^*} q^\perp) + (\epsilon^\Psi q^\perp) \epsilon_\alpha^{\gamma^*}) q_{\mu_2}^\perp \dots q_{\mu_J}^\perp F_4 \end{aligned} \quad (59)$$

For the $J^{PC} = 0^{-+}$ state, there is only one structure

(with F_1). Decompose the polarization vector of the real photon ϵ_α^Ψ into components parallel to q^\perp , parallel to P , and orthogonal to both q^\perp and P . Then:

$$\begin{aligned}\epsilon_\alpha^\Psi &= Cq_\alpha^\perp + \epsilon_\alpha^{\perp\perp} + \delta P_\alpha \\ \epsilon_{\mu_1\alpha\beta P}^\Psi &= C\epsilon_{\mu_1 q^\perp \beta P} + \epsilon_{\mu_1\alpha\beta P} \epsilon_\alpha^{\perp\perp} \\ \epsilon_{\mu_1\alpha\beta P}^\Psi \epsilon_\beta^{\gamma*} &= C\epsilon_{\mu_1 q^\perp \beta P} \epsilon_\beta^{\gamma*} + \epsilon_{\mu_1\alpha\beta P} \epsilon_\alpha^{\perp\perp} \epsilon_\beta^{\gamma*}\end{aligned}\quad (60)$$

As both $\epsilon_\alpha^{\perp\perp}, \epsilon_\beta^{\gamma*}$ are orthogonal to momentum q^\perp and P , then

$$\epsilon_{\mu_1\alpha\beta P} \epsilon_\alpha^{\perp\perp} \epsilon_\beta^{\gamma*} = Bq_{\mu_1}^\perp \quad (61)$$

Then we obtain the following structures for the vertex:

$$\begin{aligned}V_{\mu_1\dots\mu_J} &= B(q_{\mu_1}^\perp F_1 + F_2)q_{\mu_1}^\perp \dots q_{\mu_J}^\perp \\ &\quad - (CF_2 + (\epsilon^\Psi q^\perp)F_4)\epsilon_{\mu_1\beta q^\perp P} \epsilon_\beta^{\gamma*} q_{\mu_2}^\perp \dots q_{\mu_J}^\perp \\ &\quad + F_3\epsilon_{\mu_1\alpha q^\perp P} (\epsilon_\alpha^\Psi \epsilon_{\mu_2}^{\gamma*} + \epsilon_{\mu_2}^\Psi \epsilon_\alpha^{\gamma*})q_{\mu_3}^\perp \dots q_{\mu_J}^\perp\end{aligned}\quad (62)$$

The amplitudes F_1, F_2, F_4 have only two independent structures. This means that only two of them are linearly independent. This corresponds to the case $J^P = 1^+$ where the structure with F_3 cannot be produced. Thus, for $J \geq 2$, only three partial waves are linearly independent for every unnatural state. The list of the lowest partial waves is given in [Table 6](#).

B. The decay of the resonances into $4\pi^0$ final state.

Let us list the basic momenta and tensors:

$$\begin{aligned}P_{12\mu} &= (k_1 + k_2)_\mu, \quad g_{\mu\nu}^{\perp P_{12}} = g_{\mu\nu} - \frac{P_{12\mu}P_{12\nu}}{P_{12}^2}, \\ k_{12\mu}^\perp &= \frac{1}{2}(k_1 - k_2)_\nu g_{\nu\mu}^{\perp P_{12}}, \quad P_{34\mu} = (k_3 + k_4)_\mu, \\ g_{\mu\nu}^{\perp P_{34}} &= g_{\mu\nu} - \frac{P_{34\mu}P_{34\nu}}{P_{34}^2}, \quad k_{34\mu}^\perp = \frac{1}{2}(k_3 - k_4)_\nu g_{\nu\mu}^{\perp P_{34}} \\ P_{3\mu} &= (P_{12} + k_3)_\mu, \quad g_{\mu\nu}^{\perp P_3} = g_{\mu\nu} - \frac{P_{3\mu}P_{3\nu}}{P_3^2}, \\ k_{3\mu}^\perp &= \frac{1}{2}(k_3 - P_{12})_\nu g_{\nu\mu}^{\perp P_3} \\ P_\mu &= (P_3 + k_4)_\mu, \quad g_{\mu\nu}^\perp = g_{\mu\nu} - \frac{P_\mu P_\nu}{P^2}, \\ k_{4\mu}^\perp &= \frac{1}{2}(k_4 - P_3)_\nu g_{\nu\mu}^\perp \\ k_\mu^\perp &= \frac{1}{2}(k_1 + k_2 - k_3 - k_4)_\nu g_{\nu\mu}^\perp \\ O_{\mu_1\dots\mu_J}^{\nu_1\dots\nu_J} &\equiv O_{\mu_1\dots\mu_J}^{\nu_1\dots\nu_J}(P)\end{aligned}\quad (63)$$

Table 6. List of partial waves produced in radiative J/Ψ decay. The partial waves which became linear dependent from other partial waves are shown in third column. The states which are forbidden in the quark model are not listed.

0^{++}	1S_0	5D_0
2^{++}	$^1D_2 \ ^3D_2 \ ^5S_2$	$^5D_2 \ ^5G_2$
4^{++}	$^1G_4 \ ^3G_4 \ ^5D_4$	$^5G_4 \ ^5I_4$
0^{-+}	3P_0	
1^{+-}	$^3S_1 \ ^3D_1$	5D_1
2^{-+}	$^3P_2 \ ^5P_2 \ ^3F_2$	5F_2
3^{+-}	$^3D_3 \ ^3G_3 \ ^5D_3$	5G_3

Consider the decay of states into the $4\pi^0$ channel. Let us start from decay via two intermediate resonances for natural states:

$$\begin{aligned}J^{PC} &\rightarrow J_{n,\beta}^{PC}(L_4(S_{m\alpha}^{PC} \rightarrow f_{J_{12}}f_{J_{34}})) \\ 0^{++} &\rightarrow 0_{0,+}^{++}(0(0_{0,+}^{++} \rightarrow f_0f_0)) \\ V^{(0+,1)} &= 1\end{aligned}\quad (64)$$

$$\begin{aligned}0^{++} &\rightarrow 0_{2,+}^{++}(2(2_{0,+}^{++} \rightarrow f_0f_2)) \\ V^{(0+,2)} &= X_{\alpha\beta}^{(2)}(k^\perp)X_{\alpha\beta}^{(2)}(k_{34}^\perp)\end{aligned}\quad (65)$$

$$\begin{aligned}0^{++} &\rightarrow 0_{0,+}^{++}(0(0_{2,+}^{++} \rightarrow f_2f_2)) \\ V^{(0+,3)} &= X_{\alpha\beta}^{(2)}(k_{12}^\perp)X_{\alpha\beta}^{(2)}(k_{34}^\perp)\end{aligned}\quad (66)$$

$$\begin{aligned}0^{++} &\rightarrow 0_{0,+}^{++}(2(2_{1,+}^{++} \rightarrow f_2f_2)) \\ V^{(0+,4)} &= X_{\alpha\beta}^{(2)}(k^\perp)X_{\alpha\xi}^{(2)}(k_{12}^\perp)X_{\beta\xi}^{(2)}(k_{34}^\perp)\end{aligned}\quad (67)$$

$$\begin{aligned}2^{++} &\rightarrow 2_{0,+}^{++}(2(0_{0,+}^{++} \rightarrow f_0f_0)) \\ V_{\alpha\beta}^{(2+,1)} &= X_{\alpha\beta}^{(2)}(k^\perp)\end{aligned}\quad (68)$$

$$\begin{aligned}2^{++} &\rightarrow 2_{0,+}^{++}(0(2_{0,+}^{++} \rightarrow f_0f_2)) \\ V_{\alpha\beta}^{(2+,2)} &= X_{\alpha\beta}^{(2)}(k_{34}^\perp)\end{aligned}\quad (69)$$

$$\begin{aligned}2^{++} &\rightarrow 2_{1,+}^{++}(2(2_{0,+}^{++} \rightarrow f_0f_2)) \\ V_{\alpha\beta}^{(2+,3)} &= X_{\alpha\chi}^{(2)}(k^\perp)O_{\beta\chi}^{\mu\nu}X_{\mu\nu}^{(2)}(k_{34}^\perp)\end{aligned}\quad (70)$$

$$\begin{aligned}2^{++} &\rightarrow 2_{0,+}^{++}(2(0_{2,+}^{++} \rightarrow f_2f_2)) \\ V_{\alpha\beta}^{(2+,4)} &= X_{\alpha\beta}^{(2)}(k^\perp)X_{\eta\xi}^{(2)}(k_{12}^\perp)X_{\eta\xi}^{(2)}(k_{34}^\perp)\end{aligned}\quad (71)$$

$$\begin{aligned}2^{++} &\rightarrow 2_{0,+}^{++}(0(2_{1,+}^{++} \rightarrow f_2f_2)) \\ V_{\alpha\beta}^{(2+,5)} &= X_{\alpha\xi}^{(2)}(k_{12}^\perp)X_{\beta\xi}^{(2)}(k_{34}^\perp)\end{aligned}\quad (72)$$

$$2^{++} \rightarrow 2_{1,+}^{++}(2(2_{1,+}^{++} \rightarrow f_2 f_2))$$

$$V_{\alpha\beta}^{(2+,6)} = X_{\alpha\chi}^{(2)}(k^\perp) O_{\chi\beta}^{\mu\nu} X_{\mu\zeta}^{(2)}(k_{12}^\perp) X_{\nu\zeta}^{(2)}(k_{34}^\perp) \quad (73)$$

$$4^{++} \rightarrow 4_{0,+}^{++}(2(2_{0,+}^{++} \rightarrow f_0 f_2))$$

$$V_{\alpha\beta\mu\nu}^{(4+,1)} = X_{\alpha\beta}^{(2)}(k^\perp) O_{\mu\nu}^{\eta\zeta} X_{\eta\zeta}^{(2)}(k_{34}^\perp) \quad (74)$$

$$4^{++} \rightarrow 4_{0,+}^{++}(2(2_{1,+}^{++} \rightarrow f_2 f_2))$$

$$V_{\alpha\beta\mu\nu}^{(4+,2)} = X_{\alpha\beta}^{(2)}(k^\perp) O_{\mu\nu}^{\eta\zeta} X_{\eta\chi}^{(2)}(k_{12}^\perp) X_{\chi\zeta}^{(2)}(k_{34}^\perp) \quad (75)$$

$$4^{++} \rightarrow 4_{0,+}^{++}(0(4_{0,+}^{++} \rightarrow f_2 f_2))$$

$$V_{\alpha\beta\mu\nu}^{(4+,3)} = X_{\alpha\beta}^{(2)}(k_{12}^\perp) X_{\mu\nu}^{(2)}(k_{34}^\perp) \quad (76)$$

$$4^{++} \rightarrow 4_{1,+}^{++}(2(4_{0,+}^{++} \rightarrow f_2 f_2))$$

$$V_{\alpha\beta\mu\nu}^{(4+,4)} = X_{\alpha\eta}^{(2)}(k^\perp) O_{\eta\beta\mu\nu}^{\zeta\delta\gamma} X_{\zeta\chi}^{(2)}(k_{12}^\perp) X_{\delta\gamma}^{(2)}(k_{34}^\perp) \quad (77)$$

Here we provide the amplitudes with $L_4 \leq 2$ and resonances decaying into two pions up to $J = 2$. The unnatural states can only decay into two resonances with $J \geq 2$. In the case of two tensor states, we get the following:

$$0^{-+} \rightarrow 0_{1,+}^{-+}(1(1_{1,-}^{++} \rightarrow f_2 f_2))$$

$$V^{(0-,1)} = X_\chi^{(1)}(k^\perp) \varepsilon_{\chi\mu\nu\rho} X_{\mu\zeta}^{(2)}(k_{12}^\perp) X_{\nu\zeta}^{(2)}(k_{34}^\perp) \quad (78)$$

$$2^{-+} \rightarrow 2_{0,+}^{-+}(1(1_{1,-}^{++} \rightarrow f_2 f_2))$$

$$V_{\alpha\beta}^{(2-,1)} = X_\alpha^{(1)}(k^\perp) \varepsilon_{\beta\mu\nu\rho} X_{\mu\zeta}^{(2)}(k_{12}^\perp) X_{\nu\zeta}^{(2)}(k_{34}^\perp) \quad (79)$$

For the cascade decays we provide examples of amplitudes or cascade decays for the orbital momentum $L_4, L_3 \leq 2$:

$$J^{PC} \rightarrow J_{n,\beta}^{PC}(L_4(\pi J_{3,m\alpha}^{PC} \rightarrow L_3(\pi f_{J_{12}})))$$

$$0^{++} \rightarrow 0_{0,+}^{++}(0(\pi 0_{0,+}^{++} \rightarrow 0(\pi f_0)))$$

$$V^{(0+,5)} = 1 \quad (80)$$

$$0^{++} \rightarrow 0_{1,+}^{++}(1(\pi 1_{0,+}^{++} \rightarrow 1(\pi f_0)))$$

$$V^{(0+,6)} = X_\nu^{(1)}(k_4^\perp) X_\nu^{(1)}(k_3^\perp) \quad (81)$$

$$0^{++} \rightarrow 0_{2,+}^{++}(2(\pi 2_{0,+}^{++} \rightarrow 2(\pi f_0)))$$

$$V^{(0+,7)} = X_{\nu\mu}^{(2)}(k_4^\perp) X_{\nu\mu}^{(2)}(k_3^\perp) \quad (82)$$

$$0^{++} \rightarrow 0_{2,+}^{++}(2(\pi 2_{0,+}^{++} \rightarrow 0(\pi f_2)))$$

$$V^{(0+,8)} = X_{\alpha\beta}^{(2)}(k_4^\perp) O_{\alpha\beta}^{\nu\mu}(P_3) X_{\nu\mu}^{(2)}(k_{12}^\perp) \quad (83)$$

$$0^{++} \rightarrow 0_{1,+}^{++}(1(\pi 1_{1,+}^{++} \rightarrow 1(\pi f_2)))$$

$$V^{(0+,9)} = X_\mu^{(1)}(k_4^\perp) X_\nu^{(1)}(k_3^\perp) O_{\mu\nu}^{\xi\zeta}(P_3) \times X_{\xi\zeta}^{(2)}(k_{12}^\perp) \quad (84)$$

$$0^{++} \rightarrow 0_{0,+}^{++}(0(\pi 0_{2,+}^{-+} \rightarrow 2(\pi f_2)))$$

$$V^{(0+,10)} = X_{\nu_1\nu_2}^{(2)}(k_3^\perp) X_{\nu_1\nu_2}^{(2)}(k_{12}^\perp) \quad (85)$$

$$0^{++} \rightarrow 0_{2,+}^{++}(2(\pi 2_{1,+}^{-+} \rightarrow 2(\pi f_2)))$$

$$V^{(0+,11)} = X_{\alpha\beta}^{(2)}(k_4^\perp) O_{\alpha\beta}^{\eta\zeta}(P_3) X_{\eta\mu}^{(2)}(k_3^\perp) O_{\xi\nu}^{\mu\zeta}(P_3) X_{\xi\nu}^{(2)}(k_{12}^\perp) \quad (86)$$

$$2^{++} \rightarrow 2_{0,+}^{++}(2(\pi 0_{0,+}^{-+} \rightarrow 0(\pi f_0)))$$

$$V_{\alpha\beta}^{(2+,7)} = X_{\mu_1\mu_2}^{(2)}(k_4^\perp) \quad (87)$$

$$2^{++} \rightarrow 2_{0,+}^{++}(1(\pi 1_{0,+}^{++} \rightarrow 1(\pi f_0)))$$

$$V_{\alpha\beta}^{(2+,8)} = X_\alpha^{(1)}(k_4^\perp) X_\beta^{(1)}(k_3^\perp) \quad (88)$$

$$2^{++} \rightarrow 2_{0,+}^{++}(0(\pi 2_{0,+}^{-+} \rightarrow 2(\pi f_0)))$$

$$V_{\alpha\beta}^{(2+,9)} = X_{\alpha\beta}^{(2)}(k_3^\perp) \quad (89)$$

$$2^{++} \rightarrow 2_{1,+}^{++}(2(\pi 2_{0,+}^{-+} \rightarrow 2(\pi f_0)))$$

$$V_{\alpha\beta}^{(2+,10)} = X_{\alpha\nu}^{(2)}(k_4^\perp) X_{\nu\beta}^{(2)}(k_3^\perp) \quad (90)$$

$$2^{++} \rightarrow 2_{0,+}^{++}(0(\pi 2_{0,+}^{-+} \rightarrow 0(\pi f_2)))$$

$$V_{\alpha\beta}^{(2+,11)} = O_{\alpha\beta}^{\mu\nu}(P_3) X_{\mu\nu}^{(2)}(k_{12}^\perp) \quad (91)$$

$$2^{++} \rightarrow 2_{1,+}^{++}(2(\pi 2_{0,+}^{-+} \rightarrow 0(\pi f_2)))$$

$$V_{\alpha\beta}^{(2+,12)} = X_{\alpha\nu}^{(2)}(k_4^\perp) O_{\xi\zeta}^{\nu\beta}(P_3) X_{\xi\zeta}^{(2)}(k_{12}^\perp) \quad (92)$$

$$2^{++} \rightarrow 2_{0,+}^{++}(1(\pi 1_{1,+}^{++} \rightarrow 1(\pi f_2)))$$

$$V_{\alpha\beta}^{(2+,13)} = X_\alpha^{(1)}(k_4^\perp) X_{\nu_1}^{(1)}(k_3^\perp) \times O_{\beta\nu_1}^{\rho_1\rho_2}(P_3) X_{\rho_1\rho_2}^{(2)}(k_{12}^\perp) \quad (93)$$

$$2^{++} \rightarrow 2_{0,-}^{++}(1(\pi 2_{0,-}^{++} \rightarrow 1(\pi f_2)))$$

$$V_{\alpha\beta}^{(2+,14)} = \varepsilon_{\nu_1\rho_2\alpha\rho} X_{\nu_1}^{(1)}(k_4^\perp) O_{\rho_3\rho_4}^{\rho_2\beta}(P_3) \times \varepsilon_{\nu_2\nu_3\rho_3\rho_3} X_{\nu_2}^{(1)}(k_3^\perp) X_{\nu_3\rho_4}^{(2)}(k_{12}^\perp) \quad (94)$$

$$2^{++} \rightarrow 2_{0,+}^{++}(2(\pi 0_{2,+}^{-+} \rightarrow 2(\pi f_2)))$$

$$V_{\alpha\beta}^{(2+,15)} = X_{\alpha\beta}^{(2)}(k_4^\perp) X_{\nu_1\nu_2}^{(2)}(k_3^\perp) X_{\nu_1\nu_2}^{(2)}(k_{12}^\perp) \quad (95)$$

$$2^{++} \rightarrow 2_{0,+}^{++} (2 (\pi 0_{2,+}^{++} \rightarrow 2 (\pi f_2)))$$

$$V_{\alpha\beta}^{(2+,15)} = X_{\alpha\beta}^{(2)} (k_4^\perp) X_{\nu_1\nu_2}^{(2)} (k_3^\perp) X_{\nu_1\nu_2}^{(2)} (k_{12}^\perp) \quad (96)$$

$$2^{++} \rightarrow 2_{0,+}^{++} (0 (\pi 2_{1,+}^{++} \rightarrow 2 (\pi f_2)))$$

$$V_{\alpha\beta}^{(2+,16)} = O_{\alpha\beta}^{\rho_1\rho_2} (P_3) X_{\rho_1\nu_1}^{(2)} (k_3^\perp) O_{\mu\nu}^{\nu_1\rho_2} (P_3) \times X_{\mu\nu}^{(2)} (k_{12}^\perp) \quad (97)$$

$$2^{++} \rightarrow 2_{1,+}^{++} (2 (\pi 2_{1,+}^{++} \rightarrow 2 (\pi f_2)))$$

$$V_{\alpha\beta}^{(2+,17)} = X_{\rho_1\alpha}^{(2)} (k_4^\perp) O_{\rho_3\rho_2}^{\beta\rho_1} (P_3) \times X_{\rho_3\nu_1}^{(2)} (k_3^\perp) O_{\nu_1\rho_2}^{\rho_4\rho_5} (P_3) X_{\rho_4\rho_5}^{(2)} (k_{12}^\perp) \quad (98)$$

$$2^{-+} \rightarrow 2_{0,-}^{-+} (2 (\pi 1_{0,+}^{++} \rightarrow 1 (\pi f_0)))$$

$$V_{\alpha\beta}^{(2-,2)} = \varepsilon_{\nu_1\nu_2\alpha\rho} X_{\nu_1\beta}^{(2)} (k_4^\perp) X_{\nu_2}^{(1)} (k_3^\perp)$$

$$2^{-+} \rightarrow 2_{0,-}^{-+} (1 (\pi 2_{0,+}^{++} \rightarrow 2 (\pi f_2)))$$

$$V_{\alpha\beta}^{(2-,3)} = \varepsilon_{\nu_1\nu_2\alpha\rho} X_{\nu_1}^{(1)} (k_4^\perp) X_{\nu_2\beta}^{(2)} (k_3^\perp) \quad (99)$$

$$2^{-+} \rightarrow 2_{0,-}^{-+} (1 (\pi 2_{0,+}^{++} \rightarrow 0 (\pi f_2)))$$

$$V_{\alpha\beta}^{(2-,4)} = \varepsilon_{\nu_1\nu_2\alpha\rho} X_{\nu_1}^{(1)} (k_4^\perp) O_{\nu_2\beta}^{\rho_1\rho_2} (P_3) \times X_{\rho_1\rho_2}^{(2)} (k_{12}^\perp) \quad (100)$$

$$2^{-+} \rightarrow 2_{0,-}^{-+} (2 (\pi 1_{1,+}^{++} \rightarrow 1 (\pi f_2)))$$

$$V_{\alpha\beta}^{(2-,5)} = \varepsilon_{\nu_1\nu_2\alpha\rho} X_{\nu_1\beta}^{(2)} (k_4^\perp) X_{\nu_3}^{(1)} (k_3^\perp) \times O_{\nu_3\nu_2}^{\rho_1\rho_2} (P_3) X_{\rho_1\rho_2}^{(2)} (k_{12}^\perp) \quad (101)$$

$$2^{-+} \rightarrow 2_{0,+}^{-+} (0 (\pi 2_{0,-}^{++} \rightarrow 1 (\pi f_2)))$$

$$V_{\alpha\beta}^{(2-,6)} = \varepsilon_{\nu_1\nu_2\alpha\rho} X_{\nu_1}^{(1)} (k_3^\perp) \times O_{\nu_2\beta}^{\rho_1\rho_2} (P_3) X_{\rho_1\rho_2}^{(2)} (k_{12}^\perp) \quad (102)$$

$$2^{-+} \rightarrow 2_{1,+}^{-+} (2 (\pi 2_{0,-}^{++} \rightarrow 1 (\pi f_2)))$$

$$V_{\alpha\beta}^{(2-,7)} = X_{\rho_2\alpha}^{(2)} (k_4^\perp) O_{\rho_3\rho_1}^{\rho_2\beta} (P_3) \times \varepsilon_{\nu_1\nu_2\rho_3\rho_4} X_{\nu_1}^{(1)} (k_3^\perp) X_{\nu_2\rho_1}^{(2)} (k_{12}^\perp) \quad (103)$$

$$2^{-+} \rightarrow 2_{0,-}^{-+} (1 (\pi 2_{1,+}^{++} \rightarrow 2 (\pi f_2)))$$

$$V_{\alpha\beta}^{(2-,8)} = X_{\nu_1}^{(1)} (k_4^\perp) \varepsilon_{\nu_1\rho_2\alpha\rho_3} O_{\rho_3\rho_1}^{\rho_2\beta} (P_3) \times X_{\nu_2\rho_3}^{(2)} (k_3^\perp) O_{\nu_2\rho_1}^{\rho_4\rho_5} (P_3) X_{\rho_4\rho_5}^{(2)} (k_{12}^\perp) \quad (104)$$

$$0^{-+} \rightarrow 0_{2,+}^{-+} (2 (\pi 2_{0,-}^{++} \rightarrow 1 (\pi f_2)))$$

$$V_{\alpha\beta}^{(0-,2)} = X_{\rho_1\rho_2}^{(2)} (k_4^\perp) O_{\rho_3\rho_4}^{\rho_1\rho_2} (P_3) \varepsilon_{\nu_1\nu_2\rho_3\rho_4} \times X_{\nu_1}^{(1)} (k_3^\perp) O_{\nu_2\rho_4}^{\rho_5\rho_6} (P_3) X_{\rho_5\rho_6}^{(2)} (k_{12}^\perp) \quad (105)$$

$$1^{++} \rightarrow 1_{0,-}^{++} (1 (\pi 1_{0,+}^{++} \rightarrow 1 (\pi f_0)))$$

$$V_{\alpha}^{(1+,1)} = \varepsilon_{\alpha\nu_1\nu_2\rho} X_{\nu_1}^{(1)} (k_4^\perp) X_{\nu_2}^{(1)} (k_3^\perp) \quad (106)$$

$$1^{++} \rightarrow 1_{1,-}^{++} (2 (\pi 2_{0,+}^{++} \rightarrow 2 (\pi f_0)))$$

$$V_{\alpha}^{(1+,2)} = \varepsilon_{\alpha\nu_1\nu_2\rho} X_{\nu_1\nu_3}^{(2)} (k_4^\perp) X_{\nu_2\nu_3}^{(2)} (k_3^\perp) \quad (107)$$

$$1^{++} \rightarrow 1_{1,-}^{++} (2 (\pi 2_{0,+}^{++} \rightarrow 0 (\pi f_2)))$$

$$V_{\alpha}^{(1+,3)} = \varepsilon_{\alpha\nu_1\nu_2\rho} X_{\nu_1\nu_3}^{(2)} (k_4^\perp) O_{\nu_2\nu_3}^{\rho_1\rho_2} (P_3) \times X_{\rho_1\rho_2}^{(2)} (k_{12}^\perp) \quad (108)$$

$$1^{++} \rightarrow 1_{0,-}^{++} (1 (\pi 1_{1,+}^{++} \rightarrow 1 (\pi f_2)))$$

$$V_{\alpha}^{(1+,4)} = \varepsilon_{\alpha\nu_1\nu_2\rho} X_{\nu_1}^{(1)} (k_4^\perp) X_{\nu_3}^{(1)} (k_3^\perp) \times O_{\nu_3\nu_2}^{\rho_1\rho_2} (P_3) X_{\rho_1\rho_2}^{(2)} (k_{12}^\perp) \quad (109)$$

$$1^{++} \rightarrow 1_{1,+}^{++} (1 (\pi 2_{0,+}^{++} \rightarrow 1 (\pi f_2)))$$

$$V_{\alpha}^{(1+,5)} = X_{\rho_1}^{(1)} (k_4^\perp) O_{\alpha\rho_1}^{\rho_3\rho_2} (P_3) \varepsilon_{\nu_1\nu_2\rho_2\rho_3} \times X_{\nu_1}^{(1)} (k_3^\perp) O_{\nu_2\rho_3}^{\rho_4\rho_5} (P_3) X_{\rho_4\rho_5}^{(2)} (k_{12}^\perp) \quad (110)$$

$$1^{++} \rightarrow 1_{1,-}^{++} (2 (\pi 2_{1,+}^{++} \rightarrow 2 (\pi f_2)))$$

$$V_{\alpha}^{(1+,6)} = \varepsilon_{\alpha\nu_1\rho_1\rho} X_{\nu_1\rho_2}^{(2)} (k_4^\perp) O_{\rho_3\rho_4}^{\rho_1\rho_2} (P_3) \times X_{\nu_2\rho_3}^{(2)} (k_3^\perp) O_{\nu_2\rho_4}^{\rho_5\rho_6} (P_3) X_{\nu_2\rho_4}^{(2)} (k_{12}^\perp) \quad (111)$$

$$4^{++} \rightarrow 4_{0,+}^{++} (2 (\pi 2_{0,+}^{++} \rightarrow 2 (\pi f_0)))$$

$$V_{\alpha\beta\mu\nu}^{(4+,5)} = X_{\alpha\beta}^{(2)} (k_4^\perp) X_{\mu\nu}^{(2)} (k_3^\perp) \quad (112)$$

$$4^{++} \rightarrow 4_{0,+}^{++} (2 (\pi 2_{0,+}^{++} \rightarrow 0 (\pi f_2)))$$

$$V_{\alpha\beta\mu\nu}^{(4+,6)} = X_{\alpha\beta}^{(2)} (k_4^\perp) O_{\mu\nu}^{\rho_1\rho_2} (P_3) X_{\rho_1\rho_2}^{(2)} (k_{12}^\perp) \quad (113)$$

$$4^{++} \rightarrow 4_{0,+}^{++} (2 (\pi 2_{1,+}^{++} \rightarrow 2 (\pi f_2)))$$

$$V_{\alpha\beta\mu\nu}^{(4+,7)} = X_{\alpha\beta}^{(2)} (k_4^\perp) O_{\mu\nu}^{\rho_3\rho_4} (P_3) \times X_{\nu_1\rho_3}^{(2)} (k_3^\perp) O_{\nu_1\rho_4}^{\rho_1\rho_2} (P_3) X_{\rho_1\rho_2}^{(2)} (k_{12}^\perp) \quad (114)$$

The code for the generation of these tensors can be found in <https://pwa.hiskp.uni-bonn.de/version/jpsi.html>.

C. The transition amplitudes for the central production due to pomeron-pomeron ($f_0 f_0$) collision.

In the pomeron-pomeron central collision (or in $f_0 f_0$ collision) only isoscalar states with even spin, positive P, C -parity are produced. The production vertices are described by the orbital momentum tensors. Therefore, we

obtain the following expressions for the partial-wave amplitudes:

$$\begin{aligned} A^{(0+,i)} &= V^{(0+,i)} & i &= 1-11, \\ A^{(2+,i)} &= X_{\mu\nu}^{(2)}(q^\perp) O_{\alpha\beta}^{\mu\nu} V_{\alpha\beta}^{(2+,i)} & i &= 1-17, \\ A^{(4+,i)} &= X_{\mu\nu\chi\xi}^{(4)}(q^\perp) O_{\mu\nu\chi\xi}^{\alpha\beta\eta\zeta} V_{\alpha\beta\eta\zeta}^{(4+,i)} & i &= 1-7 \end{aligned} \quad (115)$$

D. The radiative J/Ψ decay into $4\pi^0$ final state.

In the case of the radiative J/ψ it is usually assumed that after emission of the photon the $c\bar{c}$ system is annihilated to a resonance which decays into hadron final states. From this point of view, all states that are allowed in the quark model can be produced. These states have isospin $I = 0$ and charged parity $C = +1$. Let us give the list of the amplitudes up to $J = 4$:

$$\begin{aligned} A^{(0+,i)} &= S^0 V^{(0+,i)} & i &= 1-11, \\ A_1^{(2+,i)} &= S^0 X_{\mu\nu}^{(2)}(q^\perp) O_{\alpha\beta}^{\mu\nu} V_{\alpha\beta}^{(2+,i)} & i &= 1-17, \\ A_2^{(2+,i)} &= \varepsilon_{\mu\eta\xi\rho} S^1 X_{\xi\nu}^{(2)}(q^\perp) O_{\alpha\beta}^{\mu\nu} V_{\alpha\beta}^{(2+,i)} & i &= 1-17, \\ A_3^{(2+,i)} &= S_{\mu\nu}^2 O_{\alpha\beta}^{\mu\nu} V_{\alpha\beta}^{(2+,i)} & i &= 1-17, \\ A_1^{(4+,i)} &= S^0 X_{\mu\nu\alpha\beta}^{(4)}(q^\perp) O_{\eta\chi\xi\varrho}^{\mu\nu\alpha\beta} V_{\eta\chi\xi\varrho}^{(4+,i)} & i &= 1-7, \\ A_2^{(4+,i)} &= \varepsilon_{\mu\eta\xi\rho} S^1 X_{\xi\nu\alpha\beta}^{(4)}(q^\perp) O_{\chi\xi\varrho\delta}^{\mu\nu\alpha\beta} V_{\chi\xi\varrho\delta}^{(4+,i)} & i &= 1-7, \\ A_3^{(4+,i)} &= S_{\mu\nu}^2 X_{\alpha\beta}^{(2)}(q^\perp) O_{\eta\chi\xi\varrho}^{\mu\nu\alpha\beta} V_{\eta\chi\xi\varrho}^{(4+,i)} & i &= 1-7, \\ A^{(0-,i)} &= S^1 X_{\eta}^{(1)}(q^\perp) V^{(0-,i)} & i &= 1-2, \\ A_1^{(1+,i)} &= S_{\mu}^1 V_{\mu}^{(1+,i)} & i &= 1-6, \\ A_2^{(1+,i)} &= S_{\eta}^1 X_{\eta\mu}^{(2)}(q^\perp) V_{\mu}^{(1+,i)} & i &= 1-6, \\ A_1^{(2-,i)} &= S_{\mu}^1 X_{\nu}^{(1)}(q^\perp) O_{\mu\nu}^{\alpha\beta} V_{\alpha\beta}^{(2-,i)} & i &= 1-8, \\ A_2^{(2-,i)} &= S_{\eta}^1 X_{\eta\mu\nu}^{(3)}(q^\perp) O_{\mu\nu}^{\alpha\beta} V_{\alpha\beta}^{(2-,i)} & i &= 1-8, \\ A_3^{(2-,i)} &= \varepsilon_{\mu\eta\xi\rho} S^2 X_{\eta\nu}^{(1)}(q^\perp) O_{\mu\nu}^{\alpha\beta} V_{\alpha\beta}^{(2-,i)} & i &= 1-8 \end{aligned} \quad (116)$$

References

- [1] D. Alde *et al.* (GAMS), *Eur. Phys. J. A* **3**, 361 (1998)
- [2] G. Grayer *et al.*, *Nucl. Phys. B* **75**, 189 (1974)
- [3] R. S. Longacre *et al.*, *Phys. Lett. B* **177**, 223 (1986)
- [4] M. Ablikim *et al.* (BESIII), *Phys. Rev. D* **92**, 052003 (2015), [Erratum: *Phys. Rev. D* **93**, 039906 (2016)], 1506.00546.
- [5] M. Ablikim *et al.* (BESIII), *Phys. Rev. D* **98**, 072003 (2018), arXiv: 1808.06946
- [6] D. V. Bugg, V. V. Anisovich, A. Sarantsev, and B. S. Zou, *Phys. Rev. D* **50**, 4412 (1994)
- [7] C. Amsler *et al.*, *Phys. Lett. B* **342**, 433 (1995)
- [8] C. Amsler *et al.* (Crystal Barrel), *Phys. Lett. B* **353**, 571 (1995)
- [9] J. Z. Bai *et al.* (BES), *Phys. Lett. B* **472**, 207 (2000), arXiv: hep-ex/9909040
- [10] D. Coffman *et al.* (MARK-III), *Phys. Rev. D* **45**, 2196 (1992)
- [11] M. Artuso *et al.* (CLEO), *Phys. Rev. D* **85**, 122002 (2012), arXiv: 1201.5716
- [12] R. Aaij *et al.* (LHCb), *JHEP* **01**, 054 (2025), arXiv: 2407.12475
- [13] U. Thoma (Crystal Barrel), *AIP Conf. Proc.* **432**, 322 (1998)
- [14] A. P. Szczepaniak and E. S. Swanson, *Phys. Lett. B* **577**, 61 (2003), arXiv: hep-ph/0308268
- [15] A. V. Sarantsev, I. Denisenko, U. Thoma, and E. Klempt, *Phys. Lett. B* **816**, 136227 (2021), arXiv: 2103.09680
- [16] E. Klempt and A. V. Sarantsev, *Phys. Lett. B* **826**, 136906 (2022), arXiv: 2112.04348
- [17] A. Rodas, A. Pilloni, M. Albaladejo, C. FernandezRamirez, V. Mathieu, and A. P. Szczepaniak (Joint Physics Analysis Center), *Eur. Phys. J. C* **82**, 80 (2022), arXiv: 2110.00027
- [18] A. Athenodorou and M. Teper, *JHEP* **11**, 172 (2020), arXiv: 2007.06422
- [19] E. Gregory, A. Irving, B. Lucini, C. McNeile, A. Rago, C. Richards, and E. Rinaldi, *JHEP* **10**, 170 (2012), arXiv: 1208.1858
- [20] E. Klempt, K. V. Nikonov, A. V. Sarantsev, and I. Denisenko, *Phys. Lett. B* **830**, 137171 (2022), arXiv: 2205.07239
- [21] C. Zemach, *Phys. Rev.* **140**, B109 (1965)
- [22] S. U. Chung, *Phys. Rev. D* **57**, 431 (1998)
- [23] B. S. Zou and D. V. Bugg, *Eur. Phys. J. A* **16**, 537 (2003), arXiv: hep-ph/0211457
- [24] A. V. Anisovich, V. V. Anisovich, V. N. Markov, M. A. Matveev, and A. V. Sarantsev, *J. Phys. G* **28**, 15 (2002), arXiv: hep-ph/0105330
- [25] A. Anisovich, E. Klempt, A. Sarantsev, and U. Thoma, *Eur. Phys. J. A* **24**, 111 (2005), arXiv: hep-ph/0407211
- [26] A. V. Anisovich, K. V. Nikonov, and A. V. Sarantsev, *Phys. Atom. Nucl.* **87**, 498 (2024), arXiv: 2406.01172