

# Observation of the $X(2370)$ in $J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$ and $J/\psi \rightarrow \gamma \pi^0 \pi^0 \eta^*$

M. Ablikim (麦迪娜)<sup>1</sup>  M. N. Achasov<sup>4,c</sup>  P. Adlarson<sup>83</sup>  X. C. Ai (艾小聪)<sup>89</sup>  C. S. Akondi<sup>31A,31B</sup>   
R. Aliberti<sup>39</sup>  A. Amoroso<sup>82A,82C</sup>  Q. An (安琪)<sup>78,65</sup>  Y. H. An<sup>89</sup>  M. S. Anderson<sup>39</sup>  Y. Bai (白羽)<sup>63</sup>   
O. Bakina<sup>40</sup>  H. R. Bao (包浩然)<sup>71</sup>  X. L. Bao (保先岚)<sup>50</sup>  M. Barbagiovanni<sup>82C</sup>  V. Batozskaya<sup>1,49</sup>   
K. Begzsuren<sup>35</sup>  N. Berger<sup>39</sup>  M. Berlowski<sup>49</sup>  M. B. Bertani<sup>30A</sup>  D. Bettoni<sup>31A</sup>  F. Bianchi<sup>82A,82C</sup>   
E. Bianco<sup>82A,82C</sup>  A. Bortone<sup>82A,82C</sup>  I. Boyko<sup>40</sup>  R. Briere<sup>5</sup>  A. Brueggemann<sup>75</sup>  D. Cabiati<sup>82A,82C</sup>   
H. Cai (蔡浩)<sup>84</sup>  M. H. Cai (蔡铭航)<sup>42,k,l</sup>  X. Cai (蔡啸)<sup>1,65</sup>  A. Calcaterra<sup>30A</sup>  G. F. Cao (曹国富)<sup>1,71</sup>   
N. Cao (曹宁)<sup>1,71</sup>  S. A. Cetin<sup>69A</sup>  X. Y. Chai (柴新宇)<sup>51,h</sup>  J. F. Chang (常劲帆)<sup>1,65</sup>  T. T. Chang (常甜甜)<sup>48</sup>   
G. R. Che (车国荣)<sup>48</sup>  Y. Z. Che (车逾之)<sup>1,65,71</sup>  C. H. Chen (陈春卉)<sup>10</sup>  C. Chen (陈超)<sup>1</sup>   
G. Chen (陈刚)<sup>1</sup>  H. S. Chen (陈和生)<sup>1,71</sup>  H. Y. Chen (陈弘扬)<sup>20</sup>  M. L. Chen (陈玛丽)<sup>1,65,71</sup>   
S. J. Chen (陈申见)<sup>47</sup>  S. M. Chen (陈少敏)<sup>68</sup>  T. Chen (陈通)<sup>1,71</sup>  W. Chen (陈旺)<sup>50</sup>   
X. R. Chen (陈旭荣)<sup>34,71</sup>  X. T. Chen (陈肖婷)<sup>1,71</sup>  X. Y. Chen (陈心宇)<sup>12,g</sup>  Y. B. Chen (陈元柏)<sup>1,65</sup>   
Y. Q. Chen<sup>16</sup>  Z. K. Chen (陈梓康)<sup>66</sup>  J. Cheng (程捷)<sup>50</sup>  L. N. Cheng (程利娜)<sup>48</sup>  S. K. Choi<sup>11</sup>   
X. Chu (初晓)<sup>12,g</sup>  G. Ciminetto<sup>31A</sup>  F. Cossio<sup>82C</sup>  J. Cottee-Meldrum<sup>70</sup>  H. L. Dai (代洪亮)<sup>1,65</sup>   
J. P. Dai (代建平)<sup>87</sup>  X. C. Dai (戴鑫琛)<sup>68</sup>  A. Dbeyssi<sup>19</sup>  R. E. de Boer<sup>3</sup>  D. Dedovich<sup>40</sup>   
C. Q. Deng (邓创旗)<sup>80</sup>  Z. Y. Deng (邓子艳)<sup>1</sup>  A. Denig<sup>39</sup>  I. Denisenko<sup>40</sup>  M. Destefanis<sup>82A,82C</sup>   
F. De Mori<sup>82A,82C</sup>  E. Di Fiore<sup>31A,31B</sup>  X. X. Ding (丁晓萱)<sup>51,h</sup>  Y. Ding (丁勇)<sup>44</sup>  Y. X. Ding (丁玉鑫)<sup>32</sup>   
J. Dong (董静)<sup>1,65</sup>  L. Y. Dong (董燎原)<sup>1,71</sup>  M. Y. Dong (董明义)<sup>1,65,71</sup>  X. Dong (董翔)<sup>84</sup>   
Z. J. Dong (董焯娟)<sup>66</sup>  M. C. Du (杜蒙川)<sup>1</sup>  S. X. Du (杜书先)<sup>89</sup>  S. X. Du (杜少旭)<sup>12,g</sup>   
X. L. Du (杜小龙)<sup>12,g</sup>  Y. Q. Du (杜一琦)<sup>84</sup>  Y. Y. Duan (段尧予)<sup>61</sup>  Z. H. Duan (段宗欢)<sup>47</sup>  P. Egorov<sup>40,a</sup>   
G. F. Fan (樊高峰)<sup>47</sup>  J. J. Fan (樊俊杰)<sup>20</sup>  Y. H. Fan (范宇晗)<sup>50</sup>  J. Fang (方建)<sup>1,65</sup>  J. Fang (方进)<sup>66</sup>   
S. S. Fang (房双世)<sup>1,71</sup>  W. X. Fang (方文兴)<sup>1</sup>  Y. Q. Fang (方亚泉)<sup>1,65</sup>  L. Fava<sup>82B,82C</sup>  F. Feldbauer<sup>3</sup>   
G. Felici<sup>30A</sup>  C. Q. Feng (封常青)<sup>78,65</sup>  J. H. Feng (冯俊华)<sup>16</sup>  Q. X. Feng (冯千禧)<sup>42,k,l</sup>   
Y. T. Feng (冯筠潼)<sup>78,65</sup>  M. Fritsch<sup>3</sup>  C. D. Fu (傅成栋)<sup>1</sup>  J. L. Fu (傅金林)<sup>71</sup>  Y. W. Fu (傅亦威)<sup>1,71</sup>   
H. Gao (高涵)<sup>71</sup>  Xu Gao (高旭)<sup>38</sup>  Y. Gao (高扬)<sup>78,65</sup>  Y. N. Gao (高原宁)<sup>51,h</sup>  Y. Y. Gao (高洋洋)<sup>32</sup>   
Y. N. Gao (高语浓)<sup>20</sup>  Z. Gao (高枝)<sup>48</sup>  S. Garbolino<sup>82C</sup>  I. Garzia<sup>31A,31B</sup>  L. Ge (葛玲)<sup>63</sup>   
P. T. Ge (葛潘婷)<sup>20</sup>  Z. W. Ge (葛振武)<sup>47</sup>  C. Geng (耿聪)<sup>66</sup>  A. Gilman<sup>76</sup>  K. Goetzen<sup>13</sup>  J. Gollub<sup>3</sup>   
J. B. Gong (龚家宝)<sup>1,71</sup>  J. D. Gong (龚家鼎)<sup>38</sup>  L. Gong (龚丽)<sup>44</sup>  W. X. Gong (龚文煊)<sup>1,65</sup>  W. Gradl<sup>39</sup>   
M. Greco<sup>82A,82C</sup>  M. D. Gu (顾萌冬)<sup>56</sup>  M. H. Gu (顾旻皓)<sup>1,65</sup>  C. Y. Guan (关春懿)<sup>1,71</sup>   
A. Q. Guo (郭爱强)<sup>34</sup>  H. Guo (郭昊)<sup>55</sup>  J. N. Guo (郭嘉宁)<sup>12,g</sup>  L. B. Guo (郭立波)<sup>46</sup>  M. J. Guo (国梦娇)<sup>55</sup> 

Received 3 June 2026; Accepted 15 July 2026; Accepted manuscript online 16 July 2026

\* This work is supported in part by the National Key R&D Program of China under Contracts Nos. 2025YFA1613900, 2023YFA1606000, 2023YFA1606704; National Natural Science Foundation of China (NSFC) under Contracts Nos. 11635010, 11935015, 11935016, 11935018, 12025502, 12035009, 12035013, 12061131003, 12192260, 12192261, 12192262, 12192263, 12192264, 12192265, 12221005, 12225509, 12235017, 12342502, 12361141819, 12535005; the Chinese Academy of Sciences (CAS) Large-Scale Scientific Facility Program; the Strategic Priority Research Program of Chinese Academy of Sciences under Contract No. XDA0480600; CAS under Contract No. YSBR-101; 100 Talents Program of CAS; China Postdoctoral Science Foundation under Contracts Nos. GZC20252256, 2025M773423; The Institute of Nuclear and Particle Physics (INPAC) and Shanghai Key Laboratory for Particle Physics and Cosmology; Agencia Nacional de Investigación y Desarrollo de Chile (ANID), Chile under Contract No. ANID CCTVal CIA250027; ERC under Contract No. 758462; German Research Foundation DFG under Contract No. FOR5327; Istituto Nazionale di Fisica Nucleare, Italy; Knut and Alice Wallenberg Foundation under Contracts Nos. 2021.0174, 2021.0299, 2023.0315; Ministry of Development of Turkey under Contract No. DPT2006K-120470; National Research Foundation of Korea under Contract No. NRF-2022R1A2C1092335; National Science and Technology fund of Mongolia; Polish National Science Centre under Contract No. 2024/53/B/ST2/00975; STFC (United Kingdom); Swedish Research Council under Contract No. 2019.04595; U. S. Department of Energy under Contract No. DE-FG02-05ER41374

<sup>†</sup> E-mail: besiii-publications@ihep.ac.cn

©2026 Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd. All rights, including for text and data mining, AI training, and similar technologies, are reserved.

R. P. Guo (郭如盼)<sup>54</sup>  X. Guo (郭旭)<sup>55</sup>  Y. P. Guo (郭玉萍)<sup>12,g</sup>  Z. Guo (郭震)<sup>78,65</sup>  A. Guskov<sup>40,a</sup>   
J. Gutierrez<sup>29</sup>  J. Y. Han (韩靖宜)<sup>78,65</sup>  T. T. Han<sup>1</sup>  X. Han (韩许)<sup>78,65</sup>  F. Hanisch<sup>3</sup>   
K. D. Hao (郝科迪)<sup>78,65</sup>  X. Q. Hao (郝喜庆)<sup>20</sup>  F. A. Harris<sup>72</sup>  C. Z. He (何承芝)<sup>51,h</sup>  K. K. He (何凯凯)<sup>17,47</sup>   
K. L. He (何康林)<sup>1,71</sup>  F. H. Heinsius<sup>3</sup>  C. H. Heinz<sup>39</sup>  Y. K. Heng (衡月昆)<sup>1,65,71</sup>  C. Herold<sup>67</sup>   
P. C. Hong (洪鹏程)<sup>38</sup>  G. Y. Hou (侯国一)<sup>1,71</sup>  X. T. Hou (侯贤涛)<sup>1,71</sup>  Y. R. Hou (侯颖锐)<sup>71</sup>   
Z. L. Hou (侯治龙)<sup>1</sup>  H. M. Hu (胡海明)<sup>1,71</sup>  J. F. Hu (胡继峰)<sup>62,j</sup>  Q. P. Hu (胡启鹏)<sup>78,65</sup>   
S. L. Hu (胡圣亮)<sup>12,g</sup>  T. Hu (胡涛)<sup>1,65,71</sup>  Y. Hu (胡誉)<sup>1</sup>  Y. X. Hu (胡宇轩)<sup>84</sup>  Z. M. Hu (胡忠敏)<sup>66</sup>   
G. S. Huang (黄光顺)<sup>78,65</sup>  K. X. Huang (黄凯旋)<sup>66</sup>  L. Q. Huang (黄麟钦)<sup>34,71</sup>  P. Huang (黄盼)<sup>47</sup>   
X. T. Huang (黄性涛)<sup>55</sup>  Y. P. Huang (黄燕萍)<sup>1</sup>  Y. S. Huang (黄永盛)<sup>66</sup>  T. Hussain<sup>81</sup>  N. Hüsken<sup>39</sup>   
N. in der Wiesche<sup>75</sup>  J. Jackson<sup>29</sup>  Q. Ji (纪全)<sup>1</sup>  Q. P. Ji (姬清平)<sup>20</sup>  W. Ji (季旺)<sup>1,71</sup>  X. B. Ji (季晓斌)<sup>1,71</sup>   
X. L. Ji (季筱璐)<sup>1,65</sup>  Y. Y. Ji (吉钰瑶)<sup>1</sup>  L. K. Jia (贾乐康)<sup>71</sup>  X. Q. Jia (贾晓倩)<sup>55</sup>  D. Jiang (姜地)<sup>1,71</sup>   
S. J. Jiang (蒋思婧)<sup>10</sup>  X. S. Jiang (江晓山)<sup>1,65,71</sup>  Y. Jiang (蒋艺)<sup>71</sup>  J. B. Jiao (焦健斌)<sup>55</sup>   
J. K. Jiao (焦俊坤)<sup>38</sup>  Z. Jiao (焦铮)<sup>25</sup>  L. C. L. Jin (金梁程龙)<sup>1</sup>  S. Jin (金山)<sup>47</sup>  Y. Jin (金毅)<sup>73</sup>   
M. Q. Jing (荆茂强)<sup>56</sup>  X. M. Jing (景新娟)<sup>71</sup>  T. Johansson<sup>83</sup>  S. Kabana<sup>36</sup>  X. L. Kang (康晓琳)<sup>10</sup>   
X. S. Kang (康晓坤)<sup>44</sup>  B. C. Ke (柯百谦)<sup>89</sup>  V. Khachatryan<sup>29</sup>  A. Khoukaz<sup>75</sup>  O. B. Kolcu<sup>69A</sup>  B. Kopf<sup>2</sup>   
L. Kröger<sup>75</sup>  L. Krümmel<sup>3</sup>  Y. Y. Kuang (匡洋洋)<sup>80</sup>  X. Kui (奎贤)<sup>1,71</sup>  N. Kumar<sup>28</sup>  A. Kupsc<sup>49,83</sup>   
W. Kühn<sup>41</sup>  Q. Lan (兰强)<sup>80</sup>  W. N. Lan (兰文宁)<sup>20</sup>  T. T. Lei (雷天天)<sup>78,65</sup>  M. Lellmann<sup>39</sup>  T. Lenz<sup>39</sup>   
C. Li (李翠)<sup>52</sup>  C. H. Li (李春花)<sup>46</sup>  C. K. Li (李春凯)<sup>48</sup>  Chunkai Li (李春凯)<sup>21</sup>  Cong Li (李聪)<sup>48</sup>   
D. M. Li (李德民)<sup>89</sup>  F. Li (李飞)<sup>1,65</sup>  G. Li (李刚)<sup>1</sup>  H. B. Li (李海波)<sup>1,71</sup>  H. J. Li (李惠静)<sup>20</sup>   
H. L. Li (李昊霖)<sup>89</sup>  H. N. Li (李衡讷)<sup>62,j</sup>  H. P. Li (李昊鹏)<sup>48</sup>  H. Li (李慧)<sup>48</sup>  J. N. Li (李佳宁)<sup>32</sup>   
J. S. Li (李静舒)<sup>66</sup>  J. W. Li (李井文)<sup>55</sup>  K. Li (李科)<sup>1</sup>  K. L. Li (李凯璐)<sup>42,k,l</sup>  L. J. Li (李林健)<sup>1,71</sup>   
L. K. Li (李龙科)<sup>26</sup>  L. Li (李蕾)<sup>53</sup>  M. H. Li (李明浩)<sup>48</sup>  M. R. Li (李明润)<sup>1,71</sup>  M. T. Li (李梦婷)<sup>55</sup>   
P. L. Li (李佩莲)<sup>71</sup>  P. R. Li (李培荣)<sup>42,k,l</sup>  Q. M. Li (李启铭)<sup>1,71</sup>  Q. X. Li (李起鑫)<sup>55</sup>  R. Li (李燃)<sup>18,34</sup>   
S. Li (李硕)<sup>89</sup>  S. X. Li (李素娴)<sup>89</sup>  S. Y. Li (李淑雨)<sup>89</sup>  S. S. Li (李珊珊)<sup>27,i</sup>  T. Li (李腾)<sup>55</sup>   
T. Y. Li (李天佑)<sup>48</sup>  W. D. Li (李卫东)<sup>1,71</sup>  W. G. Li (李卫国)<sup>1</sup>  X. Li (李旭)<sup>1,71</sup>  X. H. Li (李旭红)<sup>78,65</sup>   
X. K. Li (李小康)<sup>51,h</sup>  X. L. Li (李晓玲)<sup>55</sup>  X. Y. Li (李晓宇)<sup>78,65</sup>  X. Z. Li (李绪泽)<sup>66</sup>  Y. Li (李洋)<sup>20</sup>   
Y. H. Li (黎昱宏)<sup>48</sup>  Y. B. Li (李郁博)<sup>85</sup>  Y. C. Li (李钰程)<sup>66</sup>  Y. G. Li (李彦谷)<sup>71</sup>  Y. P. Li (李雁鹏)<sup>38</sup>   
Z. H. Li (李宗豪)<sup>42</sup>  Z. J. Li (李志军)<sup>66</sup>  Z. L. Li (李芝兰)<sup>89</sup>  Z. X. Li (李振轩)<sup>48</sup>  Z. Y. Li (李紫阳)<sup>87</sup>   
C. Liang (梁畅)<sup>47</sup>  H. Liang (梁昊)<sup>78,65</sup>  Y. F. Liang (梁勇飞)<sup>60</sup>  Y. T. Liang (梁羽铁)<sup>34,71</sup>   
Z. Z. Liang (梁振卓)<sup>66</sup>  G. R. Liao (廖广睿)<sup>14</sup>  L. B. Liao (廖立波)<sup>66</sup>  M. H. Liao (廖明华)<sup>66</sup>   
Y. P. Liao (廖一朴)<sup>1,71</sup>  J. Libby<sup>28</sup>  A. Limphirat<sup>67</sup>  C. C. Lin (蔺长城)<sup>61</sup>  C. X. Lin (林创新)<sup>34</sup>   
D. X. Lin (林德旭)<sup>34,71</sup>  T. Lin (林韬)<sup>1</sup>  B. J. Liu (刘北江)<sup>1</sup>  B. X. Liu (刘宝鑫)<sup>84</sup>  C. Liu (刘成)<sup>38</sup>   
C. X. Liu (刘春秀)<sup>1</sup>  F. Liu (刘芳)<sup>1</sup>  F. H. Liu (刘福虎)<sup>59</sup>  F. Liu (刘峰)<sup>6</sup>  G. M. Liu (刘国明)<sup>62,j</sup>   
H. Liu (刘昊)<sup>42,k,l</sup>  H. B. Liu (刘宏邦)<sup>15</sup>  H. M. Liu (刘怀民)<sup>1,71</sup>  H. H. Liu (刘汇慧)<sup>22</sup>   
J. B. Liu (刘建北)<sup>78,65</sup>  J. J. Liu (刘佳佳)<sup>21</sup>  K. Liu (刘凯)<sup>42,k,l</sup>  K. Y. Liu (刘魁勇)<sup>44</sup>  K. Liu (刘珂)<sup>23</sup>   
K. Liu (刘坤)<sup>80</sup>  L. Liu (刘亮)<sup>42</sup>  L. C. Liu (刘良辰)<sup>48</sup>  Lu Liu (刘露)<sup>48</sup>  M. H. Liu<sup>38</sup>   
P. L. Liu (刘佩莲)<sup>55</sup>  Q. Liu (刘倩)<sup>71</sup>  S. B. Liu (刘树彬)<sup>78,65</sup>  T. Liu (刘桐)<sup>1</sup>  W. M. Liu (刘卫民)<sup>78,65</sup>   
W. T. Liu (刘婉婷)<sup>43</sup>  X. Liu (刘翔)<sup>42,k,l</sup>  X. K. Liu (刘筱康)<sup>42,k,l</sup>  X. L. Liu (刘兴淋)<sup>12,g</sup>   
X. P. Liu (刘新平)<sup>12,g</sup>  X. T. Liu (刘欣桐)<sup>21</sup>  X. Y. Liu (刘雪吟)<sup>84</sup>  Y. Liu (刘英)<sup>42,k,l</sup>  Y. B. Liu (刘玉斌)<sup>48</sup>   
Y. Liu (刘义)<sup>89</sup>  Z. A. Liu (刘振安)<sup>1,65,71</sup>  Z. D. Liu (刘宗德)<sup>85</sup>  Z. L. Liu (刘智青)<sup>80</sup>  Z. Q. Liu (刘钟秀)<sup>55</sup>   
Z. X. Liu (刘正扬)<sup>1</sup>  Z. Y. Liu (刘子强)<sup>42</sup>  X. C. Lou (娄辛丑)<sup>1,65,71</sup>  H. J. Lu (吕海江)<sup>25</sup>   
J. G. Lu (吕军光)<sup>1,65</sup>  X. L. Lu (陆小玲)<sup>16</sup>  Y. Lu (卢宇)<sup>7</sup>  Y. H. Lu (卢泱宏)<sup>1,71</sup>  Y. P. Lu (卢云鹏)<sup>1,65</sup>   
Z. H. Lu (卢泽辉)<sup>1,71</sup>  C. L. Luo (罗成林)<sup>46</sup>  J. R. Luo (罗家瑞)<sup>66</sup>  J. S. Luo (罗家顺)<sup>1,71</sup>   
M. X. Luo (罗民兴)<sup>88</sup>  T. Luo (罗涛)<sup>12,g</sup>  X. L. Luo (罗小兰)<sup>1,65</sup>  Z. Y. Lv (吕在裕)<sup>23</sup>  X. R. Lyu (吕晓睿)<sup>71,o</sup>   
Y. F. Lyu (吕翌丰)<sup>48</sup>  Y. H. Lyu (吕云鹤)<sup>89</sup>  F. C. Ma (马凤才)<sup>44</sup>  H. L. Ma (马海龙)<sup>1</sup>  H. Ma (马衡)<sup>27,i</sup>   
J. L. Ma (马俊力)<sup>1,71</sup>  L. L. Ma (马连良)<sup>55</sup>  L. R. Ma (马立瑞)<sup>73</sup>  Q. M. Ma (马秋梅)<sup>1</sup> 

- R. Q. Ma (马润秋)<sup>1,71</sup>  R. Y. Ma (马若云)<sup>20</sup>  T. Ma (马腾)<sup>78,65</sup>  X. T. Ma (马晓天)<sup>1,71</sup>  X. Y. Ma (马晓妍)<sup>1,65</sup>   
F. E. Maas<sup>19</sup>  I. MacKay<sup>76</sup>  M. Maggiora<sup>82A,82C</sup>  S. Maity<sup>34</sup>  S. Malde<sup>76</sup>  L. M. Mansur<sup>39</sup>   
Y. J. Mao (冒亚军)<sup>51,h</sup>  Z. P. Mao (毛泽普)<sup>1</sup>  S. Marcello<sup>82A,82C</sup>  A. Marshall<sup>70</sup>  F. M. Melendi<sup>31A,31B</sup>   
Y. H. Meng (孟琰皓)<sup>71</sup>  Z. X. Meng (孟召霞)<sup>73</sup>  G. Mezzadri<sup>31A</sup>  H. Miao (妙晗)<sup>1,71</sup>  T. J. Min (闵天觉)<sup>47</sup>   
R. E. Mitchell<sup>29</sup>  X. H. Mo (莫晓虎)<sup>1,65,71</sup>  B. Moses<sup>29</sup>  N. Yu. Muchnoi<sup>4,e</sup>  J. Muskalla<sup>39</sup>  Y. Nefedov<sup>40</sup>   
F. Nerling<sup>19,e</sup>  H. Neuwirth<sup>75</sup>  Z. Ning (宁哲)<sup>1,65</sup>  S. Nisar<sup>33</sup>  Q. L. Niu (牛祺乐)<sup>42,k,l</sup>   
W. D. Niu (牛文迪)<sup>12,g</sup>  Y. Niu (牛艳)<sup>55</sup>  C. Normand<sup>70</sup>  S. L. Olsen<sup>11,71</sup>  Q. Ouyang (欧阳群)<sup>1,65,71</sup>   
I. V. Ovtin<sup>4</sup>  S. Pacetti<sup>30B,30C</sup>  Y. Pan (潘越)<sup>63</sup>  C. Y. Pang (庞彩莹)<sup>14</sup>  A. Pathak<sup>11</sup>  Y. P. Pei (裴宇鹏)<sup>78,65</sup>   
M. Pelizaeus<sup>3</sup>  G. L. Peng (彭高乐)<sup>78,65</sup>  H. P. Peng (彭海平)<sup>78,65</sup>  X. J. Peng (彭宣嘉)<sup>42,k,l</sup>   
Y. Y. Peng (彭云翊)<sup>42,k,l</sup>  K. Peters<sup>13,e</sup>  K. Petridis<sup>70</sup>  J. L. Ping (平加伦)<sup>46</sup>  R. G. Ping (平荣刚)<sup>1,71</sup>   
S. Plura<sup>39</sup>  V. Prasad<sup>38</sup>  L. Pöpping<sup>3</sup>  F. Z. Qi (齐法制)<sup>1</sup>  H. R. Qi (漆红荣)<sup>68</sup>  S. Qian (钱森)<sup>1,65</sup>   
W. B. Qian (钱文斌)<sup>71</sup>  C. F. Qiao (乔从丰)<sup>71</sup>  J. H. Qiao (乔佳辉)<sup>20</sup>  J. J. Qin (秦佳佳)<sup>80</sup>   
J. L. Qin (覃嘉良)<sup>61</sup>  L. Q. Qin (秦丽清)<sup>14</sup>  L. Y. Qin (秦龙宇)<sup>78,65</sup>  P. B. Qin (秦鹏勃)<sup>80</sup>   
X. P. Qin (覃潇萍)<sup>43</sup>  X. S. Qin (秦小帅)<sup>55</sup>  Z. H. Qin (秦中华)<sup>1,65</sup>  J. F. Qiu (邱进发)<sup>1</sup>  Z. H. Qu (屈子皓)<sup>80</sup>   
J. Rademacker<sup>70</sup>  K. Ravindran<sup>74</sup>  C. F. Redmer<sup>39</sup>  A. Rivetti<sup>82C</sup>  M. Rolo<sup>82C</sup>  G. Rong (荣刚)<sup>1,71</sup>   
S. S. Rong (荣少石)<sup>1,71</sup>  F. Rosini<sup>30B,30C</sup>  Ch. Rosner<sup>19</sup>  M. Q. Ruan (阮曼奇)<sup>1,65</sup>  W. R. Ruangyoo<sup>67</sup>   
N. Salone<sup>79</sup>  A. Sarantsev<sup>40,d</sup>  Y. Schelhaas<sup>39</sup>  M. Schernau<sup>36</sup>  K. Schoenning<sup>83</sup>  M. Scodreggio<sup>31A</sup>   
W. Shan (单葳)<sup>26</sup>  X. Y. Shan (单心钰)<sup>78,65</sup>  Z. J. Shang (尚子杰)<sup>42,k,l</sup>  J. F. Shangguan (上官剑锋)<sup>17</sup>   
L. G. Shao (邵立港)<sup>1,71</sup>  M. Shao (邵明)<sup>78,65</sup>  C. P. Shen (沈成平)<sup>12,g</sup>  H. F. Shen (沈宏飞)<sup>1,9</sup>   
W. H. Shen (沈文涵)<sup>71</sup>  X. Y. Shen (沈肖雁)<sup>1,71</sup>  B. A. Shi (施伯安)<sup>71</sup>  Ch. Y. Shi (石超逸)<sup>87,b</sup>   
H. Shi (史华)<sup>78,65</sup>  J. L. Shi (石家磊)<sup>8,p</sup>  J. Y. Shi (石京燕)<sup>1</sup>  M. H. Shi (时明辉)<sup>89</sup>  S. Shi (石硕)<sup>80</sup>   
S. Y. Shi (史书宇)<sup>80</sup>  X. Shi (史欣)<sup>1,65</sup>  H. L. Song (宋海林)<sup>78,65</sup>  J. J. Song (宋娇娇)<sup>20</sup>   
M. H. Song (宋铭昊)<sup>42</sup>  T. Z. Song (宋天资)<sup>66</sup>  W. M. Song (宋维民)<sup>38</sup>  Y. X. Song (宋昀轩)<sup>51,h,m</sup>   
Z. R. Song (宋子荣)<sup>27,i</sup>  S. Sosio<sup>82A,82C</sup>  S. Spataro<sup>82A,82C</sup>  S. Stansilaus<sup>76</sup>  F. Stieler<sup>39</sup>  M. Stolte<sup>3</sup>   
S. S. Su (苏闪闪)<sup>44</sup>  G. B. Sun (孙光豹)<sup>84</sup>  G. X. Sun (孙功星)<sup>1</sup>  H. Sun (孙昊)<sup>71</sup>  H. K. Sun (孙浩凯)<sup>1</sup>   
J. F. Sun (孙俊峰)<sup>20</sup>  K. Sun (孙开)<sup>68</sup>  L. Sun (孙亮)<sup>84</sup>  R. Sun (孙睿)<sup>78</sup>  S. S. Sun (孙胜森)<sup>1,71</sup>  T. Sun<sup>57,f</sup>   
W. Y. Sun (孙文玉)<sup>56</sup>  Y. C. Sun (孙雨长)<sup>84</sup>  Y. H. Sun (孙益华)<sup>32</sup>  Y. J. Sun (孙勇杰)<sup>78,65</sup>   
Y. Z. Sun (孙永昭)<sup>1</sup>  Z. Q. Sun (孙泽群)<sup>1,71</sup>  Z. T. Sun (孙振田)<sup>55</sup>  H. Tabaharizato<sup>1</sup>  N. T. Tagsinsit<sup>67</sup>   
C. J. Tang (唐昌建)<sup>60</sup>  G. Y. Tang (唐光毅)<sup>1</sup>  J. Tang (唐健)<sup>66</sup>  J. J. Tang (唐嘉骏)<sup>78,65</sup>   
L. F. Tang (唐林发)<sup>43</sup>  Y. A. Tang (唐迎澳)<sup>84</sup>  Z. H. Tang (唐梓涵)<sup>1,71</sup>  L. Y. Tao (陶璐燕)<sup>80</sup>  M. Tat<sup>76</sup>   
J. X. Teng (滕佳秀)<sup>78,65</sup>  J. Y. Tian (田济源)<sup>78,65</sup>  W. H. Tian (田文辉)<sup>66</sup>  Y. Tian (田野)<sup>34</sup>   
Z. F. Tian (田喆飞)<sup>84</sup>  K. Yu. Todyshev<sup>4</sup>  I. Uman<sup>69B</sup>  E. van der Smagt<sup>3</sup>  B. Wang (王博)<sup>66</sup>   
B. Wang (王斌)<sup>1</sup>  B. Wang (王博)<sup>78,65</sup>  C. Wang (王程)<sup>42,k,l</sup>  C. Wang (王超)<sup>20</sup>  C. Wang (王聪)<sup>23</sup>   
D. Y. Wang (王大勇)<sup>51,h</sup>  F. K. Wang (王福凯)<sup>66</sup>  H. J. Wang (王泓鉴)<sup>42,k,l</sup>  H. R. Wang (王皓冉)<sup>86</sup>   
J. Wang (王杰)<sup>10</sup>  J. J. Wang (王家驹)<sup>84</sup>  J. P. Wang (王吉鹏)<sup>37</sup>  K. Wang (王科)<sup>1,65</sup>  L. L. Wang (王亮亮)<sup>1</sup>   
L. W. Wang (王璐怡)<sup>38</sup>  M. Wang (王萌)<sup>55</sup>  Mi Wang (王咪)<sup>78,65</sup>  N. Y. Wang (王南洋)<sup>71</sup>   
P. Wang (汪品)<sup>21</sup>  S. Wang (王石)<sup>42,k,l</sup>  S. Wang (王顺)<sup>64</sup>  T. Wang (王婷)<sup>12,g</sup>  W. Wang (王为)<sup>66</sup>   
W. P. Wang (王维平)<sup>39</sup>  X. F. Wang (王雄飞)<sup>42,k,l</sup>  X. L. Wang (王小龙)<sup>12,g</sup>  X. N. Wang (王新南)<sup>1,71</sup>   
X. Wang (王心)<sup>27,i</sup>  Y. Wang (王亦)<sup>1</sup>  Y. D. Wang (王雅迪)<sup>50</sup>  Y. F. Wang (王贻芳)<sup>1,9,71</sup>   
Y. H. Wang (王英豪)<sup>42,k,l</sup>  Y. J. Wang (王祎景)<sup>78,65</sup>  Y. L. Wang (王艺龙)<sup>20</sup>  Y. N. Wang (王亚男)<sup>50</sup>   
Y. N. Wang (王燕宁)<sup>84</sup>  Y. Q. Wang (王亚乾)<sup>18</sup>  Y. Wang (王义)<sup>68</sup>  Y. Wang (王源)<sup>18,34</sup>   
Z. Wang (王铮)<sup>1,65</sup>  Z. L. Wang (王治浪)<sup>2</sup>  Z. Q. Wang (王紫祺)<sup>12,g</sup>  Z. Y. Wang (王至勇)<sup>1,71</sup>   
Z. Wang (王旨)<sup>48</sup>  Ziyi Wang (王子一)<sup>71</sup>  D. Wei (魏栋)<sup>48</sup>  D. H. Wei (魏代会)<sup>14</sup>  D. J. Wei (魏德杰)<sup>73</sup>   
H. R. Wei<sup>48</sup>  F. Weidner<sup>75</sup>  H. R. Wen (文焕然)<sup>34</sup>  S. P. Wen (文硕频)<sup>1</sup>  U. Wiedner<sup>3</sup>  G. Wilkinson<sup>76</sup>   
J. F. Wu (吴金飞)<sup>1,9</sup>  L. H. Wu (伍灵慧)<sup>1</sup>  L. J. Wu (吴连近)<sup>20</sup>  L. J. Wu (武廉杰)<sup>20</sup>   
S. G. Wu (吴韶光)<sup>1,71</sup>  S. M. Wu (吴蜀明)<sup>71</sup>  X. W. Wu (吴潇)<sup>80</sup>  Z. Wu (吴智)<sup>1,65</sup>  H. L. Xia (夏惠亮)<sup>78,65</sup> 

L. Xia (夏磊)<sup>78,65</sup> B. H. Xiang (向本后)<sup>1,71</sup> D. Xiao (肖栋)<sup>42,k,l</sup> G. Y. Xiao (肖光延)<sup>47</sup> H. Xiao (肖浩)<sup>80</sup>  
 Y. L. Xiao (肖云龙)<sup>12,g</sup> Z. J. Xiao (肖振军)<sup>46</sup> C. Xie (谢陈)<sup>47</sup> K. J. Xie (谢凯吉)<sup>1,71</sup> Y. Xie (谢勇)<sup>55</sup>  
 Y. G. Xie (谢宇广)<sup>1,65</sup> Y. H. Xie (谢跃红)<sup>6</sup> Z. P. Xie (谢智鹏)<sup>78,65</sup> T. Y. Xing (邢天宇)<sup>1,71</sup>  
 D. B. Xiong (熊东波)<sup>1</sup> G. F. Xu (许国发)<sup>1</sup> H. Y. Xu (许皓月)<sup>2</sup> Q. J. Xu (徐庆君)<sup>17</sup> Q. N. Xu<sup>32</sup>  
 T. D. Xu (徐腾达)<sup>80</sup> X. P. Xu (徐新平)<sup>61</sup> Y. Xu (徐月)<sup>12,g</sup> Y. C. Xu (胥英超)<sup>86</sup> Z. S. Xu (许昭棠)<sup>71</sup>  
 F. Yan (严芳)<sup>24</sup> L. Yan (严亮)<sup>12,g</sup> W. B. Yan (鄢文标)<sup>78,65</sup> W. C. Yan (闫文成)<sup>89</sup> W. H. Yan (闫文昊)<sup>6</sup>  
 W. P. Yan (闫文鹏)<sup>20</sup> X. Q. Yan (严薛强)<sup>12,g</sup> Y. Y. Yan<sup>67</sup> H. J. Yang (杨海军)<sup>57,f</sup> H. L. Yang (杨昊霖)<sup>38</sup>  
 H. X. Yang (杨洪勋)<sup>1</sup> J. H. Yang (杨君辉)<sup>47</sup> R. J. Yang (杨润佳)<sup>20</sup> X. Y. Yang (杨晓艺)<sup>73</sup>  
 Y. Yang (杨莹)<sup>12,g</sup> Y. G. Yang (杨雅鸽)<sup>56</sup> Y. H. Yang (杨耘赫)<sup>48</sup> Y. M. Yang (杨一鸣)<sup>89</sup>  
 Y. Q. Yang (杨永强)<sup>10</sup> Y. Z. Yang (杨颖喆)<sup>20</sup> Y. H. Yang (杨友华)<sup>47</sup> Z. Y. Yang<sup>80</sup>  
 W. J. Yao (姚文静)<sup>6</sup> Z. P. Yao (姚志鹏)<sup>55</sup> M. Ye (叶梅)<sup>1,65</sup> M. H. Ye (叶铭汉)<sup>9</sup> Z. J. Ye (叶子健)<sup>62,i</sup>  
 K. Yi (易凯)<sup>46</sup> J. H. Yin (殷俊昊)<sup>48</sup> Q. Q. Yin (尹琪钦)<sup>48</sup> Z. Y. You (尤郑昀)<sup>66</sup> B. X. Yu (俞伯祥)<sup>1,65,71</sup>  
 C. X. Yu (喻纯旭)<sup>48</sup> G. Yu<sup>13</sup> J. S. Yu (俞洁晟)<sup>27,i</sup> L. W. Yu (喻丽雯)<sup>12,g</sup> T. Yu (于涛)<sup>80</sup>  
 X. D. Yu (余旭东)<sup>51,h</sup> Y. C. Yu (郁烨淳)<sup>89</sup> Y. C. Yu (于勇超)<sup>42</sup> C. Z. Yuan (苑长征)<sup>1,71</sup>  
 H. Yuan (袁昊)<sup>1,71</sup> J. Yuan (袁菁)<sup>38</sup> J. Yuan (袁杰)<sup>50</sup> L. Yuan (袁丽)<sup>2</sup> M. K. Yuan (袁铭宽)<sup>12,g</sup>  
 S. H. Yuan (袁升豪)<sup>80</sup> Y. Yuan (袁野)<sup>1,71</sup> C. X. Yue (岳崇兴)<sup>43</sup> Ying Yue (岳颖)<sup>20</sup> A. A. Zafar<sup>81</sup>  
 F. R. Zeng (曾凡蕊)<sup>55</sup> S. H. Zeng<sup>70</sup> X. Zeng (曾鑫)<sup>12,g</sup> Y. J. Zeng (曾溢嘉)<sup>1,71</sup> Y. J. Zeng (曾宇杰)<sup>66</sup>  
 Y. C. Zhai (翟云聪)<sup>55</sup> Y. H. Zhan (詹永华)<sup>66</sup> B. L. Zhang (张伯伦)<sup>1,71</sup> B. X. Zhang (张丙新)<sup>1</sup>  
 D. H. Zhang (张丹昊)<sup>48</sup> G. Y. Zhang (张广义)<sup>20</sup> G. Y. Zhang (张耕源)<sup>1,71</sup> H. Zhang (张豪)<sup>78,65</sup>  
 H. C. Zhang (张航畅)<sup>1,65,71</sup> H. H. Zhang (张宏浩)<sup>66</sup> H. L. Zhang (张瀚林)<sup>48</sup> H. Q. Zhang (张华桥)<sup>1,65,71</sup>  
 H. R. Zhang (张浩然)<sup>78,65</sup> H. Y. Zhang (章红宇)<sup>1,65</sup> H. Zhang (张晗)<sup>89</sup> J. Zhang (张晋)<sup>66</sup>  
 J. J. Zhang (张进军)<sup>58</sup> J. L. Zhang (张杰磊)<sup>21</sup> J. Q. Zhang (张敬庆)<sup>46</sup> J. S. Zhang (张家声)<sup>12,g</sup>  
 J. W. Zhang (张家文)<sup>1,65,71</sup> J. X. Zhang (张景旭)<sup>42,k,l</sup> J. Y. Zhang (张建勇)<sup>1</sup> J. Z. Zhang (张景芝)<sup>1,71</sup>  
 J. Y. Zhang<sup>49</sup> J. Zhang (张进)<sup>53</sup> J. Y. Zhang (张纪元)<sup>12,g</sup> L. M. Zhang (张黎明)<sup>68</sup> L. Zhang (张雷)<sup>47</sup>  
 N. Zhang<sup>38</sup> P. Zhang (张鹏)<sup>1,9</sup> Q. Zhang (张强)<sup>20</sup> Q. Y. Zhang (张秋岩)<sup>38</sup> Q. Z. Zhang (张铨正)<sup>71</sup>  
 R. Y. Zhang (张若愚)<sup>42,k,l</sup> S. H. Zhang (张水涵)<sup>1,71</sup> S. N. Zhang<sup>76</sup> S. L. Zhang (张书磊)<sup>27,i</sup>  
 X. M. Zhang (张晓梅)<sup>1</sup> X. Y. Zhang (张学尧)<sup>55</sup> Y. T. Zhang (张亚腾)<sup>89</sup> Y. H. Zhang (张银鸿)<sup>1,65</sup>  
 Y. P. Zhang (张越鹏)<sup>78,65</sup> Y. Zhang (张瑶)<sup>1</sup> Y. Zhang (张宇)<sup>80</sup> Y. Zhang (张雨)<sup>66</sup> Z. Zhang (张哲)<sup>34</sup>  
 Z. D. Zhang (张正德)<sup>1</sup> Z. H. Zhang (张泽恒)<sup>1</sup> Z. L. Zhang (张兆领)<sup>38</sup> Z. X. Zhang (张泽祥)<sup>20</sup>  
 Z. Y. Zhang (张振宇)<sup>84</sup> Z. Z. Zhang<sup>20</sup> Z. K. Zhang (张兆轲)<sup>61</sup> Z. L. Zhang (张志龙)<sup>50</sup>  
 Z. Y. Zhang (张子扬)<sup>48</sup> G. Zhao (张子羽)<sup>1</sup> J.-P. Zhao (赵光)<sup>71</sup> J. Y. Zhao (赵俊鹏)<sup>1,71</sup>  
 J. Z. Zhao (赵静宜)<sup>1,65</sup> L. Zhao (赵京周)<sup>1</sup> L. Zhao (赵玲)<sup>78,65</sup> M. G. Zhao (赵雷)<sup>48</sup> R. P. Zhao (赵明刚)<sup>71</sup>  
 S. J. Zhao (赵若平)<sup>89</sup> Y. B. Zhao (赵豫斌)<sup>1,65</sup> Y. L. Zhao (赵艳琳)<sup>61</sup> Y. P. Zhao (赵祎鹏)<sup>50</sup>  
 Y. X. Zhao (赵宇翔)<sup>34,71</sup> Z. G. Zhao (赵政国)<sup>78,65</sup> A. Zhemchugov<sup>40,a</sup> B. Zheng (郑波)<sup>80</sup>  
 B. M. Zheng (郑变敏)<sup>38</sup> J. P. Zheng (郑建平)<sup>1,65</sup> W. J. Zheng (郑文静)<sup>1,71</sup> W. Q. Zheng (郑稳倩)<sup>10</sup>  
 X. R. Zheng (郑心如)<sup>20</sup> Y. H. Zheng (郑阳恒)<sup>71,o</sup> B. Zhong (钟彬)<sup>46</sup> C. Zhong (钟翠)<sup>20</sup>  
 X. Zhong (钟鑫)<sup>45</sup> H. Zhou (周航)<sup>39,55,n</sup> J. Q. Zhou (周嘉奇)<sup>38</sup> S. Zhou (周帅)<sup>6</sup> X. Zhou (周详)<sup>84</sup>  
 X. K. Zhou (周晓康)<sup>6</sup> X. R. Zhou (周小蓉)<sup>78,65</sup> X. Y. Zhou (周兴玉)<sup>43</sup> Y. X. Zhou (周亦雄)<sup>86</sup>  
 Y. Z. Zhou (周祎卓)<sup>20</sup> A. N. Zhu (朱傲男)<sup>71</sup> J. Zhu (朱江)<sup>48</sup> K. Zhu (朱凯)<sup>1</sup> K. J. Zhu (朱科军)<sup>1,65,71</sup>  
 K. S. Zhu (朱康帅)<sup>12,g</sup> L. X. Zhu (朱琳萱)<sup>71</sup> L. Zhu (朱林)<sup>20</sup> S. H. Zhu (朱世海)<sup>77</sup> T. J. Zhu (朱腾蛟)<sup>12,g</sup>  
 W. D. Zhu (朱稳定)<sup>12,g</sup> W. J. Zhu (朱文静)<sup>1</sup> W. Z. Zhu (朱文卓)<sup>20</sup> Y. C. Zhu (朱莹春)<sup>78,65</sup>  
 Z. A. Zhu (朱自安)<sup>1,71</sup> X. Y. Zhuang (庄新宇)<sup>48</sup> M. Zhuge (诸葛铭)<sup>55</sup> J. H. Zou (邹佳恒)<sup>1</sup> J. Zu<sup>34</sup>

(BESIII Collaboration)<sup>†</sup><sup>1</sup>Institute of High Energy Physics, Beijing 100049, China<sup>2</sup>Beihang University, Beijing 100191, China<sup>3</sup>Bochum Ruhr-University, D-44780 Bochum, Germany<sup>4</sup>Budker Institute of Nuclear Physics SB RAS (BINP), Novosibirsk 630090, Russia<sup>5</sup>Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA

- <sup>6</sup>Central China Normal University, Wuhan 430079, China  
<sup>7</sup>Central South University, Changsha 410083, China  
<sup>8</sup>Chengdu University of Technology, Chengdu 610059, China  
<sup>9</sup>China Center of Advanced Science and Technology, Beijing 100190, China  
<sup>10</sup>China University of Geosciences, Wuhan 430074, China  
<sup>11</sup>Chung-Ang University, Seoul, 06974, Republic of Korea  
<sup>12</sup>Fudan University, Shanghai 200433, China  
<sup>13</sup>GSI Helmholtzcentre for Heavy Ion Research GmbH, D-64291 Darmstadt, Germany  
<sup>14</sup>Guangxi Normal University, Guilin 541004, China  
<sup>15</sup>Guangxi University, Nanning 530004, China  
<sup>16</sup>Guangxi University of Science and Technology, Liuzhou 545006, China  
<sup>17</sup>Hangzhou Normal University, Hangzhou 310036, China  
<sup>18</sup>Hebei University, Baoding 071002, China  
<sup>19</sup>Helmholtz Institute Mainz, Staudinger Weg 18, D-55099 Mainz, Germany  
<sup>20</sup>Henan Normal University, Xinxiang 453007, China  
<sup>21</sup>Henan University, Kaifeng 475004, China  
<sup>22</sup>Henan University of Science and Technology, Luoyang 471003, China  
<sup>23</sup>Henan University of Technology, Zhengzhou 450001, China  
<sup>24</sup>Hengyang Normal University, Hengyang 421002, China  
<sup>25</sup>Huangshan College, Huangshan 245000, China  
<sup>26</sup>Hunan Normal University, Changsha 410081, China  
<sup>27</sup>Hunan University, Changsha 410082, China  
<sup>28</sup>Indian Institute of Technology Madras, Chennai 600036, India  
<sup>29</sup>Indiana University, Bloomington, Indiana 47405, USA  
<sup>30A</sup>INFN Laboratori Nazionali di Frascati, INFN Laboratori Nazionali di Frascati, I-00044, Frascati, Italy  
<sup>30B</sup>INFN Laboratori Nazionali di Frascati, INFN Sezione di Perugia, I-06100, Perugia, Italy  
<sup>30C</sup>INFN Laboratori Nazionali di Frascati, University of Perugia, I-06100, Perugia, Italy  
<sup>31A</sup>INFN Sezione di Ferrara, INFN Sezione di Ferrara, I-44122, Ferrara, Italy  
<sup>31B</sup>INFN Sezione di Ferrara, University of Ferrara, I-44122, Ferrara, Italy  
<sup>32</sup>Inner Mongolia University, Hohhot 010021, China  
<sup>33</sup>Institute of Business Administration, University Road, Karachi 75270, Pakistan  
<sup>34</sup>Institute of Modern Physics, Lanzhou 730000, China  
<sup>35</sup>Institute of Physics and Technology, Mongolian Academy of Sciences, Peace Avenue 54B, Ulaanbaatar 13330, Mongolia  
<sup>36</sup>Instituto de Alta Investigación, Universidad de Tarapacá, Casilla 7D, Arica 1000000, Chile  
<sup>37</sup>Jiangsu Ocean University, Lianyungang 222005, China  
<sup>38</sup>Jilin University, Changchun 130012, China  
<sup>39</sup>Johannes Gutenberg University of Mainz, Johann-Joachim-Becher-Weg 45, D-55099 Mainz, Germany  
<sup>40</sup>Joint Institute for Nuclear Research, 141980 Dubna, Moscow region, Russia  
<sup>41</sup>Justus-Liebig-Universität Giessen, II. Physikalisches Institut, Heinrich-Buff-Ring 16, D-35392 Giessen, Germany  
<sup>42</sup>Lanzhou University, Lanzhou 730000, China  
<sup>43</sup>Liaoning Normal University, Dalian 116029, China  
<sup>44</sup>Liaoning University, Shenyang 110036, China  
<sup>45</sup>Longyan University, Longyan 364000, China  
<sup>46</sup>Nanjing Normal University, Nanjing 210023, China  
<sup>47</sup>Nanjing University, Nanjing 210093, China  
<sup>48</sup>Nankai University, Tianjin 300071, China  
<sup>49</sup>National Centre for Nuclear Research, Warsaw 02-093, Poland  
<sup>50</sup>North China Electric Power University, Beijing 102206, China  
<sup>51</sup>Peking University, Beijing 100871, China  
<sup>52</sup>Qufu Normal University, Qufu 273165, China  
<sup>53</sup>Renmin University of China, Beijing 100872, China  
<sup>54</sup>Shandong Normal University, Jinan 250014, China  
<sup>55</sup>Shandong University, Jinan 250100, China  
<sup>56</sup>Shandong University of Technology, Zibo 255000, China  
<sup>57</sup>Shanghai Jiao Tong University, Shanghai 200240, China  
<sup>58</sup>Shanxi Normal University, Linfen 041004, China  
<sup>59</sup>Shanxi University, Taiyuan 030006, China  
<sup>60</sup>Sichuan University, Chengdu 610064, China  
<sup>61</sup>Soochow University, Suzhou 215006, China  
<sup>62</sup>South China Normal University, Guangzhou 510006, China  
<sup>63</sup>Southeast University, Nanjing 211100, China  
<sup>64</sup>Southwest University of Science and Technology, Mianyang 621010, China  
<sup>65</sup>State Key Laboratory of Particle Detection and Electronics, Beijing 100049, Hefei 230026, China  
<sup>66</sup>Sun Yat-Sen University, Guangzhou 510275, China  
<sup>67</sup>Suranaree University of Technology, University Avenue 111, Nakhon Ratchasima 30000, Thailand  
<sup>68</sup>Tsinghua University, Beijing 100084, China  
<sup>69A</sup>Turkish Accelerator Center Particle Factory Group, Istinye University, 34010, Istanbul, Turkey  
<sup>69B</sup>Turkish Accelerator Center Particle Factory Group, Near East University, Nicosia, North Cyprus, 99138, Mersin 10, Turkey

<sup>70</sup>University of Bristol, H H Wills Physics Laboratory, Tyndall Avenue, Bristol, BS8 1TL, UK<sup>71</sup>University of Chinese Academy of Sciences, Beijing 100049, China<sup>72</sup>University of Hawaii, Honolulu, Hawaii 96822, USA<sup>73</sup>University of Jinan, Jinan 250022, China<sup>74</sup>University of La Serena, Av. Raúl Bitrán 1305, La Serena, Chile<sup>75</sup>University of Muenster, Wilhelm-Klemm-Strasse 9, 48149 Muenster, Germany<sup>76</sup>University of Oxford, Keble Road, Oxford OX13RH, United Kingdom<sup>77</sup>University of Science and Technology Liaoning, Anshan 114051, China<sup>78</sup>University of Science and Technology of China, Hefei 230026, China<sup>79</sup>University of Silesia in Katowice, Institute of Physics, 75 Pulku Piechoty 1, 41-500 Chorzow, Poland<sup>80</sup>University of South China, Hengyang 421001, China<sup>81</sup>University of the Punjab, Lahore-54590, Pakistan<sup>82A</sup>University of Turin and INFN, University of Turin, I-10125, Turin, Italy<sup>82B</sup>University of Turin and INFN, University of Eastern Piedmont, I-15121, Alessandria, Italy<sup>82C</sup>University of Turin and INFN, INFN, I-10125, Turin, Italy<sup>83</sup>Uppsala University, Box 516, SE-75120 Uppsala, Sweden<sup>84</sup>Wuhan University, Wuhan 430072, China<sup>85</sup>Xi'an Jiaotong University, No.28 Xianning West Road, Xi'an, Shaanxi 710049, P.R. China<sup>86</sup>Yantai University, Yantai 264005, China<sup>87</sup>Yunnan University, Kunming 650500, China<sup>88</sup>Zhejiang University, Hangzhou 310027, China<sup>89</sup>Zhengzhou University, Zhengzhou 450001, China<sup>a</sup>Also at the Moscow Institute of Physics and Technology, Moscow 141700, Russia<sup>b</sup>Also at the Functional Electronics Laboratory, Tomsk State University, Tomsk, 634050, Russia<sup>c</sup>Also at the Novosibirsk State University, Novosibirsk, 630090, Russia<sup>d</sup>Also at the NRC "Kurchatov Institute", PNPI, 188300, Gatchina, Russia<sup>e</sup>Also at Goethe University Frankfurt, 60323 Frankfurt am Main, Germany<sup>f</sup>Also at Key Laboratory for Particle Physics, Astrophysics and Cosmology, Ministry of Education; Shanghai Key Laboratory for Particle Physics and Cosmology; Institute of Nuclear and Particle Physics, Shanghai 200240, China<sup>g</sup>Also at Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) and Institute of Modern Physics, Fudan University, Shanghai 200443, China<sup>h</sup>Also at State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, China<sup>i</sup>Also at School of Physics and Electronics, Hunan University, Changsha 410082, China<sup>j</sup>Also at Guangdong Provincial Key Laboratory of Nuclear Science, Institute of Quantum Matter, South China Normal University, Guangzhou 510006, China<sup>k</sup>Also at MOE Frontiers Science Center for Rare Isotopes, Lanzhou University, Lanzhou 730000, China<sup>l</sup>Also at Lanzhou Center for Theoretical Physics, Lanzhou University, Lanzhou 730000, China<sup>m</sup>Also at Ecole Polytechnique Federale de Lausanne (EPFL), CH-1015 Lausanne, Switzerland<sup>n</sup>Also at Helmholtz Institute Mainz, Staudinger Weg 18, D-55099 Mainz, Germany<sup>o</sup>Also at Hangzhou Institute for Advanced Study, University of Chinese Academy of Sciences, Hangzhou 310024, China<sup>p</sup>Also at Applied Nuclear Technology in Geosciences Key Laboratory of Sichuan Province, Chengdu University of Technology, Chengdu 610059, China

**Abstract:** Based on  $(10087 \pm 44) \times 10^6$   $J/\psi$  events collected with the BESIII detector, the  $J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$  and  $J/\psi \rightarrow \gamma \pi^0 \pi^0 \eta$  processes are studied. The  $X(2370)$  is observed in both the  $K_S^0 K_S^0 \pi^0$  and  $\pi^0 \pi^0 \eta$  invariant mass spectra, with statistical significances greater than  $14\sigma$  and  $20\sigma$ , respectively. By combining measurements from these processes with those from the previously reported  $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$  process, the mass and width of the  $X(2370)$  are determined to be  $2359_{-14}^{+13}$  MeV/ $c^2$  and  $170_{-29}^{+44}$  MeV, respectively. In addition, the decay  $X(2370) \rightarrow a_0(980)^0 \pi^0$  with  $a_0(980)^0 \rightarrow \pi^0 \eta$  is observed with a statistical significance exceeding  $9\sigma$ . The properties of the  $X(2370)$ —decay pattern similarities to that of  $\eta_c$ , are consistent with those of a pseudoscalar glueball.

**Keywords:** glueball,  $X(2370)$ , QCD

**DOI:** 10.1088/1674-1137/ae8acd **CSTR:** 32044.14.ChinesePhysicsC.50101001

Quantum Chromodynamics (QCD), a non-Abelian  $SU(3)$  gauge theory, predicts the existence of glueballs, which are uniquely formed by self-interacting gluons—the gauge bosons of the strong interactions. The search for glueballs in experiments is crucial for testing and developing QCD. Predictions of the  $0^{++}$  glueball mass from different lattice QCD (LQCD) groups range from 2.3 to 3.0 GeV/ $c^2$  [1–6]. However, although  $J/\psi$  radiative de-

cays are considered an ideal place for glueball searches [7–9], no  $0^{++}$  glueball candidates had been observed in  $J/\psi$  radiative decays in the mass range above 2.3 GeV/ $c^2$  until 2011.

In 2011, based on a sample of 0.2 billion  $J/\psi$  events collected with the BESIII detector, the  $X(2370)$  particle was first observed in the process  $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$  with a statistical significance greater than  $6.4\sigma$ . Its mass and

width were measured to be  $2376.3 \pm 8.7(\text{stat})_{-4.3}^{+3.2}(\text{syst})$  MeV/ $c^2$  and  $83 \pm 17(\text{stat})_{-6}^{+44}(\text{syst})$  MeV [10], respectively. The observation is particularly interesting because the measured mass is consistent with the range predicted by LQCD for the lightest pseudoscalar glueball. Moreover,  $J/\psi$  radiative decays are generally considered a favorable environment for glueball production, and a pseudoscalar glueball is expected to decay into channels such as  $\pi^+ \pi^- \eta'$  [10]. In subsequent studies, using 1.3 billion  $J/\psi$  events, the  $X(2370)$  was confirmed in the decays  $J/\psi \rightarrow \gamma K^+ K^- \eta'$  and  $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$  with a statistical significance of  $8.3\sigma$  [11]. Furthermore, using 10 billion  $J/\psi$  events, the spin-parity quantum numbers of the  $X(2370)$  were determined to be  $0^{++}$  with a significance greater than  $9.8\sigma$  in the  $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$  process [12]. Its mass and width were measured to be  $2395 \pm 11(\text{stat})_{-94}^{+26}(\text{syst})$  MeV/ $c^2$  and  $188_{-17}^{+18}(\text{stat})_{-33}^{+124}(\text{syst})$  MeV, respectively. The product branching fraction of  $J/\psi \rightarrow \gamma X(2370) \rightarrow \gamma f_0(980) \eta' \rightarrow \gamma K_S^0 K_S^0 \eta'$  was measured to be  $(1.31 \pm 0.22(\text{stat})_{-0.84}^{+2.85}(\text{syst})) \times 10^{-5}$ . The measured mass of the pseudoscalar  $X(2370)$  is within the range predicted by LQCD for the lightest pseudoscalar glueball [5, 6]. In addition to the glueball interpretation, other interpretations and phenomenological studies of the  $X(2370)$  have also been proposed [13–17]. To understand the nature of the  $X(2370)$ , more decay modes should be searched for and studied using 10 billion  $J/\psi$  decays collected at BESIII.

For glueball decay properties, no theoretical predictions are available, including from LQCD. The idea that glueball decays and charmonium decays exhibit qualitative similarities, since both proceed predominantly via gluons, was first proposed in Ref. [18]. In this picture, the decays of the  $0^{++}$  and  $2^{++}$  glueballs should be similar to those of  $\chi_{c0}$  and  $\chi_{c2}$ , respectively [18]. By analogy, the decays of a  $0^{++}$  glueball are expected to resemble those of the  $\eta_c$  [19–21], as illustrated in Fig. 1. As a conventional  $c\bar{c}$  charmonium state, the  $\eta_c$  can also decay electromagnetically via photons, although its hadronic decays via gluons dominate. Furthermore, since the  $0^{++}$  glueball and the  $\eta_c$  may have different masses, the resulting phase-space differences make a direct quantitative comparison of their decay properties difficult.

The  $KK\pi$  and  $\pi\pi\eta$  modes are two typical decay channels of the  $\eta_c$ . In addition, the decay  $\eta_c \rightarrow a_0(980)\pi$  has been observed, where the  $a_0(980)$  dominantly decays into  $\pi\eta$  [22]. Therefore, it is essential to search for corre-

sponding decay modes of the  $X(2370)$ , such as  $KK\pi$  and  $\pi\pi\eta$ , including the  $a_0(980)\pi$  intermediate state, in order to investigate whether it can be interpreted as a pseudoscalar glueball. In this work, the  $J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$  and  $J/\psi \rightarrow \gamma \pi^0 \pi^0 \eta$  processes are studied. These channels benefit from very low background contributions because the processes  $J/\psi \rightarrow \pi^0 K_S^0 K_S^0 \pi^0$  and  $J/\psi \rightarrow \pi^0 \pi^0 \pi^0 \eta$  are forbidden by  $C$ -parity conservation and the exchange symmetry of identical particles ( $K_S^0 K_S^0$  and  $\pi^0 \pi^0$ ). This study is based on  $(10087 \pm 44) \times 10^6$   $J/\psi$  events [23] collected with the BESIII detector [24].

A detailed description of the design and performance of the BESIII detector can be found in Ref. [24]. Monte Carlo (MC) simulated data samples produced with a GEANT4-based [25] package, which includes the geometric description of the BESIII detector [26] and the detector response, are used to determine detection efficiencies and estimate backgrounds. The simulation models the beam energy spread and initial-state radiation in  $e^+e^-$  annihilations with the generator KKMC [27, 28]. The inclusive MC sample includes both the production of the  $J/\psi$  resonance and the continuum processes incorporated in KKMC [27, 28]. All particle decays are modeled with EVTGEN [29, 30] using branching fractions either taken from the Particle Data Group (PDG) [22], when available, or otherwise estimated with LUNDCHARM [31, 32]. Final-state radiation from charged particles is incorporated using the PHOTOS package [33]. MC simulation samples of the processes  $J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$  and  $J/\psi \rightarrow \gamma \pi^0 \pi^0 \eta$ , including the subsequent decays  $K_S^0 \rightarrow \pi^+ \pi^-$ ,  $\pi^0 \rightarrow \gamma\gamma$ , and  $\eta \rightarrow \gamma\gamma$ , are generated uniformly in phase-space (PHSP).

Charged tracks detected in the multi-layer drift chamber (MDC) are required to be within a polar-angle ( $\theta$ ) range of  $|\cos\theta| < 0.93$ , where  $\theta$  is defined with respect to the  $z$ -axis, the symmetry axis of the MDC. No requirement is imposed on the distance of closest approach to the interaction point for charged tracks since all charged tracks originate from  $K_S^0$  decays. All charged tracks are assumed to be pions. To reconstruct  $K_S^0$  candidates, the tracks of each  $\pi^+ \pi^-$  pair are fitted to a secondary vertex. To suppress background events, all  $K_S^0$  candidates are required to satisfy  $|M_{\pi^+ \pi^-} - m_{K_S^0}| < 11$  MeV/ $c^2$ , where  $m_{K_S^0}$  is the nominal mass of the  $K_S^0$  from the PDG [22]. To further suppress background, the decay length of the  $K_S^0$  candidate is required to be greater than twice the vertex resolution away from the interaction point. With these selections, the miscombination rate between pions from different  $K_S^0$  candidates is 0.2%. The reconstructed  $K_S^0$  candidates are used as input for the subsequent kinematic fit.

Photon candidates are identified using isolated showers in the electromagnetic calorimeter (EMC). The deposited energy of each shower must be greater than 25 MeV in the barrel region ( $|\cos\theta| < 0.80$ ) and greater than





Fig. 1. (color online) Decays of (a) the  $0^{++}$  glueball and (b) the  $\eta_c$ .

50 MeV in the end cap region ( $0.86 < |\cos\theta| < 0.92$ ). To exclude showers that originate from charged tracks, the opening angle between the shower position and charged tracks extrapolated to the EMC must be greater than 10 degrees. The EMC time of a shower is required to be within 700 ns after the  $e^+e^-$  annihilation (the time determined by charged tracks), or, in the case of no charged tracks, within 500 ns before and after the time of the highest-energy shower.

For the  $J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$  process, each event candidate is required to have at least two  $K_S^0$  candidates, at least three photons, and no additional charged tracks from the interaction point. A four-constraint (4C) kinematic fit under the  $J/\psi \rightarrow \gamma\gamma\gamma K_S^0 K_S^0$  hypothesis is performed by enforcing energy-momentum conservation. If there are multiple  $\gamma\gamma K_S^0 K_S^0$  combinations, the one with the smallest  $\chi_{4C}^2$  is chosen. The resulting  $\chi_{4C}^2$  is required to be less than 40. To reconstruct the  $\pi^0$  candidate, a five-constraint (5C) kinematic fit is performed to further constrain the invariant mass of the two photons to  $m_{\pi^0}$ , where  $m_{\pi^0}$  is the nominal mass of the  $\pi^0$  [22]. Among the three  $\gamma\gamma$  combinations, the one with the smallest  $\chi_{5C}^2$  is chosen as the  $\pi^0$  candidate. The two photons from the  $\pi^0$  candidate are required to satisfy  $|M_{\gamma\gamma} - m_{\pi^0}| < 22 \text{ MeV}/c^2$ , with each photon having an energy greater than 100 MeV. To suppress miscombinations of the radiative photon ( $\gamma_{\text{rad}}$ ) and the photons from the  $\pi^0$  candidate ( $\gamma_{\pi^0}$ ), events with  $|M_{\gamma_{\text{rad}}\gamma_{\pi^0}} - m_{\pi^0}| < 22 \text{ MeV}/c^2$  are rejected. These photon-related requirements reduce the miscombination rate between the  $\gamma_{\text{rad}}$  and  $\gamma_{\pi^0}$  to 0.1%. Events with  $|M_{\gamma_{\text{rad}}\gamma_{\pi^0}} - m_{\eta}| < 25 \text{ MeV}/c^2$  or  $|M_{\gamma_{\text{rad}}\pi^0} - m_{\omega}| < 40 \text{ MeV}/c^2$  are rejected to suppress background containing an  $\eta$  or  $\omega$ , where  $m_{\eta}$  and  $m_{\omega}$  are the nominal masses of the  $\eta$  and  $\omega$  [22], respectively. After applying the above selection criteria, a clear signal of the  $X(2370)$  around  $2.3 \text{ GeV}/c^2$ , along with the  $\eta_c$  peak around  $2.9 \text{ GeV}/c^2$ , is observed in the  $K_S^0 K_S^0 \pi^0$  invariant-mass spectrum using the momenta obtained from the 5C kinematic fit, as shown in Fig. 2(a), Fig. 2(b), and Fig. 2(c). These two-dimensional distributions indicate several possible intermediate processes, for example,  $a_0(980)^0 \pi^0$  and  $K_0^*(1430) K_S^0 \rightarrow K_S^0 K_S^0 \pi^0$ .

An inclusive MC sample of 10 billion  $J/\psi$  decays is used to study potential background contributions. After applying the above selection criteria to the inclusive MC sample, no peaking background is found in the  $K_S^0 K_S^0 \pi^0$  invariant-mass spectrum. The contributions of non- $K_S^0$  and non- $\pi^0$  background processes are found to be negligible in the data by fitting the corresponding  $M_{\pi^+\pi^-}$  and  $M_{\gamma\gamma}$  distributions.

An unbinned maximum-likelihood fit is performed on the  $K_S^0 K_S^0 \pi^0$  invariant-mass spectrum between 1.95 and  $2.75 \text{ GeV}/c^2$ . The  $X(2370)$  signal is described by an efficiency-corrected and PHSP-weighted Breit-Wigner (BW) function convoluted with a detector resolution function.



**Fig. 2.** (color online) Two-dimensional distributions of (a)  $M_{K_S^0 K_S^0}$  versus  $M_{K_S^0 K_S^0 \pi^0}$  and (b)  $M_{K_S^0 \pi^0}$  versus  $M_{K_S^0 K_S^0 \pi^0}$ . (c) The  $K_S^0 K_S^0 \pi^0$  invariant mass spectrum. The solid red histogram shows PHSP MC with arbitrary normalization. (d) The fit results within the range  $[1.95, 2.75] \text{ GeV}/c^2$ . The dashed red line is the  $X(2370)$  signal component, the dashed green line represents the remaining processes described by a third-order polynomial function, and the blue line is the total fit result.

A third-order polynomial function is used to describe the remaining contributions. From the fit shown in Fig. 2(d), the mass and width of the  $X(2370)$  are measured to be  $2311 \pm 4(\text{stat}) \text{ MeV}/c^2$  and  $177 \pm 20(\text{stat}) \text{ MeV}$ , respectively. The significance of the  $X(2370)$  is greater than  $14\sigma$ , as determined from the changes in the log-likelihood value and degrees of freedom between fits with and without the  $X(2370)$  signal hypothesis, and includes systematic variations.

For the  $J/\psi \rightarrow \gamma\pi^0\pi^0\eta$  process, each event candidate is required to have no charged tracks and at least seven photons. To identify photon pairs originating from  $\pi^0$  decays, a one-constraint (1C) kinematic fit is performed by constraining the invariant mass of each photon pair to  $m_{\pi^0}$ . At least two pairs with  $\chi_{1C}^2 < 10$  are required and selected as  $\pi^0$  candidates, which are then used as inputs to the subsequent kinematic fit. A six-constraint (6C) kinematic fit under the  $J/\psi \rightarrow \gamma\gamma\pi^0\pi^0$  hypothesis is performed by enforcing energy-momentum conservation and constraining the invariant masses of the two  $\pi^0$  candidates to  $m_{\pi^0}$ . If there are multiple  $\gamma\gamma\pi^0\pi^0$  combinations, the one with the smallest  $\chi_{6C}^2$  is chosen. The resulting  $\chi_{6C}^2$  is required to be less than 40. To reconstruct the  $\eta$  candidate, a seven-constraint (7C) kinematic fit is performed to further constrain the invariant mass of the two photons to  $m_{\eta}$ . Among the three  $\gamma\gamma$  combinations, the one with the


smallest  $\chi_{7C}^2$  is chosen. To suppress multi-photon backgrounds, the  $\chi_{4C}^2$  from the kinematic fit under the  $J/\psi \rightarrow 7\gamma$  hypothesis is required to be less than that from the kinematic fits under the  $J/\psi \rightarrow 8\gamma$  and  $J/\psi \rightarrow 9\gamma$  hypotheses. The two photons from the  $\eta$  candidate are required to satisfy  $|M_{\gamma\gamma} - m_\eta| < 27 \text{ MeV}/c^2$ . To suppress miscombination between the radiative photon and other selected photons, events satisfying any of the following requirements are rejected:  $|M_{\gamma_{\text{rad}}\gamma_{\pi^0}} - m_{\pi^0}| < 20 \text{ MeV}/c^2$ ,  $|M_{\gamma_{\text{rad}}\gamma_\eta} - m_{\pi^0}| < 20 \text{ MeV}/c^2$ ,  $|M_{\gamma_{\text{rad}}\gamma_{\pi^0}} - m_\eta| < 30 \text{ MeV}/c^2$ , and  $|M_{\gamma_{\text{rad}}\gamma_\eta} - m_\eta| < 50 \text{ MeV}/c^2$ , where  $\gamma_\eta$  denotes the photons from the  $\eta$  candidate. With these requirements, the miscombination rate among the seven photons is 1.4%. To suppress background containing an  $\omega$ , events with  $|M_{\gamma_{\text{rad}}\pi^0} - m_\omega| < 40 \text{ MeV}/c^2$  are rejected. After applying the above selection criteria, clear  $X(2370)$  and  $\eta_c$  peaks are observed in the  $\pi^0\pi^0\eta$  invariant mass spectrum using the momenta obtained from the 7C kinematic fit, as shown in Fig. 3(a), Fig. 3(b), and Fig. 3(c). These two-dimensional distributions also indicate several possible intermediate processes, such as  $f_0(1500)\eta$  and  $a_0(980)^0\pi^0 \rightarrow \pi^0\pi^0\eta$ .

After applying the above selection criteria, no peaking background is found in the  $\pi^0\pi^0\eta$  invariant mass spectrum of the inclusive MC sample. The background contri-


bution from non- $\eta$  processes is estimated by fitting the corresponding  $M_{\gamma\gamma}$  distribution in data, yielding a non- $\eta$  background fraction of 3.5%. A validation using  $\eta$  mass sideband events indicates that the non- $\eta$  background processes produce no peaking structure. Therefore, it is described by a polynomial function in the subsequent fit.

A strategy analogous to that described for the  $J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$  process is employed in the fit to the  $\pi^0\pi^0\eta$  invariant mass spectrum between 2.0 and 2.7  $\text{GeV}/c^2$ , as shown in Fig. 3(d). The mass and width of the  $X(2370)$  are measured to be  $2366 \pm 2(\text{stat}) \text{ MeV}/c^2$  and  $127 \pm 9(\text{stat}) \text{ MeV}$ , respectively. The significance of the  $X(2370)$  is greater than  $20\sigma$ , including systematic variations.

Using the selection criteria for the  $J/\psi \rightarrow \gamma \pi^0 \pi^0 \eta$  process described above, an  $a_0(980)^0$  candidate is reconstructed from the  $\eta$  and the lower-energy  $\pi^0$ . To suppress contributions from the decay  $X(2370) \rightarrow f_0(1500)\eta$ , events with  $|M_{\pi^0\pi^0} - 1.5 \text{ GeV}/c^2| < 0.15 \text{ GeV}/c^2$  are rejected. The resulting mass spectrum of the  $\eta$  and the lower-energy  $\pi^0$  is shown in Fig. 4(a). An additional requirement of  $|M_{\pi^0\eta} - 0.98 \text{ GeV}/c^2| < 0.05 \text{ GeV}/c^2$  is applied to select the  $a_0(980)^0$  signal region. After applying these requirements, the  $\pi^0\pi^0\eta$  mass spectrum, shown in Fig. 4(b), exhibits clear  $X(2370)$  and  $\eta_c$  peaks. The  $\pi^0\pi^0\eta$  mass spectrum between 2.00 and 2.70  $\text{GeV}/c^2$  is fitted using a strategy



**Fig. 3.** (color online) The two-dimensional distributions of (a)  $M_{\pi^0\pi^0}$  versus  $M_{\pi^0\pi^0\eta}$  and (b)  $M_{\pi^0\eta}$  versus  $M_{\pi^0\pi^0\eta}$ . (c) The  $\pi^0\pi^0\eta$  invariant-mass spectrum. The solid red histogram corresponds to the PHSP MC sample with arbitrary normalization. (d) The fit results in the range  $[2.00, 2.70] \text{ GeV}/c^2$ . The dashed red line represents the  $X(2370)$  signal component, the dashed green line represents the remaining processes described by a third-order polynomial function, and the blue line indicates the total fit result.



**Fig. 4.** (color online) (a) The invariant mass spectrum of the  $\eta$  and the lower-energy  $\pi^0$ . (b) The  $\pi^0\pi^0\eta$  invariant mass spectrum for selected events in the  $a_0(980)^0$  signal region. The PHSP MC simulations are shown with arbitrary normalization. (c) and (d) show the fit results in the range  $[2.00, 2.70] \text{ GeV}/c^2$  for the  $a_0(980)^0$  signal region and the  $a_0(980)^0$  sideband region, respectively. The dashed green line represents the contribution of the remaining processes, described by a second-order polynomial plus an exponential function.

analogous to that described in the previous paragraph, as shown in Fig. 4(c). In particular, the PHSP parametrization in this fit includes an intermediate  $a_0(980)^0$  state, which is parametrized using dispersion integrals [34]. The remaining processes are described by the sum of a polynomial and an exponential function. The mass and width of the  $X(2370)$  are measured to be  $2358 \pm 5(\text{stat}) \text{ MeV}/c^2$  and  $180 \pm 20(\text{stat}) \text{ MeV}$ , respectively. The significance of the  $X(2370)$  is greater than  $9\sigma$ , including systematic variations. In contrast, there is no evidence for an  $X(2370)$  signal in the  $a_0(980)^0$  mass sideband region of  $0.20 \text{ GeV}/c^2 < |M_{\pi^0\eta} - 0.98 \text{ GeV}/c^2| < 0.25 \text{ GeV}/c^2$ , as shown in Fig. 4(d).

Systematic uncertainties associated with the invariant-mass spectrum fits are evaluated for the three decay modes and arise primarily from the PHSP parametrization, the background model, and the treatment of an additional resonance. For each source, alternative fits are performed, and the largest deviations of the fitted mass and width of the  $X(2370)$  from the nominal fit are taken as the corresponding systematic uncertainties. The uncertainty associated with the PHSP parametrization is studied by including an additional Blatt-Weisskopf barrier factor and by considering PHSP variations among different intermediate resonances. For the  $K_S^0 K_S^0 \pi^0$  mode, the alternative intermediate resonances  $K_0^*(1430)$  and  $a_0(980)^0$  are considered. For the  $\pi^0 \pi^0 \eta$  mode, the alternative intermediate resonances  $a_0(980)^0$  and  $f_0(1500)$  are considered. The  $a_0(980)^0$  is described using both the Flatté [34, 35] and dispersion-integral parametrizations [34], while all other intermediate resonances are described by BW functions with masses and widths fixed to the PDG values [22]. The uncertainty related to the background model is estimated by varying the fit range and adopting alternative background parametrizations. The uncertainty related to an additional resonance is estimated by varying its treatment. For the  $K_S^0 K_S^0 \pi^0$  mode, the uncertainty includes a contribution from the additional resonance  $X(2260)$ . This inclusion is motivated by the study of the charged channel  $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$  [36], where the  $X(2260)$  is observed in the  $K_S^0 K^\pm \pi^\mp$  mass spectrum with a significance greater than  $10\sigma$ , and its fitted mass and width are  $2256 \pm 3(\text{stat}) \text{ MeV}/c^2$  and  $74 \pm 10(\text{stat}) \text{ MeV}$ , respectively.


The significance of the  $X(2260)$  in the  $K_S^0 K_S^0 \pi^0$  mass spectrum is  $2.3\sigma$ , where the low significance is due to the available statistics being more than ten times smaller here than in the charged channel. The uncertainty related to the impact of the  $X(2260)$  is estimated by incorporating its contribution to the  $K_S^0 K_S^0 \pi^0$  mass spectrum fit, and the mass and width of the  $X(2260)$  are fixed to the values measured in the  $K_S^0 K^\pm \pi^\mp$  mass spectrum. For the  $\pi^0 \pi^0 \eta$  mode, a potential additional resonance can be added in the mass spectrum fit with a statistical significance of  $4\sigma$ , and its mass and width are fitted to be  $2142 \pm 7(\text{stat}) \text{ MeV}/c^2$  and  $186 \pm 48(\text{stat}) \text{ MeV}$ , respectively. The uncertainty related to the impact of this resonance is estimated by incorporating its contribution to the  $\pi^0 \pi^0 \eta$  mass spectrum fit. For the  $a_0(980)^0 \pi^0$  mode, since no evidence of the additional resonance is observed, the uncertainty related to the additional resonance is not included. All sources of systematic uncertainties are summarized in Table 1.

To further improve the measurement precision, the  $X(2370)$  masses and widths measured in different decay modes are combined following the procedure recommended by the PDG [22]. All  $X(2370)$  measurements included in our combination are based on the data sample of 10 billion  $J/\psi$  events collected with the BESIII detector, including the studies of the  $J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$  and  $J/\psi \rightarrow \gamma \pi^0 \pi^0 \eta$  processes presented in this paper, as well as the previously reported results for the  $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$  process [12]. Considering the statistical and systematic uncertainties, the combined mass and width of the  $X(2370)$  are  $2359^{+13}_{-14} \text{ MeV}/c^2$  and  $170^{+44}_{-29} \text{ MeV}$ , respectively. The uncertainty of the combined mass is scaled by a factor of 1.48 because  $\chi^2/N_{\text{dof}} = 2.20$ , following the procedure recommended by the PDG [22]. Figure 5 presents a comparison of the  $X(2370)$  resonance parameters measured in the individual channels and the combined result discussed in this paper. The resulting  $\chi^2/N_{\text{dof}}$  values, shown in each figure and defined in the PDG [22], indicate reasonable consistency among the input measurements.

In summary, using a sample of  $(10087 \pm 44) \times 10^6$   $J/\psi$  events [23] collected with the BESIII detector, the  $J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$  and  $J/\psi \rightarrow \gamma \pi^0 \pi^0 \eta$  processes are invest-

**Table 1.** Systematic uncertainties on the mass (in  $\text{MeV}/c^2$ ), width (in  $\text{MeV}$ ), and significance (in  $\sigma$ ) of the  $X(2370)$  in different processes, denoted by  $\Delta M$ ,  $\Delta \Gamma$ , and  $N_\sigma$ , respectively.

| Source               | $J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$ |                 |            | $J/\psi \rightarrow \gamma \pi^0 \pi^0 \eta$ |                 |            | $J/\psi \rightarrow \gamma \pi^0 \pi^0 \eta$ ( $a_0(980)^0$ signal region) |                 |            |
|----------------------|-----------------------------------------------|-----------------|------------|----------------------------------------------|-----------------|------------|----------------------------------------------------------------------------|-----------------|------------|
|                      | $\Delta M$                                    | $\Delta \Gamma$ | $N_\sigma$ | $\Delta M$                                   | $\Delta \Gamma$ | $N_\sigma$ | $\Delta M$                                                                 | $\Delta \Gamma$ | $N_\sigma$ |
| PHSP parametrization | $\pm 7$                                       | $\pm 2$         | $> 18$     | $\pm 3$                                      | $\pm 1$         | $> 20$     | $\pm 2$                                                                    | $\pm 3$         | $> 12$     |
| Background model     | $\pm 10$                                      | $\pm 37$        | $> 14$     | $\pm 9$                                      | $\pm 24$        | $> 20$     | $\pm 9$                                                                    | $\pm 58$        | $> 9$      |
| Additional resonance | $\pm 22$                                      | $\pm 62$        | $> 14$     | $\pm 4$                                      | $\pm 56$        | $> 20$     | —                                                                          | —               | —          |
| Total                | $\pm 25$                                      | $\pm 72$        | $> 14$     | $\pm 10$                                     | $\pm 61$        | $> 20$     | $\pm 9$                                                                    | $\pm 58$        | $> 9$      |



**Fig. 5.** (color online) Combined (a) mass and (b) width of the  $X(2370)$ . The bars on each black dot indicate the total uncertainty; the horizontal red shaded bands indicate the statistical uncertainties; and the vertical blue lines and blue shaded bands indicate the combined results and their total uncertainties.

igated to search for the  $X(2370)$ . The  $X(2370)$  is observed in the  $K_S^0 K_S^0 \pi^0$  and  $\pi^0 \pi^0 \eta$  invariant mass spectra with statistical significances greater than  $14\sigma$  and  $20\sigma$ , respectively. By further selecting the  $a_0(980)^0$  mass region in the latter process, the decay  $X(2370) \rightarrow a_0(980)^0 \pi^0$  with  $a_0(980)^0 \rightarrow \pi^0 \eta$  is observed with a statistical significance greater than  $9\sigma$ . In the above decay modes, the mass and width of the  $X(2370)$  are determined to be

$$\begin{aligned} X(2370) &\rightarrow K_S^0 K_S^0 \pi^0 : \\ M_{X(2370)} &= 2311 \pm 4(\text{stat}) \pm 25(\text{syst}) \text{ MeV}/c^2, \\ \Gamma_{X(2370)} &= 177 \pm 20(\text{stat}) \pm 72(\text{syst}) \text{ MeV}; \end{aligned} \quad (1)$$

$$\begin{aligned} X(2370) &\rightarrow \pi^0 \pi^0 \eta : \\ M_{X(2370)} &= 2366 \pm 2(\text{stat}) \pm 10(\text{syst}) \text{ MeV}/c^2, \\ \Gamma_{X(2370)} &= 127 \pm 9(\text{stat}) \pm 61(\text{syst}) \text{ MeV}; \end{aligned} \quad (2)$$

$$\begin{aligned} X(2370) &\rightarrow a_0(980)^0 \pi^0 : \\ M_{X(2370)} &= 2358 \pm 5(\text{stat}) \pm 9(\text{syst}) \text{ MeV}/c^2, \\ \Gamma_{X(2370)} &= 180 \pm 20(\text{stat}) \pm 58(\text{syst}) \text{ MeV}. \end{aligned} \quad (3)$$

Based on the results of the first two decay modes observed in this paper, together with the previously reported results in  $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$  [12], a combined determin-

ation of the  $X(2370)$  mass and width is performed, yielding

$$\begin{aligned} M_{X(2370)} &= 2359^{+13}_{-14} \text{ MeV}/c^2, \\ \Gamma_{X(2370)} &= 170^{+44}_{-29} \text{ MeV}. \end{aligned} \quad (4)$$

The combined mass of the  $X(2370)$  lies within the mass range predicted by LQCD calculations for the lightest pseudoscalar glueball [5, 6]. The observed decay modes of  $\pi^+ \pi^- \eta'$ ,  $K_S^0 K_S^0 \eta'$ ,  $K_S^0 K_S^0 \pi^0$ , and  $\pi^0 \pi^0 \eta$ , including  $a_0(980)^0 \pi^0$ , show similarities between the  $X(2370)$  and  $\eta_c$  decays. In addition, as shown in the scatter plot of  $M_{K_S^0 K_S^0 \eta}$  versus  $M_{K_S^0 K_S^0 \eta}$  from a previous study of the  $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$  channel [37], similar behavior is observed in the  $K_S^0 K_S^0 \eta$  decay mode. The event yields in both the  $X(2370)$  and  $\eta_c$  mass regions are enhanced in the  $f_0(1500)$  and  $f_0(1710)$  mass regions, whereas they are suppressed in the  $f_0(980)$  mass region. The properties of the  $X(2370)$ —decay pattern similarities to that of  $\eta_c$ , are consistent with those of a pseudoscalar glueball. To further identify the nature of the  $X(2370)$ , more studies are needed both experimentally and theoretically.

## ACKNOWLEDGMENTS

The BESIII Collaboration thanks the staff of BEPCII (<https://cstr.cn/31109.02.BEPC>) and the IHEP computing center for their strong support.

## References

- [1] G. S. Bali *et al.* (UKQCD Collaboration), *Phys. Lett. B* **309**, 378 (1993)
- [2] C. J. Morningstar and M. J. Peardon, *Phys. Rev. D* **60**, 034509 (1999)
- [3] Y. Chen, A. Alexandru, S. J. Dong, *et al.*, *Phys. Rev. D* **73**, 014516 (2006)
- [4] E. Gregory, A. Irving, B. Lucini *et al.*, *JHEP* **10**, 170 (2012)
- [5] L. C. Gui, J. M. Dong, Y. Chen, *et al.*, *Phys. Rev. D* **100**, 054511 (2019)
- [6] D. Vadacchino, PoS LATTICE2022, 236 (2023), arXiv: 2305.04869

- [7] L. Kopke and N. Wermes, *Phys. Rept.* **174**, 67 (1989)
- [8] M. B. Çakır and G. R. Farrar, *Phys. Rev. D* **50**, 3268 (1994)
- [9] F. E. Close, G. R. Farrar, and Z. Li, *Phys. Rev. D* **55**, 5749 (1997)
- [10] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **106**, 072002 (2011)
- [11] M. Ablikim *et al.* (BESIII Collaboration), *Eur. Phys. J. C* **80**, 746 (2020)
- [12] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **132**, 181901 (2024)
- [13] X. Sun, L. Y. Dai, S. Q. Kuang *et al.*, *Phys. Rev. D* **105**, 034010 (2022)
- [14] J. S. Yu, Z. F. Sun, X. Liu *et al.*, *Phys. Rev. D* **83**, 114007 (2011)
- [15] R. R. Dong, N. Su, H. X. Chen *et al.*, *Eur. Phys. J. C* **80**, 749 (2020)
- [16] Q. N. Wang, D. K. Lian, and W. Chen, *Phys. Rev. D* **112**, 034010 (2025)
- [17] N. Su and H. X. Chen, *Phys. Rev. D* **106**, 014023 (2022)
- [18] K. T. Chao, *Commun. Theor. Phys.* **24**, 373 (1995)
- [19] T. Huang, S. Jin, D. H. Zhang, *et al.*, *Phys. Lett. B* **380**, 189 (1996)
- [20] Y. Huang, S. Jin, and P. Zhang, *Int. J. Mod. Phys. A* **40**, 2530007 (2025)
- [21] C. Morningstar, PoS **LATTICE2024**, 004 (2025)
- [22] S. Navas *et al.* (Particle Data Group), *Phys. Rev. D* **110**, 030001 (2024)
- [23] M. Ablikim *et al.* (BESIII Collaboration), *Chin. Phys. C* **46**, 074001 (2022)
- [24] M. Ablikim *et al.* (BESIII Collaboration), *Nucl. Instrum. Meth. A* **614**, 345 (2010)
- [25] S. Agostinelli *et al.* (GEANT4 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003)
- [26] K. X. Huang, Z. J. Li, Z. Qian, *et al.*, *Nucl. Sci. Tech.* **33**, 142 (2022)
- [27] S. Jadach, B. F. L. Ward, and Z. Was, *Phys. Rev. D* **63**, 113009 (2001)
- [28] S. Jadach, B. F. L. Ward, and Z. Was, *Comput Phys Commun* **130**, 260 (2000)
- [29] D. J. Lange, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001)
- [30] R. G. Ping, *Chin. Phys. C* **32**, 599 (2008)
- [31] J. C. Chen, G. S. Huang, X. R. Qi *et al.*, *Phys. Rev. D* **62**, 034003 (2000)
- [32] R. L. Yang, R. G. Ping, and H. Chen, *Chin. Phys. Lett.* **31**, 061301 (2014)
- [33] E. Barberio and Z. Was, *Comput. Phys. Commun.* **79**, 291 (1994)
- [34] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **95**, 032002 (2017)
- [35] S. M. Flatte, *Phys. Lett. B* **63**, 224 (1976)
- [36] M. Ablikim *et al.* (BESIII Collaboration), *Observation of the  $X(2370)$  in  $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$* , publication in preparation.
- [37] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **115**, 091803 (2015)