The AME2026 Atomic Mass Evaluation *

(I). Evaluation of input data and adjustment procedures

W.J. Huang (黄文嘉)[1,2,†], R. Chakma[3], Meng Wang (王猛)[2,4,‡], F.G. Kondev[3], and S. Naimi[5]

[1] Advanced Energy Science and Technology Guangdong Laboratory, Huizhou 516007, China
[2] State Key Laboratory of Heavy Ion Science and Technology, Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China
[3] Physics Division, Argonne National Laboratory, Lemont, IL 60439, USA
[4] School of Nuclear Science and Technology, University of Chinese Academy of Sciences, Beijing 100049, China
[5] Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France

Abstract This is the first of two papers (Part I and Part II) presenting the results of the new Atomic Mass Evaluation, AME2026. It provides detailed information on the experimental input data used to derive the recommended values tabulated in Part II. It describes the evaluation philosophy and procedures used in the selection of specific nuclear reaction, decay, and mass-spectrometric data for a least-squares adjustment to determine the recommended mass values and their uncertainties. All input data, both accepted and rejected, are tabulated and compared with the adjusted values obtained from the least-squares adjustment. Differences from the previous AME2020 evaluation are discussed, and specific examples are presented for several nuclides that may be of interest to AME users.

Keywords: atomic mass evaluation, trends in the mass surface (TMS), atomic-mass tables, least-squares adjustment

DOI: 10.1088/1674-1137/aebcec **CSTR:** 32044.14.ChinesePhysicsC.50120002

1 Introduction

Atomic masses are fundamental properties of nuclei. They reflect the combined effects of the strong, electromagnetic, and weak interactions among nucleons, and provide essential input for studies of nuclear structure, nuclear astrophysics, weak-interaction physics, atomic physics, and a wide range of applications. A reliable and comprehensive table of atomic masses, derived from all available experimental information and evaluated within a consistent framework, is therefore indispensable to the research community. The Atomic Mass Evaluation (AME) was established in the 1950s to produce a reliable atomic mass table derived from experimental data. Successive AME mass tables have served the research community by providing the most reliable and comprehensive information on atomic masses [1]. The last complete evaluation of experimental atomic mass data AME2020 [2, 3] was published in March 2021. Since then, the experimental knowledge of atomic masses has continuously expanded, and a substantial body of data relevant to atomic masses has been published in the scientific literature.

This paper is the first of two AME2026 articles (Part I and Part II). It describes the evaluation procedures and the main AME2026 evaluation table, Table I, where all accepted and rejected experimental data are presented and compared with the adjusted values obtained from a least-squares adjustment. In Part II (the next article of this issue), the recommended mass values and their uncertainties are given, followed by tables and graphs of derived quantities and detailed bibliographical information. As in previous AME versions, AME2026 is accompanied by the NUBASE2026 evaluation [4], which provides information about the basic properties of the known ground and excited isomeric states that are used in the AME.

There is no strict literature cut-off date for the present AME2026 evaluation: all data available to the evaluators through the end of June 2026 were included. The final mass-adjustment calculations were performed on October 24, 2026. The present publication includes updated information from the previous AME tables, including data that were not used in the final adjustment for various reasons, e.g., data with excessively large uncertainties.

Received 28 October 2026

∗ This work has been undertaken with the endorsement of the IUPAP Commission on Symbols, Units, Nomenclature, Atomic Masses and Fundamental Constants (SUNAMCO).

†Email: huangwj@impcas.ac.cn

‡Email: wangm@impcas.ac.cn


Content from this work may be used under the terms of the Creative Commons Attribution 4.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Table A. Constants used in this work or resulting from the present evaluation.

1 u	=	$m(^{12}\mathrm{C})/12$	=	atomic mass unit				
1 u	=	1 660 539.068 92	±	0.000 52	$\times 10^{-33}$ kg	0.31	ppb	*a*
1 u	=	931 494.103 72	±	0.000 29	keV	0.31	ppb	*a*
1 MeV	=	1 073 544.100 83	±	0.000 33	nu	0.31	ppb	*a*
m_e	=	548 579.909 044 1	±	0.000 009 7	nu	0.02	ppb	*a*
	=	510 998.950 69	±	0.000 16	eV	0.31	ppb	*a*
m_p	=	1 007 276 466.597	±	0.008	nu	0.008	ppb	*b*
M_α	=	4 001 506 179.66	±	0.04	nu	0.010	ppb	*c*
$m_n - m_H$	=	839 884.00	±	0.47	nu	560	ppb	*c*
	=	782 346.99	±	0.44	eV			

a) adopted from the work of CODATA2022 [11].
b) derived from this work combined with m_e and the ionization energy [11, 12].
c) this work.

Remark: In the following text, several results of general interest are discussed. References to articles listed in Table I are generally not repeated in the text. However, when it is necessary to provide specific references, those are given using the NSR key-number style [5] (e.g., [2020Fu05]) and are listed at the end of Part II, under 'References used in the AME2026 and the NUBASE2026 evaluations'.

As in previous AME versions, all uncertainties are quoted as one standard deviation (1σ).

2 Units and energy recalibration

The unit used in the AME is the '*unified atomic mass unit*' u, which is defined as one-twelfth of the mass of one free atom of ^{12}C, at rest and in its atomic and nuclear ground states, $1\,\mathrm{u} = m(^{12}\mathrm{C})/12$. The 'unified mass unit' represents the unification of the physical and chemical units in 1960 following the proposal of Kohman, Mattauch, and Wapstra [6]. It is worth noting that the chemical mass-spectrometry community (e.g., biochemistry and polymer chemistry) widely uses the dalton unit (symbol Da, named after John Dalton*). The dalton is an alternative name for the same unit and is convenient for expressing molecular masses.

Until May 20, 2019, the mole, the SI (International System of Units) unit of amount of substance, was defined by fixing the molar mass of carbon-12 to be exactly $M(^{12}\mathrm{C}) = 0.012\mathrm{kg/mol}$. At that time the kilogram was defined as exactly the mass of the international prototype of the kilogram, an artifact made of a platinum-iridium alloy. In the current SI, the base units are defined by assigning exact numerical values to the Planck constant h, elementary charge e, Boltzmann constant k, and Avogadro constant N_A [7]. The mole is now defined by fixing the Avogadro constant N_A to $6.022\,140\,76 \times 10^{23}\ \mathrm{mol}^{-1}$, and $M(^{12}\mathrm{C})$ is no longer a defining constant. The kilogram is defined by fixing the Planck constant at $h = 6.626\,070\,15 \times 10^{-34}$ Js, where $\mathrm{J\,s} = \mathrm{kg \cdot m^2 \cdot s^{-1}}$; the metre and the second are defined by the fixed values of the speed of light in vacuum and the unperturbed ground-state hyperfine-transition frequency of ^{133}Cs [8].

Atom-recoil experiments determine the ratio $h/m(X)$ for an atom X. Because the Planck constant has an exact numerical value in the revised SI, such a measurement now yields the mass $m(X)$ in kilograms without an additional uncertainty from h. Using atom interferometry, the ratio $h/m(^{87}\mathrm{Rb})$ was measured with a relative uncertainty of 142 ppt [9], where the mass of ^{87}Rb in kilograms has the same relative uncertainty. The AME, however, determines relative atomic masses $A_\mathrm{r}(X) = m(X)/m_\mathrm{u}$, where $m_\mathrm{u} = m(^{12}\mathrm{C})/12$. The recoil result is not included in the AME, since the CODATA adjustment already incorporates this recoil measurement and the relative atomic masses from the AME to determine m_u. The inclusion of recoil results would therefore require a consistent treatment of the shared input data and their covariances. In general, the atomic mass of a particular nuclide can be obtained as an inertial mass from the motion of an ionized atom in an electromagnetic field, using other nuclides as references. The mass is then derived from a ratio of masses and expressed in 'u'. ^{12}C serves as the ultimate reference in the mass determination. Alternatively, the atomic mass of a particular nuclide can be determined by establishing an energy relation between the mass of interest and one or more well-known masses with a nuclear

*John Dalton (1766-1844) first speculated that elements combine in proportions following simple laws, and was the first to create a table of (very approximate) atomic weights.

Table B. Definition of Volt unit, and resulting mass-energy conversion constants.

	$2e/h$			u		
1983	483594.21	(1.34)	GHz/V	931501.2	(2.6)	keV
1983	483594	(exact)	GHz/V_{86}	931501.6	(0.3)	keV_{86}
1986	483597.67	(0.14)	GHz/V	931494.32	(0.28)	keV
1990	483597.9	(exact)	GHz/V_{90}	931493.86	(0.07)	keV_{90}
1999	483597.9	(exact)	GHz/V_{90}	931494.009	(0.007)	keV_{90}
2010	483597.9	(exact)	GHz/V_{90}	931494.0023	(0.0007)	keV_{90}
2012	483597.9	(exact)	GHz/V_{90}	931494.0038	(0.0004)	keV_{90}
2018	483597.8484	(exact)	GHz/V	931494.10242	(0.00028)	keV
2022	483597.8484	(exact)	GHz/V	931494.10372	(0.00029)	keV

decay or reaction measurement. This energy relation is then expressed in electron-volts (eV).

In earlier AME evaluations, the choice of volt for high-precision energy data required special care. From AME1993 through AME2016, the unit of the *international* volt (V) was not adopted, because at that time $2e/h$ had to be determined experimentally with a definite uncertainty for the *international* volt V. Instead, the *maintained* volt V_{90} was used, which was defined by $2e/h = 483597.9\,(\text{exact})\,\text{GHz}/V_{90}$ in 1990. The γ-ray energies determined in wavelength measurements can be expressed in eV_{90} without loss in precision, since the conversion coefficient is an exact quantity. These energies served as standards in other energy measurements, so the reaction and decay energies used in AME could be more accurately expressed in the *maintained* volt, as discussed by Cohen and Wapstra [10]. At that time, the precision of the conversion factor between mass units and *maintained* volt (V_{90}) was also higher than that between the former and *international* volt. Since the 2019 redefinition of the SI, the Josephson constant $K_J = 2e/h$ is exact in SI units, and the *maintained* volt is no longer needed for the present evaluation. In AME2026, all energy input values are expressed in SI electron-volts. In the most recent CODATA2022 evaluation [11], the relation between the old *maintained* volt and the *international* volt is given by the exact relation $V_{90} = 1.000\,000\,106\,66\,\text{V}$, with a difference of 106.66 ppb. This difference is much smaller than the uncertainties of almost all energy data used in the AME; the few input data for which this difference is relevant, such as the high-precision γ-ray wavelength results of [2006De21], have been recalibrated from eV_{90} to eV and marked accordingly in Table I. The detailed historical discussion and recalibrations were given in previous evaluations, in particular AME2020 [2].

Table A presents the relations between several constants of interest, obtained from the CODATA2022 evaluation [11]. The ratio of mass unit to electron-volt is also given. In addition, values for the masses of the proton, neutron, and alpha particle, as derived from the present evaluation, are also given, together with the mass difference between the neutron and the hydrogen atom.

The recommended $2e/h$ values and the mass-energy conversion factor are given in Table B.

For α-particle energies, Rytz took this change into account when updating his earlier evaluation of α-particle energies [13]. Proton energies were also recalibrated in AME2003. These recalibrated data are used in the present work and marked by 'Z' behind the reference keynumber in Table I. When it was not possible to do so, for example, when that position in the table was already used to indicate a remark, the same 'Z' symbol was added to the uncertainty value mentioned in the remark.

Until the end of the last century, the relative precision of the mass excess $M-A$ expressed in keV was, for several nuclides, poorer than that of the same quantity expressed in mass units. Due to the improved precision of the fundamental constants, the conversion between mass units and energy units is now sufficiently accurate that this difference is no longer significant. In the earlier mass tables (e.g., AME1993), values were given for the total binding energy, ZM_H+NM_n-M, mainly because its uncertainty in keV_{90} was then larger than that of the mass excess $M-A$. This consideration is no longer important, owing in particular to the improved precision of the neutron mass. Since AME2003, the tables instead give the binding energy per nucleon for pedagogical reasons, as this quantity is more useful for illustrating the Aston curve and the maximum stability near the 'iron peak', which is of importance in nuclear astrophysics.

3 Input data

As mentioned in the preceding section, atomic mass measurements are usually based on two types of experimental methods: the mass-spectrometric method (often

called the 'direct method'), where the inertial mass is determined from the trajectory of an ion — from its cyclotron frequency in a magnetic field or its time-of-flight; and the so-called 'indirect method', where a reaction or decay energy, i.e., the mass difference between two or more nuclides, is determined. In both cases, the experimental information normally provides a relation between atomic masses rather than an isolated absolute mass. For this reason, the AME data treatment makes every effort to enter the primary experimental quantities rather than only the derived absolute mass values reported in the literature. In this way, the masses can be recalibrated automatically for any future changes, and the original correlation information can be properly preserved.

In the present work, all available experimental data related to atomic masses, including both energy and mass-spectrometric data, are considered. The input data are extracted from the available literature, compiled in an appropriate format, and carefully evaluated. The data are compared with other results and with the trends in the mass surface (TMS) in the region, and calibrations are checked when necessary. As a result, some of the input data are accepted directly, or after appropriate corrections, and used in the atomic mass adjustment. Other data are not used due to large uncertainties or other reasons, as explained in the next section. The complete set of input data is given in Table I, and the data used in the mass adjustment are also presented graphically in Fig. 1.

3.1 Connection diagram

Figure 1 displays the connection diagram of the input data in the $(N-Z)$ versus $(N+Z)$ plane. Each dot represents a nuclide, and each line between two dots represents an experimentally established relation between atomic masses. This representation is straightforward for many data types, such as mass-doublet spectroscopy and decay-energy measurements. For example, Penning-trap mass measurements usually determine the ratio of two ion masses from the inverse ratio of their cyclotron frequencies in the trap. Such measurements can therefore be represented as a connection between two nuclides. For reaction energy measurements that involve several atomic masses, the diagram shows only the connection corresponding to the dominant mass difference between two nuclides. Most lines in Fig. 1 link two nuclides, with the exception of three mass-spectrometric 'triplet' measurements on Rb and Cs isotopes [1986Au02], which are best represented as linear combinations of masses of three isotopes with non-integer coefficients.

However, not all experimental results can be represented explicitly in Fig. 1. For example, results from Smith-type storage-ring, time-of-flight, and multi-reflection time-of-flight mass spectrometers involving two reference ions are often published as absolute mass values. These results are entered as absolute masses in Table I: ^{A}El–u, and they are shown as open squares in Fig. 1.

Atomic masses are extracted from the network represented in Fig. 1. Absolute mass values in the unit 'u' can be determined for nuclides that are ultimately connected to ^{12}C, the anchor point of the network. As can be seen from Fig. 1, the accepted data may determine the mass of a nuclide through several different routes; such a nuclide is called *primary*. Its mass value is then obtained by the least-squares method. In other cases, the mass of a nuclide is derived from only one connection to another nuclide; such a nuclide is called *secondary*. This classification is of importance for the calculation procedure.

The diagrams in Fig. 1 also show cases where the relation between two atomic masses is experimentally known, but neither nuclide is connected ultimately to ^{12}C, as happens for many of the heaviest nuclides. Since the AME policy is to include all available experimental results, estimated mass values based on trends in the mass surface (TMS) in the neighboring region are introduced in such cases. In addition, unmeasured nuclides located near the experimentally known mass region are included with the TMS estimates in order to complete the local representation of the mass surface. Estimates of unknown masses are further discussed in Section 5.

Excited isomeric states may be involved in experiments, and the related information is included in this work. Although some isomeric states are indicated in Fig. 1, the diagram itself is two-dimensional and cannot represent all the experimental information involving isomers. Instead, all relevant information is given in Table I.

3.2 Representation of the input data

The input data are given as linear equations in Table I. The equations are constructed from the original experimental quantities reported in the literature whenever possible. For example, Penning-trap mass spectrometers measure the cyclotron frequency of a charged ion in a magnetic field, which is compared to that of a well-known reference ion in order to determine the ratio of two masses. These mass ratios are converted to linear combinations of masses of electrically neutral atoms without any loss of accuracy. The procedure for converting frequency ratios to linear equations can be found in Appendix A. In some publications, such linear equations are also reported directly [2005Ra34]. Results from nuclear reaction or decay energy measurements are likewise entered as mass-difference relations.

When an experimental result cannot be readily expressed as a simple equation, the absolute mass value is given in Table I as a difference from the mass unit: ^{A}El-u, usually accompanied by a comment explaining the ori-




Figure 1: Diagrams of the input-data connections used in the present evaluation, shown in panels (a)–(j).

For *primary data* (those used in the least-squares analysis):

□ absolute mass-doublet nuclide (or nuclide connected by a unique secondary relative mass-doublet to a remote reference nuclide);

○ other primary nuclide;

⊡ ⊙ primary nuclide with relevant isomer;

// mass-spectrometric connection;

——— other primary reaction or decay connection.

Primary connections are drawn with two different thicknesses. Thicker lines represent the highest-precision data in the given mass region

(limits: 1 keV for $A < 36$,
2 keV for $A = 36$ to 165 and
3 keV for $A > 165$).

For *secondary data* (cases where masses are known from one type of data and are therefore not checked by a different connection):

■ secondary experimental nuclide determined from mass spectrometry;

• secondary experimental nuclide determined by a reaction or a decay;

○ nuclide for which mass is estimated from trends in the Mass Surface (TMS);

– – – – – connection to a secondary nuclide. Note that an experimental connection may exist between two estimated TMS nuclides when neither of them is connected to the network of primaries.




Figure 1 (b). Diagram of the input-data connections used in the present evaluation — continued.

ginal measurement. Such comments are not provided for results from time-of-flight and storage-ring mass spectrometry, where the mass of interest is obtained through complex calibrations involving many reference masses and is published as an absolute mass value.

In Table I, values of the equations and the corresponding uncertainties are given in 'μu' for results from mass spectrometry and in 'keV' for energy measurements.

3.3 Main types of input data

A total of 1490 new experimental data entries are included in AME2026 relative to AME2020. Among them, 1023 data entries are from mass spectrometry measurements, and 467 from energy measurements. Modern input data for the AME are dominated by direct mass spectrometry, while reaction and decay energy measurements remain essential in several mass regions, especially near the driplines and in decay chains connected to newly identified nuclides. The following paragraphs summarize the main types of measurements and how they are treated in the evaluation.

Updated values for the masses of π^+ and π^- are included from the latest Particle Data Group publication [14]. These two mesons are included because they are involved in some atomic mass experiments, although they are not considered 'nuclides' themselves.

3.3.1 High-precision mass spectrometry near β-stability

The most precise mass values are generally obtained with Penning traps, which determine the cyclotron-frequency ratio of an ion of interest to a reference ion in a uniform magnetic field B. For stable or long-lived nuclides, such measurements can reach ultrahigh precision and provide an important *backbone* for the evaluation network. In the AME, the reported frequency ratios are converted to differences between neutral atomic masses,




Figure 1 (c). Diagram of the input-data connections used in the present evaluation — continued.

with corrections for the missing electrons and their binding energies. High-precision measurements near stability are provided, for example, by dedicated Penning-trap setups such as LIONTRAP, PENTATRAP, and FSU-MIT-TRAP.

For measurements at the highest precision that involve molecules, molecular binding energies may also need to be included in the data treatment. As described, for example, in Ref. [2021Fi.1], even the rotational energy of H_2^+ has to be taken into account. For most molecules used in mass experiments, the binding energy correction is typically only a few parts in 10^{10}, and its uncertainty limits the neutral atomic mass only at the level of a few parts in 10^{12}. For measurements with relative precision no better than about 10^{-9} (100 eV/100 u), molecular binding-energy corrections can usually be neglected without loss of accuracy. In cases where the precision is better than 10^{-10} (10 eV/100 u), e.g., at MIT [1995Di08] and FSU [2015My03], it is necessary to take into account the molecular binding energy. More details on binding-energy corrections can be found in Appendix A.

3.3.2 Mass spectrometry far from β-stability

The region of the nuclear chart for which experimental masses are known has expanded substantially through the development of radioactive ion beam facilities and modern mass spectrometers. As first emphasized in AME2012 [30], the masses of short-lived nuclides, previously derived mainly from Q_β end-point measurements, are now determined predominantly by mass spectrometry. The increased accuracy and precision of direct mass measurements have significantly refined the atomic mass surface and, consequently, our understanding of nuclear interactions.




Figure 1 (d). Diagram of the input-data connections used in the present evaluation — continued.

In the past, several types of mass spectrometry provided some important results for radioactive nuclides, but are no longer used. These include classical double-focusing mass spectrometry, Smith-type radio-frequency mass spectrometry MITRAL at CERN, and cyclotron time-of-flight mass spectrometry at SARA-Grenoble and at CSS2-GANIL, all of which were described in AME2016 and will not be discussed here.

Currently, mass-spectrometric results are obtained using four main methods: Penning-trap mass spectrometry and multi-reflection time-of-flight (MR-TOF) mass spectrometry at low-energy radioactive ion beam facilities, and magnetic-rigidity time-of-flight ($B\rho$-TOF) and storage-ring based isochronous (IMS) or Schottky mass spectrometry at high-energy facilities.

Penning trap spectrometers: Currently, Penning traps remain the most accurate direct mass spectrometers for radioactive nuclides that can be delivered to the trap with sufficient yield. Examples of Penning-trap facilities that have provided important input data on short-lived nuclides in the recent AME evaluation include: ISOLTRAP-CERN, CPT-Argonne, JYFLTRAP-Jyväskylä, LEBIT-East-Lansing, SHIPTRAP-Darmstadt, TITAN-Vancouver, and TRIGA-TRAP-Mainz. For short-lived nuclides with low production rates, the Phase-Imaging Ion-Cyclotron-Resonance (PI-ICR) method [2013El01] is now widely used because of its high efficiency and resolving power. For example, the isomeric state in ^{104}Nb with an excitation energy of about 10 keV, requiring a resolving power of approximately 10^7, was identified and measured by both CPT [2018Or.A] and JYFLTRAP [2024Hu16]. PI-ICR can




Figure 1 (e). Diagram of the input-data connections used in the present evaluation — continued.

be applied to nuclides with half-lives down to a few tens of milliseconds, provided the production yield and preparation time are favorable. Thanks to the high resolving power of Pi-Icr, many low-lying isomeric states can be clearly identified. The conventional Time-of-Flight Ion-Cyclotron-Resonance (ToF-Icr) method also remains an important technique for determining the cyclotron frequency from ion motion in the trap.

For Penning-trap measurements, the linear functions are constructed using the frequency ratios reported in the literature. When inconsistencies are identified in the reported values, the authors are contacted to clarify the issues. One such example involves the results from [2017Er01]: an erratum was published following discussions with the authors.

Multi-Reflection Time-of-Flight spectrometers: The Multi-Reflection Time-of-Flight spectrometer (Mr-Tof) was initially developed for beam purification for Isoltrap and was subsequently used as an independent mass spectrometer. This method began providing mass results for short-lived nuclides in 2013, and results from Isolde-Cern, Ribf-Riken, Frs-Gsi, and Titan-Triumf are included in the current evaluation. Related devices are also operated, commissioned, or under development at other facilities, including Ganil, Jyfl, Raon, Frib, Argonne, Notre Dame, and IMP-Lanzhou.

The Mr-Tof mass measurement is based on time-of-flight: ions are reflected back and forth in a static electric field to extend the flight path and thereby improve the mass resolving power. With this method, a resolving power of 10^5 is routinely achieved within approxim-




Figure 1 (f). Diagram of the input-data connections used in the present evaluation — continued.

ately 10 milliseconds, and the typical relative precision is 10^{-7}. Careful optimization of the ion-mirror voltages using stable ions enabled the ZeroDegree Mr-Tof (ZD Mr-Tof) system to achieve a resolving power above 10^6 within a total time of flight of 12.5 ms [15]. Reaching such performance is more challenging for short-lived nuclides because of their limited yields and lifetimes, but remains feasible owing to the short measurement cycle of Mr-Tof.

For mass determination with Mr-Tof, the general relation between mass-to-charge ratio (m/q) and time-of-flight t is [2013Wi06]:

$$t = \alpha\sqrt{m/q} + \beta, \tag{1}$$

where α and β are parameters determined by the experimental setup and calibration procedure.

At Riken, Gsi, and Triumf, the parameter β can be determined independently; thus, only one reference nuclide is needed for mass calibration. In this way, the ratio of time-of-flight between the ion of interest and the reference ion is used to extract the linear equation, as is the case for Penning-trap measurements. However, at Isolde-Cern, two reference masses are used to determine the mass of interest via the so-called C_{tof} method. When using this method, we express the linear equation in terms of absolute mass.

Magnetic-Rigidity Time-of-Flight spectrometers: Projectile fragmentation or in-flight fission reactions are powerful tools for producing exotic nuclides far from stability. In magnetic-rigidity time-of-flight ($B\rho$-ToF) mass spectrometry, the masses of nearly undecelerated




Figure 1 (g). Diagram of the input-data connections used in the present evaluation — continued.

fragments produced in a thin target bombarded by heavy ions can be determined from a combination of magnetic deflection and time-of-flight measurements. Nuclides spanning an extended region in A/Z and Z are analyzed simultaneously, and even very short-lived (down to $1\,\mu$s) ions can be measured. In this way, mass values with relative precisions of 3×10^{-6} to 5×10^{-5} can be obtained for a large number of neutron-rich nuclides with mass number up to $A=115$. The limited resolving power, around 10^4, and cross-contamination can cause significant systematic shifts in the measured masses.

The main evaluation issues for $B\rho$-ToF data are calibration, extrapolation away from calibrant nuclides, and possible isomeric mixtures. Because the calibration is often empirical, newly available high-precision mass values from other methods can sometimes improve the calibration of earlier measurements. In the AME, such results are examined with particular attention to the calibration function, the calibrant set, and possible systematic effects. The MSU group re-evaluated their earlier experimental data [2015Me01, 2015Me08, 2016Me07] by incorporating recently published high-precision Penning-trap mass values [2018Le03, 2018Mo14] as additional calibration points, thereby reducing the systematic uncertainty and expanding the number of nuclides for which masses were determined [2020Me06].

Storage-ring Mass Spectrometry: Storage-ring mass spectrometry determines masses from the revolution frequencies or revolution times of stored ions. Since beam cooling is not required in the isochronous mode, the IMS method is suitable for mass measurements of nuclei with half-lives as short as tens of microseconds.




Figure 1 (h). Diagram of the input-data connections used in the present evaluation — continued.

The first implementation of this type was at GSI-ESR in Darmstadt, and a second was developed at the Cooler Storage Ring for Experiments (CSRE) at IMP-LANZHOU. At the Rare-RI Ring (R3) at RIKEN RIBF, the IMS has been realized with the individual-ion injection of rare nuclei [2022Li20]. At CSRE, the recently developed $B\rho$-defined IMS technique measures both the revolution time and the velocity of each ion. This technique improves the efficiency, sensitivity, and mass accuracy compared to conventional IMS over a broad range of mass-to-charge ratios [2022Wa39].

In classical Schottky mass spectrometry, the revolution frequencies of cooled stored ions are measured nondestructively. This method can provide very high precision, but the time required for beam cooling and observation generally limits its application to nuclides with relatively long half-lives. One advantage of Schottky detection is that many stored ions can be observed simultaneously, which is particularly useful when the nuclides of interest are produced together with a large number of contaminants. A recent experiment at GSI-ESR used a combined Schottky and isochronous mass-spectrometry technique to determine high-precision masses of neutron-rich Pt and Au nuclei near $N = 126$ [2026Fe.1]. Storage-ring techniques continue to provide important input data in regions that are difficult to access with low-energy mass spectrometers.

3.3.3 Energy measurement

Although mass spectrometry now dominates the field of mass measurements, reaction and decay-energy measurements remain indispensable in discovering new particle-unbound nuclides and determining their masses. Such measurements provide mass differences through Q values and separation energies, and are often the only source of information for particle-unbound and newly




Figure 1 (i). Diagram of the input-data connections used in the present evaluation — continued.

synthesized nuclides at the limits of stability. In Table I, these data are entered as mass-difference relations involving the parent, daughter, and emitted particles.

For very short-lived unbound nuclides, the lifetime may be too short for the nucleus to acquire atomic electrons and form a neutral atom (a process that typically takes approximately 10^{-14} s). Nevertheless, we express the result as an atomic mass so that it can be treated consistently with other nuclides. Mass information in such cases may come from complete-kinematics measurements of decay products, invariant-mass spectra, direct proton or neutron emission spectroscopy, or β-delayed particle emission studies. For example, the mass of the unbound nuclide ^{16}Be was deduced from the invariant mass of the ^{14}Be+n+n decay products [2024Mo02]. It is experimentally challenging to study these unbound nuclides far from stability, and usually only very few events can be observed. Frequently, theoretical input is required to interpret the experimental data and extract mass information. In heavier neutron-deficient regions, decay spectroscopy remains an important source of mass information. Characteristic α-decay chains can be used to identify new nuclides and provide Q_α values from which their masses are deduced, as in recent studies of ^{160}Os and ^{156}W [2023Br10, 2024Ya04].

3.4 Corrections of the input data

In most cases, the values and uncertainties quoted in the original publication are accepted as input to the AME. Corrections are made only when the published result is not the quantity needed for the adjustment, or when additional uncertainty is required to represent an unresolved experimental ambiguity. Such changes are documented in remarks accompanying the data in Table I. The main cases are summarized below.

3.4.1 Decay energies and level assignments

For an α decay, the quantity used in the mass adjustment is usually the ground-state-to-ground-state decay energy, unless a transition involving an excited state is explicitly required. In several heavy nuclides, however, the observed α line cannot be assigned unambiguously to a ground-state transition. When the daughter excitation energy is known or can be estimated, the corresponding correction is applied to the decay energy, and the estimated quantity is flagged with "#". When no reliable




Figure 1 (j). Diagram of the input-data connections used in the present evaluation — continued.

correction can be made, an additional 50 keV is added in quadrature to the quoted uncertainty. This accounts for possible unobserved low-lying excited states in the daughter or along the α-decay chain.

3.4.2 Isomeric mixtures

The resolving power of modern mass spectrometers is often sufficient to separate low-lying isomers, but this is not always the case. Some measurements determine the mass of an unresolved mixture of the ground state and one or more isomeric states. Since the ground-state mass is the primary goal of the AME, extracting it from an unresolved mixture requires information on the excitation energies and production rates of the isomers. The detailed prescription is given in Appendix B, and information about these changes is explicitly documented in remarks accompanying the data in Table I.

If the excitation energy E_1 of a single isomer and the relative production rates of the two states are known, the measured mixture is converted directly to a ground-state mass [2020Ca08]. If E_1 is known but the production ratio is not, all relative intensities are treated as equally probable. For a two-state mixture, this gives the ground-state mass $M_0 = M_{exp} - E_1/2$, with an uncertainty of $0.29E_1$ arising from the unknown admixture (see Appendix B). This policy was first defined and tested for the GSI-ESR cooled-beam cyclotron-frequency data, in consultation with the authors of the measurements. If more than one isomer may contribute, the same principle is extended to the allowed set of states as described in Appendix B.

A further complication arises when E_1 is not known. In such a case, we use an estimated value for E_1, taken from the NUBASE evaluation and flagged with '#'. When estimating the E_1 values, we first consider experimental information that may provide a lower limit, for example, whether one of two isomers decays to the other or whether γ rays of known energy are observed in the decay of the isomer. If not, we interpolate between E_1 values in neighboring nuclides that can be expected to have the same spin and configuration assignments (for odd A: isotones when Z is even, or isotopes when Z is odd). If such a comparison does not yield useful results, indications from theory are sometimes accepted, including upper limits on transition energies derived from the measured half-lives. Values estimated in this way are assigned conservative uncertainties, which are propagated into the final results.

For bare or highly charged ions, particular care is needed because internal conversion is suppressed and the lifetime of an isomer can be much longer than that of a corresponding neutral atom. This can affect whether an isomer survives long enough to contribute to a mass-spectrometric peak, as illustrated by the fully stripped $^{94m}Ru^{44+}$ case at CSRE [2017Ze02].

3.4.3 Particle energy versus decay energy

In the literature, the terms particle energy and decay energy are sometimes used inconsistently. Some authors report the particle energy E, while others quote the decay energy Q, and the distinction is not always clear. More precisely, the decay energy is the sum of the kinetic energies of the emitted particle and the recoiling daughter nuclide. In general, the α particle carries approximately 97–98% of the total kinetic energy released, with the recoiling daughter nuclide accounting for the remaining 2–3%. The literature contains many ambiguous cases, especially in proton-decay experiments, where Q_p and E_p are numerically closer to each other. In some cases, the ambiguity can be resolved through a careful inspection of the publication and discussions with the authors, but in many cases it remains unresolved.

Some authors derive a quantity which they call Q_α from the measured α-particle energy by applying both a recoil energy and an electron screening correction intended to obtain the energy release for bare nuclei. In the AME, the latter correction is removed to recover the Q_α value defined from neutral-atom masses. This should be distinguished from the electronic-energy terms retained in the treatment of high-precision α-decay energies (see Appendix C).

Finally, some measured α-particle energies are affected by coincidence summing between the α particle that feeds an excited level of the daughter nuclide and the conversion electrons that follow the decay of this level. This is sometimes apparent from the reported α spectra, since the width of the observed line is larger than that of other lines. In some cases, this summing can also produce spurious α peaks. When deriving the corresponding Q_α values, appropriate corrections are applied for the energy carried away by X-rays and γ rays. These corrections are mentioned in a remark added to the corresponding entry in Table I.

3.4.4 Recalibration of decay energies in implantation experiments

In decay-spectroscopy experiments, the parent nucleus is often implanted in a silicon detector, and the subsequent α or proton decay is measured in the same detector. In α-decay measurements, authors sometimes assume that only the α-particle energy is recorded, while in proton-decay measurements, both the proton and the heavy recoil are sometimes assumed to be detected simultaneously. Neither of these assumptions is generally correct. In practice, α particles and protons with energies of a few MeV are detected with nearly 100% efficiency. This is not the case for the heavy recoil, where only a fraction of the recoil energy contributes to the detected signal. If this partial recoil energy is not treated consistently for the calibration lines and for the decay of interest, the derived decay energy can be shifted by an amount comparable to the quoted uncertainty.

The correction procedure used in the AME was described in Ref. [16]. For a given decay, the deposited energy is written as the kinetic energy of the emitted particle plus the detected fraction of the recoil energy. The recoil detection efficiency is calculated from Lindhard's theory and has been checked against implantation measurements [17–19]. This correction is particularly relevant for superheavy and neutron-deficient nuclei measured after implantation in a double-sided silicon strip detector (DSSSD), where the calibrants and the nuclide of interest may have significantly different masses and thus different recoil energies. Some recent publications already take this recoil contribution into account. For example, the ^{256}No study of Ref. [2021Ke10] explicitly refers to this correction when treating implantation energies. When the authors' correction is documented and consistent with the AME procedure, the published value is used directly.

4 Determination of the adjusted mass values

The atomic mass evaluation differs from many other data evaluations [20] because nearly all mass determinations are relative rather than absolute measurements. Each experimental datum establishes a relation between two or more masses. The ensemble of these relations forms a highly interconnected network, as represented in Fig. 1. All the input data are expressed as linear equations in Table I.

To determine the mass values, we use a least-squares method in which each datum is weighted according to its quoted uncertainty. In the first step, we pre-average the identical and parallel data to reduce the size of the system of equations, and then separate primary data from secondary ones. The primary data are used in the least-squares calculations. Finally, we provide the recommended atomic masses (Part II, Table I), the nuclear reaction and separation energies (Part II, Table III), the adjusted values for the input data (Table I), the *influences* of data on the primary nuclides (Table I), the *influences* received by each primary nuclide (Part II, Table II), information on the inversion errors, the correlation coefficients (Part II, Table B), the χ^2 values, and the distribution of the v_i (see below), among others.

4.1 Compacting the set of data

4.1.1 Pre-averaging

Two or more measurements of the same physical quantity can be replaced without loss of information by their average value and uncertainty, thereby reducing the system of equations to be treated. Extend-

Table C. Largest discrepancies among pre-averaged data. The quantity n is the number of data in the pre-average, χ_n is the Birge ratio, and σ_e is the external error.

Item	n	χ_n	σ_e	Item	n	χ_n	σ_e
^{117}Ag^{m}(IT)^{117}Ag	2	2.48	0.37	^{15}N(p,n)^{15}O	2	2.03	1.28
^{101}Rb–u	2	2.47	44.36	^{58}Fe(t,p)^{60}Fe	4	2.03	7.38
^{186}W(n,γ)^{187}W	2	2.44	0.11	^{204}Tl(β^-)^{204}Pb	2	2.03	0.39
^{144}Ce(β^-)^{144}Pr	2	2.44	2.18	^{257}Rf^{m}(α)^{253}No	3	2.02	6.59
^{75}As(n,γ)^{76}As	2	2.32	0.17	^{106}Ag(ε)^{106}Pd	2	1.98	6.63
^{110}In(β^+)^{110}Cd	3	2.29	28.4	^{78}Se(n,γ)^{79}Se	3	1.96	0.28
^{40}Cl(β^-)^{40}Ar	2	2.21	76.10	^{31}Mg–O1.938	2	1.93	5.01
^{153}Gd(n,γ)^{154}Gd	2	2.16	0.39	^{145}Sm(ε)^{145}Pm	2	1.92	7.92
^{61}Ga–u	2	2.16	24.42	^{242}Pu(α)^{238}U	2	1.89	2.08
^{113}Cs(p)^{112}Xe	3	2.10	5.03	^{230}Th(d,t)^{229}Th	2	1.89	8.58

ing this procedure, we also consider *parallel* data: reaction data that give essentially the mass difference between the same two nuclides, except in rare cases where the precision is comparable to the uncertainty of the masses of the reaction particles. Examples include: ^{14}C(^{7}Li,^{7}Be)^{14}B and ^{14}C(^{14}C,^{14}N)^{14}B; or ^{22}Ne(t,^{3}He)^{22}F and ^{22}Ne(^{7}Li,^{7}Be)^{22}F. In such cases, one of the two links is replaced by an equivalent value for the other, and the pre-averaging proceeds as for the identical quantities.

If the n data to be pre-averaged are strongly conflicting, i.e., if the consistency factor (or Birge ratio) $\chi_n = \sqrt{\chi^2/(n-1)}$ obtained in the calculation of the pre-average is greater than 2.5, the (internal) error σ_i of the average is multiplied by the Birge ratio ($\sigma_e = \sigma_i \times \chi_n$). The quantity σ_e is often called the 'external error'. In some cases, an unweighted average is used, as indicated in the corresponding remarks in Table I. The pre-averaged data with the largest discrepancies are listed in Table C. Most of these items already appeared in a similar table in AME2020. Table D lists cases that appeared among the largest discrepancies in the previous evaluation but are no longer retained in Table C, together with the reason for each change.

In the present evaluation, we have replaced 3175 data with 1249 averages. Of these averages, 21.0% have values of χ_n beyond unity, 1.1% beyond two, and none beyond three, giving a very satisfactory overall distribution for our treatment. In a complex system such as this one, many values of χ_n exceeding 1 or 2 are expected to occur, and if the uncertainties were multiplied by χ_n in all these cases, the χ^2-test on the total adjustment would have been invalidated. This explains our choice of a relatively high threshold ($\chi_n^0 = 2.5$) for departing from the rule of 'internal error' of the weighted average, compared for example with $\chi_n^0 = 2$ recommended by Woods and Munster [21] or $\chi_n^0 = 1$ used in a different context by the Particle Data Group [14]. In addition to the automated pre-averaging, results from various capture ratios were combined manually in some β^+-decay cases. These cases are ^{109}Cd(ε)^{109}Ag (average of 3 entries), and ^{195}Au(ε)^{195}Pt (average of 5 entries); they are detailed in Table I. Five additional cases (^{139}Ce, ^{147}Tb, ^{152}Ho, ^{166}Yb, and ^{207}Bi) appear in our list but carry no weight in the adjustment and are labeled with 'U' in Table I.

4.1.2 Policies for treating parallel data

When averaging β- (or α-) decay energies derived from different branches observed in the same experiment, corresponding to transitions to or from different levels in the daughter nuclide, the uncertainty adopted for further evaluation is not the one resulting from the weighted average, but rather the smallest individual experimental uncertainty. In this way, we avoid artificially reducing the systematic component of the uncertainty. In some cases, however, when it is obvious that the uncertainty is dominated by poor statistics, we do not follow the above rule (e.g., ^{23}Al^{i}(p)^{22}Mg of [1997Bl04]). Some quantities have been reported more than once by the same experimental group. If the results are obtained by the same method in different experiments and are published in regular refereed journals, only the most recent one is usually retained in the calculation. There are two reasons for this policy. The first is that one might expect that the authors, who presumably consider their results to be of comparable quality, would have averaged them in their latest publication. The second is that averaging the two results would provide no control over the systematic component of the uncertainty. Our policy is different if the newer result is published in a secondary reference (non-refereed abstract, preprint, private communication, conference, thesis, or annual report). In such cases, the

Table D. Cases listed among the largest discrepancies of pre-averaged data in AME2020 but removed in the current work. The last column gives reasons for each change.

Item	n	χ_n	σ_e	Reason for change
^{220}Fr(α)^{216}At	3	1.70	3.34	new data available for ^{220}Fr
^{27}P^i(2p)^{25}Al	3	1.55	22.72	new data available for ^{27}P^i
^{177}Pt(α)^{173}Os	3	0.55	2.2	new data available for ^{177}Pt
^{181}Pb(α)^{177}Hg	2	0.07	0.6	re-analysis by the evaluator

older result is used in the calculation, except when the newer one is an update of the previous value. In some cases, when the systematic uncertainties are discussed explicitly, all the independent results are used.

4.1.3 Insignificant data

To preserve the interpretability of the final χ^2, the least-squares procedure excludes data whose weight is more than a factor of 10 smaller than that of other data, or of any combination of *all* other data determining the same quantity. Such data are labeled with 'U' in the list of input data; comparison with the output values allows us to check our judgment. Earlier, data were labeled 'U' if their weight was 10 times smaller than that of a *simple* combination of other data. Since AME1993, this criterion has been extended to data that weigh 10 times less than the combination of *all* other accepted data. Until AME2003, data labeled 'U' that had already appeared in a previous table were not reprinted, in order to reduce the size of the data table. This policy was changed starting with AME2012, and we now aim to include all relevant data, including those deemed insignificant. This is because conflicts may arise among recent results, and access to older data can help inform the evaluation of new measurements.

4.2 Separating secondary data

In Section 3, while examining the diagrams of connections in Fig. 1, we noted that the masses of *secondary* nuclides can be determined uniquely from a chain of secondary connections leading to a *primary* nuclide. Only the *primary* nuclides belong to the complex network of mutually constraining data for which the least-squares method is required. In terms of equations and parameters, if a parameter appears in only one equation of the system, that equation and parameter can be removed without affecting the fit result for all remaining data. Thus, we can redefine more precisely what was called *secondary* in Section 3: the parameter above is a *secondary* parameter (or mass) and its related equation is a *secondary* equation. This removal is applied recursively until no parameter appearing in only one equation remains. After the reduced set has been solved, the *secondary* equations can be used to determine the final values and uncertainties for the *secondary* parameters. The equations and parameters remaining after removing all secondaries are called *primary*.

Therefore, only the system of *primary* data is overdetermined, and only *primary* masses can be improved by the adjustment through all available information in the network. The values of *secondary* data are not adjusted by the least-squares procedure; they are used afterwards to propagate masses and uncertainties from the primary network, and they do not contribute to χ^2. The diagrams in Fig. 1 show that many *secondary* data exist. Thus, removing them considerably simplifies the system. More importantly, if a better value is later found for a *secondary* datum, the mass of the *secondary* nuclide can easily be improved. The procedure is more complicated for new *primary* data.

We define DEGREES for *secondary* nuclides and data, reflecting their distance along the chains connecting them to the network of primaries. The first secondary nuclide connected to a primary one is assigned degree 2, as is the connecting datum. Degree 1 is reserved for primary nuclides and data. Degrees for secondary nuclides and data range from 2 to 18. The heaviest nuclide, ^{295}Og, has the highest degree, 18. Its mass is determined through a long α-decay chain, although many of the intermediate values are estimates. In Table I, the degree of each datum is indicated in column 'Dg'. In the table of atomic masses (Part II, Table I), each *secondary* nuclide is marked with a label in column 'Orig.' to indicate the parent nuclide from which its mass value is determined.

To summarize, separating secondary nuclides and secondary data from primaries significantly reduces the size of the system that will be treated by the least-squares method (described below). After treatment of the primary data alone, the adjusted masses for primary nuclides can be easily combined with the secondary data to yield masses of secondary nuclides. This treatment not only decreases computational time (which is no longer a limiting factor), but also provides clearer insight into the relations between data and masses, because the secondary chains do not introduce additional overdetermined least-squares correlations.

Remark: The word *primary* used for these nuclides and for the data connecting them does not mean that they are more important than the others, but only that they are subject to the least-squares treatment above. The labels *primary* and *secondary* are not intrinsic properties of data or nuclides; they may change when new information becomes available.

4.3 Least-squares method

Each piece of data has a value q_i with an uncertainty dq_i (one standard deviation) and defines a relation between two, three, or four masses, whose unknown values are denoted m_μ. An overdetermined system relating Q data to M masses ($Q > M$) can be represented by a system of Q linear equations with M parameters:

$$\sum_{\mu=1}^{M} k_i^\mu m_\mu = q_i \pm dq_i \,, \tag{2}$$

e.g., for a nuclear reaction $A(a,b)B$ with q_i, the energy balance is written as:

$$m_{\rm A} + m_{\rm a} - m_{\rm b} - m_{\rm B} = q_i \pm dq_i \,. \tag{3}$$

Thus, $k_i^{\rm A} = +1$, $k_i^{\rm a} = +1$, $k_i^{\rm b} = -1$ and $k_i^{\rm B} = -1$.

In matrix notation, $\mathbf{K}$ being the (Q,M) matrix of coefficients, Eq. (2) is written: $\mathbf{K}|m\rangle = |q\rangle$. Most elements of the matrix $\mathbf{K}$ are zero: e.g., for $A(a,b)B$, Eq. (3) yields a row of $\mathbf{K}$ with only four nonzero elements.

We define the diagonal weight matrix $\mathbf{W}$ by its elements $w_i^i = 1/(dq_i)^2$. The solution of the least-squares method leads to a very simple construction:

$${}^{\rm t}\mathbf{KWK}|m\rangle = {}^{\rm t}\mathbf{KW}|q\rangle \,. \tag{4}$$

The NORMAL matrix $\mathbf{A} = {}^{\rm t}\mathbf{KWK}$ is a square matrix of order M, positive-definite, symmetric, and regular, and hence invertible [22]. Thus the vector $|\overline{m}\rangle$ for the adjusted masses is:

$$|\overline{m}\rangle = \mathbf{A^{-1}}\ {}^{\rm t}\mathbf{KW}|q\rangle \qquad \text{or} \qquad |\overline{m}\rangle = \mathbf{R}|q\rangle \,. \tag{5}$$

The rectangular (M,Q) matrix $\mathbf{R}$ is called the RESPONSE matrix.

The diagonal elements of $\mathbf{A^{-1}}$ are the variances of the adjusted masses, and the off-diagonal elements $(a^{-1})_\mu^\nu$ are the covariances between masses m_μ and m_ν.

Correlation coefficients for the most precise nuclides are given in Table B of Part II. Following the advice of B. N. Taylor, we now provide, on the website of the AMDC [23], the full list of correlation coefficients, allowing users to perform exact calculations of any combination of masses.

One of the most powerful tools in the least-squares calculation described above is the flow-of-information matrix, discovered in 1984 [24]. This matrix allows one to trace the contribution of each individual piece of data to each of the parameters (here the atomic masses). The AME has used this method since 1993. The flow-of-information matrix $\mathbf{F}$ is defined as follows: $\mathbf{K}$, the matrix of coefficients, is a rectangular (Q,M) matrix. The transpose of the response matrix ${}^{\rm t}\mathbf{R}$ is also a (Q,M) rectangular one. The (i,μ) element of $\mathbf{F}$ is defined as the product of the corresponding elements of ${}^{\rm t}\mathbf{R}$ and of $\mathbf{K}$. In Ref. [24], it is demonstrated that such an element represents the *"influence"* of datum i on parameter (mass) m_μ. A column of $\mathbf{F}$ thus represents all the contributions brought by all data to a given mass m_μ, and a row of $\mathbf{F}$ represents all the influences given by a single piece of data. The sum of influences along a row is the *"significance"* of that datum.

It has also been shown [24] that the influences and significances have all the expected properties, namely that the sum of all the influences on a given mass (along a column) is unity, that the significance of a datum does not exceed unity, and that it will decrease when additional independent data are added. The significance defined in this way is exactly $(d\overline{q}/dq)^2$, the squared ratio of the adjusted uncertainty to the input uncertainty. This quantity was used to assess the relative importance of each datum before the introduction of the $\mathbf{F}$ matrix.

A simple interpretation of influences and significances can be obtained by calculating, from the adjusted masses and Eq. (2), the adjusted data:

$$|\overline{q}\rangle = \mathbf{KR}|q\rangle. \tag{6}$$

The ith diagonal element of $\mathbf{KR}$ then represents the contribution of datum i to the determination of $\overline{q_i}$ (same datum): this quantity is exactly what was called above the *significance* of datum i. This ith diagonal element of $\mathbf{KR}$ is the sum of the products of row i of $\mathbf{K}$ and column i of $\mathbf{R}$. The individual terms in this sum are precisely the *influences* defined above.

The flow-of-information matrix $\mathbf{F}$ thus provides insight into how the information from datum i flows into each of the masses m_μ. The flow-of-information matrix cannot be given in full in a printed table. It can be observed along rows, displaying for each datum, the nuclides influenced by this datum and the values of these *influences*. It can also be observed along columns to display for each primary mass all contributing data with their *influence* on that mass.

The first display is partly given in the table of input data (Table I) in column 'Signf.' for the *significance* of primary data and 'Main infl.' for the largest *influence*. Since in the large majority of cases only two nuclides are concerned in each piece of data, the second largest *influence* can easily be deduced. It is therefore unnecessary to give a table of all *influences* for each primary datum.

The second display is given in Part II, Table II for up

to three of the most important data with their *influence* in the determination of each primary mass.

4.4 Consistency of data

The strongly overdetermined system of equations ($Q \gg M$) offers the evaluator several interesting possibilities to examine and judge the data. One may, for example, examine all data for which the adjusted values deviate significantly from the input ones. This helps to locate erroneous input data. One can also examine a group of data from one experiment to check whether the uncertainties assigned in the experimental paper were underestimated.

If the uncertainties dq_i assigned to the data q_i were all reliable, the normalized deviations v_i between the adjusted values $\overline{q}_i$ in Eq. (6) and the input q_i data, $v_i = (\overline{q}_i - q_i)/dq_i$, would be distributed as a Gaussian with standard deviation $\sigma = 1$, and the quantity χ^2:

$$\chi^2 = \sum_{i=1}^{Q} \left(\frac{\overline{q}_i - q_i}{dq_i} \right)^2 \qquad \text{or} \qquad \chi^2 = \sum_{i=1}^{Q} v_i^2 \qquad (7)$$

should have expectation $Q - M$, the number of degrees of freedom, with a standard deviation of $\sqrt{2(Q-M)}$.

One can then define the NORMALIZED CHI, χ_n (also called the 'consistency factor' or 'Birge ratio'): $\chi_n = \sqrt{\chi^2/(Q-M)}$ for which the expected value is $1 \pm 1/\sqrt{2(Q-M)}$.

Another quantity of interest for the evaluator is the PARTIAL CONSISTENCY FACTOR, χ_n^p, defined for a (homogeneous) group of p data points as:

$$\chi_n^p = \sqrt{\frac{Q}{Q-M} \; \frac{1}{p} \; \sum_{i=1}^{p} v_i^2}\,. \qquad (8)$$

By construction, χ_n^p reduces to χ_n when the sum is taken over all the input data.

One can also consider groups of data related to a given laboratory and obtained with a given method of measurement, and examine χ_n^p of each group. Since various methods are used in mass determination, systematic uncertainties associated with a specific method may become apparent when the adjustment is considered as a whole.

There are presently 307 groups of data in Table I (among which 208 have at least one measurement used in determining the masses), identified by the column 'Lab'.

As in previous evaluations, partial consistency factors are estimated for groups of data related to a given laboratory and, where relevant, to a specific method of measurement. A large value of χ_n^p indicates unrecognized systematic effects, underestimated uncertainties, or calibration problems within the group; it does not by itself identify which datum is responsible, and must be followed by examination of the original publications.

When such effects cannot be traced to individual input data, each group of mass-spectrometric data is assigned a factor F according to its partial consistency factor χ_n^p, because its statistical uncertainties and internal systematic errors alone may not reflect the real experimental situation. Based on comparison with all other data, and especially with combinations of reaction and decay energy measurements, we have assigned factors F of 1.0, 1.5, 2.5, or 4.0 to different laboratories. For example, the group of data H25 (see Table I) has been assigned $F = 2.5$, meaning that the total uncertainties assigned to this group are multiplied by 2.5.

One should note that the values obtained for data not used in the least-squares adjustment are particularly informative. Because such data do not contribute to the adjusted masses, their deviations provide a more independent estimate of the consistency of the group with the whole data set.

4.5 Treatment of questionable data

The strong interdependence of most data, as illustrated by the connection diagrams (Figs. 1a–1j), allows local and general consistency tests. These can indicate that certain input values may be problematic. We follow the policy of checking all significant data that differ by more than two (sometimes 1.5) standard deviations from the adjusted values. In many cases, examination of the experimental results shows that a correction is necessary. For example, a particular decay may have been assigned to the wrong final level, a reported decay energy may correspond to an isomer rather than the ground state, or the mass number assigned to a decay may be incorrect. In such cases, the values are corrected, and remarks are added below the corresponding A-group of data in Table I, in order to explain the reasons for the corrections.

It also happens that a careful examination of a particular paper leads to serious doubts about the validity of the results within the reported precision, but does not permit a specific correction. Doubts can also be expressed by the authors themselves. The results are given in Table I and compared with the adjusted values. They are labeled 'F', and not used in the final adjustment, but are always followed by a comment to explain the reason. In several cases, the reader may note that the difference between the experimental and adjusted values is small relative to the experimental uncertainty. This does not invalidate the 'F' assignment, which is based on the evaluation of the original experimental information, rather than only on the residual. Frequently, two (or more) pieces of data are discrepant, leading to a significant contribution to χ^2. A detailed examination of the papers may not allow correction or rejection, indicating that the results in at least one of them cannot be trusted within the given uncertainties. In such cases, based




Figure 2: Changes in ground-state mass uncertainties from AME2020 to AME2026. Grey squares indicate ground-state nuclides included in AME2026. Colored squares show experimentally known nuclides with reduced mass uncertainties; the color represents the ratio σ(AME2020)/σ(AME2026). Red open circles mark nuclides whose masses are experimentally known for the first time in AME2026.

on past experience, we use in the calculations the value that appears most reliable, while the other is labeled 'B' if published in a regular refereed journal, or 'C' otherwise.

In some cases, detailed analysis of strongly conflicting data does not provide grounds for assuming that one result is more dependable than the others, or for rejecting a particular datum. Moreover, poor agreement with other data is not the only reason to question the correctness of the reported data. As in previous AME evaluations, and as discussed in Section 5, we use the property of regularity of the mass surface to help make a choice, as well as to perform further checks on the other data. When a mass is determined by only a single experimental result, but the result is strongly doubted, the policy of assigning a 'D' label, as defined for 'irregular masses' in Section 5.1, may apply.

We also do not accept experimental results when other quantities (e.g., half-lives) derived in the same experiment and for the same nuclide are in strong contradiction with well-established values. To summarize, data assigned any of the labels 'F', 'B', 'C', or 'D' are excluded from the calculations. The least-squares adjustment is then repeated iteratively until consistency is achieved and no further relabeling is required.

4.6 Results of the calculation

In this evaluation, we have 15384 experimental data of which 6780 are labeled 'U' (see above), 1168 are labeled 'O' (old result from the same group), and 1026 are not accepted and labeled 'B': 620, 'C': 158, 'D': 60, or 'F': 188. In the calculation, we thus retain 6410 valid input data, compressed to 4484 through the preaveraging procedure (see Section 4.1.1), after which the separation of secondary data leaves a system of 2582 primary data, representing 1138 primary reactions and decays, and 1444 primary mass-spectrometric measurements. These primary data are used to determine 1493 primary masses (excluding ^{12}C). Thus, we solve a system of 2582 equations with 1493 parameters. Theoretically, the expectation value for χ^2 should be 1089$\pm$47 (and the theoretical $\chi_n = 1 \pm 0.021$). The total χ^2 of the adjustment is actually 1123 ($\chi_n = 1.016$), thus showing that the ensemble of evaluated data was of good quality, and that the adopted criteria for data selection and exclusion were adequate. In the present AME, there are 2685 masses for ground states (excluding ^{12}C) and 512 masses for excited isomers derived from experimental results. Among the 2685 experimental ground-state masses, 137 nuclides have a precision better than 0.1 keV (122 in AME2020), 559 better than 1 keV (402), 1745 better than 10 keV (1532), and 2597 better than 100 keV (2407). Conversely, there are 88 nuclides known with uncertain-

Table E. Experimental data that were replaced by estimated values in AME2020, along with new experimental values and corresponding references.

Item	Value in AME2020	New experimental value	Reference
^{49}Ar−u	−17060(400)#	−16974(5)	2026Fu08
^{65}Cr−u	−28310(200)#	−28208(45)	2022Si20
^{68}Mn−u	−28920(300)#	−29020(30)	2025Ch42
^{68}Fe−u	−43960(200)#	−44100.9(5.6)	2022Po02
^{69}Fe−u	−39200(200)#	−39281.7(9.1)	2022Po02
^{70}Fe−u	−36890(300)#	−37053(12)	2022Po02
^{74}Ni−u	−48700(200)#	−48451.5(3.4)	2022Gi08
		−48441.7(4.7)	2025Xi16
^{80}Zr−u	−54750(300)#	−54355.8(1.5)	2026Er.A
^{103}Sn(β^+)^{103}In	7540(100)#	7507(9)	a
^{106}Zr−u	−58750(200)#	−58582(5)	2024Hu16
^{118}In^{m}(β^-)^{118}Sn	4525(50)#	4741(1)	a
^{120}In^{m}(β^-)^{120}Sn	5420(50)#	5474(4)	2023Ne13
^{138}Sb−u	−54650(300)#	−54381(67)	2026Si.A
^{154}Ce−u	−52220(200)#	−52069(3)	2022Or02
^{189}W(β^-)^{189}Re	2170(200)#	2337(34)	a

a From least-squares analysis in the present work.

ties larger than 100 keV (143 in AME2020).

Fig. 2 summarizes the changes in mass precision from AME2020 to AME2026 for ground-state nuclides. In the present evaluation, the mass uncertainties of 1442 experimentally known nuclides are improved with respect to AME2020. In addition, 142 nuclides whose masses were previously estimated, or absent from the AME2020 mass table, are now known experimentally. The gray squares indicate all ground-state nuclides included in AME2026, providing an overview of the evaluated mass surface. The colored squares indicate nuclides with improved experimental precision, with the color representing the improvement factor, defined as the ratio σ(AME2020)/σ(AME2026). Red open circles mark the new experimental masses in AME2026.

5 Mass estimation

We provide estimated mass values for some nuclides whose masses have not been determined experimentally, as well as for nuclides whose experimental values are considered unreliable. The estimates are based on the trends in the mass surface (TMS), under the assumption that the mass surface is generally smooth. For this purpose, we use an interactive graphical program [20, 25], which allows the simultaneous display of four graphs, showing either particle energies or differences from a mass model, as a function of any of the variables N, Z, A, $N-Z$ or $N-2Z$. Isolines connecting nuclides with the same value of any of these quantities can also be drawn. The mass value of a nuclide can be modified, or a new entry created, in any view, allowing the evaluator to judge how much freedom remains in assigning a value to this mass. At the same time, interdependencies through secondary connections are taken into account. In cases where odd-even staggering produces two alternating trends, data points of one parity (proton or neutron number) may be deselected to allow a clearer view of each trend separately.

The values substituted for data yielding 'irregular masses', as well as the estimates for unknown masses (see below), are thus derived by requiring continuity in several views of the mass surface, including the consequences for connected masses in the same chain. Comparisons with the predictions of 16 nuclear mass models are presently available in this program. All values not derived from experimental data alone are marked with the '#' symbol in all tables, here and in Part II. Since publication of AME1983 [26], the uncertainties are also given for the estimated data derived from TMS. These uncertainties are obtained by considering several graphs of TMS and judging how much the estimated mass may be varied before the extrapolated surface becomes visibly distorted in any of the graphical views. This procedure is unavoidably subjective, but its reliability is tested by comparing the estimates with masses measured after the evaluation.

5.1 Irregular mass values

As mentioned in Section 4.5, some questioned results are labeled 'D' and are not used in the mass calculation. When a single mass deviates significantly from regularity, with no similar pattern for nuclides with the same N or Z, the correctness of the data determining this mass may be questioned. To avoid strongly fluctuating plots, such *irregular* masses are replaced by values derived from the TMS, provided they are based on a single mass relation and have not been verified by an independent experimental method. In most cases, only one measurement is reported for the mass relation in question. In some cases, however, two or more measurements are treated in the same way, since the measurement using the same method and the same type of relation are likely to share the same systematic uncertainty and therefore do not constitute independent verification.

In the last evaluation AME2020, there were 77 such physical quantities, as listed in Table F of Ref. [2]. Of those, 25 items have been superseded by new experimental data or by reanalysis performed by the evaluators in this work. The new results validate our earlier treatment, as shown in Table E. In the present evaluation, 55 irregular experimental values are replaced by estimated ones, as shown in Table F. Such data, as well as the other local irregularities visible in the figures in Part II, could motivate re-measurement of the masses of the nuclides involved, preferably by different methods, in order to remove any doubt and possibly reveal true irregularities due to physical properties.

As a result of this treatment, the masses of 44 nuclides that would otherwise be classified as experimentally known in AME2026 are replaced by estimated values. These nuclides are ^{20}Al, ^{28}Cl, ^{29}Ar, ^{29}Cl, ^{31}K, ^{38}Al, ^{39}Al, ^{41}Si, ^{42}Si, ^{43}P, ^{44}P, ^{45}S, ^{47}Cl, ^{57}Ca, ^{59}Ti, ^{60}Sc, ^{62}Ti, ^{64}V, ^{90}Rh, ^{91}Pd, ^{94}Br, ^{94}Ag, ^{95}Cd, ^{98}In, ^{99}Sn, ^{105}Y, ^{107}Zr, ^{110}Nb, ^{113}Mo, ^{115}Tc, ^{116}Cs, ^{118}Ru, ^{124}Pd, ^{126}Ag, ^{129}Nd, ^{135}Pm^{m}, ^{140}Eu, ^{142}Dy, ^{143}I, ^{166}Eu, ^{168}Re, ^{188}Ta, ^{188}At, and ^{228}Np. Other irregular experimental data do not change the number of known masses, because they involve only nuclides whose masses remain unknown, i.e., none of them is ultimately connected to ^{12}C. Notably, the experimental proton-decay energies of nine proton emitters, namely ^{116}La, ^{117}La, ^{130}Eu, ^{135}Tb, ^{140}Ho, ^{144}Tm, ^{146}Tm, ^{150}Lu, and ^{176}Tl, deviate from the TMS. The deviation is likely due in part to the Thomas-Ehrman shift, which makes proton-emitting nuclides more bound than suggested by the TMS. In addition, ^{130}Eu, ^{135}Tb, and ^{140}Ho are known to be strongly deformed proton emitters whose decays are described by deformed Nilsson configurations [2004Da04, 2004Wo07, 1999Ry04]. Their deviations from the TMS may therefore also contain a deformation-dependent contribution, rather than arising from the Thomas-Ehrman shift alone. Nuclides at the proton dripline have wave functions that differ from those of neighboring particle-bound nuclides, and the assumption of regularity underlying the TMS may therefore not be applicable in these cases. Thus, we do not replace these experimental results.

The irregular experimental values are replaced in all tables and graphs by the values recommended in the present evaluation. In AME1993, a different policy was adopted where the original irregular experimental values were given in all main tables, and the 'recommended' values were given separately in secondary tables. This policy caused confusion among many users of our tables. Since AME1995, we provide only what we consider the *best recommended values*, using when necessary the *values derived from* TMS, as explained above. Data that are not used following this policy can be easily located in Table I where they are flagged 'D' and always accompanied by a comment explaining in which direction the value has been changed and by which amount.

5.2 Estimates for unknown masses

Estimates for unknown masses are made by using trends in the mass surface and by requiring all graphs to remain as smooth as possible, except where effects due to shell closures or nuclear deformation are expected. We caution users that these extrapolations, based on trends in known masses, may be unreliable if unexpected regions of deformation or new (semi-)magic numbers occur.

In addition to the rather severe constraints imposed by the requirement of simultaneous REGULARITY of all graphs, all available experimental information is considered in the estimates. Knowledge of stability or instability against particle emission, or limits on proton or α emission, yields upper or lower limits on the separation energies. For example, before direct mass information became available, the proton separation energy S_p of ^{64}As was estimated to be $-100(200)$ keV from its experimental half-life. It is now experimentally determined to be $-57(30)$ keV in AME2026.

For proton-rich nuclides with $N < Z$, mass estimates can also be guided by mirror symmetry. The usual TMS estimates based on nucleon-separation energies are very useful over a broad regime of the mass surface, but their smooth behavior deteriorates when approaching the proton dripline. In this region, weak binding, Coulomb effects, and the Thomas-Ehrman shift can produce local deviations that are not well represented by a direct extrapolation of the separation-energy trends. For the present evaluation, we use a mirror-energy difference (MED) indicator as an additional guide for very proton-rich nuclides. For odd-Z nuclides, the quantity $S_n(\text{mirror}) - S_p$ is inspected, while for even-Z nuclides, the corresponding quantity $[S_{2n}(\text{mirror}) - S_{2p}]/2$ is used [27]. Fig. 3 displays

the MED indicator for different isotopic chains ranging from $Z = 13$ to $Z = 19$. These quantities remain comparatively smooth across neighboring proton-rich nuclides, making the extrapolation less sensitive to the rapid variation of the proton-separation energies. The recent precise masses of proton-rich nuclides illustrate both the usefulness of mirror-energy differences and the need to allow for mirror-symmetry breaking near the proton dripline [2024Yu13].

Table F. Experimental data replaced by estimates based on trends in the mass surface (TMS) in AME2026. Units for both mass-spectrometric and energy values are given in keV.

Item	Reference	Experimental value		Recommended value	
^{20}Al(3p)^{17}Ne	2025Xu03	1940	110	3570#	200#
^{28}Cl(p)^{27}S	2018Mu18	1600	80	2200#	300#
^{29}Cl(p)^{28}S	2018Xu04	1800	100	2260#	100#
^{29}Ar(2p)^{27}S	2018Mu18	5500	180	4100#	200#
^{31}K–u	2019Ko18	30645	200	32700#	400#
^{38}Al–u	2007Ju03	16210	250	16470#	200#
^{39}Al–u	2007Ju03	20170	630	21490#	300#
^{41}Si–u	2007Ju03	12120	370	13200#	300#
^{42}Si–u	2007Ju03	15160	580	17500#	300#
^{43}P–u	2007Ju03	4680	370	5500#	300#
^{44}P–u	2007Ju03	9380	900	11550#	400#
^{45}S–u	2007Ju03	–3990	690	–3340#	300#
^{47}Cl–u	2007Ju03	–8920	1000	–9580#	200#
^{57}Ca–u	2018Mi08	–7370	990	–6560#	400#
^{59}Ti–u	2020Me06	–25200	270	–25880#	300#
	2020Mi13	–25200	240		
^{60}Sc–u	2020Mi13	–3550	1040	–4550#	300#
^{62}Ti–u	2020Mi13	–13300	450	–12200#	400#
^{64}V–u	2020Mi13	–16690	510	–16320#	400#
^{90}Rh(ϵ)^{90}Ru	2019Pa16	13410	1330	13540#	200#
^{91}Pd(ϵ)^{91}Rh	2018Pa20	11800	2200	12240#	300#
^{94}Br–u	2016Kn03	–46800	400	–47650#	200#
^{94}Ag(ϵ)^{94}Pd	2019Pa16	13400	650	13700#	400#
^{95}Cd(ϵ)^{95}Ag	2018Pa20	10200	1700	12850#	400#
^{98}In(ϵ)^{98}Cd	2019Pa16	12930	400	13730#	300#
^{99}Sn(ϵ)^{99}In	2018Pa20	14700	3600	13400#	500#
^{105}Y–u	2016Kn03	–51270	535	–50570#	200#
^{107}Zr–u	2016Kn03	–54380	449	–53800#	200#
^{110}Nb–u	2016Kn03	–52310	335	–52190#	200#
^{113}Mo-u	2016Kn03	–53390	314	–52550#	300#
^{115}Tc–u	2016Kn03	–56320	316	–55790#	200#
	2024Wa19	–55099	874		
^{116}Cs($\epsilon\alpha$)^{112}Te	1977Bo28	12300	400	13100#	100#
	1976Jo.A	12400	900		

Table F. Experimental data replaced by estimates based on trends in the mass surface (TMS) in AME2026—continued.

Item	Reference	Experimental value		Recommended value	
^{118}Ru–u	2016Kn03	–57640	183	–57180#	200#
^{124}Pd–u	2016Kn03	–60190	372	–58290#	300#
^{126}Ag–u	2016Kn03	–61410	306	–60720#	200#
^{129}Nd(ϵp)^{128}Ce	1978Bo.A	5300	300	5870#	200#
^{135}Pm^{m}(β^+)^{135}Nd	1995Ve08	6040	150	6386#	50#
^{140}Eu(β^+)^{140}Sm	1995Ve08	8470	50	8660#	100#
^{142}Dy(β^+)^{142}Tb	1991Fi03	7100	200	6580#	200#
^{143}I–u	2016Kn03	–50160	461	–50790#	200#
^{166}Eu(β^-)^{166}Gd	2014Ha38	7322	300	7622#	100#
^{168}Re(α)^{164}Ta	1992Me10	5063	13	5200#	100#
^{184}Bi(α)^{180}Tl	2003An27	7830	15	8040#	100#
^{188}Ta–u	2013Sh30	–33612	55	–33910#	200#
^{188}At(p)^{187}Po	2025Ko13	1510	40	2080#	300#
^{228}Np(α)^{224}Pa	2003Ni10	7308	36	7538#	100#
^{236}Bk(α)^{232}Am	2020Po07	7447	14	7700#	200#
^{268}Hs(α)^{264}Sg	2010Ni14	9623	16	9760#	100#
^{270}Db(α)^{266}Lr	2014Kh04	8019	30	8310#	200#
^{270}Mt(α)^{266}Bh	2012Mo25	10414	71	10180#	100#
^{275}Ds(α)^{271}Hs^{m}	2024Og02	11366	20	11200#	300#
^{281}Rg(α)^{277}Mt	2017Og01	9410	50	10020#	400#
^{282}Mt(α)^{278}Bh	2016Ho09	8960	180	8660#	200#
^{284}Cn(α)^{280}Ds	2017Ka66	9869	193	9670#	150#
	2021Sa01	9463	10		
^{286}Cn(α)^{282}Ds	2023Co04	9432	41	9280#	100#
^{290}Nh(α)^{286}Rg	2016Ho09	9846	45	9250#	300#

Another useful estimate can be obtained from the observation that masses of nuclear states belonging to an isobaric multiplet are often well approximated by a quadratic function of the proton number Z (or of the third component of the isospin, $T_3 = \frac{1}{2}(N-Z)$): the Isobaric Multiplet Mass Equation (IMME) [28]. Use of this relation is attractive since, unlike the TMS extrapolation, it incorporates experimental information, namely excitation energies of isobaric analogue states. Up to AME1983, we used the IMME for deriving mass values for nuclides for which little or no information was available. This policy was reconsidered because a quantity derived partly from a calculation should not be presented as 'experimental'. Since AME1993, we have decided not to present any IMME-derived mass values in our evaluation, but rather use the IMME as a guideline when estimating masses of unknown nuclides. We continue this policy here, and do not replace experimental values by estimates from the IMME, even when the latter are orders of magnitude more precise.

Our estimates ensure that every nuclide for which there is an experimental Q-value is connected to the main group of primary nuclides through a continuous chain of known or estimated masses. In addition, the evaluators seek to maintain continuity of the mass surface. Therefore, an estimated value is included for any nuclide if it is between two experimentally studied nuclides on a line defined by either $Z =$ constant (isotopes), $N =$ constant (isotones), $N-Z =$ constant (isodiaphers), or, in a few cases, $N+Z =$ constant (isobars). It would be desirable to provide estimates for all unknown nuclides within reach of current accelerator and mass-separator technologies. However, such an ensemble is not straightforward to define in practice. Instead, we estimate mass values for all nuclides for which at least one piece of experimental information is available, such as identification, a half-life measurement, or evidence for stability towards proton or neutron emission. Then, the ensemble of experimental and estimated masses has the same boundary as in the NUBASE2026 evaluation.




Figure 3: Mirror-energy-difference indicators used to guide proton-rich mass estimates. Filled symbols represent values derived from experimental masses in the present evaluation, whereas open circles represent values involving TMS estimates.

6 Special cases

Special cases have been discussed in the AME series to highlight the issues encountered during the evaluation and to motivate further efforts to resolve them. Some of the special cases discussed in AME2020 have been resolved and have therefore been removed from the current list. Meanwhile, several new cases have been added.

6.1 Evaluation in the lightest mass region

The atomic masses of the lightest nuclides serve as fundamental constants and have important applications in fundamental physics. The local adjustment of ^{1}H, D, ^{3}H, and ^{3}He in AME2016 had a consistency factor $\chi_n = 3.4$, showing that at least one set of systematic uncertainties was underestimated. The additional link available for AME2020 established a consistent network between the FSU Penning trap and LIONTRAP, whereas the measurements from the Washington group were incompatible with this network. The Washington results were assigned a partial consistency factor $F = 2.5$ in AME2020. The connection between hydrogen and deuterium has since been strengthened by the measurement of the H_2^+/D^+ cyclotron-frequency ratio at FSU [2021Fi.1]. In the present evaluation, the reported primary cyclotron-frequency ratio, after applying the molecular binding-energy correction, gives $H_2-D=$ 1548.28597(1) μu. It agrees with the earlier FSU determinations [2018Sm03,2020Fi05], and with the proton and deuteron measurements from LIONTRAP [2017He14, 2020Ra.1]. The FSU group also measured three cyclotron-frequency ratios: $HD^+/^3He^+$, HD^+/T^+, and $T^+/^3He^+$ [2023Me17]. The direct $T^+/^3He^+$ comparison gives the atomic mass difference $m(^3H) - m(^3He) =$ 19.959 41(2) μu, or a tritium β-decay energy of 18.592 07(2) keV, improving the AME2020 result by a factor of three. The three frequency ratios, rather than the atomic masses derived in that work using an adopted HD^+ mass, are entered into the least-squares adjustment.

A direct carbon reference for ^{3}He, the missing link in this mass region in AME2020, was supplied by the $^3He^+/^{12}C^{4+}$ measurement at LIONTRAP [2025Be.1]. It gives $m(^3He) = 3.016\,029\,321\,011(35)$ u, with a relative precision of 12 ppt. This result agrees with the FSU ratios when the current proton and deuteron masses are used, but differs from the Washington result [2015Za13] by 7σ. The new link is used in the present evaluation, whereas the results from [2015Za13] remain excluded. Thus, the former 'light-mass puzzle' is resolved in favor of the mutually consistent LIONTRAP and FSU results.

The corresponding discrepancy for ^{4}He has also been resolved. The direct $^4He^{2+}/^{12}C^{6+}$ measurement [2023Sa.1] gives $m(^4He) = 4.002\,603\,254\,653(48)$ u and also disagrees strongly with the earlier Washington value [2006Va22]. An independent FSU measurement of $^4He^+/D_2^+$ and $^4He^+/H_2D^+$ ratios [2025Re.1] gives 4.002 603 254 670(80) u, supporting [2023Sa.1]. For the adjustment, the molecular mass-difference equations supplied by the authors are used after corrections for molecular binding, electronic binding, residual rotational energy, and polarizability.

All of the new primary links discussed above are retained in the present adjustment, and none exhibits a deviation larger than 0.7σ. The corresponding atomic-mass uncertainties for ^{1}H, D, ^{3}H, ^{3}He, and ^{4}He are 8, 13, 26, 24, and 41 pu (1 pu= 10^{-12} u), respectively. The Washington measurements remain unused with $F = 2.5$. In particular, their link between carbon and ^{16}O is no longer used. As a consequence, the uncertainty of ^{16}O increased from 300 pu in AME2020 to 442 pu in the present evaluation. Further independent measurements remain valuable, especially those that avoid contamination from, or can directly identify, molecular rotational states.

6.2 The mass of ^{70}Br

The mass of the 2.2-s 9^+ isomer in ^{70}Br was measured at LEBIT by Penning-trap mass spectrometry. Using the excitation energy of 2292.3(8) keV, the authors deduced the ground-state mass excess of $-51425(15)$ keV [2009Sa12]. This value gives $Q_{EC} =$ 10504(15) keV for the superallowed $0^+ \to 0^+$ transition and a corrected $\mathcal{F}t$ value of 4090(80) s, which disagrees with the world-average $\mathcal{F}t$ value by 13σ. An independent estimate based on the smooth variation of the Coulomb-energy difference (CED) gives a mass excess of $-51934(16)$ keV, $Q_{EC} = 9996(16)$ keV, and $\mathcal{F}t =$ 3140(60) s [2023Hu18]. The estimated Q_{EC} agrees with the early β-endpoint measurements and restores the regular trends of the CED and the $A = 70$ isobaric-multiplet

mass-equation coefficient. In the present evaluation, the Penning-trap result is therefore not used, and the CED estimate is adopted. Although this choice leaves a residual irregularity in the local mass surface, it provides the most consistent description of the available information. A new direct measurement is required to resolve this case.

6.3 The mass of ^{149}La

The mass of ^{149}La was measured directly for the first time at JYFLTRAP using the PI-ICR method [2025Ja05], reporting a mass excess of −59988.3(28) keV. Together with the other lanthanum masses measured in the same work, this result implied a pronounced bump in the two-neutron separation energies at $N = 93$–94. A subsequent measurement at CPT [2026Li.A], using doubly charged ^{149}La ions, with molecular ions serving as references, yielded −60209.9(16) keV, about 200 keV more bound than the JYFLTRAP value. The mass of ^{149}La, produced by spontaneous fission sources, was measured using the RIKEN MR-TOF mass spectrometer [2026Ki04], yielding a value of −60210(5) keV, which agrees with the CPT value. In addition, the TOF-event-correlated β decays observed at RIKEN gave a half-life of 0.90(13) s, consistent with the known half-life. In the present evaluation, the JYFLTRAP result is not used, while the CPT and RIKEN results are combined to give the mass excess of ^{149}La. With this assignment, the previously reported bump in S_{2n} values disappears and is replaced by a kink similar to that observed in the cerium isotopes. As the neighboring odd-A La isotopes are not known to support a comparable long-lived isomeric state, an unidentified contaminant may have affected the measurement reported in [2025Ja05].

6.4 The Q-value for ^{99}Rh(β^+)^{99}Ru

This case was discussed in AME2012, AME2016, and AME2020. The Q_{β^+} value for ^{99}Rh(β^+)^{99}Ru was adjusted to 2044(7) keV in AME2016, based mainly on the β-endpoint measurements of [1952Sc11] and [1959To25]. A later measurement [1974An23] gave $Q_{\beta^+} = 2170(30)$ keV. Independent decay-scheme information, however, requires $Q_{\beta^+} > 2059.34$ keV, since a level at this excitation energy in ^{99}Ru is populated in the β decay of ^{99}Rh. In AME2020, the incompletely documented result of [1952Sc11] was discarded and the unweighted average of [1959To25] and [1974An23] was used. The adjusted value, 2041(19) keV, remained marginally below the required decay energy, and a direct mass measurement of ^{99}Rh was requested.

The mass of ^{99}Rh has now been measured directly with MR-TOF at RIKEN [2026Ho.A]. The reported mass excess, -85513(23) keV, is 72(30) keV higher (i.e., less bound) than the AME2020 value derived mainly from the decay data. In the present evaluation, the measured ^{99}Rh–^{99}Cd mass difference from the MR-TOF measurement is retained. Together with the independent connection through ^{99}Pd(β^+)^{99}Rh and the well-established mass of ^{99}Ru, the adjustment gives $Q_{\beta^+}[^{99}\mathrm{Rh}(\beta^+)^{99}\mathrm{Ru}] = 2070(15)$ keV. This value is consistent with the lower limit of 2059.34 keV from the observed β-decay feeding. The unweighted average of [1959To25] and [1974An23] is also retained, but contributes only 7% to the adjusted mass. Thus, the requirement from the decay scheme is now satisfied, and this case is considered resolved.

6.5 The masses of ^{171}Os and ^{167}W

A re-examination of the mass surface around ^{171}Os reveals an irregularity in the S_{2n} values when the mass value reported from the Schottky measurement at GSI-ESR is assigned to the ground state. A $(13/2^+)$ isomer has been established in ^{171}Os [2023Zh03] at an excitation energy of 186.32(13) keV, with a measured half-life of 0.79(2) s. If the Schottky result is instead assigned to this isomer, the ground-state mass is lowered by the isomeric excitation energy. In the present adjustment, this reassignment makes the ^{171}Os ground state about 190 keV more bound and removes most of the local irregularity in the mass surface. Such an assignment is plausible because, as noted in [2005Li24], isomers separated by only a few hundred keV could not be resolved in the Schottky spectra. Moreover, the de-excitation of the isomer in ^{171}Os is inferred to proceed through one or more highly converted transitions [2023Zh03]. The suppression of internal conversion in highly charged ions may increase the survival time of the isomer in the storage ring. The Schottky spectrum cannot, however, separate the two states if both are populated.

A similar reassignment is suggested for ^{167}W. The mass excesses reported in [2005Li24] are −34275(28) keV for the state observed in ^{171}Os and −42080(28) keV for that observed in ^{167}W. If the state observed in ^{167}W is assigned to the $(13/2^+)$ isomer at 127.2(17) keV, the corresponding ground-state mass excess is −42207(28) keV. These values give $Q_\alpha = 5507(40)$ keV for the α decay from the ^{171}Os isomer to the ^{167}W ground state. This value is consistent within 1.3σ with $Q_\alpha = 5560(4)$ keV deduced from the 5306(4) keV α decay of the $(13/2^+)$ isomer in ^{171}Os to the $(13/2^+)$ state in ^{167}W [2023Zh03]. Separating such low-lying isomeric states from their ground states requires a mass resolving power of about 1×10^6, making direct mass measurements particularly demanding. Among currently available high-precision techniques, Penning-trap mass spectrometry using the PI-ICR method appears to offer the most realistic way to resolve these states, provided that the preparation and measurement time can be kept short enough for the iso-

mer to survive. Such a measurement would provide a decisive test of the proposed isomeric assignment.

7 General information

The complete data and tables of the present issue are available online at the AMDC website [23], together with additional graphs of the mass surface that supplement those presented in Part II.

Tables of masses (Part II, Table I) and nuclear reaction and separation energies (Part II, Table III) are available in ASCII format to facilitate their use in computer programs by end users. The headers of these files describe the data formats. The first file, named **mass.mas26**, contains the table of masses. The next two files correspond to the table of reaction and separation energies, in two parts of six entries each, as in Part II, Table III: **rct1.mas26** for S_{2n}, S_{2p}, Q_{α}, $Q_{2\beta}$, $Q_{\varepsilon p}$, and $Q_{\beta n}$ (odd pages in this issue); and **rct2.mas26** for S_n, S_p, $Q_{4\beta}$, $Q_{d,\alpha}$, $Q_{p,\alpha}$, and $Q_{n,\alpha}$ (facing even pages).

A cross-platform program, NUCLEUS++, has been developed to provide a unified graphical interface to data from the AME and NUBASE evaluations [29]. In addition to displaying evaluated masses and basic nuclear properties, it also gives access to the original input data associated with each mass determination. The program is available from the AMDC website.

8 Acknowledgments

This work is supported in part by the Guangdong Major Project of Basic Research (2021B0301030006), the Science and Technology Leading Talent Foundation of Gansu Province (25RCKA025), and the National Natural Science Foundation of China (12121005). Work at Argonne National Laboratory was supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under Contract Nos. DE-AC02-06CH11357 and DE-SC0023688. W.J.H. acknowledges support from the Talent Support Project of Guangdong Program (2023TQ07A872). S.N. acknowledges the support of the IJCLab-CNRS/IN2P3.

Appendix A Converting frequency ratios to linear equations

Penning-trap mass measurements are generally reported as cyclotron-frequency ratios. For inclusion in the AME, these ratios are converted into linear relations between atomic masses. A Fortran program implementing this conversion was described in AME2016 [31]. The underlying relations and the treatment of molecular binding energies are summarized below.

In the following, quantities with the subscript r refer to the reference ion, and those without a subscript refer to the ion of interest. For positive ions, the relation used in AME2012 [30] is:

$$R=\frac{f_r}{f}=\frac{\mathcal{M}-m_e\ q+B}{\mathcal{M}_r-m_e\ q_r+B_r}\ \frac{q_r}{q}\,, \tag{9}$$

where f is the cyclotron frequency, q is the positive integer charge number, $\mathcal{M}$ is the neutral-atom mass, B is the ionization energy, and m_e is the electron mass. For measurements involving molecular ions, the relation was generalized in AME2016 [31] to:

$$R=\frac{f_r}{f}=\frac{\mathcal{M}-D-m_e\ q+B}{\mathcal{M}_r-D_r-m_e\ q_r+B_r}\ \frac{q_r}{q}\,. \tag{10}$$

In this expression, $\mathcal{M}$ is the sum of the masses of the constituent neutral atoms and D is the molecular binding energy (dissociation energy). Throughout this derivation, masses are represented by their numerical values in u, and energies by the numerical values of their mass equivalents, E/c^2, in u. Let $M=\mathcal{M}-A$ denote the mass excess (in u) expressed in this convention; Eq. (10) then becomes:

$$A+M-D-qm_e+B=R\frac{q}{q_r}(A_r+M_r-D_r-q_rm_e+B_r) \tag{11}$$

or, alternatively:

$$\begin{aligned} M-R\frac{q}{q_r}M_r=q(1-R)m_e &+ A_r(\frac{q}{q_r}R-\frac{A}{A_r}) \\ &+ R\frac{q}{q_r}(B_r-D_r)-(B-D)\,. \end{aligned}$$

The general aim is to establish some quantity y and its associated precision dy. We define C as a truncated, three-digit decimal approximation of the ratio A to A_r, and then we can write:

$$y \;=\; M-C\ M_r \tag{12}$$

and so

$$y \;=\; y_1+y_2+y_3+y_4\,, \tag{13}$$

where

$$y_1 \;=\; M_r\left(R\ \frac{q}{q_r}-C\right), \tag{14}$$

$$y_2 \;=\; m_e\ q(1-R)\,, \tag{15}$$

$$y_3 \;=\; A_r\left(\frac{q}{q_r}\ R-\frac{A}{A_r}\right), \tag{16}$$

and

$$y_4 \;=\; R\frac{q}{q_r}(B_r-D_r)-(B-D)\,. \tag{17}$$

We now estimate the typical magnitude of each term. For a reference nuclide with mass number $A_r\approx 100$: M_r is generally smaller than 0.1 u; R–C is a few 10^{-4}; $(1-R)$ is usually smaller than unity (and typically 0.2 for a 20% mass change); $R-\frac{A}{A_r}$ varies from 1 to 100×10^{-6}; and

$(B_r - D_r)$ is generally smaller than 0.1 μu. The resulting values of y_1, y_2, y_3, and y_4 are of the order of 10 μu, 100 μu, 10 to 10000 μu, and 0.1 μu, respectively.

The associated precision dy is written:

$$dy = dy_1 + dy_2 + dy_3 + dy_4 , \tag{18}$$

where

$$dy_1 = \frac{q}{q_r}\, M_r\, dR + \left(R\, \frac{q}{q_r} - C \right) dM_r \simeq dR \times 10^5 \ \mu\text{u}, \tag{19}$$

$$dy_2 = m_e\, q\, dR \simeq dR \times 10^3 \ \mu\text{u}, \tag{20}$$

$$dy_3 = \frac{q}{q_r}\, A_r\, dR \simeq dR \times 10^8 \ \mu\text{u}, \tag{21}$$

and

$$dy_4 = \frac{q}{q_r}\, (B_r - D_r)\, dR + R\, \frac{q}{q_r} (dB_r - dD_r) - (dB - dD) \simeq dR \times 10^{-1} \ \mu\text{u}. \tag{22}$$

Consequently, only the third term contributes significantly to the precision of the measurement, and so we write: $dy = dy_3$.

If the two frequencies are each measured with a typical precision of 10^{-7} for ions at $A = 100$, then the precision of the frequency ratio R is 1.4×10^{-7}, and the precision of the mass is approximately 14 μu. Next, we illustrate how to calculate the molecular binding energy for high-precision Penning-trap experiments.

A.1 Bond Dissociation Energy

The molecular correction requires knowledge of the energy needed to dissociate the neutral molecule in its ground state into its constituent neutral atoms in their ground states. The bond dissociation energy D, which quantifies the strength of a chemical bond, is defined for a bond A–B broken through the reaction:

$$\text{AB} \rightarrow \text{A} + \text{B}$$

as the standard enthalpy change of the reaction at a specified temperature [32]:

$$D^o(\text{AB}) = \Delta H f_0^o(\text{A}) + \Delta H f_0^o(\text{B}) - \Delta H f_0^o(\text{AB}), \tag{23}$$

where $\Delta H f_0^o$ is the standard heat of formation. Values are available for a large number of atoms and compounds from the NIST Chemistry WebBook [33]. All D^o values refer to the gaseous state at either 0 K or 298 K. When data on the standard heat of formation at 0 K are unavailable, $D^o(\text{AB})$ is converted to $D^o_{298}(\text{AB})$ using the approximation:

$$D^o_{298}(\text{AB}) \approx D^o(\text{AB}) + 3.72 \text{ kJ/mol}. \tag{24}$$

Unlike diatomic molecules, which involve only one bond, polyatomic molecules have several bonds, and their total dissociation energies are the sum of all individual bond dissociation energies. For example, to obtain the total dissociation energy of CH_4, one calculates the dissociation energies of the four successive bonds CH_3–H, CH_2–H, CH–H, and C–H:

$$\begin{aligned} D^o(\text{CH}_3 - \text{H}) &= \Delta H f_0^o(\text{CH}_3) + \Delta H f_0^o(\text{H}) - \Delta H f_0^o(\text{CH}_4), \\ D^o(\text{CH}_2 - \text{H}) &= \Delta H f_0^o(\text{CH}_2) + \Delta H f_0^o(\text{H}) - \Delta H f_0^o(\text{CH}_3), \\ D^o(\text{CH} - \text{H}) &= \Delta H f_0^o(\text{CH}) + \Delta H f_0^o(\text{H}) - \Delta H f_0^o(\text{CH}_2), \\ D^o(\text{C} - \text{H}) &= \Delta H f_0^o(\text{C}) + \Delta H f_0^o(\text{H}) - \Delta H f_0^o(\text{CH}). \end{aligned} \tag{25}$$

Summing over all four bonds above, one obtains:

$$\begin{aligned} D^o(\text{CH}_4) &= D(\text{CH}_3 - \text{H}) + D(\text{CH}_2 - \text{H}) \\ &\quad + D(\text{CH} - \text{H}) + D(\text{C} - \text{H}) \\ &= \Delta H f_0^o(\text{C}) + 4 \times \Delta H f_0^o(\text{H}) - \Delta H f_0^o(\text{CH}_4) \end{aligned} \tag{26}$$

This expression can be generalized for a polyatomic molecule of the form $\text{A}_\text{n}\text{B}_\text{k}\text{C}_\text{i}$:

$$\begin{aligned} D^o(\text{A}_\text{n}\text{B}_\text{k}\text{C}_\text{i}) &= n\Delta H f_0^o(\text{A}) + k\Delta H f_0^o(\text{B}) + i\Delta H f_0^o(\text{C}) \\ &\quad - \Delta H f_0^o(\text{A}_\text{n}\text{B}_\text{k}\text{C}_\text{i}). \end{aligned} \tag{27}$$

Molecular ionization energies are available from the NIST Chemistry WebBook [33]; atomic ionization energies are tabulated in the NIST Atomic Spectra Database [12]. For multiply charged ions, the sum of the successive ionization energies is required.

A.1.1 $^{13}\text{C}_2\text{H}_2^+$ and $^{14}\text{N}_2^+$ mass doublet

The measurement reported in [2004Ra33] provides an example of the importance of molecular energy corrections. For the $^{13}\text{C}_2\text{H}_2^+$ and $^{14}\text{N}_2^+$ mass doublet, the measured cyclotron-frequency ratio is $R = 0.999\,421\,460\,888(7)$, with a relative precision of 7×10^{-12}. We first calculate $D^o(\text{C}_2\text{H}_2)$ by applying Eq. (27):

$$\begin{aligned} D^o(\text{C}_2\text{H}_2) &= 2 \times \Delta H f_0^o(\text{C}) + 2 \times \Delta H f_0^o(\text{H}) - \Delta H f_0^o(\text{C}_2\text{H}_2) \\ &= 2 \times 716.68 + 2 \times 217.998 - 227.40 \\ &= 1641.956 \text{ kJ/mol} \\ &= 17.018 \text{ eV}. \end{aligned}$$

$D^o(\text{N}_2) = 944.9$ kJ/mol $= 9.793$ eV is obtained from Ref. [34], since $\Delta H f_0^o(\text{N}_2)$ is not available in Ref. [33]. Combining the ionization energy $B(\text{C}_2\text{H}_2) = 11.400$ eV and $B(\text{N}_2) = 15.581$ eV, we obtain the mass difference:

$$^{13}\text{C} + \text{H} - ^{14}\text{N} = 8105.862995(98) \ \mu\text{u}.$$

In the original paper [2004Ra33], this equation is given as:

$$^{13}\text{C} + \text{H} - ^{14}\text{N} = 8105.86288(10) \ \mu\text{u},$$

which differs by 0.12(10) nu (1 nu = 10^{-9} u) from our calculation. The difference arises because updated molecular binding energies from Refs. [33, 34] are used in the present work. In this case, the molecular binding energy is 170 times larger than the reported mass uncertainty, and even small updates to the molecular energies have a significant impact on the deduced mass values.

Appendix B Mixtures of isomers or of isobars in mass spectrometry

When two or more components may appear as a single unresolved spectral line, and their identities or relative contributions are uncertain, a special treatment is required.

The goal is to determine the expected mass M_{exp} and the uncertainty associated with the unknown mixture composition. It is assumed that all lines may contribute, and a uniform probability density is assigned to the allowed mixture compositions. The following treatment assumes that the measured mass is represented by an intensity-weighted centroid of the unresolved components. Let M_0 denote the mass of the lowest-energy component, and M_1, M_2, M_3,...the masses of the other components. For a given composition of the mixture, the resulting mass m is given by

$$m = (1-\sum_{i=1}^{n} x_i)M_0 + \sum_{i=1}^{n} x_i M_i\,, \qquad \text{with} \begin{cases} 0 \le x_i \le 1 \\ \sum_{i=1}^{n} x_i \le 1 \end{cases} \tag{28}$$

where the unknown relative contributions x_1, x_2, x_3, ...are jointly uniform over the region defined by these constraints. If $P(m)$ is the normalized probability density of the mass m, then:

$$\overline{M} = \int P(m)\, m\, dm \tag{29}$$

$$\text{and} \quad \sigma^2 = \int P(m)\,(m-\overline{M})^2\, dm\,. \tag{30}$$

It is thus assumed that the experimentally measured mass is $M_{exp} = \overline{M}$, and that σ, which reflects the uncertainty in the composition of the mixture, is added in quadrature to the experimental uncertainty, assuming the two are independent.

B.1 Case of 2 spectral lines

In the case of two lines, one simply gets

$$m = (1-x_1)M_0 + x_1 M_1 \quad \text{with} \quad 0 \le x_1 \le 1\,. \tag{31}$$

The relation between m and x_1 is one-to-one, so that

$$P(m) = \begin{cases} 1/(M_1 - M_0) & \text{if } M_0 \le m \le M_1, \\ 0 & \text{elsewhere} \end{cases} \tag{32}$$

i.e., a rectangular (uniform) distribution, and one obtains:

$$\begin{aligned} M_{exp} &= \frac{1}{2}(M_0 + M_1)\,, \\ \sigma &= \frac{\sqrt{3}}{6}(M_1 - M_0) = 0.290\ (M_1 - M_0)\,. \end{aligned} \tag{33}$$

Therefore, the ground-state mass $M_0 \pm \sigma_0$ can be determined from the measured mass $M_{exp} \pm \sigma_{exp}$ and the excitation energy $E_1 \pm \sigma_1$ using the following equations:

$$M_0 = M_{exp} - \frac{1}{2}E_1\,,$$

$$\sigma^2 = \frac{1}{12}E_1^2 \qquad \text{or} \qquad \sigma \simeq 0.29\, E_1\,,$$

$$\sigma_0^2 = \sigma_{exp}^2 + (\frac{1}{2}\sigma_1)^2 + \sigma^2\,.$$

The uncertainty expression assumes that the experimental and excitation-energy uncertainties are independent, and evaluates the mixture variance at the adopted excitation energy.

B.2 Case of 3 spectral lines

In the case of three spectral lines, we derive from Eq. (28):

$$m = (1-x_1-x_2)M_0 + x_1 M_1 + x_2 M_2\,, \tag{34}$$

$$\text{with} \begin{cases} 0 \le x_1 \le 1 \\ 0 \le x_2 \le 1 \\ 0 \le x_1 + x_2 \le 1\,. \end{cases} \tag{35}$$

After normalization, one gets:

$$P(m) = \frac{2k}{M_2 - M_0}\,, \tag{36}$$

$$\text{with} \begin{cases} k = (m - M_0)/(M_1 - M_0) & \text{if } M_0 \le m \le M_1 \\ k = (M_2 - m)/(M_2 - M_1) & \text{if } M_1 \le m \le M_2 \end{cases} \tag{37}$$

and finally:

$$M_{exp} = \frac{1}{3}(M_0 + M_1 + M_2) \tag{38}$$

$$\sigma = \frac{\sqrt{2}}{6}\sqrt{M_0^2 + M_1^2 + M_2^2 - M_0 M_1 - M_1 M_2 - M_2 M_0}\,.$$

The ground-state mass $M_0 \pm \sigma_0$ can be determined from the measured mass $M_{exp} \pm \sigma_{exp}$, the excitation en-

ergies $E_1 \pm \sigma_1$, and $E_2 \pm \sigma_2$:

$$\begin{aligned} M_0 &= M_{exp} - \frac{1}{3}(E_1+E_2)\,, \\ \sigma^2 &= \frac{1}{18}(E_1^2+E_2^2-E_1E_2) \\ &\quad \text{or} \qquad \sigma = 0.236\sqrt{E_1^2+E_2^2-E_1E_2}\,, \\ \sigma_0^2 &= \sigma_{exp}^2 + (\frac{1}{3}\sigma_1)^2 + (\frac{1}{3}\sigma_2)^2 + \sigma^2\,. \end{aligned}$$

If the levels are equally spaced, i.e., $E_2 = 2E_1$,

$$\sigma = \frac{\sqrt{6}}{12}E_2 = 0.204\,E_2\,,$$

while for a value of E_1 very near 0 or E_2,

$$\sigma = \frac{\sqrt{2}}{6}E_2 = 0.236\,E_2\,.$$

B.3 Case of more than 3 spectral lines

For more than three lines, one may readily infer $M_{exp} = \sum_{i=0}^{n} M_i/(n+1)$, but the determination of σ requires $P(m)$. As the exact calculation of $P(m)$ becomes rather difficult, it is simpler to use Monte Carlo simulations.

The composition vectors must be sampled uniformly over the allowed region. For each set of x_i values, m is calculated, and the histogram $N_j(m_j)$ of its distribution is built. Writing *nbin* for the number of histogram bins, one gets:

$$P(m_j) = \frac{N_j}{\sum_{j=1}^{nbin} N_j}\,, \tag{39}$$

and

$$\begin{aligned} M_{exp} &= \sum_{j=1}^{nbin} P(m_j) m_j\,, \\ \sigma^2 &= \sum_{j=1}^{nbin} P(m_j)(m_j - M_{exp})^2\,. \end{aligned}$$

For a ground state and three excited isomers, the ground-state mass $M_0 \pm \sigma_0$ can be determined from the measured mass $M_{exp} \pm \sigma_{exp}$ and the excitation energies $E_1 \pm \sigma_1$, $E_2 \pm \sigma_2$, and $E_3 \pm \sigma_3$ as:

$$\begin{aligned} M_0 &= M_{exp} - \frac{1}{4}(E_1+E_2+E_3)\,, \\ \sigma_0^2 &= \sigma_{exp}^2 + (\frac{1}{4}\sigma_1)^2 + (\frac{1}{4}\sigma_2)^2 + (\frac{1}{4}\sigma_3)^2 + \sigma^2\,. \end{aligned}$$

An example of a Fortran program based on the CERN library was given for the cases of two, three, and four lines in AME2016 [31].

Appendix C Definition of decay energies

Conventionally, the total energy released in α decay, Q_α, is defined as the difference between the atomic mass of the parent nuclide and the sum of the atomic masses of the daughter nuclide and ^{4}He:

$$Q_\alpha = M_{\text{parent}} - M_{\text{daughter}} - M_{^4\text{He}}\,. \tag{40}$$

This value equals the sum of the kinetic energies of the α particle and of the recoiling nuclide (with only a minor correction for the fact that the electron cloud of the recoiling atom may be in an excited state). However, some authors quote Q_α as a value 'corrected for screening', which essentially means that they use the masses of bare nuclides for the M values in the above equation.

A similar convention has been used for some published proton-decay energies. This convention differs from that adopted in the AME; values based on nuclear masses should be clearly distinguished from the atomic-mass-based Q values used here.

High-precision α-decay energies

The most precise α-decay energies are those obtained from absolute measurements using magnetic spectrographs, as summarized in Ref. [1991Ry01] and references therein. All α-energy standards are determined using this method. The relation between the α-particle energy E_α and the decay energy Q_α is:

$$Q_\alpha = (M - m_\alpha) - \sqrt{(M-m_\alpha)^2 - 2\times M \times E_\alpha} + 79.0\ \text{eV}\,, \tag{41}$$

where M is the mass of the parent nuclide, m_α is the mass of the α particle (bare ^{4}He nucleus), and 79.0 eV is the total binding energy of the two electrons in neutral helium. This formula was discussed in AME1977 and derived in Ref. [35].

References

References identified in the text such as [25Be.1],[2014Ya.A] or [2016Xu10] are listed in Part II under "References used in the AME2026 *and the* NUBASE2026 *evaluations".*

1 G. Audi, Int. J. Mass Spectr. **251**, 85 (2006)
2 W.J. Huang, M. Wang, F.G. Kondev, G. Audi, and S. Naimi, Chin. Phys. C **45**, 030002 (2021)
3 M. Wang, W.J. Huang, F.G. Kondev, G. Audi, and S. Naimi, Chin. Phys. C **45**, 030003 (2021)
4 F.G. Kondev, R. Chakma, W.J. Huang, M. Wang, and S. Naimi, Chin. Phys. C **50**, 120001 (2026)
5 B. Pritychenko, E. Běták, M.A. Kellett, B. Singh and J. Totans, Nucl. Instrum. Methods Phys. Res. A **640**, 213 (2011)
6 T.P. Kohman, J.H.E. Mattauch and A.H. Wapstra, J. de Chimie Physique **55**, 393 (1958)
7 D.B. Newell, F. Cabiati, J. Fischer, K. Fujii, S.G. Karshenboim, H.S. Margolis, E.d. Mirandés, P.J. Mohr, F. Nez, K. Pachucki, T.J. Quinn, B.N. Taylor, M. Wang, B.M. Wood, and Z. Zhang, Metrologia **55**, L13-L16 (2018)
8 Bureau International des Poids et Mesures (BIPM) SI Brochure (2019, 9th edition)
9 L. Morel, Z. Yao, P. Cladé and S. Guellati-Khélifa, Nature(London) **588**, 61 (2020)
10 E.R. Cohen and A.H. Wapstra, Nucl. Instrum. Methods **211**, 153 (1983)
11 P.J. Mohr, D.B. Newell, B.N. Taylor and E. Tiesinga, Rev. Mod. Phys. **97**, 025002 (2025)
12 A. Kramida, Yu. Ralchenko, J. Reader and NIST ASD Team, NIST Atomic Spectra Database, version 5.12, National Institute of Standards and Technology, Gaithersburg, MD, 2024; available at `https://physics.nist.gov/asd` (accessed 30 June 2026)
13 A. Rytz, At. Data Nucl. Data Tables **47**, 205 (1991)
14 F. Takahashi et al. (Particle Data Group), Int. J. Mod. Phys. A **41**, 2630011 (2026)
15 M. Rosenbusch et al., Nucl. Instrum. Methods Phys. Res. A **1047**, 167824 (2023)
16 W.J. Huang and G. Audi, ND 2016 International Conference on Nuclear Data for Science and Technology, EPJ Web Conf. **146**, 10007 (2017)
17 J. Lindhard, V. Nielsen, M. Scharff, and P.V. Thomsen, Mat. Fys. Medd. Vid. Selsk **33**, 10 (1963)
18 A. Ratkowski, Nuclear Instruments and Methods, **130(2)**, 533 (1975)
19 S. Hofmann, G. Munzenberg, K. Valli, F. Hessberger, J.R.H. Schneider, P. Armbruster, B. Thuma, and Y. Eyal, GSI-Report, page 241, 1982
20 G. Audi, Hyperfine Interactions **132**, 7 (2001), École Internationale Joliot-Curie 2000, Spa, p.103;
21 M.J. Woods and A.S. Munster, NPL Report RS(EXT)95 (1988)
22 Y.V. Linnik, Method of Least Squares (Pergamon, New York, 1961); Méthode des Moindres Carrés (Dunod, Paris, 1963)
23 The AME2026 files in the electronic distribution and complementary documents can be retrieved from the Atomic Mass Data Center (AMDC) through the *Web*: http://amdc.impcas.ac.cn/ and the collaboration site at **ANL**: https://www.anl.gov/phy/atomic-mass-data-resources/, as well as at **IAEA**: https://www-nds.iaea.org/amdc/
24 G. Audi, W.G. Davies and G.E. Lee-Whiting, Nucl. Instrum. Methods A, **249**, 443 (1986)
25 C. Borcea and G. Audi, Rev. Roum. Phys. **38**, 455 (1993) http://amdc.impcas.ac.cn/extrapolations/bernex.pdf
26 A.H. Wapstra, G. Audi and R. Hoekstra, Nucl. Phys. A **432**, 185 (1985)
27 H.T. Fortune, Phys. Rev. C **97**, 034301 (2018)
28 M. MacCormick and G. Audi, Nucl. Phys. A, **925**, 61 (2014)
29 J.Y. Shi, W.J. Huang, M. Wang, X.-L. Yan, D. Lunney, G. Audi, F. G. Kondev, S. Naimi and R. Chakma, Nuclear Science and Techniques **35**, 186 (2024)
30 G. Audi, M. Wang, A.H. Wapstra, F.G. Kondev, M. MacCormick, X. Xu and B. Pfeiffer, Chin. Phys. C **36**, 1287 (2012)
31 W.J. Huang, G. Audi, M. Wang, F.G. Kondev, S. Naimi and X. Xu, Chin. Phys. C **41**, 030002 (2017)
32 B.D. Darwent, NSRDS-NBS **31**, 48 (1970)
33 P.J. Linstrom and W. G. Mallard, NIST Chemistry WebBook, *NIST Standard Reference Database Number* 69, National Institute of Standards and Technology, Gaithersburg MD, 20899, http://webbook.nist.gov, 2026
34 Y.R. Luo and J.A. Kerr, Bond dissociation energies. *CRC Handbook of Chemistry and Physics*. **89**, 2012
35 W.J. Huang, Direct Mass Measurements and Global Evaluation of Atomic Masses. Diss. Université Paris Saclay (COmUE), 2018

Table I. Comparison of input and adjusted values

EXPLANATION OF TABLE

The entries are arranged in groups according to the highest relevant mass number occurring in each item.

Item	K^m, Cs^m, Cs^n, In^p, Tl^q: higher isomers, see NUBASE.	In nuclear reactions: ε = electron capture.	In mass-doublet equations: H = ^{1}H, N = ^{14}N, D = ^{2}H, O = ^{16}O, C = ^{12}C, u = absolute mass-doublet.	In mass-triplet equations: Rb^x, Rb^y: different mixtures of isomers or contaminants.

Input value

Mass doublet: value and standard uncertainty in μu.
Triplet: value and standard uncertainty in keV.
Reaction: value and standard uncertainty in keV.
The value is the combination of mass excesses $\Delta(M-A)$ specified under 'item'. It represents the experimental result and uncertainty reported by the authors, except in a few cases for which comments are provided and for certain α-decay data. If an α decay cannot be clearly assigned as feeding the ground state, an additional uncertainty of 50 keV is added in quadrature to the reported uncertainty. If such energies are reported by more than one group, a further 50 keV is added in quadrature to the uncertainty of the average used in the adjustment.

Adjusted value

Result of the least-squares adjustment. For secondary data (Dg = 2–18), the adjusted value is identical to the input value and is therefore not repeated. The adjusted value is also given only once for a group of results corresponding to the same reaction or mass doublet. Values and uncertainties are rounded, but never to a precision worse than tens of keV.

- # Value and uncertainty not derived solely from experimental data, but based at least partly on trends in the mass surface (TMS).
- $*$ No mass value has been calculated for at least one of the masses involved.

v_i

Normalized deviation between the input and adjusted values, defined as their difference divided by the input uncertainty (see Section 4.4).

Dg

- 1 Primary data (see Section 4).
- 2–18 Secondary data of different degrees.
- B Well-documented data, or data published in peer-reviewed journals, that disagree with other well-documented values.
- C Data from incomplete reports that disagree with other data.
- o Data included in or superseded by later work of same group.
- D Data not verified by an independent method and strongly deviating from the TMS; replaced by an estimated value (see Section 5, p. 120002-21).
- F Critical assessment of the paper raises doubts about the validity of the data within the reported uncertainty.
- R Item replaced for computational reasons by an equivalent item yielding the same result.
- U Data carrying much less weight than a combination of other data.
- – Data to be averaged.

Signf.

Significance ($\times$100) of primary data only (see Section 4.3); the significance of secondary data is always 100%.

Main infl.

Largest *influence* ($\times$100) and the nucleus to whose mass the datum contributes the most (see Section 4.3).

Lab

Identifies the group that measured the corresponding item. For example, the laboratory key MA8 denotes Penning-trap data from the Mainz-Isolde group. The numbers distinguish different experimental conditions.

F

Multiplication factor applied to mass-spectrometric data (see Section 4.4). The standard uncertainty given in the 'Input value' column has been multiplied by this factor before being used in the least-squares adjustment.

Reference

Reference keys:
To reduce the width of the Table, the first two digits of the publication year are omitted. At the end of this volume, however, the complete reference key number is given (e.g., 2003Ba49 rather than 03Ba49).

20Ma09	Results published in peer-reviewed journals. These keys are taken from Nuclear Data Sheets. When an official key is not yet available, a provisional key of the form 20Ma.1 is used.
20Ba.A	Results from abstracts, preprints, private communications, conference proceedings, theses, or annual reports.
Ens169	References for relevant excited-state energies are cited in the remarks in the Qfile. Their reference keys refer to the "Evaluated Nuclear Structure Data Files" (ENSDF), the electronic version of Nuclear Data Sheets (NDS). A key such as Ens169 identifies the release date of the ENSDF file: '16' indicates the year 2016 and '9' the month (the letters a, b, and c denote October, November, and December, respectively).
Nub269	An excitation energy derived or estimated in NUBASE2026 is identified by 'Nubase' (see preceding entry).
Averag	Average of the data from the following lines.
AHW	(or FGK, GAu, HWJ, MMC, SAR, WgM, RCH) : comment written by one of the evaluators.
$*$	A remark on the corresponding item is given below the block of data corresponding to the same (highest) A.
Y	Recalibrations of 65Ry01 for charged particle recalibrations, and recalculated triplets for isomeric mixtures.
Z	Recalibrations of 91Ry01 for α particles, 90Wa22 for γ rays in (n,γ) and (p,γ) reactions, or 91Wa.A for protons and γ rays in (p,γ) reactions.

Remarks. For data marked with a star in the reference column, additional remarks are provided. These remarks are collected at the end of each data block sharing the same highest relevant mass number. They provide:

i) Information explaining how the values in 'Input value' column were derived when the original papers do not explicitly report, for example, mass differences obtained from measured frequency ratios, reaction energies, or transitions to excited states in the final nuclei for which improved excitation-energy values are now available.

ii) Reasons for changing values (e.g., through recalibrations) or uncertainties reported by the authors, or for rejecting the data (i.e., assigning the labels B, C, or F).

iii) Value suggested by TMS and recommended in this evaluation as the best estimate (see Section 5, p. 120002-21).

iv) Separate values for capture ratios (see Section 4.1).

Special notation in remarks:

E_{β^-}, Q_{β^-}	β^- endpoint energy, β^- decay energy
E_{β^+}, Q_{β^+}	β^+ endpoint energy, β^+ decay energy
E_p, Q_p	laboratory-frame proton energy, proton-decay energy
a_s	scattering length
T	threshold energy for a given reaction
ε	electron capture; $\beta^+ = \varepsilon + e^+$ (see NUBASE2026, p. 120001-20)
p^+, pK, pL	fractions of β^+, ε(K) or ε(L) transitions to the mentioned states
L/K , L/M	ε(L)/ε(K), ε(L)/ε(M)
IBE	internal bremsstrahlung endpoint
M-A , D_M	mass excess (in keV), mass difference (in μu)
TMS	trends in the mass surface
',Z' (after uncertainty)	recalibrated value (see 'Reference' above)

Table I. Comparison of input data and adjusted values (Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
π^+	140081.39	0.18	140081.39	0.18	0.0	1	100	100 π^+			20PaDG	*
$\pi^+(2\beta^+)\pi^-$	1021.998	0.001	1021.9980	0.0010	−0.0	1	100	100 π^-			20PaDG	*
*π^+	By convention! This is M=139570.39(0.18) keV + m(e^-)										WgM20a	**
*$\pi^+(2\beta^+)\pi^-$	By convention = 2m(e-).										WgM20a	**
*$\pi^+(2\beta^+)\pi^-$	π^+, π^- should have same mass, with relative difference determined										20PaDG	**
*	as $2(5)10^{-4}$										20PaDG	**
$H_{12}-C$	93902.7	0.4	93900.38289	0.00009	−2.3	U			M17	2.5	66Be10	
	93900.66	0.48			−0.2	U			A2	2.5	70St25	
	93900.32	0.12			0.3	U			B07	1.5	71Sm01	
	93900.391	0.012			−0.3	U			WA1	2.5	95Va38	
	93900.3804	0.0084			0.3	U			MI1	1.0	95Di08	
	93900.3865	0.0017			−0.9	U			WA1	2.5	01Va33	
	93900.3860	0.0042			−0.7	U			ST2	1.0	02Be64	
	93900.38272	0.00040			0.4	o			MZ3	1.0	17He14	
	93900.38290	0.00040			−0.0	U			MZ3	1.0	19He19	
n(β^-)^{1}H	782	13	782.3470	0.0004	0.0	U					51Ro50	
D_6-C	84610.56	0.12	84610.66707	0.00008	0.4	U			A2	2.5	70St25	
	84610.62	0.09			0.3	U			B07	1.5	71Sm01	
	84611.60	0.34			−1.1	U			J5	2.5	72Ka57	*
	84611.47	0.40			−0.8	U			J6	2.5	76Ka50	
	84610.644	0.005			1.8	U			WA1	2.5	92Va.A	
	84610.584	0.078			0.4	U			OH1	2.5	93Ma.A	
	84610.662	0.007			0.3	o			WA1	2.5	93Va.C	
	84610.6616	0.0067			0.3	o			WA1	2.5	95Va38	
	84610.6710	0.0054			−0.7	U			MI1	1.0	95Di08	
	84610.6656	0.0036			0.4	U			MI1	1.0	95Di08	
	84610.66897	0.00086			−0.9	U			WA1	2.5	06Va22	
	84610.66834	0.00024			−2.1	F			WA1	2.5	15Za13	*
	84610.66705	0.00010			0.2	1	60	60 ^{2}H	MZ3	1.0	20Ra.1	*
H_2-D	1547.77	0.28	1548.28597	0.00001	0.7	U			C1	2.5	64Mo.A	
	1548.22	0.05			0.5	o			M19	2.5	67Jo18	
	1548.08	0.08			1.0	o			J2	2.5	69Na21	
	1548.286	0.004			−0.0	o			B07	1.5	71Sm01	
	1548.222	0.063			0.4	o			J5	2.5	72Ka57	
	1548.176	0.133			0.3	o			J5	2.5	72Ka57	
	1548.298	0.008			−1.0	U			B08	1.5	75Sm02	
	1548.301	0.005			−2.0	U			B08	1.5	75Sm02	
	1548.190	0.023			1.7	U			J6	2.5	76Ka50	
	1548.28	0.05			0.0	U			M25	2.5	78Ha14	
	1548.302	0.012			−0.5	U			OH1	2.5	93Go37	
	1548.2836	0.0018			1.3	U			MI1	1.0	95Di08	
	1548.28649	0.00035			−1.5	U			ST2	1.0	08So20	
	1548.28588	0.00006			1.5	o			FS1	1.0	18Sm03	*
	1548.28597	0.00004			−0.0	o			FS1	1.0	20Fi05	*
	1548.28597	0.00001			−0.0	1	98	86 ^{1}H	FS1	1.0	21Fi.1	*
^{1}H(n,γ)^{2}H	2224.564	0.017	2224.5662	0.0004	0.1	U			BNL		80Gr02	
	2224.5	0.12			0.6	U			MMn		80Is02	
	2224.561	0.009			0.6	U			Utr		82Va13	Z
	2224.549	0.009			1.9	U					82Vy10	Z
	2224.560	0.009			0.7	U					83Ad05	Z
	2224.5756	0.008			−1.2	U			NBS		86Gr01	*
	2224.5727	0.0500			−0.1	U			PTc		97Ro26	*
	2224.5660	0.0004			0.6	o			NBS		99Ke05	*
	2224.58	0.05			−0.3	U			Bdn		06Fi.A	*
	2224.56624	0.00044			−0.0	1	100	100 1 n	NBS		06De21	*
*D_6-C	For all 72Ka57 doublets, see also reference										72Og03	**
*D_6-C	F : the other result from same paper is not trusted within error bar										GAu	**
*D_6-C	From frequency ratio $D^+/(^{12}C_6)^+$ =1.0070527379117(85)										20Ra.1	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*H_2-D$	Uncorrected frequency ratio $H_3^+/(HD)^+$ =0.999487821168(19); Reconstructed										HWJ201	**
*	from upper limit for the mass difference between ions										HWJ201	**
$*H_2-D$	From frequency ratio $H_2^+/(^2H)^+$ =0.999231660004(19)										HWJ201	**
$*H_2-D$	From frequency ratio $H_2^+/(^2H)^+$ =0.9992316600030(43)										HWJ241	**
$*^1H(n,\gamma)^2H$	Original 2224.5890(0.0022) revised in reference; error increased by evaluator										90Wa22	**
$*^1H(n,\gamma)^2H$	Original error 0.0005 increased for calibration										GAu	**
$*^1H(n,\gamma)^2H$	More precisely, H+n−D=2388170.07(0.42) nu corrected to 2388169.95(0.42) nu										99Ke05	**
$*^1H(n,\gamma)^2H$	All errors in reference increased by 20 ppm for calibration										06Fi.A	**
$*^1H(n,\gamma)^2H$	Original 2224.56610(0.00044) recalibrated with 2018 Codata										WgM211	**
$^3H_4-C$	64197.0690	0.0062	64197.12559	0.00011	3.7	B			WA1	2.5	93Va04	*
	64197.1136	0.0116			1.0	o			ST2	1.0	02Bf02	
	64197.1148	0.0100			1.1	U			ST2	1.0	06Na49	
H D−C	21926.80973	0.00006	21926.80975	0.00002	0.4	1	11	7 ^2H	MZ3	1.0	20Ra.1	
$^3He_4-C$	64117.2399	0.0039	64117.28796	0.00009	4.9	B			WA1	2.5	93Va04	
	64117.252	0.030			0.5	U			WA1	2.5	93Va04	*
	64117.294	0.011			−0.5	o			ST2	1.0	01Fr18	
	64117.2868	0.0100			0.1	o			ST2	1.0	06Na49	*
	64117.28668	0.00017			3.0	B			WA1	2.5	15Za13	
	64117.28804	0.00014			−0.6	1	45	45 ^3He	MZ3	1.0	25Be.1	*
H_3-^3He	7445.858	0.012	7445.77373	0.00002	−2.8	U			MZ1	2.5	91Ha31	*
	7445.7737	0.0001			0.3	U			FS1	1.0	18Sm03	*
$D_2-H\ ^3H$	4329.257	0.003	4329.24238	0.00002	−3.2	B			B08	1.5	75Sm02	
$^3H-H\ D$	−5877.2	0.7	−5877.52835	0.00002	−0.2	U			C1	2.5	64Mo.A	
	−5877.52837	0.00014			0.1	U			FS1	1.0	15My03	
	−5877.52836	0.00003			0.2	1	52	44 ^3H	FS1	1.0	23Me17	
$^3He-H\ D$	−5896.84	0.42	−5897.48776	0.00002	−0.6	U			C1	2.5	64Mo.A	
	−5897.512	0.005			3.2	B			B08	1.5	75Sm02	
	−5897.495	0.006			0.8	U			B09	1.5	81Sm02	
	−5897.48771	0.00014			−0.4	o			FS1	1.0	15My03	
	−5897.48780	0.00007			0.6	o			FS1	1.0	17Ha33	*
	−5897.48777	0.00003			0.3	1	45	32 ^3He	FS1	1.0	23Me17	
$^3H-^3He$	19.83	0.18	19.95941	0.00002	0.3	U			C1	2.5	64Mo.A	
	19.951	0.004			0.8	o				2.5	84Ni16	*
	19.967	0.003			−1.0	o				2.5	84Li24	
	19.967	0.002			−1.5	U				2.5	85Li02	
	19.948	0.003			1.5	U				2.5	85Ta.A	
	19.9570	0.0013			1.9	U			ST2	1.0	06Na49	
	19.95934	0.00007			1.0	o			FS1	1.0	15My03	
	19.95941	0.00002			−0.1	1	79	56 ^3H	FS1	1.0	23Me17	*
	18.59150	0.00060			911.9	U				2.5	22Ak01	
$^2H(n,\gamma)^3H$	6257.6	0.3	6257.2300	0.0004	−1.2	U					69Pr06	
	6256.96	0.25			1.1	U			ILn		79Br25	Z
$^2H(d,p)^3H$	4029	12	4032.66375	0.00002	0.3	U			CIT		49To23	Y
	4034	6			−0.2	U			MIT		64Sp12	
	4033.7	1.7			−0.6	U			NDm		67Od01	
$^2H(d,n)^3He$	3260	9	3268.9088	0.0004	1.0	U			CIT		49To23	Y
	3269	11			−0.0	U			Wis		56Do41	Y
$^3H(\beta^-)^3He$	18.645	0.016	18.59207	0.00002	−3.3	B					72Be11	*
	18.619	0.040			−0.7	U					73Pi01	*
	18.607	0.013			−1.1	o					76Tr07	
	18.614	0.013			−1.7	U					81Lu07	*
	18.562	0.020			1.5	U					83De47	*
	18.590	0.008			0.3	U					85Si07	*
	18.604	0.006			−2.0	o					85Bo34	
	18.603	0.010			−1.1	o					86Fr09	*
	18.600	0.004			−2.0	U					87Bo07	
	18.598	0.015			−0.4	o					87Bu.A	
	18.603	0.004			−2.7	B					88Ka32	
	18.589	0.003			1.0	o					89St05	
	18.595	0.006			−0.5	o					91Bu12	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{3}H(β^-)^{3}He	18.592	0.003	18.59207	0.00002	0.0	U					91Ka41	*
	18.591	0.002			0.5	U					91Ro07	*
	18.595	0.006			−0.5	U					92Bu13	*
	18.589	0.003			1.0	o					92Ot.A	
	18.593	0.003			−0.3	U					92Ho09	*
	18.591	0.003			0.4	U					93We03	
	18.597	0.014			−0.4	U					95Hi14	
	18.5895	0.0025			1.0	U					95St26	
	18.550	0.019			2.2	U					23As02	*
^{3}H(p,n)^{3}He	−764.08	0.15	−763.7549	0.0004	2.1	o			Zur		61Ry05	
	−764.39	0.37			1.7	U			NRL		64Bo10	
	−763.82	0.08			0.8	U			Zur		64Sa12	
*^{3}H$_4$−C	Item preliminarily disregarded										AHW	**
*^{3}He$_4$−C	Original changed after discussion with authors										AHW	**
*^{3}He$_4$−C	Use instead the most precise difference between ^{3}H and ^{3}He (see below)										GAu	**
*^{3}He$_4$−C	From frequency ratio $^3He^+/(^{12}C_4)^+$ =1.0053440800401(115)										25Be.1	**
*H$_3$−^{3}He	From $^3He^+/H_2^+$ =1.496441095(6) +3eV ionization										AHW	**
*H$_3$−^{3}He	Uncorrected frequency ratio $H_3^+/(^3He)^+$ =0.997536905902(14); Reconstructed										HWJ201	**
*	from upper limit for the mass difference between ions										HWJ201	**
*^{3}He−H D	Other data are related to a metastable state in ^{3}H, not used										HWJ183	**
*^{3}H−^{3}He	Atom mass difference=ion mass difference 18.573 + 0.011 keV										AHW	**
*	required correction cannot be estimated										85Au07	**
*^{3}H−^{3}He	Q_{β^-}=18.592071 0.000022 keV										23Me17	**
*^{3}H−^{3}He	rebuilt from Q value 18.57520(60) keV for molecular T_2 decay to ^{3}HeT+										22Ak01	**
*^{3}H(β^-)^{3}He	For corrections to 72Be11 see reference										82Di01	**
*^{3}H(β^-)^{3}He	For corrections to 73Pi01 and 81Lu07 see reference										85Au07	**
*^{3}H(β^-)^{3}He	Error for 83De47 increased, see reference										85Au07	**
*^{3}H(β^-)^{3}He	Original value 18580(7) corrected in reference										89Re04	**
*^{3}H(β^-)^{3}He	As calculated from their data in reference										89Re04	**
*^{3}H(β^-)^{3}He	E_{β^-}=18.5721(0.0030), SFS and recoil as in reference										88Ka32	**
*^{3}H(β^-)^{3}He	E_{β^-}=18.5705(0.0020), SFS and recoil as in reference										89St05	**
*^{3}H(β^-)^{3}He	E_{β^-}=18.556(0.006), corrections as in reference										91Bu12	**
*^{3}H(β^-)^{3}He	E_{β^-}=18.5733(0.0002+syst), SFS and recoil as in reference										88Ka32	**
*^{3}H(β^-)^{3}He	average 18553(+18−19) (bayesian) and 18548(19) (frequentist)										24As04	**
^{4}He$_3$−C	7809.706	0.009	7809.76396	0.00012	2.6	B			WA1	2.5	92Va.A	
	7809.7493	0.0030			2.0	B			WA1	2.5	95Va38	
	7809.7704	0.0039			−1.7	U			ST2	1.0	01Fr18	
	7809.7620	0.0003			2.6	U			WA1	2.5	01Va.A	
	7809.7467	0.0066			1.0	U			MZ2	2.5	01Br27	
	7809.76246	0.00019			3.2	B			WA1	2.5	04Va14	
	7809.76239	0.00019			3.3	B			WA1	2.5	06Va22	
^{4}He$_5$−C	13016.27326	0.00024	13016.27327	0.00021	0.1	1	73	73 ^{4}He	MZ3	1.0	23Sa.1	*
^{4}He−H$_4$	−28696.8747	0.0026	−28696.87297	0.00005	0.7	o			ST2	1.0	06Na49	
	−28696.8750	0.0026			0.8	U			ST2	1.0	06Na13	
D$_2$−^{4}He	25600.315	0.014	25600.30103	0.00004	−0.7	U			B08	1.5	75Sm02	
	25600.331	0.005			−2.4	U			MZ1	2.5	90Ge12	*
	25600.328	0.005			−2.2	U			MZ1	2.5	92Ke06	*
	25600.308	0.005			−0.3	U			BL1	4.0	01He36	
	25600.30105	0.00008			−0.2	1	31	23 ^{4}He	FS1	1.0	25Re.1	
H$_2$ D−^{4}He	27148.58686	0.00020	27148.58700	0.00004	0.7	1	5	4 ^{4}He	FS1	1.0	25Re.1	
H ^{3}H−^{4}He	21271.075	0.012	21271.05865	0.00005	−0.9	U			B08	1.5	75Sm02	
^{4}H(γ,n)^{3}H	5200	1700	1600	100	−2.1	U					62Ar05	
	2900	500			−2.6	U					69Mi10	*
	8000	3000			−2.1	U					79Me13	*
	2700	600			−1.8	U					81Se11	
	2600	200			−5.0	B					85Fr01	*
	3500	500			−3.8	B					86Be35	*
	2600	400			−2.5	U					86Mi14	*
	3000	200			−7.0	B					87Go25	*
	3800	300			−7.3	B					90Am04	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{4}H(γ,n)^{3}H	3100	300	1600	100	−5.0	B					91Bl05	*
	2300	300			−2.3	U					95Al31	*
	2670	310			−3.5	B					03Me11	
	1600	100				2					09Gu17	*
^{3}He(d,p)^{4}He	18380	10	18353.05439	0.00004	−2.7	U			Mex		64Ma.B	
	18382	15			−1.9	U			Mex		64Ma.B	
	18350.1	3.9			0.8	U			NDm		67Od01	
^{4}Li(p)^{3}He	3300	300	3100	210	−0.7	2					87Br.B	
*^{4}He$_5$−C	From frequency ratio $^{12}C_6^+/(^4He)^{++}$ =1.00065092119 28(119)										23Sa.1	**
*D$_2$−^{4}He	Error has to be confirmed										GAu	**
*^{4}H(γ,n)^{3}H	From ^{7}Li(π^-,t)^{4}H reaction										69Mi10	**
*^{4}H(γ,n)^{3}H	From ^{7}Li(π^-,t)^{4}H reaction										AHW	**
*^{4}H(γ,n)^{3}H	From ^{7}Li(^{3}He,^{3}He ^{3}He)^{4}H reaction										85Fr01	**
*^{4}H(γ,n)^{3}H	From ^{9}Be(^{11}B,^{16}O)^{4}H reaction										86Be35	**
*^{4}H(γ,n)^{3}H	From ^{7}Li(n,α)^{4}H reaction										86Mi14	**
*^{4}H(γ,n)^{3}H	From ^{9}Be(π^-,dt)^{4}H, same data in reference										91Go19	**
*^{4}H(γ,n)^{3}H	From ^{7}Li(π^-,t)^{4}H										90Am04	**
*^{4}H(γ,n)^{3}H	From ^{2}D(t,n)^{4}H										91Bl05	**
*^{4}H(γ,n)^{3}H	From ^{6}Li(^{6}Li,^{8}B)										95Al31	**
*^{4}H(γ,n)^{3}H	Fit with 3 resonances 1.6(0.1), 3.4(0.1), 6.0(0.1) MeV										09Gu17	**
^{5}H(γ,2n)^{3}H	1800	800	1850	90	0.1	U					68Yo06	
	11000	1500			−6.1	F					81Se.A	*
	7400	700			−7.9	F					87Go25	*
	5200	400			−8.4	F					95Al31	*
	1700	300			0.5	U					01Ko52	*
	1800	100			0.5	2					03Go11	*
	1800	200			0.2	2					04St18	
	2400	300			−1.8	2					17Wu03	*
^{4}He(n,γ)^{5}He	−890	50	−735	20	3.1	B					66La04	*
	−735	20				2					09Ak03	
^{4}He(p,γ)^{5}Li	−1965	50				2					65Ma56	*
*^{5}H(γ,2n)^{3}H	From ^{6}Li(π^-,p)^{5}H. F : private communication by author										AHW	**
*^{5}H(γ,2n)^{3}H	From ^{9}Be(π^-,pt)^{5}H, same data in reference										91Go19	**
*^{5}H(γ,2n)^{3}H	F : probably higher state										01Ko52	**
*^{5}H(γ,2n)^{3}H	From ^{7}Li(^{6}Li,^{8}B)										95Al31	**
*^{5}H(γ,2n)^{3}H	F : probably higher state										01Ko52	**
*^{5}H(γ,2n)^{3}H	From p(^{6}He,^{2}He)										01Ko52	**
*^{5}H(γ,2n)^{3}H	From t(t,p)										03Go11	**
*^{5}H(γ,2n)^{3}H	From ^{6}He(d,^{3}He)										17Wu03	**
*^{4}He(n,γ)^{5}He	Average of many reactions leading to ^{5}He										AHW	**
*^{4}He(p,γ)^{5}Li	Average of many reactions leading to ^{5}Li										AHW	**
^{6}Li$_2$−C	30246.152	0.119	30245.7750	0.0030	−0.8	F			BL1	4.0	98He.B	*
	30245.575	0.034			1.5	U			BL1	4.0	01He36	
	30245.7748	0.0031			0.0	1	100	100 ^{6}Li	FS1	1.0	10Mo30	
^{6}Li−H$_6$	−31827.302	0.040	−31827.3040	0.0016	−0.1	U			ST2	1.0	06Na13	
^{6}Li−D$_3$	−27182.500	0.040	−27182.4461	0.0016	0.3	U			BL1	4.0	01He36	
^{4}He−^{6}Li$_{.667}$	−7483.694	0.046	−7483.7112	0.0010	−0.4	U			TT1	1.0	09Br10	
^{6}He−^{7}Li$_{.857}$	5170.947	0.057	5170.95	0.06	0.0	1	100	100 ^{6}He	TT1	1.0	12Br03	
^{6}H(γ,3n)^{3}H	2700	400	2660	240	−0.1	2					84Al08	*
	2600	500			0.1	2					86Be35	*
	2800	500			−0.3	2					92Al.A	*
	2850	900			−0.2	2					08Ca22	*
	2300	640			0.6	2					25Sh16	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{6}Li(n,α)^{3}H	4794	6	4783.4712	0.0015	−1.8	U			Win		67De15	
^{6}Li(p,α)^{3}He	4017	12	4019.7162	0.0014	0.2	U			CIT		49To16	Y
	4021	5			−0.3	U			Wis		51Wi26	Y
	4023	2			−1.6	U			Bir		53Co02	Y
	4025	6			−0.9	o			MIT		64Sp12	
	4018.2	1.1			1.4	U			MIT		81Ro02	
^{6}Li(d,α)^{4}He	22396	12	22372.7706	0.0014	−1.9	U			Bir		53Co02	Y
	22376	14			−0.2	U			Ric		53Ph28	Y
	22403	12			−2.5	U			Mex		64Ma.B	
^{6}Li(p,t)^{4}Li	−18700	300	−18900	210	−0.7	R			Brk		65Ce02	
^{6}He(β^-)^{6}Li	3509.8	3.8	3505.21	0.05	−1.2	U					63Jo04	
^{6}Li(p,n)^{6}Be	−5074	13	−5071	5	0.3	2			CIT		67Ho01	
^{6}Li(^{3}He,t)^{6}Be	−4306	6	−4307	5	−0.1	2			CIT		66Wh01	
$*^{6}$Li$_2$−C	F : leak during the measurement										98He.B	**
$*^{6}$H(γ,3n)^{3}H	From ^{7}Li(^{7}Li,^{8}B)^{6}H										84Al08	**
$*^{6}$H(γ,3n)^{3}H	From ^{9}Be(^{11}B,^{14}O)^{6}H										86Be35	**
*	^{6}H not observed in ^{6}Li(π^-,π^+)										87Se.A	**
$*^{6}$H(γ,3n)^{3}H	From ^{7}Li(^{7}Li,^{8}B)^{6}H										92Al.A	**
$*^{6}$H(γ,3n)^{3}H	Symmetrized from 2910(+850–950) keV										08Ca22	**
$*^{6}$H(γ,3n)^{3}H	From ^{7}Li(ε,epπ^+)^{6}H and 2.3 (+0.5(stat) 0.4(syst)) MeV										25Sh16	**
^{7}Li−H$_7$	−38771.7889	0.0045	−38771.789	0.004	−0.0	1	100	100 ^{7}Li	ST2	1.0	06Na13	*
^{7}Li−^{6}Li$_{1.167}$	−1644.991	0.079	−1644.976	0.005	0.2	o			TT1	1.0	09Br10	
^{6}Li−^{7}Li$_{.857}$	1407.954	0.013	1407.944	0.004	−0.7	U			TT1	1.0	09Br.A	
^{7}Be−^{7}Li	925.355	0.025	925.348	0.024	−0.3	1	90	90 ^{7}Be	MS1	1.0	24Bh03	
^{3}He(α,γ)^{7}Be	1586.3	0.6	1587.075	0.023	1.3	U					82Kr05	
^{7}Li(p,α)^{4}He	17364	11	17346.243	0.004	−1.6	U			CIT		51Wh05	Y
	17352	9			−0.6	U			Bir		53Co02	Y
	17345	13			0.1	U			Ric		53Fa18	Y
	17373	6			−4.5	C			Mex		64Ma.B	
	17357	14			−0.8	U			MIT		64Sp12	
^{7}H(γ,2n)^{5}H	−1100	340	100#	1000#	3.5	F					08Ca22	*
^{7}He(γ,n)^{6}He	450	20	410	7	−2.0	3			MSU		01Ch31	
	430	20			−1.0	3					02Me07	
	360	50			1.0	U					06Sk03	
	400	10			1.0	3					08De29	
	388	20			1.1	3					09Ak03	
	380	28			1.1	U					16Re14	
	410	20			−0.0	3					25Kh09	*
^{7}Li(t,α)^{6}He	9788	30	9839.90	0.05	1.7	U			ChR		54Al35	Y
^{7}Li(d,^{3}He)^{6}He−^{19}F()^{18}O	−1981.09	0.42	−1980.36	0.05	1.7	U			MSU		78Ro01	*
^{6}Li(n,γ)^{7}Li	7250.0	0.5	7251.094	0.004	2.2	U			Utr		68Sp01	
	7250.3	0.9			0.9	U					72Op01	
	7250.98	0.09			1.3	U			Ptn		85Ko47	*
	7249.94	0.15			7.7	C			Bdn		06Fi.A	
^{6}Li(d,p)^{7}Li	5028	2	5026.528	0.004	−0.7	U			Bir		53Co02	Y
	5035	5			−1.7	U			Mex		61Ja23	
	5024	7			0.4	U			MIT		64Sp12	
^{6}Li(t,d)^{7}Li	986	7	993.864	0.004	1.1	U			ChR		54Al35	
^{7}Li(^{3}He,α)^{6}Li	13322	10	13326.527	0.004	0.5	U			Mex		64Ma.B	
^{6}Li(n,γ)^{7}Li^{i}	−3947	50	−4044	10	−1.9	U					69Pr04	*
^{6}Li(^{3}He,d)^{7}Be	136	3	113.316	0.023	−7.6	C			Mex		64Ma.B	
^{7}B(p)^{6}Be	2013	25				3					11Ch32	
^{7}Li(t,^{3}He)^{7}He	−11184	30	−11147	7	1.2	U			LAl		69St02	
^{7}Li^{i}(IT)^{7}Li	11295	10				2					23Dr02	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{7}Be(ε)^{7}Li		866	7	861.956	0.022	−0.6	U						72Pe05	
^{7}Li(p,n)^{7}Be		−1644.04	0.22	−1644.303	0.022	−1.2	U				Zur		61Ry05	Z
		−1643.68	0.26			−2.4	U				Wis		63Ga09	Y
		−1644.30	0.10			−0.0	–				Mar		70Ro07	*
		−1644.18	0.10			−1.2	–				Auc		85Wh03	*
	ave.	−1644.24	0.07			−0.9	1	10	10	^{7}Be			average	
^{7}Li(π^{+},π^{-})^{7}B		−11870	100	−11748	26	1.2	U						81Se.A	
^{7}Be^{i}(IT)^{7}Be		11000	50	10980	30	−0.4	2						67Ha08	
		10970	40			0.3	2						67Mc14	
*^{7}Li−H$_{7}$	D_M=7016003.4256(45)–7×1007825.03207(10) using Ame2003												06Na13	**
*^{7}H(γ,2n)^{5}H	From ^{7}H(γ,4n)^{3}H = 704(323), and ^{5}H(γ,2n)^{3}H = 1800(100) keV												08Ca22	**
*^{7}H(γ,2n)^{5}H	F : not confirmed in later work of reference with higher statistics												10Ni10	**
*^{7}He(γ,n)^{6}He	supersedes analysis in 24Go06 where no uncertainties were given												25Kh09	**
*^{7}Li(d,^{3}He)^{6}He−^{19}F()^{18}O	Q−Q =0.98(0.41) to 2^{+} level at 1982.07(0.09) keV in ^{18}O												Ens967	**
*^{6}Li(n,γ)^{7}Li	Original 7251.02 recalibrated using ^{35}Cl(n,γ) of reference												82Kr12	**
*^{6}Li(n,γ)^{7}Li	Typo 7250.02 in Ame1986 recalib. 7249.97 in Ame1993, 7249.98 Ame2003												GAu	**
*^{6}Li(n,γ)^{7}Li^{i}	IT=11200(50); Q rebuilt with Ame1965												MMC128	**
*^{7}Li(p,n)^{7}Be	T=1880.64(0.09,Z); error in Q increased												AHW	**
*^{7}Li(p,n)^{7}Be	T=1880.43(0.02,Z); error in Q increased												AHW	**
^{8}Li−u		22488.2	4.0	22486.24	0.05	−0.5	U				RI1	1.0	13It01	
^{8}C−u		37606	32	37643	20	1.2	1	37	37	^{8}C	1.0	1.0	11Ch32	*
^{8}He−^{6}Li$_{1.333}$		13776.88	0.72	13775.61	0.11	−1.8	o				TT1	1.0	08Ry03	
		13775.50	0.19			0.6	o				TT1	1.0	08Br.D	*
^{8}Li−^{6}Li$_{1.333}$		2327.426	0.034	2327.44	0.05	0.3	o				TT1	1.0	08Sm.A	
		2327.42	0.11			0.2	o				TT1	1.0	08Sm03	
		2327.42	0.11			0.2	1	21	21	^{8}Li	TT1	1.0	09Br10	
^{8}He−^{7}Li$_{1.143}$		15642.49	0.11	15642.49	0.11	0.0	1	100	100	^{8}He	TT1	1.0	12Br03	
^{4}He(^{18}O,^{14}O)^{8}He		−37967	25	−37975.39	0.11	−0.3	U				MIT		75Ja10	
^{4}He(^{26}Mg,^{22}Mg)^{8}He		−44962	30	−44999.33	0.20	−1.2	U				Brk		74Ce05	
^{4}He(^{64}Ni,^{60}Ni)^{8}He		−31818	15	−31810.7	0.4	0.5	U				Pri		75Ko18	
		−31796	8			−1.8	U				Tex		77Tr07	
^{8}Be(α)^{4}He		91.88	0.05	91.84	0.04	−0.8	2				Zur		68Be02	*
		91.80	0.05			0.8	2						92Wu09	*
		92.2	0.4			−0.9	U						16Re14	
^{6}Li(t,p)^{8}Li		790	11	801.92	0.05	1.1	U				ChR		54Al35	
^{6}Li(^{3}He,p)^{8}Be		16824	12	16787.46	0.04	−3.0	C				Mex		64Ma.B	
^{6}Li(d,γ)^{8}Be^{j}		−5216.5	3.0	−5213.4	2.0	1.0	1	43	43	^{8}Be^{j}			76No07	*
^{6}Li(^{3}He,n)^{8}B		−1974.8	1.0	−1974.8	1.0	−0.0	1	100	100	^{8}B	Nvl		58Du78	Y
^{7}Li(n,γ)^{8}Li		2032.78	0.15	2032.62	0.05	−1.1	–						74Ju.A	*
		2032.77	0.18			−0.8	–				ORn		91Ly01	Z
		2032.57	0.06			0.8	–				Bdn		06Fi.A	
^{7}Li(d,p)^{8}Li		−192	1	−191.95	0.05	0.1	U				Wis		51Wi26	Y
		−188	7			−0.6	U				MIT		64Sp12	
^{7}Li(n,γ)^{8}Li	ave.	2032.61	0.05	2032.62	0.05	0.1	1	79	79	^{8}Li			average	
^{7}Li(^{3}He,d)^{8}Be		11795	13	11760.93	0.04	−2.6	U				Mex		64Ma.B	
^{8}B^{i}(IT)^{8}B		10614	20	10623	7	0.4	1	14	14	^{8}B^{i}			14Br15	
*^{8}C−u	Represents ^{8}C → 4p + ^{4}He, yielding M − A=35030(30) keV												GAu	**
*^{8}He−^{6}Li$_{1.333}$	superseded by 12Br03 as per private comm. from M. Brodeur 23Br.A												FKo233	**
*^{8}Be(α)^{4}He	For atomic binding energy correction see reference												67St30	**
*^{6}Li(d,γ)^{8}Be^{j}	E_d=6962.8(3.0) keV												76No07	**
*^{7}Li(n,γ)^{8}Li	PrvCom to reference												74Aj01	**
^{9}Li−^{6}Li$_{1.500}$		4105.867	0.092	4105.86	0.20	−0.1	o				TT1	1.0	08Sm.A	
		4105.86	0.20				2				TT1	1.0	08Sm03	
^{9}Be−u		12183.0344	0.0035	12183.0340	0.0030	0.0	1	100	100	^{9}Be	1.0	1.0	24Di10	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{9}Be−^{7}Li$_{1.286}$		−8397.39	0.10	−8397.382	0.007	0.1	U			TT1	1.0	09Ri03	
^{9}Be(p,α)^{6}Li		2117	7	2125.599	0.004	1.2	U			CIT		49To16	Y
		2130	10			−0.4	U			Chi		51Ca37	Y
		2125	4			0.1	U			Wis		51Wi26	Y
		2126	2			−0.2	U			Bir		53Co02	Y
		2144	6			−3.1	B			MIT		64Sp12	
		2125.4	1.8			0.1	U			NDm		67Od01	
^{6}Li(α,p)^{9}Be		−2125.6	1.2	−2125.599	0.004	0.0	U			NDm		65Br28	
^{6}Li(α,n)^{9}B		−3974	12	−3976.0	0.9	−0.2	U			Tal		63Me08	
^{7}Li(t,p)^{9}Li		−2397	20	−2386.96	0.19	0.5	U					64Mi04	
		−2385.7	3.0			−0.4	U			MSU		75Ka18	
^{9}Be(^{15}N,^{17}F)^{7}He		−16610	30	−16575	7	1.2	U			Ber		99Bo26	
^{9}Be(d,α)^{7}Li		7162	10	7152.127	0.005	−1.0	U			CIT		51Wh05	Y
		7153	3			−0.3	U			Bir		53Co02	Y
		7162	4			−2.5	U			Mex		64Ma.B	
		7157	8			−0.6	U			MIT		64Sp12	
^{7}Li(^{3}He,p)^{9}Be		11215	15	11200.928	0.005	−0.9	U			Mex		64Ma.B	
^{9}Be(p,^{3}He)^{7}Li^{i}		−22499	50	−22496	10	0.1	o			Brk		65De08	
		−22479	40			−0.4	U			Brk		67Mc14	
^{7}Be(^{3}He,n)^{9}C		−6287	5	−6282.1	2.1	1.0	–			CIT		67Ba.A	Z
		−6275.2	3.5			−2.0	–			CIT		71Mo01	Z
	ave.	−6282.1	2.1			−0.0	1	100	100 ^{9}C			average	
^{9}He(γ,n)^{8}He		100	60	1250	50	19.2	B			MSU		01Ch31	*
		1270	100			−0.2	–			Ber		99Bo26	
		2000	200			−3.7	B					07Go24	
		1330	80			−0.9	–					10Jo06	*
	ave.	1310	60			−0.8	1	56	56 ^{9}He			average	
^{9}Be(γ,n)^{8}Be		−1665	1	−1664.57	0.04	0.4	U			Wis		50Mo56	Y
^{9}Be(p,d)^{8}Be		557	3	560.00	0.04	1.0	U			CIT		49To16	Y
		558	5			0.4	U			Chi		51Ca37	Y
		557.5	1.			2.5	U			Wis		51Wi26	Y
		560	2			0.0	U			Bir		53Co02	Y
		562	4			−0.5	U			MIT		64Sp12	
		559.0	1.1			0.9	U			Zur		66Re02	
		559.6	0.6			0.7	U			NDm		67Od01	Z
^{9}Be(d,t)^{8}Be		4602	13	4592.66	0.04	−0.7	U			MIT		64Sp12	
		4591.7	3.1			0.3	U			NDm		67Od01	
^{9}Be(^{3}He,α)^{8}Be		18931	13	18913.05	0.04	−1.4	U			Mex		64Ma.B	
^{9}N(p)^{8}C		2750	210				2					23Ch46	*
^{9}Be(π^{-},π^{+})^{9}He		−30472	100	−30610	50	−1.4	–					87Se05	
^{9}Be(^{13}C,^{13}O)^{9}He		−50200	600	−49580	50	1.0	o			Ber		88Bo20	
		−49470	80			−1.3	o			Ber		91Bo.B	
^{9}Be(^{14}C,^{14}O)^{9}He		−34580	100	−34580	50	0.0	–			Ber		95Bo.B	
^{9}Be(π^{-},π^{+})^{9}He	ave.	−30540	70	−30610	50	−0.9	1	44	44 ^{9}He			average	
^{9}Be(p,n)^{9}B		−1850.4	1.0				2			Wis		50Ri59	Z
		−1852	3	−1850.4	0.9	0.5	U			Ric		55Ma84	Z
*^{9}Be−u	derived from g-factor measurement, theory dependent											24Di10	**
*^{9}He(γ,n)^{8}He	From scattering length a_s =–10 fm; questioned in reference											10Jo06	**
*^{9}He(γ,n)^{8}He	Scattering length a_s =–3.17(66) fm											10Jo06	**
*^{9}N(p)^{8}C	broad resonant state											23Ch46	**
^{10}B ^{37}Cl−C ^{35}Cl		9987.21	0.56	9986.73	0.07	−0.3	U			H38	2.5	84El05	
^{10}Be−^{7}Li$_{1.429}$		−9334.16	0.13	−9334.24	0.05	−0.6	1	17	17 ^{10}Be	TT1	1.0	09Ri03	
^{10}B−u		12936.862	0.016	12936.862	0.016	−0.0	1	100	100 ^{10}B	MS1	1.0	16Gu02	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{10}C−^{10}B		3916.413	0.090	3916.36	0.07	−0.6	1	67	67 ^{10}C	JY1	1.0	11Er02	
^{7}Li(t,γ)^{10}Be^{i}		−3929.8	21.0				2					73Ab10	
^{10}B(n,α)^{7}Li		2801	4	2789.908	0.015	−2.8	U					67De15	
^{7}Li(α,n)^{10}B		−2787	4	−2789.908	0.015	−0.7	U			Ric		57Bi84	Y
^{10}B(p,α)^{7}Be		1147	5	1145.605	0.027	−0.3	U			CIT		49Ch35	Y
		1146	6			−0.1	U			CIT		51Br10	Y
		1146	2			−0.2	U			Wis		52Cr30	Y
		1153	4			−1.8	U			MIT		64Sp12	
^{10}B(^{3}He,^{6}He)^{7}B		−18550	100	−18287	26	2.6	U			Brk		67Mc14	
^{10}He(γ,2n)^{8}He		1200	300	1440	90	0.8	U					94Ko16	
		1420	100			0.2	2					10Jo06	
		2100	200			−3.3	B					12Si07	
		1600	250			−0.6	2					12Ko43	
		1400	300			0.1	U					15Ma54	
^{10}Be(p,^{3}He)^{8}Li^{i}		−26802.3	5.4				2			MSU		75Ro01	*
^{10}Be(p,t)^{8}Be^{j}		−27487.0	2.6	−27489.3	2.0	−0.9	1	57	57 ^{8}Be^{j}	MSU		75Ro01	*
^{10}B(d,α)^{8}Be		17829	10	17819.74	0.04	−0.9	U			Bir		54El10	Y
		17830	6			−1.7	U			Mex		64Ma.B	
		17818.6	4.1			0.3	U			NDm		67Od01	
^{10}B(p,^{3}He)^{8}Be		−535.5	2.5	−533.31	0.04	0.9	U			Wis		52Cr30	Y
^{10}Li(γ,n)^{9}Li		150	150	26	13	−0.8	U					90Am05	*
		25	15			0.1	3					95Zi03	*
		30	24			−0.1	3					08Ak03	*
^{10}Li^{p}(γ,n)^{9}Li		240	60	220	40	−0.3	3					97Bo10	*
		210	50			0.2	3					97Zi04	*
^{9}Be(^{9}Be,^{8}B)^{10}Li^{q}		−33770	260	−33750	40	0.1	U			Brk		75Wi26	*
^{9}Be(^{13}C,^{12}N)^{10}Li^{q}		−36370	50	−36390	40	−0.4	2			Ber		93Bo03	*
^{10}Be(d,^{3}He)^{9}Li		−14142.8	2.5	−14142.94	0.19	−0.1	U			MSU		75Ka18	
^{9}Be(n,γ)^{10}Be		6812.33	0.06	6812.28	0.05	−0.9	–			MMn		86Ke14	Z
		6812.10	0.14			1.3	–			Bdn		06Fi.A	
^{9}Be(d,p)^{10}Be		4583	8	4587.71	0.05	0.6	U			Ric		51Kl55	Y
		4595	4			−1.8	U			Mex		64Ma.B	
		4590	8			−0.3	U			MIT		64Sp12	
^{9}Be(n,γ)^{10}Be	ave.	6812.29	0.06	6812.28	0.05	−0.3	1	83	83 ^{10}Be			average	
^{9}Be(^{3}He,d)^{10}B		1123	5	1093.310	0.015	−5.9	C			Mex		64Ma.B	
^{10}B(d,t)^{9}B		−2189	10	−2179.9	0.9	0.9	U			MIT		64Sp12	
^{10}B(^{3}He,α)^{9}B		12130	15	12140.5	0.9	0.7	U			Ric		60Sp08	
		12171	15			−2.0	U			Mex		64Ma.B	
^{10}N(p)^{9}C		2200	200	2280	180	0.4	1	80	80 ^{10}N			17Ho10	*
^{10}Be(^{14}C,^{14}O)^{10}He		−41190	70	−41580	90	−5.5	B			Ber		94Os04	
^{10}Be(β^{-})^{10}B		560	5	556.85	0.05	−0.6	U					50Hu27	
		555	5			0.4	U					52Fe16	
^{10}C(β^{+})^{10}B		3604	16	3648.06	0.07	2.8	U					63Ba52	
^{10}B(p,n)^{10}C		−4433.7	1.5	−4430.41	0.07	2.2	U			Har		75Fr.A	
		−4430.17	0.34			−0.7	o			Auc		84Ba12	*
		−4430.17	0.09			−2.7	o			Auc		89Ba28	*
		−4430.30	0.12			−0.9	1	33	33 ^{10}C	Auc		98Ba83	*
^{10}B(^{3}He,t)^{10}C		−3667	10	−3666.65	0.07	0.0	U			Brk		68Br23	
^{10}B(^{14}N,^{14}B)^{10}N		−47550	400	−47230	180	0.8	1	20	20 ^{10}N			02Le16	
*^{10}Be(p,^{3}He)^{8}Li^{i}	Original value –26804.1(5.4) recalibrated											AHW	**
*^{10}Be(p,t)^{8}Be^{j}	Original –27487.6(2.6) recalibrated											GAu	**
*^{10}Li(γ,n)^{9}Li	From ^{11}B(π^{-},p)^{10}Li											GAu	**
*^{10}Li(γ,n)^{9}Li	Resonance less than 50 above the one neutron threshold, but											95Zi03	**
*	could also be final state interaction; then ^{10}Li would be 200 higher											97Bo10	**
*^{10}Li(γ,n)^{9}Li	Deduced from s-state scattering length a_s =–22.4(4.8) fm											08Ak03	**
*^{10}Li^{p}(γ,n)^{9}Li	From ^{10}Be(^{12}C,^{12}N)^{10}Li^{m} (1^{+} level)											GAu	**
*^{10}Li^{p}(γ,n)^{9}Li	Theoretical work: 1^{+} level above 1^{-} ground state											02Ga12	**
*^{9}Be(^{9}Be,^{8}B)^{10}Li^{q}	Q =–34060(250) to 2^{+} level 290(80) above 1^{+} level											93Bo03	**
*	revised with Breit−Wigner line shape. Probably 2^{+} level											97Bo10	**
*^{9}Be(^{13}C,^{12}N)^{10}Li^{q}	Revised with Breit−Wigner line shape (probably 2^{+} level)											97Bo10	**
*^{10}N(p)^{9}C	Q_p=2200(200) keV for 2^{-} state and Q_p=1900(200) for 1^{-} state											17Ho10	**
*^{10}B(p,n)^{10}C	T=4876.90(0.37); withdrawn by author											89Ba28	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{10}$B(p,n)^{10}C	T=4876.88(0.10,Z); original T=4876.95(0.10) keV										MMC126	**
$*^{10}$B(p,n)^{10}C	Average of two datasets; withdrawn by author										98Ba83	**
$*^{10}$B(p,n)^{10}C	T=4877.03(0.13); this is the second 89Ba28 dataset, recalibrated by author										98Ba83	**
^{11}B ^{37}Cl−^{13}C ^{35}Cl	2998.15	1.30	3000.20	0.07	0.6	U			H38	2.5	84El05	
^{11}Li−^{6}Li$_{1.833}$	16003.5	1.2	16003.3	0.7	−0.1	o			TT1	1.0	08Sm.A	
	16003.33	0.66				2			TT1	1.0	08Sm03	
^{11}Be−^{6}Li$_{1.833}$	−6059.27	0.28	−6058.7	0.6	2.1	o			TT1	1.0	08Br.C	*
^{11}Be−^{7}Li$_{1.571}$	−3479.83	0.62				2			TT1	1.0	09Ri03	
^{11}C−^{14}N$_{.786}$	9016.430	0.064	9016.43	0.06	0.0	1	100	100 ^{11}C	MS1	1.0	16Gu02	
^{11}Li−u	43780	130	43723.6	0.7	−0.3	U			TO2	1.5	88Wo09	
	43805	28			−2.9	U			P40	1.0	03Ba.A	
	43715.4	5.0			1.6	o			P40	1.0	04Ba.A	
	43714.5	5.1			1.8	U			P40	1.0	09Ga24	
^{11}Be−u	21654.0	3.6	21661.6	0.6	2.1	o			P40	1.0	04Ba.A	
	21653.5	3.5			2.3	U			P40	1.0	09Ga24	
	21658.5	3.8			0.8	U			P40	1.0	09Ga24	*
^{11}B−u	9305.167	0.013	9305.167	0.013	−0.0	1	100	100 ^{11}B	MS1	1.0	16Gu02	
^{9}Li−^{11}Li$_{.491}$ ^{6}Li$_{.600}$	−3949	175	−3494.8	0.4	1.0	U			P12	2.5	75Th08	
^{9}Li−^{11}Li$_{.409}$ ^{7}Li$_{.643}$	−1250	86	−1288.2	0.3	−0.3	U			P11	1.5	75Th08	*
	−1223	195			−0.3	U			P13	1.0	75Th08	
^{9}Li−^{11}Li$_{.273}$ ^{8}Li$_{.750}$	−1928	31	−1873.26	0.25	0.7	U			P12	2.5	75Th08	
	−1923	31			1.6	U			P13	1.0	75Th08	
^{7}Li(α,γ)^{11}B^{i}	−3885.6	20.0	−3896	9	−0.5	1	21	21 ^{11}B^{i}			66Cu02	*
^{11}B(p,α)^{8}Be	8583	15	8590.09	0.04	0.5	U			CIT		51Li26	Y
	8589	4			0.3	U			Bir		53Co02	Y
	8597	6			−1.2	U			Mex		61Ja23	
	8575	11			1.4	U			MIT		64Sp12	
^{11}B(^{3}He,^{6}He)^{8}B^{i}	−27539	8	−27538	7	0.2	1	86	86 ^{8}B^{i}	MSU		75Ro01	*
^{11}Be^{i}(γ,d)^{9}Li	3245	20				3					97Te07	*
^{9}Be(t,p)^{11}Be	−1164	15	−1168.4	0.6	−0.3	U			Ald		62Pu01	
^{11}B^{j}(2p)^{9}Li	2700	80				3					12Ch40	
^{11}B(d,α)^{9}Be	8029	4	8030.090	0.013	0.3	U			Bir		54El10	Y
	8035	9			−0.5	U			Mex		64Ma.B	
	8024	7			0.9	U			MIT		64Sp12	
	8029.7	2.8			0.1	U			NDm		67Od01	
^{9}Be(^{3}He,p)^{11}B	10344	13	10322.964	0.013	−1.6	U			Mex		64Ma.B	
	10322.1	2.3			0.4	U			NDm		67Od01	
^{11}B(p,^{3}He)^{9}Be^{i}	−24713.3	1.7				2			MSU		74Ka15	*
^{9}Be(^{3}He,p)^{11}B^{i}	−2240	12	−2237	9	0.2	–					63Gr.A	*
	−2240.6	20.			0.2	–			MSU		82Zw02	*
ave.	−2240	10			0.3	1	79	79 ^{11}B^{i}			average	
^{11}B(p,t)^{9}B^{i}	−26064.1	2.3				2			MSU		74Ka15	*
^{9}Be(^{3}He,n)^{11}C^{i}	−4612	50	−4600	40	0.2	1	50	50 ^{11}C^{i}			71Wa21	*
^{11}O(2p)^{9}C	4250	60				2					20We08	
^{10}Be(d,p)^{11}Be	−1721	7	−1723.4	0.6	−0.3	U			CIT		70Go11	
^{11}B(^{7}Li,^{8}B)^{10}Li	−32431	80	−32399	13	0.4	U			MSU		94Yo01	*
^{11}B(^{7}Li,^{8}B)^{10}Li^{q}	−32908	62	−32870	40	0.5	R			MSU		94Yo01	
^{10}Be(p,γ)^{11}B^{i}	−1322	30	−1332	9	−0.3	U					70Go04	*
^{10}B(n,γ)^{11}B	11454.1	0.2	11454.221	0.019	0.6	U			Ptn		86Ko19	Z
	11454.15	0.27			0.3	U			Bdn		06Fi.A	
^{10}B(d,p)^{11}B	9227	5	9229.654	0.019	0.5	U			Bir		54El10	Y
	9234	6			−0.7	U			Mex		64Ma.B	
	9244	11			−1.3	U			MIT		64Sp12	
	9232.9	2.			−1.6	U			NDm		66Br18	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{11}B(^{3}He,α)^{10}B	9101	20	9123.400	0.019	1.1	U			Man		60Ta12	
^{10}B(^{3}He,d)^{11}C	3174	15	3196.71	0.06	1.5	U			Man		60Fo01	
	3226	10			−2.9	C			Mex		64Ma.B	
^{11}N(p)^{10}C	1973	180	1378	5	−3.3	B			MSU		74Be20	*
	1300	40			1.9	o			Lis		96Ax01	
	1450	400			−0.2	U			MSU		98Az01	*
	1630	50			−5.0	B			Spe		00Ol01	*
	1350	120			0.2	U			Lis		00Ma62	*
	1310	50			1.4	U			INS		03Gu06	
	1540	20			−8.1	B					06Ca05	
	1378	5				2					19We11	
^{11}B(π^-,π^+)^{11}Li	−33120	50	−33082.5	0.6	0.7	U					91Ko.B	
^{11}B(^{14}C,^{14}O)^{11}Li	−37120	35	−37048.4	0.6	2.0	U			MSU		93Yo07	
^{11}B^i(IT)^{11}B	12510	50	12560	9	1.0	U					71Wa21	*
^{11}C(β^+)^{11}B	1982.8	2.6	1981.69	0.06	−0.4	U					75Be28	
^{11}B(p,n)^{11}C	−2759.7	3.	−2764.04	0.06	−1.4	U			Wis		50Ri59	Z
	−2763.2	1.4			−0.6	U			Ric		61Be13	Z
^{11}B(^{3}He,t)^{11}C	−2002.1	1.2	−2000.28	0.06	1.5	U			Str		65Go05	Z
^{11}B(^{3}He,t)^{11}C^i	−14151	50	−14160	40	−0.2	1	50	50 ^{11}C^i			71Wa21	*
*^{11}Be−^{6}Li$_{1.833}$	superseded by 09Ri03 as per private comm. from M. Brodeur 23Br.A										FKo233	**
*^{11}Be−u	Result from the “cooling” experiment										09Ga24	**
*^{9}Li−^{11}Li$_{.409}$ ^{7}Li$_{.643}$	Symmetric double-doublet 6−9 8−11 included										GAu	**
*^{7}Li(α,γ)^{11}B^i	IT=12550(30); Q rebuilt with Ame1964										GAu	**
*^{11}B(^{3}He,^{6}He)^{8}B^i	IT=10619(9); rebuilt Q =–27538.9(8.2) keV										GAu	**
*^{11}Be^i(γ,d)^{9}Li	Q(d)=Q(p+n)+2224.5660(0.0004) Q(p+n)=1020(20) keV										MMC162	**
*^{11}B(p,^{3}He)^{9}Be^i	IT=14392.2(1.8); rebuilt Q =–24715.2(1.7); recalibrated +1.87 keV										MMC121	**
*^{9}Be(^{3}He,p)^{11}B^i	IT=12565(12); Q rebuilt with Ame1961										MMC121	**
*^{9}Be(^{3}He,p)^{11}B^i	IT=12563(20), Q rebuilt with Ame1977										MMC121	**
*^{11}B(p,t)^{9}B^i	IT=14655.4(2.5); rebuilt Q =–26064.3; recalibrated +0.16 keV										MMC121	**
*^{9}Be(^{3}He,n)^{11}C^i	IT=12170(40); Q rebuilt; possibly not pure T=3/2										MMC121	**
*^{11}B(^{7}Li,^{8}B)^{10}Li	Original (>–32471) re-evaluated										GAu	**
*	existence of this level not completely certain										94Yo01	**
*^{10}Be(p,γ)^{11}B^i	IT=12550(30); Q rebuilt with Ame1964										MMC121	**
*^{11}N(p)^{10}C	From ^{14}N(^{3}He,^{6}He)^{11}N Q =–25010(100) to 250(150) level										90Aj01	**
*^{11}N(p)^{10}C	From ^{9}Be(^{12}N,^{10}Be)^{11}N										98Az01	**
*^{11}N(p)^{10}C	From ^{10}B(^{14}N,^{13}B)^{11}N										00Ol01	**
*^{11}N(p)^{10}C	From scattering ^{10}C on H; precisely, 1270(+180–50) keV										00Ma62	**
*^{11}B^i(IT)^{11}B	From ^{11}B(^{3}He,^{3}He')										AHW	**
*^{11}B(^{3}He,t)^{11}C^i	IT=12150(50); Q rebuilt; possibly not pure T=3/2										MMC121	**
^{12}Be−u	26911.3	14.2	26922.1	2.0	0.8	U			P40	1.0	09Ga24	
^{12}Be−C	26922.4	2.3			−0.1	1	79	79 ^{12}Be	TT1	1.0	10Et01	
$C_{14}-{}^{12}C_{12}$	1.2	4.9	0.00000	0.00013	−0.2	U			TG1	1.5	09Ke.A	
$C_{14}-{}^{12}C_{15}$	2.1	2.7			−0.5	U			TG1	1.5	11Ke03	
$C_{15}-{}^{12}C_{14}$	−4.0	4.7			0.6	U			TG1	1.5	11Ke03	
	−0.5	6.2			0.1	U			TG1	1.5	11Ke03	
$C_{15}-{}^{12}C_{16}$	−1.6	2.1			0.5	U			TG1	1.5	10Ke09	
	4.7	4.2			−0.7	U			TG1	1.5	11Ke03	
	3.7	4.1			−0.6	U			TG1	1.5	11Ke03	
$C_{16}-{}^{12}C_{12}$	−0.3	6.1			0.0	U			TG1	1.5	09Ke.A	
	2.6	5.0			−0.3	U			TG1	1.5	09Ke.A	
$C_{16}-{}^{12}C_{15}$	−1.3	3.9			0.2	U			TG1	1.5	11Ke03	
	−0.7	1.3			0.4	U			TG1	1.5	12Sm07	
$C_{21}-{}^{12}C_{22}$	1.6	1.3			−0.8	U			TG1	1.5	14Ei01	
$C_{23}-{}^{12}C_{22}$	1.2	1.7			−0.5	U			TG1	1.5	14Ei01	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$C_{23}-{}^{12}C_{21}$	2.2	1.8			−0.8	U			TG1	1.5	14Ei01	
^{7}Li(^{7}Li,2p)^{12}Be	−9710	100	−9841.5	1.9	−1.3	U			LAl		71Ho26	
^{12}C(α,^{8}He)^{8}C	−64520	200	−64249	18	1.4	U					74Ro17	
	−64278	26			1.1	–			Tex		76Tr01	
ave.	−64270	23			0.9	1	63	63 ^{8}C			average	
^{9}Be(^{7}Li,α)^{12}B^{i}	−2308.4	50.	−2258	19	1.0	1	14	14 ^{12}B^{i}	Phi		75Aj03	*
^{12}C(^{3}He,^{6}He)^{9}C	−31578	8	−31571.9	2.1	0.8	U			MSU		71Tr03	
	−31575.6	3.2			1.2	R			MSU		79Ka.A	
^{10}Be(t,p)^{12}Be	−4809	15	−4809.5	1.9	−0.0	U			Brk		78Al29	
	−4808.3	4.2			−0.3	1	21	21 ^{12}Be	Phi		94Fo08	
^{10}B(t,p)^{12}B	6346	6	6342.1	1.3	−0.7	U			Man		60Ja17	
^{10}B(α,d)^{12}C	1340.3	0.8	1339.804	0.015	−0.6	U			Wis		56Do41	Z
	1340.6	1.5			−0.5	U			NDm		65Br28	
^{12}C(d,α)^{10}B	−1340.1	1.2	−1339.804	0.015	0.2	U			NDm		65Br28	
^{10}B(^{3}He,p)^{12}C	19694.5	3.6	19692.858	0.015	−0.5	U			NDm		67Od01	
	19692.86	0.44			−0.0	U			Mun		83Ch08	*
^{10}B(^{3}He,p)^{12}C^{i}	4585	6	4585	3	−0.1	1	31	31 ^{12}C^{i}			62Br10	
^{10}B(^{3}He,n)^{12}N	1570	25	1572.4	1.0	0.1	U			CIT		64Fi02	
	1561	9			1.3	U			CIT		64Ka08	
	1568	20			0.2	U			LAl		66Za01	
	1574	7			−0.2	U			Har		68Ad03	
^{12}N^{i}(2p)^{10}B	2905	29	2952	4	1.6	U					12Ja11	*
^{12}O(2p)^{10}C	1770	20	1636.0	2.0	−6.7	B					95Kr03	
	1638	24			−0.1	U					12Ja11	
	1688	29			−1.8	o					19We03	
	1718	15			−5.5	B					19We11	
	1636	2				2					23Ch47	*
^{12}Li(γ,n)^{11}Li	120	15	210	30	6.0	B					08Ak03	*
	210	30				3					13Ko03	
^{11}B(d,p)^{12}B	1141	4	1145.1	1.3	1.0	1	11	11 ^{12}B	Mex		61Ja23	
	1137	5			1.6	U			MIT		64Sp12	
^{11}B(^{3}He,d)^{12}C	10436	17	10463.204	0.012	1.6	U			Man		60Fo01	
	10469.7	5.7			−1.1	U			NDm		67Od01	
^{11}B(d,n)^{12}C^{i}	−1376.2	4.0	−1376	3	0.0	1	69	69 ^{12}C^{i}			55Ma76	*
^{12}N(β^{+})^{12}C	17406	15	17338.1	1.0	−4.5	B					63Gl04	
^{12}C(p,n)^{12}N	−18119.9	4.4	−18120.4	1.0	−0.1	U			Yal		69Ov01	Z
^{12}C(π^{+},π^{-})^{12}O	−31037	48	−30890.6	2.0	3.0	B					80Bu15	
	−31014	24			5.1	B					92Iv.A	
^{12}N^{i}(IT)^{12}N	12242	4				3					19We11	
*^{9}Be(^{7}Li,α)^{12}B^{i}	IT=12770(50) using Q_{gs}=10461.6(1.6)keV										75Aj03	**
*	energy and resolution arguments for T=1, not an IAS										08Ch28	**
*^{10}B(^{3}He,p)^{12}C	Original Q =15305.45(0.3) revised by authors to 15253.95(31) keV										83Vo.A	**
*	to 2^{+} level at 4438.91(0.31) keV										Ens006	**
*^{12}N^{i}(2p)^{10}B	Q_{2p}=1165(29) to ^{10}B^{i} at 1740.05(0.04)										Nub269	**
*^{12}O(2p)^{10}C	Re-analysis of 19We11										23Ch47	**
*^{12}Li(γ,n)^{11}Li	E_{res} derived in reference from scattering length a_s =–13.7(1.6) fm										10Ha04	**
*^{11}B(d,n)^{12}C^{i}	E_{res}=1627(4) Q =–1376(4); recalibrated Q =–1376.16(4.00) keV										MMC121	**
C H−^{13}C	4500	36	4470.19669	0.00029	−0.6	U			R08	1.5	69De19	
	4470.185	0.008			1.0	U			B08	1.5	75Sm02	
	4470.10	0.05			0.8	U			M25	2.5	78Ha14	
C D−^{13}C H	2921.923	0.008	2921.91072	0.00029	−1.0	U			B08	1.5	75Sm02	
	2921.87	0.05			0.3	U			M25	2.5	78Ha14	
	2921.9086	0.0012			1.8	U			MI1	1.0	95Di08	
	2921.9074	0.0015			2.2	U			MI1	1.0	95Di08	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{13}C−u	3354.8404	0.0041	3354.83522	0.00029	−0.5	U			WA1	2.5	95Va38	
^{9}Be(α,γ)^{13}C^{i}	−4458.4	2.0	−4460.4	1.1	−1.0	2					73Ad02	
	−4461.4	1.4			0.7	2					78Hi06	
^{10}B(α,p)^{13}C	4068	12	4061.547	0.015	−0.5	U			MIT		64Sp12	
	4063.4	2.4			−0.8	U			NDm		67Od01	
^{13}Li(γ,2n)^{11}Li	1470	310	110	70	−4.4	B					08Ak03	*
	1470	350			−3.9	B					10Jo07	*
	110	70				3					13Ko03	*
^{11}B(t,p)^{13}B	−233	4	−233.4	1.0	−0.1	U			Man		60Mu07	
	−233.4	1.0				2			Str		83An15	
^{13}C(d,α)^{11}B	5169	6	5168.108	0.012	−0.1	U			CIT		51Li29	Y
	5166	5			0.4	U			Ric		53Ph28	Y
	5165	10			0.3	U			MIT		64Sp12	
	5166.6	2.5			0.6	U			NDm		70Br23	
^{11}B(^{3}He,p)^{13}C	13221	10	13184.947	0.012	−3.6	C			Mex		64Ma.B	
	13185.4	4.0			−0.1	U			NDm		67Od01	
^{11}B(^{3}He,n)^{13}N	10183	11	10182.13	0.27	−0.1	U					71Hs03	
^{13}Be(γ,n)^{12}Be	100	70	510	10	5.9	B					01Th01	*
	60	10			45.0	B					08Ch07	*
	510	10				2					10Ko17	
	400	30			3.7	B					14Ra07	
^{12}C(n,γ)^{13}C	4946.47	0.17	4946.3088	0.0005	−0.9	U					67Pr10	
	4946.03	0.15			1.9	U			Utr		68Sp01	
	4946.51	0.31			−0.6	U			ILn		79Br25	Z
	4946.321	0.024			−0.5	U					80Wa24	*
	4946.337	0.031			−0.9	U			Utr		81Va.B	*
	4946.31	0.10			−0.0	U			Bdn		06Fi.A	
^{12}C(d,p)^{13}C	2727	6	2721.74261	0.00027	−0.9	o			Ric		51Kl55	Y
	2722	4			−0.1	U			Ric		53Fa18	Y
	2720	2			0.9	U			Bir		54El10	Y
	2725	5			−0.7	U			Mex		61Ja23	
	2722	4			−0.1	U			MIT		64Sp12	
	2722.3	0.6			−0.9	o			NDm		67Od01	
	2721.9	0.8			−0.2	U			NDm		74Jo14	
	2721.80	0.50			−0.1	U			Rez		90Pi05	*
^{13}C(p,d)^{12}C	−2722	7	−2721.74261	0.00027	0.0	o			MIT		64Sp12	
^{13}C(d,t)^{12}C	1311	3	1310.92115	0.00027	−0.0	U			CIT		51Li29	Y
	1311	6			−0.0	U			Mex		64Ma.B	
	1311	6			−0.0	U			MIT		64Sp12	
	1310.9	0.7			0.0	U			NDm		67Od01	
^{12}C(p,γ)^{13}N	1943.24	0.32	1943.49	0.27	0.8	2					77Fr20	Z
	1944.1	0.5			−1.2	2					77He26	Z
^{12}C(d,n)^{13}N	−280.5	3.	−281.08	0.27	−0.2	U			Ric		49Bo67	Y
^{12}C(p,γ)^{13}N^{i}	−13121.62	0.18				2					73Hu07	
^{13}C(^{14}C,^{14}O)^{13}Be^{p}	−37020	50				2			Ber		92Os04	
^{13}N(β^{+})^{13}C	2222.3	3.8	2220.47	0.27	−0.5	U					54Ki23	
^{13}C(p,n)^{13}N	−3002.3	1.0	−3002.82	0.27	−0.5	o			Ric		61Be13	Z
	−3004.1	1.5			0.9	U			NRL		64Bo10	Z
	−3002.4	1.0			−0.4	U			Ric		66Bo20	
^{13}O(β^{+})^{13}N	17500	200	17770	10	1.3	U					65Mc09	
*^{13}Li(γ,2n)^{11}Li	Corresponds to excited state										13Ko03	**
*^{13}Li(γ,2n)^{11}Li	Symmetrized from 120(+60–80) keV										GAu	**
*^{13}Be(γ,n)^{12}Be	From scattering length a_s <–10 fm; questioned in reference										10Ko17	**
*^{13}Be(γ,n)^{12}Be	From scattering length a_s =–20 fm; questioned in reference										10Ko17	**
*	a_s =–3.4(0.6) fm in 10Ko17; a_s =–3.2(+.9–1.1) fm in 07Si24										07Si24	**
*^{12}C(n,γ)^{13}C	$Q(\gamma)$=1261.844(0.006,Z) to 3684.477(0.023,Z) level										81Va.B	**
*^{12}C(n,γ)^{13}C	$Q(\gamma)$=1261.844(0.006,Z) to 3684.493(0.030,Z) level										81Va.B	**
*^{12}C(d,p)^{13}C	Estimated systematic error 0.5 added to statistical error 0.038 keV										AHW	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{14}Be−u		42660	150	42760	70	0.5	–			TO2	1.5	88Wo09	
	ave.	42890	140			−0.9	1	28	28 ^{14}Be			average	
C D_2−^{14}C H_2		9311.498	0.006	9311.503	0.004	0.6	1	20	20 ^{14}C	B08	1.5	75Sm02	
C H_2−N		12576.22	0.10	12576.05967	0.00028	−0.6	U			J2	2.5	69Na21	
		12576.086	0.009			−2.0	U			B07	1.5	71Sm01	
		12576.0598	0.0008			−0.2	U			MI1	1.0	95Di08	
C D−N		11027.815	0.018	11027.77370	0.00028	−1.5	o			B07	1.5	71Sm01	
		11027.773	0.007			0.1	U			B08	1.5	75Sm02	
C H_4−N D		14124.17	0.14	14124.34564	0.00028	0.5	U			J6	2.5	76Ka50	
C D_2−N H_2		9479.68	0.13	9479.48773	0.00028	−0.6	U			J6	2.5	76Ka50	
^{14}N−u		3074.014	0.019	3074.00414	0.00028	−0.2	U			OH1	2.5	93Ma.A	
		3074.0056	0.0018			−0.3	U			WA1	2.5	95Va38	
^{13}C H−^{14}N		8105.86299	0.00010	8105.86299	0.00010	−0.0	1	97	75 ^{13}C	MI3	1.0	04Ra33	*
^{14}C H_2−N D		1716.269	0.003	1716.270	0.004	0.3	1	80	80 ^{14}C	B08	1.5	75Sm02	
^{14}O−^{14}N		5522.702	0.027	5522.702	0.027	−0.0	1	100	100 ^{14}O	MS1	1.0	15Va08	
^{14}N(^{3}He,^{9}Li)^{8}C		−42214	50	−42225	18	−0.2	R			MSU		76Ro04	
^{11}B(α,p)^{14}C		789	17	783.760	0.013	−0.3	U			MIT		64Sp12	
^{14}B^{i}(γ,d)^{12}Be		2515	20				2					01Ta23	*
^{14}C(^{18}O,^{20}Ne)^{12}Be		−15770	50	−15798.8	1.9	−0.6	U			ChR		74Ba15	
^{14}C(d,α)^{12}B		361.8	1.4	361.3	1.3	−0.4	1	89	89 ^{12}B	Wis		56Do41	Z
^{14}C(p,^{3}He)^{12}B^{i}		−30702.73	19.96	−30711	19	−0.4	1	86	86 ^{12}B^{i}			71Ne.A	*
^{14}C(p,t)^{12}C^{j}		−32235.9	2.4				2			MSU		78Ro08	*
^{14}N(d,α)^{12}C		13579	6	13574.22329	0.00026	−0.8	U			Mex		64Ma.B	
		13588	6			−2.3	U			MIT		64Sp12	
^{12}C(^{3}He,p)^{14}N		4779.0	1.4	4778.83111	0.00026	−0.1	U			CIT		62Ba26	Y
		4806	9			−3.0	C			Mex		64Ma.B	
		4776.3	1.5			1.7	U			NDm		67Od01	
^{14}N(p,t)^{12}N		−22135.5	1.0	−22135.5	1.0	0.0	1	100	100 ^{12}N	MSU		75No.A	
^{12}C(^{3}He,n)^{14}O		−1146.86	0.72	−1147.880	0.025	−1.4	U			Nvl		61Bu04	*
		−1148.61	0.56			1.3	U			CIT		62Ba26	*
		−1149.01	0.48			2.4	U			Mar		70Ro07	*
^{14}C(^{7}Li,^{8}B)^{13}Be		−39990	500	−38654	10	2.7	U			Dbn		83Al20	
^{14}C(^{11}B,^{12}N)^{13}Be		−39600	90	−39310	10	3.2	B			Dbn		98Be28	
^{13}C(n,γ)^{14}C		8177	2	8176.434	0.004	−0.3	U					67Th05	
		8176.61	0.24			−0.7	U			Bdn		06Fi.A	
^{13}C(d,p)^{14}C		5946	4	5951.868	0.004	1.5	U			CIT		51Li29	Y
		5952	10			−0.0	U			Nob		54Ah47	Y
		5951	10			0.1	U			Mex		64Ma.B	
		5951	8			0.1	U			MIT		64Sp12	
		5951.85	0.54			0.0	U			Rez		90Pi05	*
^{13}C(p,γ)^{14}N		7551.0	0.8	7550.56358	0.00009	−0.5	U					56Ma87	Z
		7551.1	0.5			−1.1	U					63Bo07	*
^{13}C(^{3}He,d)^{14}N		2048	14	2057.08850	0.00009	0.6	U			MIT		64Sp12	
^{14}N(^{3}He,α)^{13}N		10015	10	10024.24	0.27	0.9	U			Ric		59Yo25	
^{14}F(p)^{13}O		1560	40				3					10Go16	*
^{14}C(π^{-},π^{+})^{14}Be		−38100	170	−37840	70	1.6	R					84Gi09	*
^{14}C(^{14}C,^{14}O)^{14}Be^{p}		−43440	60				2			Ber		95Bo10	
^{14}C(^{7}Li,^{7}Be)^{14}B		−21499	30	−21507	21	−0.3	–			ChR		73Ba34	
^{14}C(^{14}C,^{14}N)^{14}B		−20494	30	−20488	21	0.2	–			Ors		81Na.A	
^{14}C(^{7}Li,^{7}Be)^{14}B	ave.	−21506	21	−21507	21	−0.0	1	100	100 ^{14}B			average	
^{14}C(β^{-})^{14}N		155.2	0.5	156.476	0.004	2.6	U					54Ki23	
		155.74	0.08			9.2	F					91Su09	*
		155.95	0.22			2.4	U					95Wi20	
		156.27	0.14			1.5	U					00Ku25	
^{14}C(p,n)^{14}N		−626.15	0.3	−625.871	0.004	0.9	U			Wis		56Sa06	
		−625.88	0.09			0.1	U			Zur		73Hi.A	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{14}N(p,n)^{14}O	−5930.7	2.8	−5926.711	0.025	1.4	U			Ric		65Ku02	
	−5927.6	1.5			0.6	U			Har		73Cl12	
	−5925.6	0.4			−2.8	F			Auc		77Wh01	*
	−5925.41	0.08			−16.3	F			Auc		81Wh03	*
	−5925.41	0.11			−11.8	F			Auc		98Ba83	*
	−5926.68	0.17			−0.2	U			Auc		03To03	
^{14}N(^{3}He,t)^{14}O	−5161.3	0.8	−5162.956	0.025	−2.1	F			Mun		77Vo02	*
*^{13}C H−^{14}N	Original 8105.86288(10) μu recalculated for molecular and ionization										HWJ153	**
*^{14}B^i(γ,d)^{12}Be	Q(d)=Q(p+n)+2224.5660(0.0004) Q(p+n)=290(20) keV										MMC162	**
*^{14}C(p,^{3}He)^{12}B^i	IT=12710(20); Q rebuilt with Ame1964										MMC129	**
*	energy and resolution arguments for T=1, not an IAS										08Ch28	**
*^{14}C(p,t)^{12}C^j	IT=27595.0(2.4); Q rebuilt										MMC121	**
*^{12}C(^{3}He,n)^{14}O	Originals T=1436.2(0.9,Bu) 1437.5(0.7,Ba) 1437.9(0.6,Ro) respectively, recalibrated										MMC126	**
*^{13}C(d,p)^{14}C	Estimated systematic error 0.5 added to statistical error 0.20										AHW	**
*^{13}C(p,γ)^{14}N	E_p=1747.06(0.53) to 2$^+$ level at 9172.25 keV										Ens01a	**
*^{14}F(p)^{13}O	Broad distribution may contain two states										23Ch22	**
*^{14}C(π^-,π^+)^{14}Be	Original error 160 increased with 60 calibration uncertainty										GAu	**
*^{14}C(β^-)^{14}N	F : find 17 keV neutrino. See also reference										91No07	**
*^{14}N(p,n)^{14}O	F : withdrawn by author										81Wh03	**
*^{14}N(p,n)^{14}O	Authors recalibrated 77Wh01 for atomic effects										81Wh03	**
*^{14}N(p,n)^{14}O	F : withdrawn by author										98Ba83	**
*^{14}N(p,n)^{14}O	Original T=6353.08(0.07) recalibrated to T=6352.99(0.12) by author										98Ba83	**
*^{14}N(p,n)^{14}O	F : withdrawn by author										03To03	**
*^{14}N(^{3}He,t)^{14}O	F : rejected in reference of same group										09Fa15	**
C D H−^{15}N	21817.9119	0.0008	21817.9115	0.0006	−0.5	1	61	61 ^{15}N	MI1	1.0	95Di08	
C H$_3$−^{15}N	23366.1979	0.0017	23366.1975	0.0006	−0.3	1	14	14 ^{15}N	MI1	1.0	95Di08	
^{15}F−u	17477	86	17785	15	1.4	F				2.5	01Ze.A	*
^{14}N D−^{15}N H	9242.29	0.11	9241.8518	0.0007	−1.6	U			C5	2.5	71Ke01	
	9241.780	0.008			6.0	B			B08	1.5	75Sm02	
^{15}Be(γ,3n)^{12}Be	330	20	333	20	0.1	1	98	98 ^{15}Be			24Ku30	
^{15}N(p,α)^{12}C	4966	6	4965.4928	0.0006	−0.1	U			CIT		51Li26	Y
	4962	4			0.9	U			Bir		53Co02	Y
	4954	8			1.4	U			Mex		64Ma.B	
	4965	7			0.1	U			MIT		64Sp12	
^{12}C(α,n)^{15}O	−8503	12	−8502.0	0.5	0.1	U			Tal		63Ne05	
^{15}N(d,α)^{13}C	7675	9	7687.2354	0.0006	1.4	U			Mex		64Ma.B	
	7689	6			−0.3	U			MIT		64Sp12	
^{15}Ne(2p)^{13}O	2522	66				3					14Wa09	
^{15}Be(γ,n)^{14}Be	1800	100	1720	70	−0.8	–					13Sn02	
	1700	130			0.1	–					21Co07	
ave.	1760	80			−0.5	1	74	72 ^{14}Be			average	
^{14}C(d,p)^{15}C	−1006.5	0.8				2			Wis		56Do41	Y
^{14}C(p,γ)^{15}N^i	−1407.8	3.5				2					59Fe99	*
^{14}N(n,γ)^{15}N	10833.2	0.6	10833.2967	0.0008	0.2	U					68Gr14	
	10833.1	0.7			0.3	U					74Sp04	
	10833.5	0.05			−4.1	B			MMn		80Is02	
	10833.314	0.012			−1.4	U					97Ju02	
	10833.2339	0.0500			1.3	U			PTc		97Ro26	*
	10833.32	0.22			−0.1	U			Bdn		06Fi.A	
	10833.3	0.5			−0.0	U					06Be33	
^{14}N(d,p)^{15}N	8629	11	8608.7305	0.0006	−1.8	U			CIT		52Mi54	Y
	8614	6			−0.9	U			Mex		64Ma.B	
	8623	3			−4.8	B			MIT		64Sp12	
	8608.83	0.50			−0.2	U			Rez		90Pi05	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{14}N(p,γ)^{15}O	7297.1	0.9	7296.8	0.5	−0.4	1	30	30 ^{15}O	CIT		72Ne05	
^{14}N(^{3}He,d)^{15}O	1803	10	1803.3	0.5	0.0	U			Ric		59Yo25	Y
	1802	15			0.1	U			Man		60Fo01	
^{15}F(p)^{14}O	1410	150	1270	14	−0.9	o					03Le26	
	1510	110			−2.2	U					03Pe23	
	1490	130			−1.7	U					04Go15	*
	1560	130			−2.2	U					04Le12	
	1230	50			0.8	U					05Gu25	
	1270	14				3					16De15	
^{15}C(β^-)^{15}N	9810	30	9771.7	0.8	−1.3	U					59Al06	
^{15}O(β^+)^{15}N	2745	5	2754.2	0.5	1.8	U					57Ki22	
^{15}N(p,n)^{15}O	−3541.7	0.9	−3536.5	0.5	5.7	B					58Jo28	Y
	−3535.1	1.0			−1.5	–			CIT		72Je02	Z
	−3537.6	0.8			1.4	–					72Sh08	Z
ave.	−3536.6	0.6			0.2	1	70	70 ^{15}O			average	
*^{15}F−u	F : results distrusted (see also ^{18}Na and ^{19}Mg)										GAu	**
*^{14}C(p,γ)^{15}N^{i}	From a parametrized fit										MMC122	**
*^{14}N(n,γ)^{15}N	Original error 0.0005 increased for calibration										GAu	**
*^{14}N(d,p)^{15}N	Estimated systematic error 0.5 added to statistical error 0.061 keV										AHW	**
*^{15}F(p)^{14}O	Symmetrized from 1450(+160–100) keV										04Go15	**
^{16}O−H$_{16}$	−130285.89116	0.00069	-130285.891	50.000	4−0.	5 1		413 9 ^{16}O	MZ3	1.0	19He19	*
C H$_2$ D−O	34837.406	0.033	34837.2226	0.0004	−3.7	B			B07	1.5	71Sm01	
	34837.202	0.020			0.7	U			B08	1.5	75Sm02	
C D$_2$−O	33289.129	0.033	33288.9367	0.0004	−3.9	B			B07	1.5	71Sm01	
	33289.061	0.038			−2.2	U			B07	1.5	71Sm01	
	33288.940	0.019			−0.1	U			B08	1.5	75Sm02	
C$_4$−O$_3$	15256.131	0.018	15256.1429	0.0013	0.3	o			WA1	2.5	92Va.A	
	15256.086	0.081			0.3	U			OH1	2.5	93Ma.A	
	15256.121	0.009			1.0	o			WA1	2.5	95Va38	
	15256.1425	0.0008			0.2	o			WA1	2.5	01Va33	
	15256.1415	0.0005			1.1	o			WA1	2.5	03Va.A	
	15256.14129	0.00054			1.2	U			WA1	2.5	06Va22	
C H$_4$−O	36387.55	0.8	36385.5086	0.0004	−1.0	U			J1	2.5	68Ma45	
	36386.01	0.24			−0.8	U			J2	2.5	69Na21	
	36385.644	0.036			−2.5	U			B07	1.5	71Sm01	
	36385.5062	0.0013			1.9	U			MI1	1.0	95Di08	
	36385.5073	0.0019			0.7	U			MI1	1.0	95Di08	
	36385.5060	0.0022			1.2	U			MI1	1.0	95Di08	
^{16}O−u	−5085.362	0.027	−5085.3810	0.0004	−0.3	U			OH1	2.5	93Ma.A	
	−5085.3798	0.0011			−0.4	U				2.5	16Ho.A	
^{14}C H$_2$−O	23977.413	0.014	23977.433	0.004	1.0	U			B08	1.5	75Sm02	
N D−O	22261.160	0.013	22261.1630	0.0003	0.2	U			B08	1.5	75Sm02	
N$_2$−C O	11233.57	0.20	11233.3893	0.0004	−0.4	U			J2	2.5	69Na21	
	11233.543	0.025			−4.1	B			B07	1.5	71Sm01	
	11233.43	0.21			−0.1	U			J6	2.5	76Ka50	
	11259	27			−0.4	U			CR1	2.5	89Sh10	
	11233.3909	0.0022			−0.7	U			MI1	1.0	95Di08	
	11233.38932	0.00042			−0.1	1	82	74 ^{14}N	MI1	1.0	04Th17	
^{16}O(α,^{8}He)^{12}O	−66020	120	−65834.4	2.0	1.5	U			Brk		78Ke06	
^{16}O(p,α)^{13}N	−5211	10	−5218.43	0.27	−0.7	U			MIT		64Sp12	
^{16}O(^{3}He,^{6}He)^{13}O	−30516	14	−30513	10	0.2	2			Brk		70Me11	*
	−30511	13			−0.2	2			MSU		71Tr03	*
^{16}Be(γ,2n)^{14}Be	1350	100	840	30	−5.1	B					12Sp01	
	840	30				2					24Mo02	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{14}C(^{14}C,^{12}N)^{16}B	−48380	60	−48410	25	−0.5	o			Ber		95Bo10	
	−48378	60			−0.5	1	17	17 ^{16}B	Ber		00Ka21	
^{14}C(t,p)^{16}C	−3015	8	−3013	4	0.2	2			MSU		77Fo09	
	−3013	4			−0.1	2			LAl		78Se04	
^{14}C(^{3}He,p)^{16}N	4983	4	4978.2	2.3	−1.2	R			BNL		66Ga08	
^{14}C(^{3}He,p)^{16}N^{i}	−4951	7				2					68He03	*
^{14}N(t,p)^{16}N	4853	10	4840.3	2.3	−1.3	U			Ald		66He10	
^{14}C(^{3}He,n)^{16}O^{j}	−8100	8	−8104	4	−0.5	1	23	23 ^{16}O^{j}			70Ad01	*
^{16}O(d,α)^{14}N	3110.	3.5	3110.38743	0.00029	0.1	U			Wis		52Cr30	Y
	3119	5			−1.7	U			Ric		53Fa18	Y
	3110	6			0.1	U			Mex		64Ma.B	
	3113	6			−0.4	U			MIT		64Sp12	
^{14}N(^{3}He,p)^{16}O^{i}	2444	6	2447	4	0.5	1	54	54 ^{16}O^{i}			64Br08	*
^{14}N(d,γ)^{16}O^{j}	−1986.3	4.4	−1985	4	0.3	1	77	77 ^{16}O^{j}			72Ne10	
^{14}N(^{3}He,n)^{16}F	−963	40	−952	5	0.3	U			LAl		65Za01	
	−970	15			1.2	R			Har		68Ad03	
^{16}Ne(2p)^{14}O	1350	80	1430	12	1.0	U					08Mu13	
	1425	4			1.2	U					21Ji15	
	1461	10			−3.1	U					25Ch72	
	1430	12				3					averag	
^{16}B(γ,n)^{15}B	85	15	83	15	−0.1	1	95	83 ^{16}B			09Le02	
^{15}N(d,p)^{16}N	286	12	264.3	2.3	−1.8	U			CIT		55Pa50	Y
	269	10			−0.5	U			Pit		57Wa01	Y
	259	6			0.9	2			Mex		64Ma.B	
	267	8			−0.3	2			MIT		64Sp12	
	270	10			−0.6	U			Pen		66He10	
^{15}N(p,γ)^{16}O^{i}	−665.3	6.6	−669	4	−0.5	1	46	46 ^{16}O^{i}			57Ha99	
^{16}O(^{3}He,α)^{15}O	4920	10	4913.7	0.5	−0.6	U			Ald		59Hi68	Y
	4907	7			1.0	U			Ric		59Yo25	Y
^{16}F(p)^{15}O	527	7	531	5	0.5	1	59	58 ^{16}F			18Ch25	
	531	22			−0.0	U					23Ch22	*
^{16}N(β^-)^{16}O	10400	20	10420.9	2.3	1.0	U					59Al06	
^{16}O(^{3}He,t)^{16}F	−15430	10	−15431	5	−0.1	–			KVI		80Ja.A	
ave.	−15436	8			0.6	1	42	42 ^{16}F			average	
^{16}O(π^+,π^-)^{16}Ne	−27766	45	−27731	12	0.8	U					80Bu15	
	−27685	23			−2.0	U					20Ne-4	
*^{16}O(^{3}He,^{6}He)^{13}O	$M-A$ increased by 7 for more recent calibrator M−A(^{9}C)=28913(2)										AHW	**
*^{16}O(^{3}He,^{6}He)^{13}O	Recalibrated using their ^{12}C(^{3}He,^{6}He) result										AHW	**
*^{14}C(^{3}He,p)^{16}N^{i}	IT=9928(7), Q rebuilt with Ame1965										MMC121	**
*^{14}C(^{3}He,n)^{16}O^{j}	IT=22717(8), Q rebuilt with Ame1964										MMC121	**
*^{14}N(^{3}He,p)^{16}O^{i}	IT=12798(6), Q rebuilt										MMC121	**
*^{16}F(p)^{15}O	A systematic error of 20 keV was added to stat. error										HWJ247	**
^{17}O$_2$−^{28}Si D$_3$	−20968.3557	0.0014	−20968.3561	0.0013	−0.3	1	85	84 ^{17}O	FS1	1.0	10Mo29	
^{17}B−u	45970	860	46930	220	0.7	U			GA1	1.5	87Gi05	
	46830	180			0.4	2			TO2	1.5	88Wo09	
	47127	250			−0.5	2			GA3	1.5	91Or01	
^{17}Ne−^{22}Ne$_{.773}$	24373.27	0.38				2			MA8	1.0	08Ge07	
^{17}Na−u	37273	64				2				1.0	17Br07	*
^{17}O−^{16}O H	−3607.8961	0.0016	−3607.8951	0.0007	0.6	1	20	16 ^{17}O	FS1	1.0	10Mo29	
^{17}O(n,α)^{14}C	1817.2	3.5	1817.744	0.004	0.2	U					01Wa50	
^{14}C(α,n)^{17}O	−1819.07	2.0	−1817.744	0.004	0.7	U			Wis		56Sa06	Y
^{17}O(p,α)^{14}N	1200	17	1191.8737	0.0006	−0.5	U			MIT		64Sp12	
^{17}O(d,α)^{15}N	9818	12	9800.6042	0.0009	−1.4	U			Nob		54Pa39	Y

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{16}O(n,γ)^{17}O		4143.24	0.23	4143.0800	0.0008	−0.7	U					77Mc05	Z
		4143.06	0.13			0.2	U			Bdn		06Fi.A	
^{16}O(d,p)^{17}O		1915	8	1918.5137	0.0007	0.4	U			Ric		51Kl55	Y
		1918	4			0.1	U			MIT		57Br82	
		1918	3			0.2	U			Mex		61Ja23	
		1920	3			−0.5	U			MIT		64Sp12	
		1918.74	0.5			−0.5	U			Rez		90Pi05	*
^{16}O(n,γ)^{17}O^{i}		−6935.70	0.17				2					81Hi01	*
^{16}O(p,γ)^{17}F		600.35	0.28	600.27	0.25	−0.3	–			CIT		75Ro05	
^{16}O(d,n)^{17}F		−1626	4	−1624.30	0.25	0.4	U			Ric		51Bo49	Y
		−1624.6	0.5			0.6	–			Nvl		60Bo21	Z
^{16}O(p,γ)^{17}F	ave.	600.27	0.25	600.27	0.25	−0.0	1	100	100 ^{17}F			average	
^{16}O(p,γ)^{17}F^{i}		−10592.8	1.9				2					76Hi09	
^{16}O(^{3}He,2n)^{17}Ne		−22420	190	−22448.9	0.4	−0.2	U			BNL		67Es02	
^{17}F(β^{+})^{17}O		2770	6	2760.47	0.25	−1.6	U					54Wo23	
*^{17}Na−u	Authors says this is an upper limit $M-A$<34720(60) keV											17Br07	**
*^{17}Na−u	Represents ^{17}Na → 3p + ^{14}O, yielding $M-A$<34720(60) keV											GAu	**
*^{16}O(d,p)^{17}O	Estimated systematic error 0.5 added to statistical error 0.062 keV											AHW	**
*^{16}O(n,γ)^{17}O^{i}	Original Q =–6934.41(0.17) does not match original T=7373.31(0.18)											MMC129	**
C D_3−^{18}O		43145.72216	0.00088	43145.7214	0.0007	−0.9	–			FS1	1.0	09Re15	*
		43145.72116	0.00136			0.2	–			FS1	1.0	09Re15	*
	ave.	43145.7219	0.0007			−0.6	1	87	87 ^{18}O			average	
C_3−$^{18}O_2$		1680.7695	0.0038	1680.7757	0.0014	1.6	1	13	13 ^{18}O	FS1	1.0	09Re15	
^{18}F−u		943	85	937.3	0.5	−0.0	U				2.5	92Ge08	
^{18}Na−u		25969	54	27020	21	7.8	F				2.5	01Ze.A	*
		26882	183			0.8	U			1.0	1.0	04Ze05	*
^{18}Mg−u		45120.0	36.5				2			1.0	1.0	21Ji15	*
^{18}Ne−^{22}Ne$_{.818}$		12755.68	0.39	12755.7	0.4	−0.1	1	100	100 ^{18}Ne	MA8	1.0	04Bl20	
^{14}C(^{7}Li,^{3}He)^{18}N		−10170	60	−10117	19	0.9	U			Str		80Kr.A	
^{18}O(^{48}Ca,^{51}V)^{15}B		−21760	50	−21762	21	−0.0	–			Hei		78Bh02	
		−21768	25			0.2	–			Can		83Ho08	
	ave.	−21766	22			0.2	1	88	88 ^{15}B			average	
^{18}O(p,α)^{15}N		3954	9	3979.8003	0.0009	2.9	U			Nob		54Mi60	Y
		3964	10			1.6	U			Mex		64Ma.B	
^{18}O(d,α)^{16}N		4235	7	4244.1	2.3	1.3	R			CIT		55Pa50	Z
		4219	20			1.3	U			Mex		64Ma.B	
		4249	15			−0.3	U			Phi		66He10	
		4244	4			0.0	R			MIT		67Sp09	Z
^{16}O(^{3}He,p)^{18}F		2033	5	2032.1	0.5	−0.2	U			Ric		59Yo25	
		2055	5			−4.6	C			Mex		64Ma.B	
^{16}O(^{3}He,n)^{18}Ne		−3205	13	−3194.7	0.4	0.8	U			Nvl		61Du02	Y
		−3198	6			0.6	U			Ald		61To03	Y
		−3194.0	1.5			−0.5	U					94Ma14	
^{18}B(γ,n)^{17}B		5	5				3					10Sp02	*
^{18}O(^{48}Ca,^{49}Ti)^{17}C		−17465	35	−17476	17	−0.3	2			Hei		77No08	
		−17479	20			0.2	2			Can		82Fi10	
^{18}O(^{207}Pb,^{208}Po)^{17}C		−26870	220	−26797	17	0.3	U			ChR		79Ba31	
^{18}O(t,α)^{17}N		3872	15				2			LAl		60Ja13	
^{17}O(n,γ)^{18}O		8043.5	1.0	8045.3701	0.0010	1.9	U			Bdn		06Fi.A	
^{17}O(d,p)^{18}O		5820	10	5820.8039	0.0009	0.1	U			Nob		54Ah37	Y
		5820	10			0.1	U			Man		65Mo16	
^{17}O(p,γ)^{18}F		5603	3	5607.1	0.5	1.4	U			Str		73Se03	
		5606.2	0.6			1.5	1	60	60 ^{18}F	CIT		75Ro05	Z
^{18}Na(p)^{17}Ne		1270	170	1380	20	0.6	U					04Ze05	
		1230	150			1.0	U					12Mu05	
		1380	20				3					25Ni11	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{18}O(π^-,π^+)^{18}C	−26712	150	−26720	30	−0.1	U					78Se07	
^{18}O(^{48}Ca,^{48}Ti)^{18}C	−21434	30				2			Can		82Fi10	
	−21331	300	−21430	30	−0.3	U			Ors		82Na04	
^{18}N(β^-)^{18}O	13860	400	13896	19	0.1	U					64Ch19	
^{18}O(d,2p)^{18}N	−15270	100	−15338	19	−0.7	U			Brk		78De.A	
^{18}O(t,^{3}He)^{18}N	−13917	60	−13877	19	0.7	U			LAl		69St07	*
^{18}O(^{7}Li,^{7}Be)^{18}N	−14761	20	−14758	19	0.2	2			Can		83Pu01	
^{18}O(^{14}C,^{14}N)^{18}N	−13720	50	−13739	19	−0.4	2			Ors		80Na14	
^{18}F(β^+)^{18}O	1657	2	1655.9	0.5	−0.5	U					64Ho28	
^{18}O(p,n)^{18}F	−2451	4	−2438.3	0.5	3.2	B			Wis		50Ri59	Y
	−2436.97	0.73			−1.8	1	40	40 ^{18}F	Nvl		64Bo13	Z
	−2440.2	2.8			0.7	U					67Pr04	
^{18}Ne(β^+)^{18}F	4438	9	4444.5	0.6	0.7	U					63Fr10	
*C D_3-^{18}O	Respectively $CD_3^+/^{18}O^+$, $C_2D_6^+/^{18}O^{2+}$ considered independent										GAu	**
*^{18}Na−u	F : results distrusted (see also ^{15}F and ^{19}Mg)										GAu	**
*^{18}Na−u	Other interpretation : D_M=25969(172) μu, $M-A$=24190(160) keV										04Ze05	**
*^{18}Mg−u	from decay energy 4865(34) keV ^{18}Mg → 4p + ^{14}O										HWJ229	**
*^{18}B(γ,n)^{17}B	Decay energy <10 keV, derived from scattering length a_s <−50 fm										10Sp02	**
*^{18}O(t,^{3}He)^{18}N	From Q =14038(30), reinterpreted as mainly to (2$^-$) level at 114.90 keV										Ens967	**
^{28}Si H_3-C ^{19}F	1998.4687	0.0022	1998.4680	0.0009	−0.3	1	18	14 ^{19}F	FS1	1.0	09Re15	
^{19}B−u	64166	376				2			GA8	1.5	12Ga45	
^{19}C−u	34680	260	34800	110	0.3	o			TO1	1.5	86Vi09	
	35370	450			−0.8	U			GA1	1.5	87Gi05	
	35180	130			−2.0	o			TO2	1.5	88Wo09	
	35506	253			−1.9	U			GA3	1.5	91Or01	
C D_4-H ^{19}F	50178.88	0.05	50178.9175	0.0009	0.5	U			B08	1.5	75Sm02	
^{19}Mg−u	35470	270	34216	11	−1.9	F				2.5	01Ze.A	*
^{13}C D_3-^{19}F	47257.00669	0.00091	47257.0068	0.0008	0.1	1	86	86 ^{19}F	FS1	1.0	09Re15	
^{19}Ne−^{22}Ne$_{.864}$	9323.92	0.33	9324.17	0.17	0.8	2			MA8	1.0	04Bl20	
	9324.26	0.20			−0.5	2			MA8	1.0	08Ge07	
^{19}F(p,α)^{16}O	8115	10	8113.6119	0.0008	−0.1	U			CIT		50Ch53	Y
	8115	10			−0.1	U			CIT		57Yo04	Y
	8122	9			−0.9	U			MIT		64Sp12	
^{17}O(t,p)^{19}O	3524	7	3519.2	2.6	−0.7	R			Man		65Mo19	
^{19}F(d,α)^{17}O	10060	12	10032.1256	0.0010	−2.3	U			MIT		64Sp12	
^{19}Mg(2p)^{17}Ne	750	50	794	10	0.9	o					07Mu15	
	760	60			0.6	U					12Mu05	
	980	170			−1.1	U					18Xu04	*
	794	10				3					24Ni16	
^{18}C(n,γ)^{19}C	530	120	580	90	0.4	3					99Na27	*
	650	150			−0.5	3					01Ma08	*
^{18}O(^{18}O,^{17}F)^{19}N	−19374	50	−19372	16	0.0	2			Ors		81Na.A	
	−19334	35			−1.1	2			Can		89Ca25	
^{18}O(^{48}Ca,^{47}Sc)^{19}N	−16540	20	−16528	16	0.6	2			Can		83Ho08	
^{18}O(^{208}Pb,^{207}Bi)^{19}N	−18440	150	−18332	17	0.7	U			ChR		79Ba31	
^{18}O(d,p)^{19}O	1727	8	1731.1	2.6	0.5	o			Nob		54Mi89	Y
	1732	8			−0.1	2			CIT		54Th30	
	1731	5			0.0	2			Nob		57Ah19	Y
	1733	6			−0.3	2			Mex		64Ma.B	
	1727	5			0.8	2			MIT		64Sp12	Z
	1734	10			−0.3	U			Man		65Mo16	
^{19}F(^{3}He,α)^{18}F	10166	15	10145.7	0.5	−1.4	U			Ald		59Hi67	Y
^{19}Na(p)^{18}Ne	160	110	323	11	1.5	U					04Ze05	
	328	22			−0.2	1	23	23 ^{19}Na			10Mu12	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{19}O(β^-)^{19}F	4800	12	4820.3	2.6	1.7	U					59Al06	
^{19}Ne(β^+)^{19}F	3262	10	3239.49	0.16	−2.3	U					60Wa04	
^{19}F(p,n)^{19}Ne	−4021.3	4.7	−4021.83	0.16	−0.1	U			Ric		55Ma84	
	−4019.6	1.4			−1.6	U			Ric		61Be13	Z
	−4021.1	1.0			−0.7	U			Zur		61Ry04	Z
	−4020.7	0.8			−1.4	U					66Ma60	
	−4019.6	0.7			−3.2	B					69Ov01	Z
^{19}F(^{3}He,t)^{19}Ne^i	−10759	9				2					98Ut02	*
*^{19}Mg−u	F : results distrusted (see also ^{15}F and ^{18}Na)										GAu	**
*^{19}Mg(2p)^{17}Ne	Symmetrized from 870(+240–70); Confirmed previous result in 12Mu05										HWJ192	**
*^{18}C(n,γ)^{19}C	From Coulomb dissociation cross sections and angular distribution										99Na27	**
*^{18}C(n,γ)^{19}C	From momentum distribution following one neutron removal										01Ma08	**
*^{19}F(^{3}He,t)^{19}Ne^i	rebuilt from E_x =7500(9); with references from Ame1995; recalibrated +1keV										MMC141	**
^{20}C−u	39940	1210	40260	250	0.2	U			GA1	1.5	87Gi05	
	40360	240			−0.3	2			TO2	1.5	88Wo09	
	40165	491			0.1	2			GA3	1.5	91Or01	
	40420	550			−0.2	2			GA5	1.5	99Sa.A	
	40108	290			0.4	2			GA8	1.5	12Ga45	
^{20}N−u	23230	280	23370	80	0.3	U			TO1	1.5	86Vi09	
	23210	150			0.7	2			GA1	1.5	87Gi05	
	23380	130			−0.1	2			TO2	1.5	88Wo09	
	23397	69			−0.3	2			GA3	1.5	91Or01	
C D_4−^{20}Ne	63966.9329	0.0026	63966.94259	0.00010	3.7	B			MI1	1.0	95Di08	
^{20}Ne−u	−7559.309	0.090	−7559.83121	0.00008	−2.3	U			OH1	2.5	93Ma.A	
	−7559.814	0.014			−1.2	U			ST2	1.0	02Bf02	
	−7559.83122	0.00008			0.1	1	100	100 ^{20}Ne	Hep	1.0	23He18	*
^{20}Al−u	43275	118	45020#	220#	5.9	D				2.5	25Xu03	*
O D_2−^{20}Ne	30677.497	0.067	30678.0059	0.0005	3.0	B			OH1	2.5	93Go38	
^{20}Mg−^{23}Na$_{.870}$	27663.8	2.0	27663.8	2.0	−0.0	1	100	100 ^{20}Mg	TT1	1.0	14Ga20	
^{20}Ne−^{22}Ne$_{.909}$	270.89	0.43	271.1144	0.0024	0.5	U			MA8	1.0	04Bl20	
	271.16	0.20			−0.2	U			MA8	1.0	08Ge07	
^{20}Ne(^{3}He,^{8}Li)^{15}F	−29960	200	−29623	14	1.7	U			MSU		78Be26	
	−29730	180			0.6	U			Brk		78Ke06	
^{20}Ne^i(α)^{16}O	5548.8	6.3	5542.6	2.0	−1.0	U					73To08	
^{20}Ne(α,^{8}He)^{16}Ne	−60150	80	−60242	12	−1.2	U			Brk		78Ke06	
	−60197	23			−2.0	R			Tex		83Wo01	
^{20}Ne(^{3}He,^{6}He)^{17}Ne	−26188	50	−26203.3	0.4	−0.3	U			Brk		70Me11	*
	−26158	32			−1.4	F					98Gu10	*
^{18}O(^{48}Ca,^{46}Sc)^{20}N	−25873	60	−25010	80	14.3	B			Can		89Or03	*
^{18}O(t,p)^{20}O	3086	15	3081.9	0.9	−0.3	U			LAl		60Ja13	
	3076	10			0.6	U			Ald		62Hi06	
	3082.4	1.9			−0.3	2			Str		82An12	
	3081.7	1.0			0.2	2			Str		85An17	
^{18}O(^{3}He,p)^{20}F	6875.2	1.5	6876.895	0.030	1.1	U			NDm		70Ro06	
^{18}O(^{3}He,p)^{20}F^i	356.0	3.0				2					76Mi01	*
^{20}Ne(d,α)^{18}F	2795	9	2795.8	0.5	0.1	U			Nob		54Mi61	Y
	2766	20			1.5	U			Mex		64Ma.B	
	2790	10			0.6	U					75Bo59	
^{20}B(γ,n)^{19}B	1560	150				3					18Le18	
^{19}F(n,γ)^{20}F	6601.1	0.3	6601.336	0.030	0.8	U			Utr		68Sp01	
	6601.29	0.14			0.3	2			ILn		83Hu12	Z
	6601.32	0.05			0.3	2			MMn		87Ke09	Z
	6601.35	0.04			−0.3	2			ORn		96Ra04	
	6601.34	0.13			−0.0	2			Bdn		06Fi.A	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{19}F(d,p)^{20}F	4377	7	4376.770	0.030	−0.0	U			MIT		64Sp12	
	4377.7	0.9			−1.0	U			NDm		70Ro06	
^{19}F(p,γ)^{20}Ne^{j}	−3889.4	2.7				2					67Bl19	
^{20}Ne(^{3}He,α)^{19}Ne	3750	13	3712.32	0.16	−2.9	U			MIT		64Sp12	
^{20}Na^{i}(p)^{19}Ne	4381	50	4307.9	1.2	−1.5	U			Brk		79Mo02	
	4332	16			−1.5	U			MSU		92Go10	
	4326	30			−0.6	U			Lis		95Pi03	
^{20}F(β^-)^{20}Ne	7053	13	7024.475	0.030	−2.2	U					54Wo23	*
	7050	15			−1.7	U					59Al06	*
	7032	6			−1.3	U					76Ge08	*
	7026.9	1.8			−1.3	U					87Va20	*
	7019.8	1.7			2.7	U					89He11	*
^{20}Ne^{i}(IT)^{20}Ne	10274	3	10272.5	2.0	−0.5	2					76In06	
	10271.2	2.7			0.5	2					77Fi08	
^{20}Na(β^+)^{20}Ne	13906	40	13892.5	1.1	−0.3	U					67Su05	
^{20}Ne(p,n)^{20}Na	−14672.1	7.	−14674.8	1.1	−0.4	U					71Wi07	Z
^{20}Ne(^{3}He,t)^{20}Na−^{36}Ar()^{36}K	−1078.06	1.06	−1078.1	1.1	−0.0	1	100	100 ^{20}Na	Mun		10Wr01	
^{20}Na^{i}(IT)^{20}Na	6498.4	0.5				2					15Gl03	
*^{20}Ne−u	From frequency ratio ^{20}Ne^{10+}/^{12}C^{6+}=1.000378141802(4)										23He18	**
*^{20}Al−u	derived from Q =1.93(+0.12−0.10) MeV for ^{20}Al->^{17}N×10^{3}p										25Xu03	**
*^{20}Al−u	Trends in Mass surface TMS suggest ^{20}Al 1600keV less bound										HWJ261	**
*^{20}Ne(^{3}He,^{6}He)^{17}Ne	Original $M-A$=16479(50) but revised calibrator M(^{9}C)=28910.2(2.1)										AHW	**
*^{20}Ne(^{3}He,^{6}He)^{17}Ne	F : calibrated with ^{24}Mg(^{3}He,^{6}He) from reference for excitation energies										92Ku02	**
*	no details given. No correction possible										GAu	**
*^{18}O(^{48}Ca,^{46}Sc)^{20}N	Probably to excited levels in ^{20}N and ^{46}Sc										GAu	**
*^{18}O(^{3}He,p)^{20}F^{i}	IT=6519.4(3.0), Q rebuilt with Ame1971										MMC122	**
*^{20}F(β^-)^{20}Ne	E_{β^-}=5419(13) 5416(15) 5398(6) 5392.3(1.8) 5386.1(1.7) respectively,										GAu	**
*	to 2$^+$ level at 1633.674 keV										Ens992	**
^{21}N−u	26580	200	27090	140	1.7	U			TO1	1.5	86Vi09	
	27060	190			0.1	2			GA1	1.5	87Gi05	
	26930	210			0.5	2			TO2	1.5	88Wo09	
	27162	131			−0.4	2			GA3	1.5	91Or01	
^{21}Na−^{39}K$_{.538}$	17180.51	0.29	17180.37	0.05	−0.5	o			MA8	1.0	04Mu26	
	17180.51	0.29			−0.3	U			Ma8	1.5	08Mu05	*
^{21}Al^{p}−u	34391	54	34410	50	0.4	1	83	83 ^{21}Al^{p}	MSU	1.0	25Ni12	*
H$_3$ ^{18}O−^{21}Ne	28787.76	0.25	28788.03	0.04	1.1	U			CP1	1.0	04Sa53	*
^{21}Na−^{23}Na$_{.913}$	6995.25	0.32	6995.10	0.05	−0.5	o			MA8	1.0	04Mu26	
	6995.25	0.32			−0.3	U			Ma8	1.5	08Mu05	
^{21}Mg−^{23}Na$_{.913}$	21046.41	0.81				2			TT1	1.0	14Ga20	
^{21}Ne−^{22}Ne$_{.955}$	2073.82	0.40	2073.91	0.04	0.2	U			MA8	1.0	04Bl20	
	2074.04	0.26			−0.5	U			MA8	1.0	08Ge07	
^{21}Na−^{21}Ne	3808.017	0.097	3807.774	0.019	−2.5	U			MS1	1.0	15Ei01	
	3807.774	0.019			0.0	1	100	100 ^{21}Na	MA8	1.0	19Ka30	*
^{21}Na−^{20}Na	−9732	50	−9699.9	1.2	0.3	U			CR1	2.5	89Sh10	
^{21}Na^{i}(α)^{17}F	2397	10	2413	4	1.6	o					15Lu12	
	2409.1	12.4			0.3	U					24Je01	*
^{18}O(^{18}O,^{15}O)^{21}O	−12574	70	−12483	12	1.3	U			Ors		78Na02	
	−12499	20			0.8	2			Can		89Ca25	
^{18}O(^{64}Ni,^{61}Ni)^{21}O	−11713	15	−11722	12	−0.6	2			Dar		85Wo01	
^{18}O(^{208}Pb,^{205}Pb)^{21}O	−6860	75	−6823	12	0.5	U			ChR		79Ba31	
^{21}B(γ,2n)^{19}B	2470	190				3					18Le18	
^{19}F(t,p)^{21}F	6221.0	1.8				2			Str		84An17	
^{19}F(^{3}He,p)^{21}Ne	11911	15	11886.58	0.04	−1.6	U			Ald		59Hi75	Y
^{20}Ne(n,γ)^{21}Ne	6760.8	1.5	6761.16	0.04	0.2	U					70Se14	
	6761.16	0.04			0.1	–			MMn		86Pr05	Z
	6761.19	0.14			−0.2	–			Bdn		06Fi.A	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{20}Ne(d,p)^{21}Ne		4531	9	4536.60	0.04	0.6	U				Nob		55Ah41	Y
		4532	6			0.8	U				Mex		64Ma.B	
		4534	7			0.4	U				MIT		64Sp12	
^{20}Ne(n,γ)^{21}Ne	ave.	6761.16	0.04	6761.16	0.04	−0.0	1	100	100	^{21}Ne			average	
^{20}Ne(p,γ)^{21}Na		2431.2	0.7	2431.90	0.04	1.0	U						69Bl03	Z
^{20}Ne(p,γ)^{21}Na^{i}		−6547.9	14.3	−6542	4	0.4	U						81Fe05	
^{21}Na^{i}(p)^{20}Ne		6543	4	6542	4	−0.1	2						73Se08	*
		6542	10			0.0	o						15Lu12	*
		6561	30			−0.6	U						18Wa20	*
		6539	10			0.3	2						24Je01	*
^{21}Al(p)^{20}Mg		1170	90	1284	14	1.3	U						24Ko22	*
		1285	14			−0.1	1	99	99	^{21}Al			25Ph02	*
^{21}O(β^-)^{21}F		8150	175	8110	12	−0.2	U						81Al07	
^{21}Na(β^+)^{21}Ne		3522	30	3546.919	0.018	0.8	U						52Sc15	
		3532	20			0.7	U						60Wa04	
^{21}Al^{p}(IT)^{21}Al		6090	110	6000	50	−0.8	1	18	17	^{21}Al^{p}	MSU		25Ni12	
*^{21}Na−^{39}K$_{.538}$	CF=1.5 for prelim. results; not trusted within given uncertainties												GAu	**
*^{21}Al^{p}−u	derived from Q =4.85(0.05) MeV for ^{21}Al^{p}->^{18}Ne+3p												25Ni12	**
*H$_3$ ^{18}O−^{21}Ne	D_M=28787.78(0.25) corrected –0.02 keV for molecular and ionization												04Sa53	**
*^{21}Na−^{21}Ne	Electron binding energies not considered by authors												HWJ201	**
*^{21}Na^{i}(α)^{17}F	decay to the gs. via s-wave emission												24Je01	**
*^{21}Na^{i}(p)^{20}Ne	average Q_p=6548(4), 4904(4) to ground state and 2$^+$ level at 1634 keV												HWJ25a	**
*^{21}Na^{i}(p)^{20}Ne	Q_p=6537(10), 4913(10) to ground state and 2$^+$ level at 1633.674 keV												15Lu12	**
*^{21}Na^{i}(p)^{20}Ne	Q_p=6580(30), 4907(25) to ground state and 2$^+$ level at 1633.674 keV												18Wa20	**
*^{21}Na^{i}(p)^{20}Ne	average E_α=6230(10), 4670(10), 2180(10), 1500(10), 870(10) to ground state and												HWJ25a	**
*^{21}Na^{i}(p)^{20}Ne	1634, 4248, 4967, 5621 keV levels												HWJ25a	**
*^{21}Al(p)^{20}Mg	symmetrized from Q_p=1150(+100−70) keV												24Ko22	**
*^{21}Al(p)^{20}Mg	from Q_p=1285(2,stat)(14,syst)												25Ph02	**
^{22}C−u		57585	408	57550	250	−0.1	U				GA8	1.5	12Ga45	
^{22}N−u		32990	790	34100	220	0.9	U				GA1	1.5	87Gi05	
		34340	250			−0.6	2				TO2	1.5	88Wo09	
		34683	389			−1.0	2				GA3	1.5	91Or01	
		34240	320			−0.3	2				GA5	1.5	99Sa.A	
		33398	279			1.7	2				GA8	1.5	12Ga45	
^{22}O−u		9842	81	9970	60	1.0	R				GA3	1.5	91Or01	
^{22}Ne−u		−8614.885	0.019	−8614.9018	0.0026	−0.9	U				ST2	1.0	02Bf02	
		−8614.9018	0.0026			−0.0	1	100	100	^{22}Ne	1.0	1.0	23He18	*
^{22}Na−^{39}K$_{.564}$		14907.33	0.30	14907.08	0.14	−0.8	o				MA8	1.0	04Mu26	
		14907.33	0.30			−0.5	1	10	10	^{22}Na	Ma8	1.5	08Mu05	
^{22}Mg−^{39}K$_{.564}$		20040.33	0.35	20040.12	0.18	−0.6	o				MA8	1.0	04Mu26	
		20040.33	0.35			−0.4	1	11	11	^{22}Mg	Ma8	1.5	08Mu05	
^{22}Al−u		19434	11	19423.09	0.29	−1.0	U				LZ2	1.0	24Su04	
		19442	8			−2.4	U				LZ2	1.0	25Ch67	*
^{22}Si−u		34168	58				2				LZ2	1.0	25Xi05	
O H−^{22}Ne$_{.773}$		9398.87	0.19	9398.9701	0.0021	0.5	U				MA8	1.0	08Ge07	
^{22}Na−^{24}Mg$_{.917}$		8153.64	0.31	8154.31	0.14	2.2	o				MA8	1.0	04Mu26	
		8153.64	0.31			1.4	1	9	9	^{22}Na	Ma8	1.5	08Mu05	
^{22}Na−^{23}Na$_{.957}$		4228.11	0.29	4228.34	0.14	0.8	o				MA8	1.0	04Mu26	
		4228.11	0.29			0.5	–				Ma8	1.5	08Mu05	
		4228.35	0.25			−0.1	–				TT1	1.0	17Re10	
	ave.	4228.29	0.22			0.2	1	43	43	^{22}Na			average	
^{22}Mg−^{23}Na$_{.957}$		9361.33	0.25	9361.37	0.18	0.2	1	49	49	^{22}Mg	TT1	1.0	17Re10	
^{22}Al−^{23}Na$_{.957}$		29215.5	1.3	29213.89	0.29	−1.2	2				MS1	1.0	24Ca05	
		29213.80	0.30			0.3	2				MS2	1.0	24Ca05	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{22}Na−^{22}Ne		3052.75	0.33	3052.44	0.14	−0.9	1	19	19 ^{22}Na	CP1	1.0	04Sa53	*
^{22}Mg−^{22}Ne		8185.79	0.73	8185.48	0.18	−0.4	U			CP1	1.0	04Sa53	*
^{22}Mg−^{22}Na		5132.99	0.34	5133.04	0.18	0.1	o			MA8	1.0	04Mu26	
		5132.99	0.34			0.1	–			Ma8	1.5	08Mu05	
		5133.05	0.26			−0.0	–			TT1	1.0	17Re10	
	ave.	5133.04	0.23			0.0	1	59	39 ^{22}Mg			average	
^{22}Ne−^{20}Ne		−1056.415	0.290	−1055.0706	0.0026	1.9	U			OH1	2.5	93Go38	
^{18}O(^{18}O,^{14}O)^{22}O		−19060	100	−18860	60	2.0	2			Can		76Hi10	
^{18}O(^{208}Pb,^{204}Pb)^{22}O		−6710	180	−6700	60	0.0	2			ChR		79Ba31	
^{22}Mg^{i}(α)^{18}Ne		5885	40	5904	4	0.5	U			Bor		97Bl03	*
		5904	8			−0.0	–			Bor		06Ac04	*
		5917	10			−1.3	–			Lzr		21Wu10	*
	ave.	5909	6			−0.9	1	43	43 ^{22}Mg^{i}			average	
^{19}F(α,p)^{22}Ne		1674	11	1673.2303	0.0026	−0.1	U			MIT		64Sp12	
^{19}F(α,n)^{22}Na		−1958	10	−1952.45	0.13	0.6	U			Duk		60Wi07	Y
^{22}C(γ,2n)^{20}C		−200	120	−35	20	1.4	o					11Ya25	*
		−110	60			1.3	U					12Fo04	*
		−35	20				3					13Mo12	*
^{20}Ne(^{3}He,n)^{22}Mg		197	25	217.97	0.16	0.8	U			Har		68Ad03	
		209	11			0.8	U			CIT		70Mc06	
^{22}Mg^{i}(2p)^{20}Ne		6098	13	6110	4	0.9	–					06Ac04	*
		6090	34			0.6	o			Lzr		18Wa24	*
		6120	11			−0.9	–			Lzr		21Wu10	*
	ave.	6111	8			−0.1	1	24	24 ^{22}Mg^{i}			average	
^{22}Ne(t,α)^{21}F		4545	10	4547.8	1.8	0.3	U			LAl		61Si03	Y
^{21}Ne(n,γ)^{22}Ne		10364.4	0.3	10364.27	0.04	−0.4	U			MMn		86Pr05	Z
		10363.9	0.5			0.7	U			Bdn		06Fi.A	
^{21}Ne(d,p)^{22}Ne		8152	11	8139.70	0.04	−1.1	U			CIT		52Mi54	Y
^{21}Ne(p,γ)^{22}Na		6739.0	0.7	6738.59	0.14	−0.6	U					70An06	*
^{22}Mg^{i}(p)^{21}Na		8547	15	8542	4	−0.3	–			Brk		82Ca16	*
		8531	8			1.4	–			Lzr		21Wu10	*
	ave.	8535	7			1.1	1	34	34 ^{22}Mg^{i}			average	
^{22}F(β^-)^{22}Ne		11000	150	10818	12	−1.2	U					73Gu05	
		10950	120			−1.1	U			ANB		74Da02	
^{22}Ne(t,^{3}He)^{22}F		−10788	33	−10799	12	−0.3	2					69St07	*
		−10794	18			−0.3	2			Dar		88Cl04	*
^{22}Ne(^{7}Li,^{7}Be)^{22}F		−11691	20	−11680	12	0.5	2			Can		89Or04	*
^{22}Na(β^+)^{22}Ne		2842.2	0.5	2843.33	0.13	2.3	U					68Be35	*
		2840.4	1.5			2.0	U					68We02	*
		2841.5	1.0			1.8	U					72Gi17	*

*^{22}Ne−u	derived from of g-factor calculation, theory dependent	23He18 **
*^{22}Al−u	combined results with 24Yu13	25Ch67 **
*^{22}Na−^{22}Ne	D_M=3052.79(0.33) μu, $M-A$=–5181.06(0.31) keV; corrected –0.04 keV for	04Sa53 **
*	ion-ion interaction	04Sa53 **
*^{22}Mg−^{22}Ne	D_M=8185.84(0.73) μu, $M-A$=–399.65(0.68) keV; corrected –0.05 keV for	04Sa53 **
*	ion-ion interaction	04Sa53 **
*^{22}Mg^{i}(α)^{18}Ne	E_α=3270(40) to 2$^+$ level at 1887.3 keV	Ens967 **
*^{22}Mg^{i}(α)^{18}Ne	Q_α=4017(8) to 2$^+$ level at 1887.3 keV	Ens967 **
*^{22}Mg^{i}(α)^{18}Ne	Q_α=4030(10) to 2$^+$ level at 1887.3 keV	21Wu10 **
*^{22}C(γ,2n)^{20}C	From upper limit S_{2n}<400	GAu **
*^{22}C(γ,2n)^{20}C	From upper limit S_{2n}<220	GAu **
*^{22}C(γ,2n)^{20}C	The two items are estimates derived from the experimental result of reference	10Ta04 **
*^{22}C(γ,2n)^{20}C	From upper limit S_{2n}<70	GAu **
*^{22}Mg^{i}(2p)^{20}Ne	Original Q_{2p}=4464(8) to 1633.674(0.015) level	06Ac04 **
*^{22}Mg^{i}(2p)^{20}Ne	Estimated systematic 10 keV uncertainty added	16Ma.A **
*^{22}Mg^{i}(2p)^{20}Ne	Q_{2p}=4460(30) to 1630(15) keV level	18Wa24 **
*^{22}Mg^{i}(2p)^{20}Ne	Q_{2p}=6106(16), 4495(11) to gs., 2$^+$ level at 1633 keV	21Wu10 **
*^{21}Ne(p,γ)^{22}Na	T=701.8(0.5) to 1$^+$ level at 7408.6(0.5) keV	Ens157 **
*^{21}Ne(p,γ)^{22}Na	Reanalysis using E(exc) for lower levels of reference	90En08 **
*^{22}Mg^{i}(p)^{21}Na	E_p=8149(21), 7839(15) to 3/2$^+$ ground state, 5/2$^+$ level at 331.90 keV	Ens04c **
*^{22}Mg^{i}(p)^{21}Na	Q_p=8516(18),8205(17),6804(19),5705(8) to ground state, 5/2$^+$ level at 331.90 keV	21Wu10 **
*	7/2$^+$ level at 1716 keV, 9/2$^+$ level at 2829 keV	21Wu10 **

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{22}$Ne(t,^{3}He)^{22}F	Original value –10834(30) re-calculated from Q to										GAu	**
*	(2^+) level at 709.1, 1^+ at 1627.1 and 1^+ at 2571.7 keV										Ens157	**
$*^{22}$Ne(t,^{3}He)^{22}F	Original value –10836(12) re-calculated										GAu	**
$*^{22}$Ne(^{7}Li,^{7}Be)^{22}F	Q =–12400(20) to (2^+) level at 709.1 keV										Ens157	**
$*^{22}$Na(β^+)^{22}Ne	E_{β^+}=545.7(0.5) 543.9(1.5) 545(1) respectively, to 2^+ level at 1274.537 keV										Ens157	**
^{23}N−u		37110	2000	39420	450	0.8	o			GA5	1.5	99Sa.A	
		39378	923			0.0	U			GA7	1.5	07Ju03	
		39421	301				2			GA8	1.5	12Ga45	
^{23}O−u		15700	320	15700	130	−0.0	o			TO1	1.5	86Vi09	
		15860	320			−0.3	o			GA1	1.5	87Gi05	
		15700	150			−0.0	2			TO2	1.5	88Wo09	
		15621	186			0.3	o			GA3	1.5	91Or01	
		15695	107			0.0	2			GA7	1.5	07Ju03	
^{23}F−u		3530	210	3530	40	−0.0	U			TO1	1.5	86Vi09	
		3553	43			−0.4	–			GT1	1.5	04Ma.A	
		3503	48			0.5	–			LZ1	1.0	15Xu14	
	ave.	3520	40			0.2	1	86	86 ^{23}F			average	
^{23}Na−u		−10230.721	0.0037	−10230.7180	0.0019	0.8	–			MI2	1.0	99Br47	
		−10230.716	0.0048			−0.4	–			MI2	1.0	99Br47	
		−10230.7172	0.0026			−0.3	–			FS1	1.0	10Mo30	
	ave.	−10230.7181	0.0019			0.0	1	100	100 ^{23}Na			average	
^{23}Si−u		25268	128	25081	6	−1.5	o			LZ2	1.0	24Yu13	
		25083	17			−0.1	1	12	12 ^{23}Si	LZ2	1.0	25Xi05	
		25122	53			−0.8	o			LZ2	1.0	25Ch67	*
^{23}Ne−^{22}Ne$_{1.045}$		3469.59	0.37	3469.46	0.11	−0.3	U			MA8	1.0	04Bl20	
^{23}Mg−^{23}Na		4354.80	0.83	4354.49	0.03	−0.4	U			JY1	1.0	09Sa38	
		4354.66	0.17			−1.0	U			TT1	1.0	14Sc09	
		4354.487	0.034				2			MA8	1.0	19Ka30	
^{23}Al−^{23}Na		17475.07	0.37				2			JY1	1.0	09Sa38	
^{23}Si−^{23}Na		35311.8	6.2	35312	6	0.0	1	88	88 ^{23}Si	MS1	1.0	25Ma41	
^{23}Na(p,α)^{20}Ne		2377	3	2376.1394	0.0018	−0.3	U			Wis		53Do04	Y
		2373	8			0.4	U			MIT		64Sp12	
^{23}Na(d,α)^{21}Ne		6911	9	6912.74	0.04	0.2	U			Mex		64Ma.B	
		6909	10			0.4	U			MIT		64Sp12	
^{23}Al^i(2p)^{21}Na		6000	64	6080	40	1.2	1	44	44 ^{23}Al^i			18Wa05	*
^{22}Ne(^{18}O,^{17}F)^{23}F		−14080	90	−14040	30	0.4	1	14	14 ^{23}F	Can		89Or04	
^{22}Ne(n,γ)^{23}Ne		5200.2	2.0	5200.65	0.10	0.2	U					70Se14	
		5200.65	0.12			−0.0	2			MMn		86Pr05	Z
		5200.64	0.20			0.0	2			Bdn		06Fi.A	
^{22}Ne(d,p)^{23}Ne		2967	8	2976.08	0.10	1.1	U			Nob		54Ah20	Y
		2971	9			0.6	o			MIT		60Fr04	
		2974	6			0.3	U			Mex		64Ma.B	
		2968	7			1.2	U			MIT		64Sp12	
^{22}Ne(p,γ)^{23}Na		8794.0	1.5	8794.0940	0.0030	0.1	U					71Pi08	Z
		8794.26	0.17			−1.0	U					89Ba42	Z
^{22}Ne(p,γ)^{23}Na^j		−10796.3	2.0				2					85Ev01	*
^{23}Al^i(p)^{22}Mg		11644	57	11580	40	−1.1	1	56	56 ^{23}Al^i	Bor		97Bl04	*
^{23}F(β^-)^{23}Ne		8510	170	8440	30	−0.4	U					74Go17	
^{23}Ne(β^-)^{23}Na		4383	8	4375.79	0.10	−0.9	U					63Ca06	
^{23}Mg(β^+)^{23}Na		4121	12	4056.18	0.03	−5.4	B					63Fr10	
^{23}Na(p,n)^{23}Mg		−4832	10	−4838.53	0.03	−0.7	U			Oak		55Ki28	Z
		−4836.5	6.			−0.3	U			Ric		58Bi41	Y
		−4848.0	7.			1.4	U			ChR		58Go77	Y
		−4835.8	2.5			−1.1	U			Har		62Fr09	Z
		−4843.2	5.1			0.9	U			Tkm		63Ok01	Z
$*^{23}$Si−u	combined results with 24Yu13											25Ch67	**
$*^{23}$Al^i(2p)^{21}Na	Also Q_{2p}=5857(66) keV to $5/2^+$ level at 331.9 keV											18Wa05	**
$*^{22}$Ne(p,γ)^{23}Na^j	Original E_{res}=–10793.6(2.0) is a typo											16Ma.A	**
$*^{23}$Al^i(p)^{22}Mg	Q_p=11620(100), 10410(70) to ground state and 2^+ level at 1247.02 keV											Ens157	**
*	also Q_{2p}=6180(100), 5860(100) to ground state and 331.90 level in ^{21}Na											97Bl04	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{24}O−u	20080	1070	19860	180	−0.1	o			GA1	1.5	87Gi05	
	20000	500			−0.2	U			TO2	1.5	88Wo09	
	20659	442			−1.2	o			GA3	1.5	91Or01	
	20460	340			−1.2	o			GA5	1.5	99Sa.A	
	19861	118				2			GA7	1.5	07Ju03	
^{24}F−u	8070	170	8100	100	0.1	U			TO1	1.5	86Vi09	
	8450	240			−1.0	U			GA1	1.5	87Gi05	
	8135	86			−0.3	2			GA3	1.5	91Or01	
	8030	120			0.4	2			TO4	1.5	91Zh24	
^{24}Ne−u	−6391.8	7.4	−6389.4	0.6	0.3	U			TR1	1.0	23Ja05	
^{24}Mg−H_{24}	−202759.080	0.014	−202759.076	0.014	0.3	1	98	98 ^{24}Mg	ST2	1.0	03Be02	
^{24}Si−u	11544.7	4.0				2			MS1	1.0	22Pu01	*
^{24}Ne−^{22}Ne$_{1.091}$	3009.49	0.55				2			MA8	1.0	04Bl20	
^{24}Mg−^{23}Na$_{1.043}$	−4287.23	0.32	−4287.672	0.014	−0.9	U			Ma8	1.5	08Mu05	
^{24}Al−^{23}Na$_{1.043}$	10618.18	0.25	10618.24	0.24	0.2	1	96	96 ^{24}Al	TT1	1.0	15Ch58	
^{24}Mg(p,^{6}He)^{19}Na	−37213	70	−37166	11	0.7	U			Brk		69Ce01	
^{24}Mg(^{3}He,^{8}Li)^{19}Na	−32876	12	−32878	11	−0.1	1	77	77 ^{19}Na	MSU		75Be38	
^{24}Mg(α,^{8}He)^{20}Mg	−60900	210	−60596.0	1.9	1.4	U					74Ro17	
	−60677	27			3.0	B			Tex		76Tr03	
^{24}Mg(^{3}He,^{6}He)^{21}Mg	−27488	40	−27498.3	0.8	−0.3	U			Brk		70Me11	
	−27512	18			0.8	U			MSU		71Tr03	
^{22}Ne(t,p)^{24}Ne	5587	10	5587.8	0.5	0.1	U			LAl		61Si03	Z
^{24}Mg(d,α)^{22}Na	1955	12	1958.63	0.13	0.3	U			MIT		64Sp12	
^{24}Mg(p,t)^{22}Mg	−21194	3	−21194.41	0.16	−0.1	U			MSU		74Ha02	
	−21198.3	1.5			2.6	U			MSU		74No07	
	−21193.9	1.0			−0.5	U			Yal		05Pa31	
^{23}Na(n,γ)^{24}Na	6959.50	0.12	6959.365	0.016	−1.1	o			BNn		74Gr37	Z
	6959.42	0.07			−0.8	2			BNn		80Gr12	*
	6959.67	0.14			−2.2	U			ILn		83Hu11	Z
	6959.38	0.08			−0.2	U			Ptn		83Ti02	
	6959.44	0.05			−1.5	2			ORn		04To03	
	6959.59	0.14			−1.6	o			Bdn		06Fi.A	
	6959.352	0.018			0.7	2			Bdn		14Fi01	
^{23}Na(d,p)^{24}Na	4735	7	4734.799	0.016	−0.0	U			CIT		52Mi54	Y
	4736	5			−0.2	U			Mex		64Ma.B	
	4736	7			−0.2	U			MIT		64Sp12	
^{23}Na(p,γ)^{24}Mg	11692.95	0.17	11692.695	0.013	−1.5	U			Wis		67Mo17	Z
	11691.2	1.1			1.4	U					72Me09	
	11692.43	0.31			0.9	U					85Uh01	Z
^{24}Mg(p,d)^{23}Mg	−14307.5	1.5	−14306.66	0.03	0.6	U			MSU		74No07	
^{24}Mg(^{3}He,α)^{23}Mg	4051	15	4046.40	0.03	−0.3	U			Man		59Ba13	Y
^{24}Mg(^{7}Li,^{8}He)^{23}Al	−37397	27	−37384.3	0.4	0.5	U					01Ca37	
^{24}Al^{i}(p)^{23}Mg	4086	9	4085	3	−0.1	3			Brk		79Ay01	
	4084.5	3.5			0.1	3			MSU		80Le18	
	4093	20			−0.4	U			Bor		98Cz01	
^{24}Ne(β^-)^{24}Na	2449	50	2466.2	0.5	0.3	U					56Dr11	
^{24}Na(β^-)^{24}Mg	5511.8	2.	5515.677	0.021	1.9	U					61De25	*
	5515.8	2.			−0.1	U					64Le09	*
	5516.8	2.			−0.6	U					65Be24	*
	5511.5	1.0			4.2	B					69Bo48	*
	5511.8	2.			1.9	U					72Gi17	*
	5512.5	1.2			2.6	U					76Ge06	*
^{24}Al(β^+)^{24}Mg	13880	50	13884.77	0.23	0.1	U					68Ar03	
^{24}Mg(p,n)^{24}Al	−14659.0	2.8	−14667.11	0.23	−2.9	B			Yal		69Ov01	Z
^{24}Mg(^{3}He,t)^{24}Al	−13880	60	−13903.36	0.23	−0.4	U			Brk		66Ma18	
^{24}Mg(^{3}He,t)^{24}Al−^{36}Ar()^{36}K	−1071.48	1.05	−1070.4	0.4	1.1	1	14	10 ^{36}K	Mun		10Wr01	
^{24}Mg(π^+,π^-)^{24}Si	−23594	52	−23665	4	−1.4	U					80Bu15	
	−23667	21			0.1	U					28Si-4	
*^{24}Si−u	From frequency ratio $^{24}SiO_2H^+/(C_3H_2F)^+$ =1.000085151(71)										22Pu01	**
*^{23}Na(n,γ)^{24}Na	Original value (,Z) increased by 0.037 for better recoil correction										AHW	**
*^{24}Na(β^-)^{24}Mg	E_{β^-}=1389(2) 1393(2) 1394(2) 1388.7(1.0) 1389(2) 1389.7(1.2) respectively,										GAu	**
*	to 4$^+$ level at 4122.889 keV										Ens07a	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{25}F−u		12010	220	12170	100	0.5	o			TO1	1.5	86Vi09	
		12010	290			0.4	o			GA1	1.5	87Gi05	
		12210	150			−0.2	2			TO2	1.5	88Wo09	
		12120	151			0.2	o			GA3	1.5	91Or01	
		11990	130			0.9	2			TO4	1.5	91Zh24	
		12249	97			−0.6	2			GA7	1.5	07Ju03	
^{25}Ne−u		−2293	32	−2173	9	3.8	F			P40	1.0	01Lu20	*
		−2166	41			−0.2	U			LZ1	1.0	15Xu14	
^{25}Ne−^{12}C ^{13}C		−5527.4	8.5				2			TR1	1.0	23Ja05	
^{25}Al−^{25}Mg		4591.342	0.048				2			JY1	1.0	16Ca22	
^{25}Mg(p,α)^{22}Na		−3151	8	−3147.33	0.14	0.5	U			MIT		59Br74	Y
^{23}Na(t,p)^{25}Na		7488.8	1.2				2			Str		84An17	
^{25}Mg(d,α)^{23}Na		7026	13	7047.88	0.05	1.7	U			MIT		64Sp12	
		7048	10			−0.0	U					67Ha17	
^{25}O(γ,n)^{24}O		776	15	757	8	−1.2	3					08Ho03	*
		740	40			0.4	U					13Ca18	*
		749	10			0.8	3					16Ko11	
^{24}Mg(n,γ)^{25}Mg		7330.5	9.99	7330.52	0.05	0.0	U					69Ha.A	
		7330.5	0.3			0.1	U			MMn		80Is02	Z
		7330.78	0.14			−1.8	U			ILn		82Hu02	Z
		7330.4	0.2			0.6	U			MMn		85Ke.A	
		7330.64	0.08			−1.5	–			MMn		90Pr02	Z
		7330.69	0.05			−3.4	B			ORn		92Wa06	
		7330.53	0.15			−0.1	–			Bdn		06Fi.A	
^{24}Mg(d,p)^{25}Mg		5098	12	5105.95	0.05	0.7	U			Har		61Hi11	Y
		5112	12			−0.5	U			Mex		61Ja23	
		5102	7			0.6	U			MIT		64Sp12	
^{24}Mg(n,γ)^{25}Mg	ave.	7330.62	0.07	7330.52	0.05	−1.3	1	45	43 ^{25}Mg			average	
^{24}Mg(p,γ)^{25}Al		2271.6	1.1	2271.37	0.06	−0.2	U					71Ev01	Z
		2271.7	0.7			−0.5	U					72Pi07	Z
		2271.4	0.8			−0.0	U					85Uh01	Z
^{24}Mg(^{3}He,d)^{25}Al		−3218.0	4.5	−3222.11	0.06	−0.9	U			NDm		73Br27	
^{24}Mg(p,γ)^{25}Al^{i}		−5629.3	5.8	−5630.3	1.1	−0.2	U					68Te01	*
^{25}Ne(β^-)^{25}Na		7380	300	7334	8	−0.2	U					73Go11	
^{25}Na(β^-)^{25}Mg		3650	250	3835.0	1.2	0.7	U					54Na18	
		4000	200			−0.8	U					55Ma63	
^{25}Al(β^+)^{25}Mg		4292	30	4276.81	0.04	−0.5	U					60Wa04	
^{25}Mg(p,n)^{25}Al		−5058	6	−5059.15	0.04	−0.2	U			Har		69Fr08	
^{25}Al^{i}(IT)^{25}Al		7901	2	7901.7	1.1	0.3	3					77Ro03	
		7902.0	1.4			−0.2	3					21Su03	
*^{25}Ne−u	F : rejected by authors: “unreliable double peak”											06Ga04	**
*^{25}O(γ,n)^{24}O	Symmetrized from 770(+20–10)											08Ho03	**
*^{25}O(γ,n)^{24}O	Symmetrized from 725(+54–29)											13Ca18	**
*^{24}Mg(p,γ)^{25}Al^{i}	IT=7916(6), Q rebuilt with Ame1964, error estimated by evaluator											GAu	**
^{26}F−u		19800	1000	20050	110	0.2	o			TO1	1.5	86Vi09	
		20940	640			−0.9	o			GA1	1.5	87Gi05	
		19820	210			0.7	–			TO2	1.5	88Wo09	
		19544	300			1.1	o			GA3	1.5	91Or01	
		19490	210			1.8	U			TO4	1.5	91Zh24	
		20054	86			−0.0	–			GA7	1.5	07Ju03	
	ave.	20020	120			0.2	1	93	93 ^{26}F			average	
^{26}Ne−u		448	90	494	8	0.3	–			GA3	1.5	91Or01	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{26}Ne−u		461	33	494	8	1.0	o			P40	1.0	01Lu20	
		518	20			−1.2	–			P40	1.0	06Ga04	
	ave.	516	20			−1.1	1	16	16 ^{26}Ne			average	
^{26}Ne−^{12}C ^{14}N		−2583.9	8.6	−2580	8	0.5	1	84	84 ^{26}Ne	TR1	1.0	23Ja05	
^{26}Na−u		−7367	7	−7365	4	0.2	o			P40	1.0	01Lu17	
		−7365	4			−0.1	2			P40	1.0	06Ga04	*
		−7368	11			0.2	2			P40	1.0	06Ga04	
^{26}Mg−H_{26}		−220857.848	0.034	−220857.86	0.03	−0.3	1	85	85 ^{26}Mg	ST2	1.0	03Be02	
^{26}P−u		11806.8	11.8	11806	11	−0.1	o			LZ2	1.0	24Yu13	
		11806	11				2			LZ2	1.0	25Ch67	*
^{26}Al−^{23}$Na_{1.130}$		−1547.46	0.24	−1547.41	0.07	0.2	U			MA8	1.0	08Ge08	
^{26}Si−^{23}$Na_{1.13}$		3895.1	2.1	3894.53	0.12	−0.3	U			MS1	1.0	10Kw02	
^{26}Al−^{25}$Mg_{1.040}$		1621.46	0.48	1621.43	0.06	−0.1	U			JY1	1.0	06Er08	
^{26}Al^m−^{25}$Mg_{1.040}$		1867.09	0.53	1866.53	0.06	−1.1	U			JY1	1.0	06Er08	
^{26}Al−^{26}Mg		4299.14	0.17	4298.91	0.07	−1.4	1	16	15 ^{26}Al	JY1	1.0	06Er08	
^{26}Al^m−^{26}Mg		4544.09	0.17	4544.00	0.07	−0.5	1	16	15 ^{26}Al^m	JY1	1.0	06Er08	
^{26}Al^m−^{26}Al		245.09	0.17	245.096	0.014	0.0	U			JY1	1.0	06Er08	
		244.91	0.14			1.3	U			JY1	1.0	09Er02	
		245.114	0.049			−0.4	U			JY1	1.0	09Er07	
^{26}Si−^{26}Al		5441.97	0.14	5441.94	0.09	−0.2	2			JY1	1.0	09Er02	
		5441.92	0.12			0.2	2			JY1	1.0	09Er02	*
^{25}Na−^{26}$Na_{.721}$ ^{22}$Na_{.284}$		−2881	33	−2939.7	2.8	−1.8	U			P13	1.0	75Th08	
		−2921	22			−0.8	U			P13	1.0	75Th08	
^{26}Al(n,α)^{23}Na		2966.5	2.5	2966.12	0.07	−0.2	U					01Wa50	
^{23}Na(α,n)^{26}Al		−2968	4	−2966.12	0.07	0.5	U			Duk		60Wi07	Y
^{26}O(γ,2n)^{24}O		90	110	18	5	−0.7	U					12Lu07	*
		18	5				3					16Ko11	
^{24}Mg(t,p)^{26}Mg		9940	12	9941.81	0.03	0.2	U			Har		61Hi11	Y
^{24}Mg(^{3}He,p)^{26}Al		5932	15	5918.81	0.07	−0.9	U			Ald		59Hi66	Y
		5922	8			−0.4	U			Phi		72Be51	
^{24}Mg(^{3}He,n)^{26}Si		85	18	67.32	0.11	−1.0	U			CIT		67Mi02	
		75	30			−0.3	U					67Mc03	
		95	15			−1.8	U			Har		68Ad03	
		65	30			0.1	U			Ber		68Ha09	
^{26}Mg(^{7}Li,^{8}B)^{25}Ne		−22050	100	−22205	8	−1.6	U			Brk		73Wi06	
^{26}Mg(^{13}C,^{14}O)^{25}Ne		−19067	50	−19074	8	−0.1	U			Can		85Wo04	
^{26}Mg(d,^{3}He)^{25}Na		−8653	10	−8652.2	1.2	0.1	U			MSU		73Be14	
^{26}Mg(t,α)^{25}Na		5664	12	5668.2	1.2	0.3	U			Ald		62Hi01	
^{25}Mg(n,γ)^{26}Mg		11092.9	0.3	11093.08	0.04	0.6	U			MMn		80Is02	
		11091.84	0.44			2.8	U			ILn		82Hu02	Z
		11093.10	0.06			−0.3	1	54	46 ^{25}Mg	MMn		90Pr02	Z
		11093.17	0.06			−1.5	o			ORn		91Ki04	Z
		11093.23	0.05			−3.0	B			ORn		92Wa06	Z
		11093.16	0.22			−0.4	U			Bdn		06Fi.A	
^{25}Mg(d,p)^{26}Mg		8865	12	8868.51	0.04	0.3	U			Ald		61Hi11	Y
		8876	12			−0.6	U			Mex		61Ja23	
		8889	12			−1.7	U			MIT		64Sp12	
^{25}Mg(p,γ)^{26}Al		6305.0	1.2	6306.33	0.06	1.1	U					74De37	
		6304.9	1.1			1.3	U					79El11	
		6306.39	0.11			−0.6	–					85Be17	Z
		6306.38	0.08			−0.7	–			Utr		91Ki04	Z
	ave.	6306.38	0.06			−0.9	1	75	63 ^{26}Al			average	
^{26}Si^i(p)^{25}Al		7563	15	7541.0	2.0	−1.5	U			Brk		83Ca06	
		7544	15			−0.2	U			Brk		83Ho23	*
^{26}Mg(π^-,π^+)^{26}Ne		−17676	72	−17697	7	−0.3	U					80Na12	
^{26}Na(β^-)^{26}Mg		9210	200	9354	4	0.7	U					73Al13	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{26}Mg(t,^{3}He)^{26}Na	−9292	20	−9335	4	−2.2	U			LAl		74Fl01	
^{26}Mg(^{7}Li,^{7}Be)^{26}Na	−10182	40	−10216	4	−0.8	U			ChR		72Ba35	*
^{26}Al(β^+)^{26}Mg	3991	8	4004.41	0.06	1.7	U					58Fe16	*
^{26}Mg(p,n)^{26}Al	−4786.7	10.	−4786.75	0.06	−0.0	U			Oak		55Ki28	*
	−4787.04	0.48			0.6	U			Utr		69De27	
	−4786.1	1.6			−0.4	U			Har		69Fr08	*
	−4785.66	0.22			−5.0	C			Auc		84Ba.B	*
	−4786.57	0.05			−3.7	C			Auc		92Ba.A	*
	−4786.25	0.12			−4.2	B			Auc		94Br11	*
^{26}Mg(^{3}He,t)^{26}Al	−4023.0	0.6	−4023.00	0.06	0.0	F			Mun		77Vo02	*
^{26}Mg(^{3}He,t)^{26}Al−^{27}Al()^{27}Si	808.2	2.0	807.95	0.12	−0.1	U			ChR		74Ha35	
^{26}Mg(^{3}He,t)^{26}Al−^{14}N()^{14}O	1139.43	0.13	1139.96	0.07	4.1	B			ChR		87Ko34	*
^{26}Al^{m}(IT)^{26}Al	228.305	0.013	228.306	0.013	0.0	1	99	85 ^{26}Al^{m}			Ens164	
^{26}Si(β^+)^{26}Al	5079	13	5069.14	0.08	−0.8	U					63Fr10	
^{26}Si^{i}(IT)^{26}Si	13026	11	13055.0	2.0	2.6	U					04Th09	*
	13055	2				3					22Li66	*
*^{26}Na−u	Result from the “Thermo” experiment. Next item from “Rilis”										06Ga04	**
*^{26}P−u	combined results with 24Yu13										25Ch67	**
*^{26}Si−^{26}Al	D_M=5196.82(0.12) μu for ^{26}Al^{m} at 228.306(0.013); $M-A$=–7141.05(0.13) keV										Nub269	**
*^{26}O(γ,2n)^{24}O	Symmetrized from 150(+50–150) keV										12Lu07	**
*^{26}O(γ,2n)^{24}O	less than 40 keV										13Ca18	**
*^{26}Si^{i}(p)^{25}Al	E_p=3699(15) to 3695.5 level; different from preceding data										Ens098	**
*^{26}Mg(^{7}Li,^{7}Be)^{26}Na	Q =–10222(30) corrected for contribution of unresolved 82.2 level										Ens164	**
*^{26}Al(β^+)^{26}Mg	E_{β^+}=1160(8) to 2$^+$ level at 1808.74 level										Ens164	**
*^{26}Mg(p,n)^{26}Al	T=5191(10,Z) to ^{26}Al^{m} at 228.306 keV										Nub269	**
*^{26}Mg(p,n)^{26}Al	T=5209.3(1.6,Z) to ^{26}Al^{m} at 228.306 keV										Nub269	**
*^{26}Mg(p,n)^{26}Al	T=5208.86(0.23) to ^{26}Al^{m} at 228.306 keV										Nub269	**
*^{26}Mg(p,n)^{26}Al	T=5209.71(0.05) to ^{26}Al^{m} at 228.306 keV										Nub269	**
*^{26}Mg(p,n)^{26}Al	T=5209.46(0.12) to ^{26}Al^{m} at 228.306 keV										Nub269	**
*^{26}Mg(^{3}He,t)^{26}Al	Q =–4251.3(0.6,Z) to ^{26}Al^{m} at 228.306 keV										Nub269	**
*^{26}Mg(^{3}He,t)^{26}Al	F : rejected in reference of same group										09Fa15	**
*^{26}Mg(^{3}He,t)^{26}Al−^{14}N()^{14}O	Q(to 1057.740(0.023) level)-^{14}N()^{14}O=81.69(0.13)										82Al19	**
*^{26}Si^{i}(IT)^{26}Si	Unweighted average of 13015(4) and 13036(4) keV										HWJ238	**
*^{26}Si^{i}(IT)^{26}Si	Derived from several proton groups.										22Li66	**
^{27}F−u	27500	700	26980	130	−0.5	U			TO2	1.5	88Wo09	
	26005	770			0.8	U			GA3	1.5	91Or01	
	27100	900			−0.1	U			TO4	1.5	91Zh24	
	26900	580			0.1	o			GA5	1.5	99Sa.A	
	26441	204			1.8	B			GA7	1.5	07Ju03	
	27322	279			−0.8	1	10	10 ^{27}F	GA8	1.5	12Ga45	
^{27}Ne−u	6010	640	7570	100	1.6	F			TO1	1.5	86Vi09	*
	7470	300			0.2	U			GA1	1.5	87Gi05	
	7567	172			0.0	o			GA3	1.5	91Or01	
	7670	130			−0.5	2			TO4	1.5	91Zh24	
	7536	75			0.3	2			GA7	1.5	07Ju03	
^{27}Na−u	−5922	11	−5924	4	−0.1	o			P40	1.0	01Lu17	*
	−5922	4			−0.4	o			P40	1.0	06Ga04	*
^{27}P−u	−735	45	−720.0	0.6	0.3	U			LZ1	1.0	18Fu11	
	−720.03	0.64				2			MSU	1.0	23Ya31	*
^{27}S−u	18699	42	18719	26	0.5	o			LZ2	1.0	24Yu13	
	18719	26				2			LZ2	1.0	25Ch67	*
^{27}Mg−^{23}Na$_{1.174}$	−3648.27	0.15	−3648.49	0.05	−1.5	1	11	11 ^{27}Mg	TT1	1.0	17Br14	
^{27}Al−^{23}Na$_{1.174}$	−6450.79	0.25	−6450.73	0.05	0.2	U			MA8	1.0	08Ge08	
	−6450.754	0.054			0.5	1	89	89 ^{27}Al	TT1	1.0	15Kw01	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{27}Na−^{27}Al		12538	4				2			P40	1.0	01Lu17	
^{24}Na−^{27}Na$_{.356}$ ^{22}Na$_{.655}$		−3006	38	−3059.8	1.3	−0.9	U			P10	1.5	75Th08	
^{26}Na−^{27}Na$_{.770}$ ^{22}Na$_{.236}$		−1437	86	−1389	5	0.6	U			P13	1.0	75Th08	
^{26}Na−^{27}Na$_{.481}$ ^{25}Na$_{.520}$		676	66	659	4	−0.2	U			P10	1.5	75Th08	
		734	86			−0.6	U			P11	1.5	75Th08	
^{23}Na(α,γ)^{27}Al		10090.0	1.3	10091.93	0.05	1.5	U			Utr		78Ma23	
^{27}O(γ,3n)^{24}O		1090	45				3					23Ko15	
^{27}Al(p,α)^{24}Mg		1601.7	0.7	1600.77	0.05	−1.3	o			Zur		63Ry04	
		1598.4	1.0			2.4	U			NDm		65Br28	
		1601.3	0.5			−1.1	U			Zur		67St30	Z
		1600.06	0.21			3.4	B			Utr		78Ma23	Z
^{24}Mg(α,p)^{27}Al		−1598.9	1.0	−1600.77	0.05	−1.9	U			NDm		65Br28	
^{25}Mg(t,p)^{27}Mg		9055	11	9054.65	0.06	−0.0	U			Tal		61Hi11	Y
^{27}Al(d,α)^{25}Mg		6699	12	6706.72	0.07	0.6	U			Ald		61Hi11	Y
		6691	11			1.4	U			Tal		62Sh01	Y
		6700	10			0.7	U			MIT		64Sp12	
^{27}Al(p,t)^{25}Al^{i}		−23843.4	4.7	−23843.4	1.1	−0.0	U			MSU		73Be14	*
^{27}P^{i}(2p)^{25}Al		6410	45	6368	14	−0.9	3			Lis		91Bo32	
		6270	50			2.0	3					01Ca60	*
		6372	15			−0.3	3					21Sh23	
^{27}F(γ,n)^{26}F		−1620	60	−1610	60	0.1	1	98	90 ^{27}F			20Re06	
^{26}Mg(^{18}O,^{17}F)^{27}Na		−13295	55	−13431	4	−2.5	F			Mun		78Pa12	*
		−13433	60			0.0	U			Can		85Fi08	
^{26}Mg(n,γ)^{27}Mg		6443.35	0.55	6443.37	0.04	0.0	U			ILn		82Hu02	
		6443.56	0.25			−0.8	o			MMn		85Ke.A	
		6443.26	0.08			1.4	–			MMn		90Pr02	Z
		6443.44	0.05			−1.4	–			ORn		92Wa06	Z
		6443.35	0.13			0.1	–			Bdn		06Fi.A	
^{26}Mg(d,p)^{27}Mg		4214	12	4218.80	0.04	0.4	U			Ald		61Hi11	Y
		4215	10			0.4	U			Mex		61Ja23	
		4211	6			1.3	U			MIT		64Sp12	
^{26}Mg(n,γ)^{27}Mg	ave.	6443.39	0.04	6443.37	0.04	−0.4	1	93	89 ^{27}Mg			average	
^{26}Mg(p,γ)^{27}Al		8270.8	0.5	8271.29	0.06	1.0	U			Utr		59An33	*
		8271.2	0.5			0.2	U					63Va24	Z
		8271.3	0.5			−0.0	U			Utr		78Ma24	*
^{27}Al(t,α)^{26}Mg		11541	12	11542.57	0.06	0.1	U			Ald		61Hi11	
^{27}Al(^{3}He,α)^{26}Al		7523	15	7519.57	0.08	−0.2	U			Ald		59Hi66	
		7519	15			0.0	U			Man		60Ta12	
^{26}Al(p,γ)^{27}Si		7464.9	0.9	7463.34	0.13	−1.7	U					84Bu09	Z
^{27}P(p)^{26}Si		−788	31	−818.7	0.6	−1.0	U			Dbp		17Ja05	*
		−807	9			−1.3	U					19Su14	*
^{27}Na(β^-)^{27}Mg		8930	150	9069	4	0.9	U					73Al13	
^{27}Mg(β^-)^{27}Al		2600	11	2610.27	0.07	0.9	U					54Da22	
^{27}Si(β^+)^{27}Al		4872	20	4812.36	0.10	−3.0	B					60Wa04	
^{27}Al(p,n)^{27}Si		−5573	10	−5594.71	0.10	−2.2	U			Oak		55Ki28	Z
		−5597.5	6.0			0.5	U			Tkm		63Ok01	
		−5593.6	4.3			−0.3	U			Ric		65Ku02	
		−5585.1	2.3			−4.2	B			Ric		66Bo20	Z
		−5592.0	1.0			−2.7	U			Yal		69Ov01	Z
		−5594.1	3.2			−0.2	U			Har		76Fr13	
		−5593.8	0.26			−3.5	F			Auc		77Na24	*
		−5594.27	0.11			−4.0	F			Auc		85Wh03	*
		−5594.72	0.10				2			Auc		94Br37	Z
^{27}Si^{i}(IT)^{27}Si		6626	3	6638.0	1.0	4.0	B					86Sc21	
		6638	1				3					16Mc09	*
*^{27}Ne−u	F : contaminated by ^{27}Na											91Zh24	**
*^{27}Na−u	Not independent of ^{27}Na−^{27}Al from isobaric method, do not use											01Lu17	**
*^{27}P−u	From frequency ratio ^{27}PO$^+$/(HCNO+)=1.000270250(16)											23Ya31	**
*^{27}S−u	combined results with 24Yu13											25Ch67	**
*^{27}Al(p,t)^{25}Al^{i}	IT=7904(5), rebuilt Q =–23847.9(4.7); recalib +4.5 keV											GAu	**
*^{27}P^{i}(2p)^{25}Al	And E_{2p}=5315(60) to 3/2$^+$ level at 944.9 keV											Ens098	**
*^{26}Mg(^{18}O,^{17}F)^{27}Na	F : shape of peak raises doubt on centroid determination											GAu	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{26}$Mg(p,γ)^{27}Al	E_p=338.65(0.12) to 8596.8(0.5) level										78Ma24	**
$*^{26}$Mg(p,γ)^{27}Al	E_p=338.21(0.30) to 8596.8(0.5) level										78Ma24	**
$*^{26}$Mg(p,γ)^{27}Al	E_p=809.90(0.05,Z) to 9050.7(0.5,Z) level										78Ma24	**
$*^{27}$P(p)^{26}Si	Qp=332(30) keV from 3/2$^+$ level at 1120(8) keV from 08Ga10										17Ja05	**
$*^{27}$P(p)^{26}Si	Qp=318(8) keV from 3/2$^+$ level at 1125(2) keV										19Su14	**
$*^{27}$Al(p,n)^{27}Si	F : rejected by same group "measurement contains error"										94Br37	**
^{28}Ne−u	11490	430	12130	140	1.0	o			TO1	1.5	86Vi09	
	12270	560			−0.2	o			GA1	1.5	87Gi05	
	11958	238			0.5	o			GA3	1.5	91Or01	
	12160	140			−0.1	2			TO4	1.5	91Zh24	
	12110	118			0.1	2			GA7	1.5	07Ju03	
^{28}Na−u	−1220	190	−1061	11	0.6	o			TO1	1.5	86Vi09	
	−1097	96			0.2	U			GA3	1.5	91Or01	
	−1062	14			0.1	o			P40	1.0	01Lu17	
	−1061	11				2			P40	1.0	06Ga04	
^{28}Si−u	−23073.43	0.30	−23073.4657	0.0006	−0.1	U			ST1	1.0	93Je06	
	−23073.4676	0.0020			0.9	U			MI1	1.0	95Di08	
	−23073.00	0.27			−0.7	U			OH1	2.5	94Go.A	
	−23073.466	0.008			0.0	U			ST2	1.0	02Be64	*
C_2 D_2−^{28}Si	51277.0224	0.0024	51277.0214	0.0006	−0.4	U			MI1	1.0	95Di08	
$^{15}N_2$−^{28}Si H_2	7641.2007	0.0024	7641.1984	0.0013	−0.9	1	30	25 ^{15}N	MI1	1.0	95Di08	
C_2 H_4−^{28}Si	54373.59360	0.00079	54373.5934	0.0006	−0.3	1	57	57 ^{28}Si	FS1	1.0	08Re16	
$^{13}C_2$ H_2−^{28}Si	45433.19986	0.00071	45433.2000	0.0005	0.2	1	54	29 ^{28}Si	FS1	1.0	08Re16	
^{28}S−u	4485.3	15.0	4488	13	0.2	o			LZ2	1.0	24Yu13	
	4488	13				2			LZ2	1.0	25Ch67	*
$^{28}Si_2$ ^{16}O−^{35}Cl ^{37}Cl	14013.07	0.70	14012.42	0.07	−0.6	U			H46	1.5	93Nx02	
^{28}Mg−$^{23}Na_{1.217}$	−3673.79	0.28				2			TT1	1.0	17Br14	
^{25}Na−$^{28}Na_{.446}$ $^{22}Na_{.568}$	−5869	75	−5974	5	−0.9	U			P10	1.5	75Th08	*
^{26}Na−$^{28}Na_{.619}$ $^{22}Na_{.394}$	−4229	613	−4208	7	0.0	U			P11	1.5	75Th08	
	−4205	128			−0.0	U			P12	2.5	75Th08	
	−4203	87			−0.1	U			P13	1.0	75Th08	
^{28}Si(p,^{6}He)^{23}Al	−38569	80	−38544.0	0.3	0.3	U			Brk		69Ce01	
^{28}Si(^{3}He,^{8}Li)^{23}Al	−34274	25	−34255.5	0.3	0.7	U			MSU		75Be38	
^{28}O(γ,4n)^{24}O	466	49				3					23Ko15	*
^{28}Si(α,^{8}He)^{24}Si	−61433	21	−61431	4	0.1	R			Tex		80Tr04	
^{28}Si(p,α)^{25}Al	−7709.3	2.6	−7712.77	0.06	−1.3	U			NDm		73Br27	
^{28}Si(^{3}He,^{6}He)^{25}Si	−27976	50	−27981	10	−0.1	U			Brk		70Me11	
	−27981	10				2			MSU		72Be12	
^{26}Mg(t,p)^{28}Mg	6474	12	6466.24	0.26	−0.6	U			Har		61Hi11	Y
^{26}Mg(^{3}He,p)^{28}Al	8285	5	8278.42	0.06	−1.3	U			Phi		74Be07	
^{28}Si(d,α)^{26}Al	1429	4	1428.15	0.07	−0.2	U			MIT		64Sp12	
^{28}Si(p,t)^{26}Si	−22009	3	−22012.64	0.11	−1.2	U			MSU		74Ha02	
	−22014.1	1.0			1.5	U			Yal		05Pa31	
^{28}F(γ,n)^{27}F	220	50	199	6	−0.4	U			MSU		12Ch02	
	199	6				2					20Re06	
^{27}Al(n,γ)^{28}Al	7725.02	0.20	7725.174	0.011	0.8	U			BNn		78St25	Z
	7725.07	0.30			0.3	U			ILn		79Br25	Z
	7725.13	0.3			0.1	U			MMn		80Is02	Z
	7725.02	0.10			1.5	U					81Su.A	Z
	7725.14	0.09			0.4	o			ILn		82Sc14	Z
	7725.17	0.15			0.0	U			Bdn		06Fi.A	
	7725.174	0.011				2			ILn		18Rz01	
^{27}Al(d,p)^{28}Al	5511	5	5500.608	0.011	−2.1	U			Mex		61Ja23	
	5503	10			−0.2	U			MIT		64Sp12	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{27}Al(p,γ)^{28}Si	11584.89	0.30	11584.90	0.05	0.0	U			Utr		78Ma23	Z
	11585.09	0.14			−1.3	1	11	11 ^{27}Al			28Sir-0	
^{27}Al(^{3}He,d)^{28}Si	6049	18	6091.43	0.05	2.4	U					60Fo01	
^{27}Al(p,γ)^{28}Si^{r}	−956.15	0.03	−956.139	0.025	0.3	2			Utr		78Ma23	Z
	−956.025	0.020			−5.7	B			Auc		94Br37	Z
	−956.13	0.05			−0.4	2					98Wa.A	Z
^{28}Si(^{3}He,α)^{27}Si	3407	15	3398.01	0.11	−0.6	U			Ald		59Hi68	Y
^{28}Si(^{7}Li,^{8}He)^{27}P	−37513	40	−37524.7	0.6	−0.3	U					01Ca37	
^{28}P^{i}(p)^{27}Si	3835	20				3			Lis		89Po10	
^{28}Cl(p)^{27}S	1600	80	2200#	300#	7.5	D					18Mu18	*
^{28}Mg(β^-)^{28}Al	1791	10	1830.77	0.27	4.0	B					53Ma23	*
	1831.8	2.0			−0.5	U					54Ol03	*
^{28}Al(β^-)^{28}Si	4644	10	4642.08	0.05	−0.2	U					52Mo22	*
	4657	14			−1.1	U					54Ol03	*
^{28}Si^{r}(IT)^{28}Si	12541.23	0.14	12541.04	0.05	−1.3	R			Utr		90En02	Z
^{28}P(β^+)^{28}Si	14290	40	14345.0	1.2	1.4	U					68Ar03	*
^{28}Si(p,n)^{28}P	−15118.3	4.1	−15127.3	1.2	−2.2	U			Yal		69Ov01	Z
	−15112.5	5.8			−2.6	U			BNL		71Go18	Z
^{28}Si(^{3}He,t)^{28}P	−14380	60	−14363.6	1.2	0.3	U			Brk		66Ma18	
^{28}Si(^{3}He,t)^{28}P−^{36}Ar()^{36}K	−1530.58	1.10	−1530.6	1.1	0.0	1	100	100 ^{28}P	Mun		10Wr01	
^{28}Si(π^+,π^-)^{28}S	−24544	160	−24651	12	−0.7	U					82Mo12	*
*^{28}Si−u	Unc. was erroneously 0.0008 in Ame2012, was correct in Ame2003										GAu	**
*^{28}S−u	combined results with 24Yu13										25Ch67	**
*^{25}Na−^{28}Na$_{.446}$ ^{22}Na$_{.568}$	Symmetric double-doublet 22−24 26−28 included										GAu	**
*^{28}Cl(p)^{27}S	Trends in Mass Surface TMS suggest ^{28}Cl 600 keV less bound										HWJ266	**
*^{28}Mg(β^-)^{28}Al	E_{β^-}=418(10) 459(2) respectively, to 1^+ level at 1372.917 keV										Ens13a	**
*^{28}Al(β^-)^{28}Si	E_{β^-}=2865(10) 2878(14) respectively, to 2^+ level at 1779.030 keV										Ens13a	**
*^{28}P(β^+)^{28}Si	E_{β^+}=11490(40) to 2^+ level at 1779.030 keV										Ens13a	**
*^{28}Si(π^+,π^-)^{28}S	Original –24603(160) recalibrated to ^{16}O(π^+,π^-)^{16}Ne Q =–27704(20) keV										GAu	**
^{29}F−u	43103	376				2			GA8	1.5	12Ga45	
^{29}Ne−u	19433	551	19750	160	0.4	o			GA3	1.5	91Or01	
	19300	400			0.8	U			TO4	1.5	91Zh24	
	19400	410			0.6	o			GA5	1.5	00Sa21	
	19753	107				2			GA7	1.5	07Ju03	
^{29}Na−u	2820	230	2877	8	0.2	U			TO1	1.5	86Vi09	
	2838	143			0.2	U			GA3	1.5	91Or01	
	2861	14			1.1	o			P40	1.0	01Lu17	
	2866	13			0.9	1	37	37 ^{29}Na	P40	1.0	06Ga04	
^{29}Na−^{39}K$_{.744}$	29885.9	9.9	29879	8	−0.6	1	63	63 ^{29}Na	TT1	1.0	13Ch49	
^{29}Mg−u	−11375	19	−11392.8	0.4	−0.9	U			P40	1.0	06Ga04	*
	−11388	16			−0.3	U			P40	1.0	06Ga04	
^{29}Al−O$_{1.812}$	−10328.8	1.0	−10332.1	0.4	−3.3	C			TT1	1.0	12Ch.A	
	−10333.5	1.7			0.8	U			TT1	1.0	15Kw01	
^{29}P ^{40}Ar−u	−55816.80	0.50	−55816.5	0.4	0.6	1	59	59 ^{29}P	MS1	1.0	15Ei01	
^{29}S−u	−3322	14				2			LZ1	1.0	18Fu11	
^{29}Mg−^{23}Na$_{1.261}$	1508.09	0.37				2			TT1	1.0	17Br14	
^{29}Al−^{23}Na$_{1.261}$	−6645.90	0.37				2			TT1	1.0	17Ga20	
^{29}Si−^{28}Si H	−8256.90198	0.00024				2			MI3	1.0	05Ra34	
^{26}Na−^{29}Na$_{.512}$ ^{22}Na$_{.506}$	−5763	91	−5611	5	1.1	U			P10	1.5	75Th08	
	−6379	293			1.7	U			P11	1.5	75Th08	
	−5252	277			−0.5	U			P12	2.5	75Th08	
	−5576	66			−0.5	U			P13	1.0	75Th08	
^{27}Na−^{29}Na$_{.466}$ ^{25}Na$_{.540}$	−1708	124	−1713	5	−0.0	U			P10	1.5	75Th08	
^{18}O(^{13}C,2p)^{29}Mg	−1456	50	−1623.4	0.3	−3.3	B					81Pa17	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{26}Mg(^{11}B,^{8}B)^{29}Mg		−19720	50	−19856.1	1.1	−2.7	U			Brk		74Sc26	
^{26}Mg(^{18}O,^{15}O)^{29}Mg		−9207	55	−9240.6	0.6	−0.6	U			Mun		78Pa12	
		−9250	45			0.2	U			Can		85Fi08	
^{26}Mg(α,p)^{29}Al		−2880	40	−2870.8	0.3	0.2	U			Yal		57Gr47	Y
		−2874	10			0.3	U			ANL		68Be13	
^{29}Si(n,α)^{26}Mg		−21	21	−34.137	0.029	−0.6	U			Ham		62An05	
^{27}Al(t,p)^{29}Al		8679.5	1.2	8671.7	0.3	−6.5	B			Str		84An17	
^{29}Si(d,α)^{27}Al		6000	11	6012.59	0.05	1.1	U			MIT		64Sp12	
^{29}Si(p,t)^{27}Si^{i}		−23802	5	−23809.4	1.0	−1.5	U			MSU		77Be13	
^{27}Al(^{3}He,n)^{29}P		6616	30	6615.9	0.4	−0.0	U			Oak		72Gr39	
^{29}Ar(2p)^{27}S		5500	180	4100#	200#	−7.8	D					18Mu18	*
^{28}Si(n,γ)^{29}Si		8473.6	0.3	8473.6026	0.0005	0.0	o			MMn		80Is02	Z
		8473.61	0.04			−0.2	U			MMn		90Is02	Z
		8473.55	0.04			1.3	U			ORn		92Ra19	Z
		8473.55	0.05			1.1	o			PTc		97Ro26	Z
		8473.54	0.17			0.4	U			Bdn		06Fi.A	
		8473.551	0.030			1.7	U			PTc		01Pa52	*
		8473.5957	0.0050			1.4	U			NBS		06De21	
^{28}Si(d,p)^{29}Si		6252	10	6249.03634	0.00022	−0.3	U			Mex		64Ma.B	
		6252	10			−0.3	U			MIT		64Sp12	
		6249.35	0.5			−0.6	U			Rez		90Pi05	*
^{28}Si(p,γ)^{29}P		2747.1	1.7	2749.0	0.4	1.1	–					73Ba35	Z
		2748.8	0.6			0.4	–					74By01	Z
^{28}Si(d,n)^{29}P		560	30	524.5	0.4	−1.2	U			Ald		60Ma21	
^{28}Si(^{3}He,d)^{29}P		−2733	12	−2744.5	0.4	−1.0	U			Ald		60Hi03	Y
^{28}Si(p,γ)^{29}P	ave.	2748.6	0.6	2749.0	0.4	0.7	1	40	40 ^{29}P			average	
^{28}Si(p,γ)^{29}P^{i}		−5630	10	−5632.8	2.5	−0.3	U			ANL		66Yo01	
		−5631.9	5.0			−0.2	1	24	24 ^{29}P^{i}			68Te01	*
^{29}Cl(p)^{28}S		1800	100	2260#	100#	4.6	D					15Mu13	
		1800	100			4.6	D					18Xu04	*
^{29}Mg(β^-)^{29}Al		7624	400	7595.4	0.5	−0.1	U					73Go34	
^{29}Al(β^-)^{29}Si		3850	100	3687.3	0.3	−1.6	U					54Na14	*
^{29}P(β^+)^{29}Si		4967	20	4942.2	0.4	−1.2	U					55Ro05	
^{29}P^{i}(IT)^{29}P		8382.1	2.8	8381.8	2.4	−0.1	1	76	76 ^{29}P^{i}			72Ba26	
*^{29}Mg−u	Result from the "Plasma" experiment. Next item from "Rilis"											06Ga04	**
*^{29}Ar(2p)^{27}S	Trends in Mass surface TMS suggests ^{29}Ar 1400 keV more bound											HWJ261	**
*^{28}Si(n,γ)^{29}Si	Original error 0.005 increased for calibration											GAu	**
*^{28}Si(d,p)^{29}Si	Estimated systematic error 0.5 added to statistical error 0.037 keV											AHW	**
*^{28}Si(p,γ)^{29}P^{i}	IT=8376(6), Q rebuilt with Ame1964, error estimated by evaluator											GAu	**
*^{29}Cl(p)^{28}S	Trends in Mass Surface TMS suggest ^{29}Cl 460 keV less bound											GAu	**
*^{29}Al(β^-)^{29}Si	E_{β^-}=1550(100) to 5/2$^+$ level at 2028.16 and 3/2$^+$ at 2425.97 keV											Ens125	**
^{30}Ne−u		23872	884	24990	270	0.8	o			GA3	1.5	91Or01	
		25660	850			−0.5	o			GA5	1.5	00Sa21	
		24734	301			0.6	–			GA7	1.5	07Ju03	
		25024	301			−0.1	–			GA8	1.5	12Ga45	
	ave.	24880	320			0.4	1	73	73 ^{30}Ne			average	
^{30}Na−u		7620	540	9098	5	1.8	F			TO1	1.5	86Vi09	*
		9200	370			−0.2	U			GA1	1.5	87Gi05	
		9126	218			−0.1	U			GA3	1.5	91Or01	
		9330	130			−1.2	U			TO4	1.5	91Zh24	
		8976	27			4.5	B			P40	1.0	01Lu17	
		8990	25			4.3	B			P40	1.0	06Ga04	
^{30}Na−O$_{1.876}$		18638.9	5.6	18638	5	−0.1	1	82	82 ^{30}Na	TT1	1.0	13Ch49	
^{30}Na−^{39}K$_{.769}$		37004	12	37008	5	0.3	1	18	18 ^{30}Na	TT1	1.0	13Ch49	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{30}Mg−O$_{1.876}$	3.0	3.7	5.6	1.4	0.7	1	14	14 ^{30}Mg	TT1	1.0	13Ch49	
^{30}Mg−u	−9700	230	−9534.5	1.4	0.5	o			TO1	1.5	86Vi09	
	−9597	98			0.4	U			GA3	1.5	91Or01	
	−9490	110			−0.3	U			TO4	1.5	91Zh24	
	−9546	14			0.8	U			P40	1.0	06Ga04	
^{30}Mg−^{39}K$_{.769}$	18375.6	1.5	18375.2	1.4	−0.3	1	86	86 ^{30}Mg	TT1	1.0	20Le06	
^{30}Al−O$_{1.875}$	−7495.1	2.8	−7495.7	2.1	−0.2	1	55	55 ^{30}Al	TT1	1.0	20Le06	
^{30}Al−^{39}K$_{.769}$	10878.1	3.1	10878.9	2.1	0.3	1	45	45 ^{30}Al	TT1	1.0	15Kw01	
^{30}P−^{30}Si	4543.353	0.066				4			JY1	1.0	16Ca22	
^{30}S−^{30}P	6593.28	0.21				5			JY1	1.0	11So11	
^{26}Na−^{30}Na$_{.433}$ ^{22}Na$_{.591}$	−7454	287	−7468	4	−0.0	U			P10	1.5	75Th08	
	−8060	641			0.6	U			P11	1.5	75Th08	
	−7045	225			−0.8	U			P12	2.5	75Th08	
	−7515	117			0.4	U			P13	1.0	75Th08	
^{27}Na−^{30}Na$_{.360}$ ^{25}Na$_{.648}$	−2750	213	−2505	4	0.8	U			P10	1.5	75Th08	
^{26}Mg(^{18}O,^{14}O)^{30}Mg	−16234	55	−16123.8	1.3	2.0	U			Mun		78Pa12	*
^{30}Si(n,α)^{27}Mg	−4193	21	−4199.97	0.05	−0.3	U			Ham		62An05	
^{30}Si(p,α)^{27}Al	−2368	10	−2372.04	0.05	−0.4	U			MIT		64Sp12	
^{27}Al(α,p)^{30}Si	2375	8	2372.04	0.05	−0.4	U			Man		59Ba13	Y
^{30}Si(d,α)^{28}Al	3123	10	3128.56	0.05	0.6	U			MIT		64Sp12	
^{28}Si(^{3}He,n)^{30}S	−573	15	−573.64	0.21	−0.0	U			CIT		67Mi02	
^{30}Ar(2p)^{28}S	2280	130	2420	80	1.1	o					15Mu13	*
	2420	80				3					18Xu04	*
^{30}F(γ,n)^{29}F	472	67				3					24Ka32	
^{29}Si(n,γ)^{30}Si	10609.6	0.3	10609.199	0.022	−1.3	o			MMn		80Is02	Z
	10609.21	0.04			−0.3	3			MMn		90Is02	Z
	10609.24	0.05			−0.8	3			ORn		92Ra19	Z
	10609.1776	0.0500			0.4	o			PTc		97Ro26	*
	10609.178	0.030			0.7	3			PTc		01Pa52	*
	10609.23	0.21			−0.1	U			Bdn		06Fi.A	
^{29}Si(d,p)^{30}Si	8413	10	8384.633	0.022	−2.8	U			Mex		61Ja23	
	8396	13			−0.9	U			MIT		64Sp12	
	8384.92	0.53			−0.5	U			Rez		90Pi05	*
^{29}Si(p,γ)^{30}P	5594.5	0.4	5594.75	0.07	0.6	U					85Re02	
	5594.5	0.5			0.5	U					96Wa33	
^{30}Cl(p)^{29}S	480	20	364	29	−5.8	B					18Mu18	*
	364	29				3					22Gi04	
^{30}Na(β^-)^{30}Mg	17167	330	17356	5	0.6	U					83De04	*
^{30}Mg(β^-)^{30}Al	6690	240	6982.7	2.3	1.2	U					83De04	*
^{30}Al(β^-)^{30}Si	8550	250	8568.8	1.9	0.1	U					61Ro12	*
^{30}Si(t,^{3}He)^{30}Al	−8520	40	−8550.3	1.9	−0.8	U					69Aj03	
	−8545	15			−0.4	U					87Pe06	
^{30}P(β^+)^{30}Si	4262	40	4232.11	0.06	−0.7	U					56Gr07	
	4267	25			−1.4	U					63Fr10	
^{30}Si(p,n)^{30}P	−5012.1	5.	−5014.45	0.06	−0.5	U			Har		75Fr.A	Z
^{30}S(β^+)^{30}P	6118	22	6141.60	0.20	1.1	U					63Fr10	*
*^{30}Na−u	F : contaminated by ^{30}Mg										91Zh24	**
*^{26}Mg(^{18}O,^{14}O)^{30}Mg	Tentative, say authors; four counts only										AHW	**
*^{30}Ar(2p)^{28}S	Symmetrized from 2250(+150–100)										GAu	**
*^{30}Ar(2p)^{28}S	Symmetrized from 2450(+50–100), supersedes 15Mu13										HWJ192	**
*^{29}Si(n,γ)^{30}Si	Original error 0.0005 increased for calibration										GAu	**
*^{29}Si(n,γ)^{30}Si	Original error 0.005 increased for calibration										GAu	**
*^{29}Si(d,p)^{30}Si	Estimated systematic error 0.5 added to statistical error 0.16 keV										AHW	**
*^{30}Cl(p)^{29}S	Might be a mixture of gs. and first excited state in ^{30}Cl										22Gi04	**
*^{30}Na(β^-)^{30}Mg	Calculated from 3 values used for calibration										GAu	**
*^{30}Mg(β^-)^{30}Al	Calculated from value used for calibration										GAu	**
*^{30}Al(β^-)^{30}Si	E_{β^-}=5050(250) to 2^+ level at 3498.49 keV										Ens109	**
*^{30}S(β^+)^{30}P	E_{β^+}=4422(22) to 0^+ level at 677.01 keV										Ens109	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{31}Ne−u	33087	1739	33470	290	0.1	U			GA7	1.5	07Ju03	
	33752	333			−0.6	1	33	33 ^{31}Ne	GA8	1.5	12Ga45	
^{31}Na−u	13440	1000	13145.9	2.8	−0.2	o			GA1	1.5	87Gi05	
	13559	327			−0.8	o			GA3	1.5	91Or01	
	13610	210			−1.5	U			TO4	1.5	91Zh24	
	13441	118			−1.7	U			GA7	1.5	07Ju03	
^{31}Na−^{12}C ^{16}O ^{1}H$_3$	−5243.8	2.8				2			TR1	1.0	25Ly01	*
^{31}Mg−O$_{1.938}$	6503.7	3.3	6499.8	2.6	−1.2	2			TT1	1.0	13Ch49	
^{31}Mg−^{12}C ^{16}O ^{1}H$_3$	−21751.8	4.2	−21745.4	2.6	1.5	2			TR1	1.0	25Ly01	*
^{31}Mg−u	−3830	220	−3355.7	2.6	1.4	o			TO1	1.5	86Vi09	
	−3520	180			0.6	o			GA1	1.5	87Gi05	
	−3458	149			0.5	U			GA3	1.5	91Or01	
	−3370	120			0.1	U			TO4	1.5	91Zh24	
	−3425	18			3.8	B			P40	1.0	06Ga04	
O$_2$−^{31}P H	8242.20819	0.00086	8242.2088	0.0007	0.7	1	69	37 ^{31}P	FS1	1.0	08Re16	
^{31}Ar−u	12120	17	12125	14	0.3	o			LZ2	1.0	24Yu13	
	12125	14				2			LZ2	1.0	25Ch67	*
^{31}K−u	32899	215	35110#	430#	10.3	D			1.0	1.0	19Ko18	*
^{31}Na−^{39}K$_{.795}$	42017	25	41999.3	2.8	−0.7	o			TT1	1.0	13Ch49	
	42000	15			−0.0	U			TT1	1.0	17Ga20	
^{31}Al−^{39}K$_{.795}$	12803.1	2.4				2			TT1	1.0	15Kw01	
^{31}P−^{28}Si H$_3$	−26639.6290	0.0056	−26639.6326	0.0007	−0.6	U			FS1	1.0	06Re19	
	−26639.63324	0.00089			0.7	1	67	63 ^{31}P	FS1	1.0	08Re16	
^{31}S−^{31}P	5794.98	0.25				2			JY1	1.0	10Ka30	
^{31}Cl−^{31}P	18686.1	3.7				2			JY1	1.0	16Ka15	
O$_2$−^{31}P	16067.228	0.096	16067.2407	0.0007	0.1	U			MS1	1.0	09Kw02	*
^{26}Na−^{31}Na$_{.373}$ ^{22}Na$_{.657}$	−7457	286	−8024	4	−0.8	U			P12	2.5	75Th08	
^{18}O(^{15}N,2p)^{31}Al	−170	90	−308.6	2.2	−1.5	U					81Pa11	
^{27}Al(α,γ)^{31}P	9667.4	1.3	9668.60	0.05	0.9	U			Utr		78Ma23	
^{31}P(p,α)^{28}Si	1912	5	1916.3073	0.0007	0.9	U			Bar		56Va14	Y
	1919	4			−0.7	U			VUn		64Sm03	
	1911	10			0.5	U			MIT		64Sp12	
	1915.8	0.2			2.5	U			Zur		67St30	
^{28}Si(α,n)^{31}S	−8135	44	−8096.64	0.23	0.9	U			Tal		63Ne05	
^{31}P(d,α)^{29}Si	8166	11	8165.3436	0.0007	−0.1	U			MIT		64Sp12	
^{31}Cl^{i}(2p)^{29}P	7700	100	7631	3	−0.7	U					90Bo24	*
	7610	60			0.3	U			Lis		91Bo32	
	7643	50			−0.2	U			Lis		92Ba01	*
	7627	15			0.3	o					98Ax02	
	7631	3				2					00Fy01	*
^{30}Ne(n,γ)^{31}Ne	190	130	170	130	−0.2	1	95	67 ^{31}Ne			14Na10	*
^{30}Si(^{18}O,^{17}F)^{31}Al	−12200	25	−12216.8	2.2	−0.7	U					88Wo02	
	−12237	35			0.6	U			Ber		89Bo.A	
^{30}Si(n,γ)^{31}Si	6589.1	0.7	6587.39	0.04	−2.4	U					70Be48	
	6587.5	0.8			−0.1	U					70Sp02	
	6588.4	0.3			−3.4	B					72Dz13	
	6587.32	0.20			0.4	U			MMn		90Is02	Z
	6587.39	0.05			0.1	4			ORn		92Ra19	Z
	6587.3970	0.0500			−0.1	o			PTc		97Ro26	*
	6587.39	0.14			0.0	U			Bdn		06Fi.A	
	6587.397	0.057			−0.1	4			PTc		01Pa52	
^{30}Si(d,p)^{31}Si	4368	7	4362.83	0.04	−0.7	U			MIT		64Sp12	
	4364.18	0.55			−2.5	U			Rez		90Pi05	*
^{30}Si(p,γ)^{31}P	7297.4	1.2	7296.553	0.022	−0.7	U					68Wo01	
^{31}S^{i}(p)^{30}P	273	10	259.79	0.29	−1.3	U			MSU		22Bu14	
	258	4			0.4	U			MSU		25Bu.A	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{31}Cl^{i}(p)^{30}S	12033	10	12026	3	−0.7	o					98Ax02	*
	12033	14			−0.5	U					00Fy01	*
^{31}Mg(β^-)^{31}Al	10150	700	11825	3	2.4	U					83De04	
^{31}Al(β^-)^{31}Si	7940	100	7998.3	2.2	0.6	U					73Go22	
^{31}Si(β^-)^{31}P	1471	8	1491.51	0.04	2.6	U					52Mo12	
	1486	12			0.5	U					52Wa12	
^{31}S(β^+)^{31}P	5412	30	5397.99	0.23	−0.5	U					60Wa04	
^{31}P(p,n)^{31}S	−6212.3	20.	−6180.34	0.23	1.7	o			ChR		58Go77	Y
	−6250	20			3.5	B			ChR		59Br06	Y
^{31}S^{i}(IT)^{31}S	6390.46	0.16				3					16Be05	
*^{31}Na−^{12}C ^{16}O ^{1}H$_3$	From updated frequency ratio ^{31}Na$^+$/(COH$_3$)$^+$ =0.999830942(91)										25Ly01	**
*^{31}Mg−^{12}C ^{16}O ^{1}H$_3$	From updated frequency ratio ^{31}Mg$^+$/(COH$_3$)$^+$ =0.999298732(137)										25Ly01	**
*^{31}Ar−u	combined results with 24Yu13										25Ch67	**
*^{31}K−u	from three-proton separation energy -4600(200) keV										HWJ201	**
*^{31}K−u	Trends in Mass Surface TMS suggest ^{31}K 2450 keV less bound										GAu	**
*O$_2$−^{31}P	For original doublet ^{31}P−O$_{1.938}$, D_M=–16382.522(0.096) μu										GAu	**
*^{31}Cl^{i}(2p)^{29}P	Large error in E_{cm} due to sequential decay kinematics										MMC122	**
*^{31}Cl^{i}(2p)^{29}P	reference also finds 3p emission at 4870 keV										92Ba01	**
*^{31}Cl^{i}(2p)^{29}P	Q_{2p}=7620(5), 6245(2), 5679(3), 5223(5) keV										00Fy01	**
*	to ground state and levels 3/2$^+$ at 1383.55, 5/2$^+$ at 1953,91, 3/2$^+$ 2422.7 keV										Ens125	**
*^{30}Ne(n,γ)^{31}Ne	Symmetrized from 150(+160–100) keV										14Na10	**
*^{30}Si(n,γ)^{31}Si	Original error 0.0005 increased for calibration										GAu	**
*^{30}Si(d,p)^{31}Si	Estimated systematic error 0.5 added to statistical error 0.23 keV										AHW	**
*^{31}Cl^{i}(p)^{30}S	Average of 3 branches										AHW	**
*^{31}Cl^{i}(p)^{30}S	E_p=11654(28), 9493(20), 8347(15), 8092(14) keV										00Fy01	**
^{32}Na−u	19720	636	19913	6	0.2	o			GA3	1.5	91Or01	
	19900	1100			0.0	U			TO4	1.5	91Zh24	
	20980	500			−1.4	o			GA5	1.5	00Sa21	
	20193	129			−1.4	U			GA7	1.5	07Ju03	
^{32}Mg−O$_2$	9280.9	3.5	9280.7	1.7	−0.1	1	25	25 ^{32}Mg	TT1	1.0	13Ch49	
^{32}Mg−u	−800	260	−890.1	1.7	−0.2	o			TO1	1.5	86Vi09	
	−890	270			−0.0	U			GA1	1.5	87Gi05	
	−924	214			0.1	U			GA3	1.5	91Or01	
	−820	130			−0.4	U			TO4	1.5	91Zh24	
	−1142	113			2.2	o			P40	1.0	01Lu20	
	−966	38			2.0	U			P40	1.0	06Ga04	*
	−983	22			4.2	B			P40	1.0	06Ga04	
^{32}Al−O$_2$	−1744	13	−1745	8	−0.1	o			TT1	1.0	12Ch.A	
	−1744.9	7.7				2			TT1	1.0	15Kw01	
^{32}Al−u	−12160	220	−11916	8	0.7	U			TO1	1.5	86Vi09	
	−11870	200			−0.2	U			GA1	1.5	87Gi05	
	−11877	104			−0.2	U			GA3	1.5	91Or01	
^{32}Si O$_2$−C$_5$ H$_4$	−67319.35	0.32				2			MS1	1.0	09Kw02	*
O$_2$−^{32}S	17754.2	1.0	17758.0647	0.0015	1.5	U			J1	2.5	68Ma45	
C$_2$ H$_8$−^{32}S	90531.3	1.4	90529.0819	0.0014	−0.6	U			J1	2.5	68Ma45	
^{32}S−O$_2$	−17758.0663	0.0020	−17758.0647	0.0015	0.8	1	53	43 ^{32}S	FS1	1.0	05Sh38	
^{32}S−C$_2$ D$_4$	−84335.9367	0.0019	−84335.9381	0.0014	−0.7	1	57	57 ^{32}S	FS1	1.0	05Sh38	
^{32}S−H C F	−34156.50	0.57	−34157.0205	0.0016	−0.9	U			MS1	1.0	09Kw02	*
^{12}C ^{32}S$_2$−^{152}Pr$_{.500}$	−21613.9	2.4	−21626.0	1.5	−5.0	B			RI1	1.0	24Ki16	*
^{1}H ^{12}C ^{32}S$_2$−^{154}Nd$_{.500}$	−12842.3	10.0	−12830.9	0.5	1.1	U			RI1	1.0	24Ki16	*
^{32}Na−^{39}K$_{.821}$	49808	40	49710	6	−2.5	B			TT1	1.0	17Ga20	*
^{32}Ar−^{39}K$_{.821}$	27434.8	1.9				2			MA8	1.0	03Bl17	
^{32}Na−^{32}S	47841.7	5.9				2			TR1	1.0	25Ly01	*
^{32}Na^{m}−^{32}S	47968.0	3.5				2			TR1	1.0	25Ly01	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{32}Mg−^{32}S	27038.7	2.0	27038.8	1.7	0.0	1	75	75 ^{32}Mg	TR1	1.0	25Ly01	*
C F_3−^{32}S O_2 H	25483.43	0.34	25484.0430	0.0030	1.8	U			MS1	1.0	09Kw02	*
C F_3−^{32}S O_2	33310.02	0.59	33309.0740	0.0030	−1.6	U			MS1	1.0	09Kw02	*
^{26}Na−^{32}Na$_{.325}$ ^{22}Na$_{.709}$	−8569	354	−9216	4	−0.7	U			P12	2.5	75Th08	
^{32}S(^{3}He,^{8}Li)^{27}P	−31277	35	−31359.4	0.6	−2.4	B			MSU		77Be13	
^{32}S(p,α)^{29}P	−4171	20	−4198.6	0.4	−1.4	U			Tky		64Ej05	
^{32}S(^{3}He,^{6}He)^{29}S	−25520	50	−25582	13	−1.2	U			MSU		73Be09	
^{30}Si(t,p)^{32}Si	7307	1	7305.57	0.30	−1.4	U			Str		80An.A	
^{32}S(d,α)^{30}P	4892	10	4896.13	0.07	0.4	U			MIT		64Sp12	
^{32}S(p,t)^{30}S	−19614	3	−19617.12	0.21	−1.0	U			MSU		74Ha02	
^{31}Si(n,γ)^{32}Si	9203.2180	0.0500	9200.0	0.3	−65.0	B			PTc		97Ro26	*
	9203.22	0.76			−4.3	B			PTc		01Pa52	
^{31}P(n,γ)^{32}P	7935.73	0.16	7935.65	0.04	−0.5	U			MMn		85Ke11	Z
	7935.65	0.04				2			ILn		89Mi16	Z
	7935.60	0.16			0.3	U			Bdn		06Fi.A	
^{31}P(d,p)^{32}P	5712	8	5711.08	0.04	−0.1	U			MIT		64Sp12	
^{31}P(p,γ)^{32}S	8864.9	0.9	8863.9637	0.0014	−1.0	U					72Co13	
	8862.5	3.			0.5	U					73Ve06	*
	8865.6	1.0			−1.6	U					73Ve08	Z
	8865.1	0.9			−1.3	U					74Vi02	
^{31}P(^{3}He,d)^{32}S	3356	13	3370.4886	0.0014	1.1	U			MIT		68Gr17	
^{32}S(p,d)^{31}S	−12817.8	1.5	−12819.73	0.23	−1.3	U			MSU		73Mo23	
^{32}S(^{3}He,α)^{31}S	5415	15	5533.32	0.23	7.9	B					66Gr26	
	5515	15			1.2	U			MIT		67Sp09	
	5486	20			2.4	U			Ors		67Ro17	
	5538	6			−0.8	U			CIT		70Mo08	
^{32}Cl^i(p)^{31}S	3462.8	3.1	3465.6	1.2	0.9	U					85Bj01	
	3457.6	2.1			3.8	B					93Sc16	
	3458.8	1.2			5.7	B					99Ad10	
	3465	7			0.1	o					20Ar04	
	3461.7	4.1			1.0	U					21Bl02	
^{32}Na(β^-)^{32}Mg	18300	1400	19378	6	0.8	U					83De04	
^{32}Si(β^-)^{32}P	213	7	227.2	0.3	2.0	U					64Br09	
	221.4	1.2			4.8	B					84Po09	
^{32}P(β^-)^{32}S	1707.6	0.7	1710.66	0.04	4.4	B					61Ni02	
	1710.1	0.7			0.8	U					68Fi04	
^{32}Cl(β^+)^{32}S	12720	30	12681.4	1.2	−1.3	U					68Ar03	*
^{32}S(p,n)^{32}Cl	−13470	14	−13463.7	1.2	0.4	U			Yal		69Ov01	Z
	−13470	9			0.7	U			BNL		71Go18	Z
^{32}S(^{3}He,t)^{32}Cl	−12699	15	−12700.0	1.2	−0.1	U					89Je07	
^{32}S(^{3}He,t)^{32}Cl−^{36}Ar()^{36}K	133.01	1.10	133.0	1.1	−0.0	1	100	100 ^{32}Cl	Mun		10Wr01	
^{32}S(π^+,π^-)^{32}Ar	−22813	50	−22793.2	1.8	0.4	U					80Bu15	
^{32}Cl^i(IT)^{32}Cl	5046.17	0.24				2					08Bh08	*
*^{32}Mg−u	Result from the "Plasma" experiment. Next item from "Rilis"										06Ga04	**
*^{32}Si O_2−C_5 H_4	For original doublet ^{32}Si O_2 H_3−C_5 H_7										GAu	**
*^{32}S−H C F	For original doublet ^{32}S O_2 H−H_2 C O_2 F										GAu	**
*^{12}C ^{32}S_2−^{152}Pr$_{.500}$	Reported $M-A$=–52024(10) keV, author confirmed uncertainty error										24Ki16	**
*^{1}H ^{12}C ^{32}S_2−^{154}Nd$_{.500}$	Reported $M-A$=–44752(16) keV, author confirmed uncertainty error										24Ki16	**
*^{32}Na−^{39}K$_{.821}$	could be a mixture with the iso. at 118 keV										HWJ258	**
*^{32}Na−^{32}S	From updated frequency ratio ^{32}Na^+/(^{32}$S)^+$ =1.001496385(184)										25Ly01	**
*^{32}Na^m−^{32}S	From updated frequency ratio ^{32}Nam^+/(^{32}$S)^+$ =1.001500334(109)										25Ly01	**
*^{32}Mg−^{32}S	From updated frequency ratio ^{32}Mg^+/(^{32}$S)^+$ =1.000845711(100)										25Ly01	**
*C F_3−^{32}S O_2 H	For original doublet ^{32}S O_2 H−(C F_3)$_{0.942}$, D_M=–25761.27(0.34) μu										GAu	**
*C F_3−^{32}S O_2	For original doublet ^{32}S O_2–(C F_3)$_{0.928}$, D_M=–33654.93(0.59) μu										GAu	**
*^{31}Si(n,γ)^{32}Si	Original error 0.0005 increased for calibration										GAu	**
*^{31}P(p,γ)^{32}S	T=3289(3) Q =–3185.3(3.) to ^{32}S^j at 12047.84(0.20) keV										Nub269	**
*^{32}Cl(β^+)^{32}S	E_{β^+}=9470(30) to 2^+ level at 2230.57 keV										Ens119	**
*^{32}Cl^i(IT)^{32}Cl	E(γ)=3877.5(0.3) to the level at 1168.5(0.2) keV										08Bh08	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{33}Na−u		27386	1601	25334	5	−0.9	o			GA3	1.5	91Or01	
		26370	1160			−0.6	o			GA5	1.5	00Sa21	
		25142	376			0.3	o			GA7	1.5	07Ju03	
		25529	322			−0.4	U			GA8	1.5	12Ga45	
^{33}Na−^{1}H O_2		27680.1	4.6				2			TR1	1.0	25Ly01	*
^{33}Mg−$O_{2.062}$		15813.3	3.1	15810.6	2.3	−0.9	–			TT1	1.0	13Ch49	
^{33}Mg−^{1}H $^{16}O_2$		7664.1	3.9	7670.3	2.3	1.6	–			TR1	1.0	25Ly01	*
^{33}Mg−$O_{2.062}$	ave.	15809.9	2.4	15810.6	2.3	0.3	1	90	90 ^{33}Mg			average	
^{33}Mg−u		5460	900	5324.5	2.3	−0.1	o			GA1	1.5	87Gi05	
		5203	318			0.3	U			GA3	1.5	91Or01	
		5710	180			−1.4	U			TO4	1.5	91Zh24	
		5311	24			0.6	U			P40	1.0	06Ga04	
^{33}Al−u		−9490	250	−9122	8	1.0	o			TO1	1.5	86Vi09	
		−9250	160			0.5	o			GA1	1.5	87Gi05	
		−9167	142			0.2	U			GA3	1.5	91Or01	
		−9020	120			−0.6	U			TO4	1.5	91Zh24	
		−9125	64			0.0	o			GT1	1.5	04Ma.A	
		−8957	100			−0.7	o			GT2	2.5	08Kn.A	
		−8915	128			−0.6	U			GT2	2.5	08Su19	
		−9209	62			1.4	U			LZ1	1.0	15Xu14	
^{33}Si O_2−^{13}C C_4 H_4		−66848.76	0.75				2			MS1	1.0	09Kw02	*
^{33}Cl−u		−22536.9	7.5	−22548.0	0.4	−1.5	U			LZ1	1.0	11Tu09	
^{33}Mg−^{39}K$_{.846}$		36035.7	7.4	36028.9	2.3	−0.9	1	10	10 ^{33}Mg	MA8	1.0	19As04	
^{33}Al−^{39}K$_{.846}$		21582.0	7.5				2			TT1	1.0	15Kw01	
^{33}Ar−^{39}K$_{.846}$		20629.86	0.43				2			MA8	1.0	03Bl17	
^{33}Ar−^{36}Ar$_{.917}$		19689.2	4.5	19686.7	0.4	−0.6	U			MA6	1.0	01He29	
^{33}S−^{32}S H		−8437.29682	0.00030	−8437.2968	0.0003	0.0	1	100	100 ^{33}S	MI3	1.0	05Ra34	
^{30}Si(α,p)^{33}P		−2965	10	−2959.7	1.1	0.5	U			ANL		68Be13	
^{33}S(n,α)^{30}Si		3497.6	5.	3493.506	0.022	−0.8	U			ILL		81Wa31	
		3496.9	5.0			−0.7	U					01Wa50	
^{31}P(^{3}He,p)^{33}S		9787	15	9787.5616	0.0015	0.0	U					71Gr04	
^{32}S(n,γ)^{33}S		8641.5	0.3	8641.6392	0.0005	0.5	o			MMn		80Is02	Z
		8641.82	0.10			−1.8	U			ORn		83Ra04	Z
		8641.60	0.03			1.3	U			MMn		85Ke08	Z
		8641.81	0.17			−1.0	U			Bdn		06Fi.A	
		8641.6398	0.0033			−0.2	U			NBS		06De21	
^{32}S(d,p)^{33}S		6420	6	6417.07299	0.00028	−0.5	U			MIT		64Sp12	
^{32}S(p,γ)^{33}Cl		2276.4	0.9	2276.8	0.4	0.4	–					59Ku79	
		2276.8	0.5			−0.1	–					76Al01	
^{32}S(d,n)^{33}Cl		62	9	52.2	0.4	−1.1	U					72El03	
^{32}S(^{3}He,d)^{33}Cl		−3218	15	−3216.7	0.4	0.1	U					66Gr26	
		−3217	5			0.1	U			CIT		70Mo08	
^{32}S(p,γ)^{33}Cl	ave.	2276.7	0.4	2276.8	0.4	0.2	1	80	80 ^{33}Cl			average	
^{32}S(p,γ)^{33}Cl^{i}		−3267.0	1.0	−3271.7	0.5	−4.7	B					70Ab15	
		−3271.4	2.0			−0.1	U					82Wi.A	
		−3271.5	0.8			−0.3	1	37	37 ^{33}Cl^{i}			02Py01	
^{33}Cl^{i}(p)^{32}S		3266.8	1.0	3271.7	0.5	4.9	B					87Bo21	
^{33}Si(β^-)^{33}P		5768	50	5823.0	1.3	1.1	U					73Go33	
^{33}P(β^-)^{33}S		249	2	248.5	1.1	−0.2	2					54Ni06	
		248.3	1.3			0.2	2					84Po09	
^{33}Cl(β^+)^{33}S		5532	50	5582.5	0.4	1.0	U					60Wa04	
^{33}Cl^{i}(IT)^{33}Cl		5548.5	0.4	5548.4	0.4	−0.1	1	83	63 ^{33}Cl^{i}			06Tr10	
*^{33}Na−^{1}H O_2	From updated frequency ratio $^{33}Na^+/(HO_2)^+$ =1.000838864(138)											25Ly01	**
*^{33}Mg−^{1}H $^{16}O_2$	From updated frequency ratio $^{33}Mg^+/(HO_2)^+$ =1.000232267(118)											25Ly01	**
*^{33}Si O_2−^{13}C C_4 H_4	For original doublet ^{33}Si O_2 H_3−^{13}C C_4 H_7											GAu	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{34}Na−u		34010	429				2			GA8	1.5	12Ga45	
^{34}Mg−$O_{2.126}$		19747	31	19739.5	2.4	−0.2	U			TT1	1.0	13Ch49	
^{34}Mg−$^{1}H_2$ $^{16}O_2$		3445.4	3.1	3448.7	2.4	1.1	1	60	60 ^{34}Mg	TR1	1.0	25Ly01	*
^{34}Mg−u		8855	476	8928.0	2.4	0.1	o			GA3	1.5	91Or01	
		9190	350			−0.5	U			TO4	1.5	91Zh24	
		9900	350			−1.9	U			GA5	1.5	00Sa21	
		9190	97			−1.8	U			GA7	1.5	07Ju03	
^{34}Mg−^{10}B $^{12}C_2$		−4004.9	4.4	−4008.9	2.4	−0.9	1	30	30 ^{34}Mg	TR1	1.0	25Ly01	*
^{34}Al−u		−3760	430	−3218.1	2.3	0.8	o			TO1	1.5	86Vi09	*
		−3400	250			0.5	o			GA1	1.5	87Gi05	*
		−3262	218			0.1	o			GA3	1.5	91Or01	*
		−2940	120			−1.5	U			TO4	1.5	91Zh24	*
		−3199	97			−0.1	U			GT1	1.5	04Ma.A	*
		−3328	86			0.9	U			GA7	1.5	07Ju03	*
^{34}S ^{16}O−^{100}Cd$_{.500}$		2616	13	2607.3	0.9	−0.7	U			RI1	1.0	26Ro.A	*
^{34}Mg−^{39}K$_{.872}$		40583.4	7.4	40575.9	2.4	−1.0	1	10	10 ^{34}Mg	MA8	1.0	19As04	
^{34}Al−^{39}K$_{.872}$		28438.0	7.8	28429.9	2.3	−1.0	o			TT1	1.0	15Kw01	
		28427.0	3.3			0.9	2			TT1	1.0	17Ga20	
		28432.4	3.1			−0.8	2			MA8	1.0	19As04	*
^{34}Si−^{39}K$_{.872}$		10185.99	0.86				2			MA8	1.0	19As04	
^{34}Ar−^{39}K$_{.872}$		11919.02	0.36	11918.04	0.08	−2.7	U			MA8	1.0	02He23	
^{34}Ar−^{36}Ar$_{.944}$		10907.4	3.8	10907.51	0.09	0.0	U			MA6	1.0	01He29	
^{34}Cl−^{34}S		5895.548	0.058	5895.48	0.04	−1.2	1	49	31 ^{34}Cl	JY1	1.0	09Er07	
^{34}Cl^m−^{34}S		6052.575	0.068	6052.60	0.04	0.4	1	41	31 ^{34}Cl^m	JY1	1.0	09Er07	
^{34}S−^{34}Ar		−12403.19	0.20	−12403.08	0.08	0.5	1	14	13 ^{34}Ar	JY1	1.0	11Er02	
^{34}Cl^m−^{34}Cl		157.05	0.11	157.124	0.029	0.7	U			JY1	1.0	09Er07	
		157.30	0.27			−0.7	U			JY1	1.0	11Er02	
^{34}Ar−^{34}Cl		6507.627	0.092	6507.60	0.07	−0.3	1	54	52 ^{34}Ar	JY1	1.0	11Er02	
^{34}Cl^m−^{34}Ar		−6350.41	0.11	−6350.48	0.07	−0.6	1	39	35 ^{34}Ar	JY1	1.0	11Er02	
C_4 H_3−^{34}P O		54914.59	0.87				2			MS1	1.0	09Kw02	*
^{30}Si(^{7}Li,^{3}He)^{34}P		100	40	91.6	0.8	−0.2	U					77Pe17	
^{31}P(α,p)^{34}S		629.9	2.9	627.09	0.04	−1.0	U			Har		73Ry01	
^{31}P(α,n)^{34}Cl		−5632	10	−5646.86	0.05	−1.5	U			Tal		70Um01	
		−5641.5	3.7			−1.4	U			Har		73Ry01	
^{34}S(d,α)^{32}P		5096	10	5083.99	0.06	−1.2	U					78Ba30	
^{32}S(^{3}He,n)^{34}Ar		−759	15	−777.34	0.08	−1.2	U			CIT		67Mi02	
^{34}S(^{13}C,^{14}O)^{33}Si		−14243	75	−14300.1	0.7	−0.8	U			Can		86Fi06	
^{33}S(n,γ)^{34}S		11417.12	0.10	11417.15	0.04	0.3	–			ORn		83Ra04	Z
		11417.22	0.23			−0.3	–			Bdn		06Fi.A	
^{33}S(d,p)^{34}S		9202	10	9192.58	0.04	−0.9	U			MIT		64Sp12	
		9195	6			−0.4	U			Utr		71Va21	
^{33}S(n,γ)^{34}S	ave.	11417.14	0.09	11417.15	0.04	0.1	1	24	24 ^{34}S			average	
^{33}S(p,γ)^{34}Cl		5142.42	0.20	5143.20	0.05	3.9	B			Oak		83Ra04	*
		5142.4	0.3			2.7	U			Utr		83Wa27	Z
		5143.29	0.07			−1.3	1	48	48 ^{34}Cl	Auc		94Li20	
^{34}K(p)^{33}Ar		608	17				3					24Dr01	
^{34}Si(β^-)^{34}P		4700	300	4557.0	1.1	−0.5	U					77Na05	
^{34}P(β^-)^{34}S		5383	45	5383.0	0.8	−0.0	U			ANB		73Go33	
^{34}S(t,^{3}He)^{34}P		−5368	20	−5364.4	0.8	0.2	U			LAl		77Aj01	
^{34}S(^{7}Li,^{7}Be)^{34}P		−6224	40	−6244.9	0.8	−0.5	U			Can		85Dr06	
^{34}Cl(β^+)^{34}S		5522	30	5491.60	0.04	−1.0	U					56Gr07	
^{34}S(p,n)^{34}Cl		−6252	10	−6273.95	0.04	−2.2	U			Tal		70Um01	
		−6271.9	1.9			−1.1	U			Har		75Fr.A	
		−6274.27	0.56			0.6	U			Auc		77Ba16	
		−6273.11	0.25			−3.4	F			Auc		92Ba.A	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{34}S(^{3}He,t)^{34}Cl	−5510.8	0.4	−5510.20	0.04	1.5	F			Mun		77Vo02	*
^{34}S(^{3}He,t)^{34}Cl−^{27}Al()^{27}Si	−678.7	2.3	−679.25	0.10	−0.2	U			ChR		74Ha35	
^{34}Cl^{m}(IT)^{34}Cl	146.36	0.03	146.360	0.027	−0.0	1	84	65 ^{34}Cl^{m}			Ens126	
*^{34}Mg−^{1}H$_2$ ^{16}O$_2$	From updated frequency ratio $^{34}Mg^{+}/(H_2O_2)^{+}$ =1.000101321(91)										25Ly01	**
*^{34}Mg−^{10}B ^{12}C$_2$	From updated frequency ratio $^{34}Mg^{+}/(^{10}BC_2)^{+}$ =0.999882252(130)										25Ly01	**
*^{34}Al−u	Possible isomeric mixture 26(1) ms, E=46.5 keV										12Ro25	**
*^{34}S ^{16}O−^{100}Cd$_{.500}$	from frequency ratio $^{34}S^{+}_{160}/(^{100}Cd)^{++}$ =1.000052352(140)										26Ro.A	**
*^{34}S ^{16}O−^{100}Cd$_{.500}$	Original error 5.8; added systematic error 10 keV										SNa265	**
*^{34}Al−^{39}K$_{.872}$	Authors stated only ground state										19As04	**
*C$_4$ H$_3$−^{34}P O	For original doublet ^{34}P H$_2$ O−C$_4$ H$_5$										GAu	**
*^{33}S(p,γ)^{34}Cl	E_p=974.76(0.15,Z) to 6088.20(0.10,Z) level										83Ra04	**
*^{34}S(p,n)^{34}Cl	F : disturbed by resonance; at least 0.5 keV uncertain										94Li20	**
*^{34}S(^{3}He,t)^{34}Cl	F : rejected in reference of same group										09Fa15	**
^{35}Mg−u	18669	1721	16672	6	−0.8	o			GA3	1.5	91Or01	
	18830	1070			−1.3	o			GA5	1.5	00Sa21	
	16790	193			−0.4	U			GA7	1.5	07Ju03	
^{35}Mg−^{11}B ^{12}C$_2$	7371	12	7366	6	−0.4	1	27	27 ^{35}Mg	TR1	1.0	25Ly01	*
^{35}Al−u	−340	460	−240	8	0.1	o			GA1	1.5	87Gi05	
	−296	298			0.1	o			GA3	1.5	91Or01	
	80	190			−1.1	U			TO4	1.5	91Zh24	
	−236	75			−0.0	U			GA7	1.5	07Ju03	
C$_3$−^{35}Cl H	23320.8	0.3	23322.27	0.04	2.0	U			M17	2.5	66Be10	
	23322.239	0.034			0.6	1	56	56 ^{35}Cl	B07	1.5	71Sm01	
	23321.83	0.63			0.3	U			J5	2.5	72Ka57	
	23322.328	0.325			−0.1	U			J5	2.5	72Ka57	
C$_5$ H$_{10}$−^{35}Cl$_2$	140549.37	2.98	140544.93	0.08	−0.6	U			C2	2.5	65De09	
	140545.01	0.13			−0.4	1	15	15 ^{35}Cl	B07	1.5	71Sm01	
C$_4$ H$_6$ O−^{35}Cl$_2$	104153.75	3.45	104159.42	0.08	0.7	U			C2	2.5	65De09	
C$_2$ D$_6$−^{35}Cl H	107934.90	0.54	107932.94	0.04	−1.5	U			J5	2.5	72Ka57	
	107933.422	0.538			−0.4	U			J5	2.5	72Ka57	
C$_3$ H−D ^{35}Cl	24871.92	0.75	24870.56	0.04	−0.7	U			C2	2.5	65De09	
C$_8$ H$_9$−^{35}Cl$_3$	163867.25	0.90	163867.20	0.11	−0.0	U			A2	2.5	70St25	
^{35}Ar−u	−24747.3	4.3	−24742.3	0.7	1.2	U			LZ1	1.0	11Tu09	
^{35}Al−^{39}K$_{.897}$	32315.1	7.9				2			TT1	1.0	17Ga20	
^{35}K−^{39}K$_{.897}$	20560.69	0.55				2			MA8	1.0	07Ya08	
^{35}Mg−^{35}Cl	47817.2	7.3	47819	6	0.2	1	73	73 ^{35}Mg	TR1	1.0	25Ly01	*
^{35}Cl(p,α)^{32}S	1862	5	1866.06	0.04	0.8	U			Bar		57Va03	Y
	1865	8			0.1	U			MIT		64Sp12	
^{32}S(α,p)^{35}Cl	−1862	17	−1866.06	0.04	−0.2	U			MIT		64Sp12	
^{32}S(α,n)^{35}Ar	−8751	18	−8614.7	0.7	7.6	B			Tal		63Ne05	
^{35}Cl(d,α)^{33}S	8285	10	8283.14	0.04	−0.2	U			MIT		64Sp12	
^{33}S(^{3}He,n)^{35}Ar	3335	16	3321.3	0.7	−0.9	U					75Da14	
^{35}K^{i}(2p)^{33}Cl	4311	40				2					85Ay01	
^{34}S(^{18}O,^{17}F)^{35}P	−7796	40	−7808.4	1.9	−0.3	U			Can		88Or01	
^{34}S(n,γ)^{35}S	6986.00	0.10	6985.84	0.04	−1.6	–			ORn		83Ra04	Z
	6985.84	0.05			−0.1	–			MMn		85Ke08	Z
	6986.09	0.14			−1.8	U			Bdn		06Fi.A	
^{34}S(d,p)^{35}S	4762	10	4761.27	0.04	−0.1	U			MIT		64Sp12	
	4757	5			0.9	U			Utr		71Va18	
^{34}S(n,γ)^{35}S ave.	6985.87	0.04	6985.84	0.04	−0.8	1	75	46 ^{34}S			average	
^{34}S(p,γ)^{35}Cl	6367.4	1.6	6370.81	0.04	2.1	U					72Hu10	
	6370.7	0.4			0.3	U					76Sp08	Z
	6370.70	0.20			0.6	U			Oak		83Ra04	*
^{35}Cl(γ,n)^{34}Cl	−12660	40	−12644.76	0.05	0.4	U					61Sa11	
^{35}Ar(p,d)^{34}Ar	−10491	25	−10515.7	0.7	−1.0	U			MT2		21La04	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{35}P(β^-)^{35}S	3909	75	3988.4	1.9	1.1	U					72Go31	
^{35}S(β^-)^{35}Cl	167.4	0.2	167.321	0.026	−0.4	U					57Co62	
	166.80	0.15			3.5	B					85Al11	
	167.288	0.100			0.3	U					85Ap01	*
	166.93	0.2			2.0	o					85Ma59	
	167.4	0.1			−0.8	U					85Oh06	*
	166.7	0.2			3.1	B					89Si04	*
	167.56	0.03			−8.0	B					92Ch27	*
	167.35	0.10			−0.3	U					93Ab11	*
	167.23	0.10			0.9	U					93Be21	*
	167.27	0.10			0.5	U					93Mo01	*
	167.334	0.027			−0.5	1	91	71 ^{35}S			00Ho13	
^{35}Cl(n,p)^{35}S	612	4	615.026	0.026	0.8	U			BNL		68Sc01	
^{35}Ar(β^+)^{35}Cl	5980	40	5966.2	0.7	−0.3	U					56Ki29	
	5950	50			0.3	U					60Wa04	
^{35}Cl(p,n)^{35}Ar	−6747.2	1.6	−6748.6	0.7	−0.9	2			Har		75Fr.A	Z
	−6747.9	1.0			−0.7	2			Auc		77Wh03	Z
	−6750.4	1.2			1.5	2			Mtr		78Az01	*
*^{35}Mg−^{11}B ^{12}C$_2$	From updated frequency ratio ^{45}Mg$^+$/(^{11}BC$_2$)$^+$ =1.000210554(342)										25Ly01	**
*^{35}Mg−^{35}Cl	From updated frequency ratio ^{35}Mg$^+$/(^{35}Cl)$^+$ =1.001367445(208)										25Ly01	**
*^{34}S(p,γ)^{35}Cl	E_p=1264.97(0.13,Z) to 7598.91(0.15,Z) level										83Ra04	**
*^{35}S(β^-)^{35}Cl	Original error (0.030) increased to 0.100										AHW	**
*^{35}Cl(p,n)^{35}Ar	Original T=6942.2(2.2) recalibrated 6945.5(1.2)										GAu	**
^{36}Mg−u	24930	1610	21880	740	−1.3	o			GA5	1.5	00Sa21	
	21879	494				2			GA7	1.5	07Ju03	
^{36}Al−u	6187	421	6390	160	0.3	o			GA3	1.5	91Or01	
	6500	400			−0.2	U			TO4	1.5	91Zh24	
	6140	310			0.5	o			GA5	1.5	00Sa21	
	6388	107				2			GA7	1.5	07Ju03	
^{36}Si−u	−13850	640	−13350	80	0.5	U			TO1	1.5	86Vi09	
	−13490	320			0.3	o			GA1	1.5	87Gi05	
	−13578	191			0.8	o			GA3	1.5	91Or01	
	−13110	150			−1.1	2			TO4	1.5	91Zh24	
	−13376	75			0.2	2			GT1	1.5	04Ma.A	
	−13280	118			−0.4	2			GA7	1.5	07Ju03	
	−13484	163			0.8	2			LZ1	1.0	15Xu14	
^{36}Ar−u	−32454.895	0.015	−32454.894	0.029	0.1	o			ST2	1.0	02Bf02	
	−32454.895	0.029			0.0	1	100	100 ^{36}Ar	ST2	1.0	03Fr08	
^{36}Ar ^{1}H−^{39}K$_{.949}$	9812.47	0.40	9812.688	0.029	0.5	U			RKM	1.0	25Xi16	
^{36}Ca−u	−6971	18	−6960	6	0.6	2			MS1	1.0	21Su04	*
	−6959.1	6.4			−0.2	2			MS1	1.0	21Su04	*
^{36}K−^{39}K$_{.923}$	14800.99	0.38	14800.9	0.4	−0.4	1	90	90 ^{36}K	MA8	1.0	07Ya08	
^{36}Ar(^{3}He,^{8}Li)^{31}Cl	−29180	50	−29212	3	−0.6	U			MSU		77Be13	
^{36}S(^{48}Ca,^{52}V)^{32}Al	−12651	370	−12347	7	0.8	o			Dar		87Ch.A	
^{36}S(^{48}Ca,^{51}V)^{33}Al	−14150	140	−14189	7	−0.3	U			Dar		86Wo07	
^{36}S(^{14}C,^{17}O)^{33}Si	−6380	20	−6321.2	0.7	2.9	U			Mun		84Ma49	
^{36}S(^{11}B,^{14}N)^{33}Si	−4311	30	−4345.5	0.7	−1.2	U			Can		85Fi03	
^{36}Ar(^{3}He,^{6}He)^{33}Ar	−23512	30	−23508.1	0.4	0.1	U			MSU		74Na07	
^{36}S(^{11}B,^{13}N)^{34}Si	−7327	25	−7350.2	0.9	−0.9	U			Can		85Fi03	
^{36}S(^{14}C,^{16}O)^{34}Si	−2989	20	−2915.6	0.8	3.7	B			Mun		84Ma49	
^{36}S(^{64}Ni,^{66}Zn)^{34}Si	−8890	41	−8872.1	1.1	0.4	o			Dar		85Wo07	*
	−8903	33			0.9	U			Dar		86Sm05	*
^{36}S(d,α)^{34}P	4604.4	5.	4595.4	0.8	−1.8	U					82So.A	*
^{36}Ar(p,t)^{34}Ar	−19513	3	−19514.09	0.08	−0.4	U			MSU		74Ha02	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{36}Ar(p,t)^{34}Ar^{i}	−27473	50	−27448	5	0.5	U					69Br21	*
	−27448	5				2					72Pa02	*
^{36}S(^{14}C,^{15}O)^{35}Si	−16184	50	−16110	40	1.5	2			Mun		84Ma49	
^{36}S(^{13}C,^{14}O)^{35}Si	−21122	60	−21160	40	−0.6	2			Can		86Fi06	
^{36}S(^{64}Ni,^{65}Zn)^{35}Si	−17250	100	−17460	40	−2.1	2			Dar		86Sm05	*
^{36}S(d,^{3}He)^{35}P	−7607	5	−7601.8	1.9	1.0	2			BNL		84Th08	
	−7601	2			−0.4	2			Hei		85Kh04	
^{36}S(^{14}C,^{15}N)^{35}P	−2927	10	−2887.9	1.9	3.9	B			Mun		84Ma49	*
^{36}S(^{6}Li,^{7}Be)^{35}P	−7521	17	−7488.5	1.9	1.9	U			Can		85Dr06	
^{36}S(^{64}Ni,^{65}Cu)^{35}P	−5659	34	−5641.5	2.0	0.5	U			Dar		85Wo.A	
^{36}S(p,d)^{35}S^{i}	−16791	7	−16801	6	−1.4	1	67	67 ^{35}S^{i}			26Jo01	
^{35}Cl(n,γ)^{36}Cl	8579.73	0.20	8579.794	0.005	0.3	U			BNn		78St25	Z
	8579.7	0.3			0.3	o			MMn		80Is02	Z
	8579.81	0.20			−0.1	U			MMn		81Ke02	Z
	8579.66	0.10			1.3	U					81Su.A	Z
	8579.61	0.09			2.0	U			ILn		82Kr12	Z
	8579.67	0.17			0.7	U			Bdn		06Fi.A	
	8579.7945	0.0048			−0.0	1	100	99 ^{36}Cl	NBS		06De21	
^{35}Cl(d,p)^{36}Cl	6360	8	6355.228	0.005	−0.6	U			MIT		64Sp12	
^{35}Cl(p,γ)^{36}Ar	8506.1	0.5	8506.98	0.04	1.8	U					72Ho40	Z
^{35}Cl(p,γ)^{36}Ar^{j}	−2346.8	1.5	−2345.2	1.2	1.1	2					76Hu01	
	−2342.5	1.9			−1.4	2					76Ma40	
^{36}Ar(d,t)^{35}Ar	−9007	10	−8998.3	0.7	0.9	U			Yal		70Wh04	
^{36}K^{i}(p)^{35}Ar	2592	21	2623.8	2.3	1.5	U			Brk		81Ay01	
	2623.8	2.3				3					95Ga16	
^{36}S(^{7}Li,^{7}Be)^{36}P	−11277	27	−11275	13	0.1	2			Can		85Dr06	
^{36}S(^{14}C,^{14}N)^{36}P	−10256	15	−10257	13	−0.0	2			Mun		84Ma49	
^{36}Cl(β^{+})^{36}S	1137	18	1142.13	0.19	0.3	U					68Pi03	
^{36}Cl(ε)^{36}S	1180	15			−2.5	U					64Li10	
	1160	18			−1.0	U					65Be19	
^{36}S(p,n)^{36}Cl	−1924.64	0.31	−1924.48	0.19	0.5	1	37	36 ^{36}S			01Wa50	
^{36}Cl(β^{-})^{36}Ar	708.7	0.6	709.54	0.04	1.4	U					67Sp06	
^{36}Ar(p,n)^{36}K	−13588.3	8.	−13596.8	0.3	−1.1	U			BNL		71Go18	Z
	−13618	23			0.9	U					71Ja09	
^{36}Ar(^{3}He,t)^{36}K	−12930	40	−12833.0	0.3	2.4	U			Duk		70Dz04	
*^{36}Ca−u	Represents frequency ratio $^{36}Ca^{+}/(^{12}C_3)^{+}$ =1.00019369(50)										21Su04	**
*^{36}Ca−u	Represents frequency ratio $^{36}Ca^{++}/(H_2O)^{+}$ =1000780.41(18)										21Su04	**
*^{36}S(^{64}Ni,^{66}Zn)^{34}Si	Calibrated with ^{36}S(^{64}Ni,^{62}Ni) $M-A$=–26862(12) now−26861(7)										AHW	**
*^{36}S(d,α)^{34}P	Original error 1.2 judged too small										GAu	**
*^{36}Ar(p,t)^{34}Ar^{i}	IT=7950(50); Q rebuilt, estimated with 72Pa02 Q =–19523 for ground state										MMC12a	**
*^{36}Ar(p,t)^{34}Ar^{i}	IT=7925(5); Q rebuilt with author's Q =–19523 for ground state										MMC128	**
*^{36}S(^{64}Ni,^{65}Zn)^{35}Si	$M-A$=–14482(59) for average of ground state and 54, 114, 207 levels										86Sm05	**
*^{36}S(^{14}C,^{15}N)^{35}P	Original report –2693 is a typo										GAu	**
^{37}Al−u	10310	579	10530	190	0.3	o			GA3	1.5	91Or01	
	10900	450			−0.5	o			GA5	1.5	00Sa21	
	10531	129				2			GA7	1.5	07Ju03	
^{37}Si−u	−7550	1410	−7050	120	0.2	o			GA1	1.5	87Gi05	
	−7310	305			0.6	o			GA3	1.5	91Or01	
	−6930	150			−0.6	2			TO4	1.5	91Zh24	
	−7107	97			0.4	2			GA7	1.5	07Ju03	
^{37}P−u	−20740	430	−20390	40	0.5	U			TO1	1.5	86Vi09	
	−19910	190			−1.7	o			GA1	1.5	87Gi05	
	−20442	200			0.2	U			GA3	1.5	91Or01	
C_3 H−^{37}Cl	41924.73	1.09	41922.46	0.06	−0.8	U			C2	2.5	65De09	
	41922.2	0.2			0.5	U			M17	2.5	66Be10	
	41922.176	0.305			0.4	U			J5	2.5	72Ka57	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$C_2\ D_8-{}^{37}Cl\ H_3$	123436.51	0.12	123436.56	0.06	0.3	U			B07	1.5	71Sm01	
$C_3\ H_6\ O_2-{}^{37}Cl_2$	104974.24	0.08	104974.29	0.11	0.5	1	86	86 ^{37}Cl	B07	1.5	71Sm01	
$C_3\ H_5-D_2\ {}^{37}Cl$	45020.96	1.14	45019.04	0.06	−0.7	U			C2	2.5	65De09	
$C_8\ H_{15}-{}^{37}Cl_3$	219665.80	0.90	219667.78	0.17	0.9	U			A2	2.5	70St25	
$C_3\ H_3-D\ {}^{37}Cl$	43473.27	1.33	43470.75	0.06	−0.8	U			C2	2.5	65De09	
$^{37}K-u$	−26632.5	6.4	−26624.11	0.10	1.3	U			LZ1	1.0	11Tu09	
$H_3\ O-{}^{37}Ca_{.514}$	25638.22	0.35				2			MS1	1.0	07Ri08	*
$D_2\ {}^{35}Cl-H_2\ {}^{37}Cl$	15505.41	0.71	15503.62	0.07	−1.0	U			C2	2.5	65De09	
	15503.80	0.09			−0.8	U			H31	2.5	77So02	
$C_5\ H_{12}-{}^{35}Cl\ {}^{37}Cl$	159145.17	0.12	159145.12	0.07	−0.3	1	14	9 ^{37}Cl	B07	1.5	71Sm01	
$H_2\ {}^{35}Cl-{}^{37}Cl$	18600.0	0.4	18600.19	0.07	0.2	U			M17	2.5	66Be10	
$D\ {}^{35}Cl-{}^{37}Cl$	17052.95	1.02	17051.91	0.07	−0.4	U			C2	2.5	65De09	
	17051.816	0.185			0.2	U			J5	2.5	72Ka57	
$^{37}Cl-{}^{39}K_{.949}$	350.2	6.6	345.12	0.06	−0.8	U			RKM	1.0	25Xi16	
$^{37}K-{}^{39}K_{.949}$	7817.98	0.33	7818.44	0.10	1.4	U			MA8	1.0	07Ya08	
$^{37}Cl(p,\alpha)^{34}S$	3030	6	3034.19	0.07	0.7	U			Bar		57Va03	Y
	3029	8			0.6	U			MIT		64Sp12	
$^{37}Ar(n,\alpha)^{34}S$	4630	7	4630.41	0.21	0.1	U			ILL		78As06	
$^{34}S(\alpha,n)^{37}Ar$	−4625	90	−4630.41	0.21	−0.1	U			Tal		63Ne05	
$^{37}Cl(d,\alpha)^{35}S$	7791	12	7795.46	0.07	0.4	U			MIT		64Sp12	
$^{37}Cl(p,{}^3He)^{35}S^i$	−19713	10	−19693	6	2.0	1	33	33 $^{35}S^i$			75Gu15	
$^{35}Cl({}^3He,p)^{37}Ar$	9582	15	9576.40	0.21	−0.4	U			MIT		67Sp09	
$^{37}K(p,t)^{35}K$	−21256	110	−21288.1	0.5	−0.3	U			MT2		23La09	
$^{37}Ca(p,t)^{35}Ca$	−25573	105				3			MT2		23La09	*
$^{36}Mg(n,\gamma)^{37}Mg$	240	110				3					14Ko14	*
$^{36}S(^{18}O,{}^{17}F)^{37}P$	−14410	40	−14400	40	0.2	2			Can		88Or.A	*
$^{36}S(^{48}Ca,{}^{47}Sc)^{37}P$	−11490	120	−11560	40	−0.6	2			Dar		88Fi04	*
$^{36}S(n,\gamma)^{37}S$	4303.52	0.12	4303.60	0.06	0.7	2			ORn		84Ra09	Z
	4303.61	0.09			−0.1	2			Bdn		06Fi.A	
$^{36}S(d,p)^{37}S$	2079.12	0.13	2079.04	0.06	−0.6	2					84Pi03	
$^{36}S(^{14}C,{}^{13}C)^{37}S$	−3874	7	−3872.83	0.06	0.2	U			Mun		84Ma49	
$^{37}Cl(^{13}C,{}^{14}O)^{36}P$	−16433	50	−16393	13	0.8	U			Can		88Or01	
$^{36}S(p,\gamma)^{37}Cl$	8386.47	0.23	8386.39	0.19	−0.3	1	65	64 ^{36}S	Utr		84No05	Z
$^{36}S(p,\gamma)^{37}Cl^i$	−1835.5	0.3				2					84No05	
$^{36}Ar(n,\gamma)^{37}Ar$	8791.1	1.0	8787.46	0.21	−3.6	B					68Wi25	Z
	8788.8	1.2			−1.1	U					70Ha56	Z
	8789.9	0.9			−2.7	U			Bdn		06Fi.A	
$^{36}Ar(p,\gamma)^{37}K$	1857.3	1.0	1857.63	0.09	0.3	U					64Ar17	
	1857.63	0.09				2			Utr		88De03	Z
$^{36}Ar(d,n)^{37}K$	−320	100	−366.94	0.09	−0.5	U			Yal		61Ya01	
$^{36}Ar(p,\gamma)^{37}K^i$	−3192.6	0.8				2					88De03	
$^{37}Ca(p,d)^{36}Ca$	−12508	42	−12499	6	0.2	U			MT2		21La04	
$^{36}Ca(p,\gamma)^{37}Sc$	−2690	410				3					24Dr01	
$^{37}S(\beta^-)^{37}Cl$	4750	40	4865.14	0.20	2.9	U					67Wi14	
$^{37}Cl(t,{}^3He)^{37}S$	−4854	30	−4846.54	0.20	0.2	U			LAl		70Aj01	
$^{37}Ar(\varepsilon)^{37}Cl$	818	15	813.87	0.20	−0.3	U					53An01	
$^{37}Cl(p,n)^{37}Ar$	−1595.5	4.0	−1596.22	0.20	−0.2	U			Wis		50Ri59	Y
	−1595.4	1.0			−0.8	U			MIT		52Sc09	Z
	−1596.9	2.4			0.3	U			Oak		64Jo11	
	−1596.8	1.0			0.6	U			Duk		66Pa18	Z
	−1596.22	0.20				2			PTB		98Bo30	
	−1596.3	1.0			0.1	U					01Wa50	
$^{37}K(\beta^+)^{37}Ar$	6120	70	6147.48	0.23	0.4	U					58Su60	
	6170	70			−0.3	U					60Wa04	
$*H_3\ O-{}^{37}Ca_{.514}$	Error in Table II : $M-A$=13135.7(1.4) corrected to –13136.06(0.64) keV										07Ri08	**
$*^{37}Ca(p,t)^{35}Ca$	Q value reconstructed from mass excess										23La09	**
$*^{36}Mg(n,\gamma)^{37}Mg$	Symmetrized from 220(+120–90) keV										14Ko14	**
$*^{36}S(^{18}O,{}^{17}F)^{37}P$	And Q =–13650(40), $M-A$=–19750(40) if other peak is ground state one										88Or.A	**
$*^{36}S(^{48}Ca,{}^{47}Sc)^{37}P$	And Q =–11569(80), $M-A$=–18980(80) if other peak due to ^{47}Sc 807.89 level										88Fi04	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{38}Al−u	15240	1500	17680#	220#	1.1	o			GA4	1.5	00Sa21	
	17980	920			−0.2	o			GA5	1.5	00Sa21	
	17402	268			0.7	D			GA7	1.5	07Ju03	*
^{38}Si−u	−4510	180	−4480	110	0.1	o			GA4	1.5	00Sa21	
	−4020	290			−1.1	U			TO4	1.5	91Zh24	
	−4100	320			−0.8	o			GA5	1.5	00Sa21	
	−4477	75				2			GA7	1.5	07Ju03	
^{38}P−u	−14420	620	−15700	80	−1.4	U			GA1	1.5	87Gi05	
	−15910	140			1.0	2			GA4	1.5	00Sa21	
	−15530	150			−0.7	2			TO4	1.5	91Zh24	
	−16110	310			0.9	U			GA5	1.5	00Sa21	
	−15717	75			0.2	o			GT1	1.5	04Ma.A	
	−15660	100			−0.1	2			GT2	2.5	08Kn.A	
	−15688	97			−0.1	2			LZ1	1.0	15Xu14	
^{38}Ar ^{1}H−^{78}Zn$_{.500}$	1412.8	3.6	1412.5	1.1	−0.1	U			RKM	1.0	25Xi16	
^{38}Ar ^{1}H−^{39}K	6845.4	7.4	6850.65	0.21	0.7	U			RKM	1.0	25Ki14	
^{38}Ca−H$_6$ O$_2$	−60460.24	0.30	−60460.21	0.21	0.1	o			MS1	1.0	06Bo11	
	−60460.24	0.30			0.1	1	48	48 ^{38}Ca	MS1	1.0	07Ri08	
^{38}Ar−^{39}K$_{.974}$	−1917.88	0.37	−1918.01	0.21	−0.4	1	32	32 ^{38}Ar	MA8	1.0	02He23	
	−1915.2	1.5			−1.9	U			RKM	1.0	25Xi16	
^{38}K−^{39}K$_{.974}$	4430.88	0.44	4431.00	0.21	0.3	1	23	23 ^{38}K	MA8	1.0	07Ya08	
^{38}Ca ^{19}F−^{39}K$_{1.462}$	27783.80	0.63	27783.51	0.21	−0.5	U			MA8	1.0	07Ge07	
^{38}K−^{38}Ar	6348.974	0.068	6349.01	0.05	0.6	1	50	27 ^{38}K	JY1	1.0	09Er07	
^{38}K^{m}−^{38}Ar	6488.743	0.049	6488.73	0.04	−0.2	1	72	45 ^{38}K^{m}	JY1	1.0	09Er07	
^{38}Ar−^{38}Ca	−13587.18	0.12	−13587.12	0.07	0.5	1	32	17 ^{38}Ar	JY1	1.0	11Er02	
^{38}K^{m}−^{38}K	139.698	0.065	139.72	0.05	0.3	–			JY1	1.0	09Er07	
	139.78	0.14			−0.4	–			JY1	1.0	11Er02	
ave.	139.71	0.06			0.1	1	60	34 ^{38}K^{m}			average	
^{38}Ca−^{38}K	7238.04	0.10	7238.11	0.07	0.7	1	45	25 ^{38}K	JY1	1.0	11Er02	
^{38}K^{m}−^{38}Ca	−7098.43	0.11	−7098.39	0.07	0.4	1	37	21 ^{38}K^{m}	JY1	1.0	11Er02	
^{24}Mg(^{16}O,2n)^{38}Ca	−12727	30	−12754.71	0.19	−0.9	U					72Zi02	*
^{35}Cl(α,p)^{38}Ar	837.2	2.4	837.25	0.20	0.0	U			Har		75Sq01	
^{35}Cl(α,n)^{38}K	−5862.1	1.5	−5859.17	0.20	2.0	U			Mun		76Sh24	Z
	−5858.7	2.9			−0.2	U			Har		75Sq01	*
^{36}S(t,p)^{38}S	3838	30	3858	7	0.7	U					85Da15	
^{36}S(^{14}C,^{12}C)^{38}S	−781	10	−783	7	−0.2	R			Mun		84Ma49	
^{36}Ar(^{3}He,n)^{38}Ca	−1365	21	−1313.14	0.20	2.5	U			CIT		69Sh04	
^{37}Cl(n,γ)^{38}Cl	6107.84	0.30	6107.88	0.08	0.1	U					73Sp06	Z
	6107.95	0.10			−0.7	2			MMn		81Ke02	Z
	6107.73	0.15			1.0	2			Bdn		06Fi.A	
^{37}Cl(d,p)^{38}Cl	3885	8	3883.32	0.08	−0.2	U			MIT		64Sp12	
	3883.28	0.50			0.1	U			Rez		90Pi05	*
^{37}Cl(p,γ)^{38}Ar	10243.0	1.0	10242.24	0.20	−0.8	U					68En01	Z
^{38}Ca(p,d)^{37}Ca	−14764	13	−14769.2	0.7	−0.4	U			MT2		21La04	
^{37}Ca(p,γ)^{38}Sc	−1191	14				3					24Dr01	
^{38}S(β^-)^{38}Cl	2947	20	2937	7	−0.5	3					71En01	
	2936	12			0.1	3					72Vi11	
^{38}Cl(β^-)^{38}Ar	4913	5	4916.71	0.22	0.7	U					68Va06	
^{38}K(β^+)^{38}Ar	5870	30	5914.07	0.04	1.5	U					56Gr07	*
	5790	50			2.5	U					67Va27	*
^{38}Ar(p,n)^{38}K	−6695.5	4.	−6696.41	0.04	−0.2	U			Har		75Sq01	
	−6695.65	0.70			−1.1	U					78Ja06	Z
^{38}Ar(p,n)^{38}K^{m}	−6826.73	0.12	−6826.56	0.04	1.4	U			Auc		98Ha36	Z
*^{38}Al−u	Trends in Mass Surface TMS suggest ^{38}Al 260 keV less bound										HWJ266	**
*^{24}Mg(^{16}O,2n)^{38}Ca	E(^{16}O)=24880(30) to 2$^+$ level at 2213.13(0.10) keV										Ens082	**
*^{35}Cl(α,n)^{38}K	Q =–5989.1(2.9,Z) to ^{38}K^{m} at 130.15(0.04) keV										Nub269	**
*^{37}Cl(d,p)^{38}Cl	Estimated systematic error 0.5 added to statistical error 0.064 keV										AHW	**
*^{38}K(β^+)^{38}Ar	E_{β^+}=2680(30) 2600(50) respectively, to 2$^+$ level at 2167.64 keV										Ens082	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{39}Al−u	22970	1580	23070#	320#	0.0	o			GA5	1.5	00Sa21	
	21653	676			1.4	D			GA7	1.5	07Ju03	*
^{39}Si−u	1900	540	2490	150	0.7	o			GA4	1.5	00Sa21	
	2210	490			0.4	o			GA5	1.5	00Sa21	
	2491	97				2			GA7	1.5	07Ju03	
^{39}P−u	−13890	140	−13710	120	0.8	2			GA4	1.5	00Sa21	
	−13580	160			−0.6	2			TO4	1.5	91Zh24	
	−13870	280			0.4	2			GA5	1.5	00Sa21	
	−13602	140			−0.5	2			GT1	1.5	04Ma.A	
^{39}K−^{23}Na$_{1.696}$	−18942.88	0.58	−18942.222	0.006	0.8	U			Ma8	1.5	08Mu05	
^{39}Ca−u	−29278.8	6.4	−29289.2	0.6	−1.6	U			LZ1	1.0	11Tu09	
^{39}K−^{36}Ar$_{1.083}$	−1144.65	0.44	−1144.87	0.03	−0.5	U			MA8	1.0	02He23	
	−1144.83	0.40			−0.1	U			MA8	1.0	03Bl17	
^{39}K−^{37}K$_{1.054}$	−8231.29	0.53	−8231.71	0.11	−0.5	U			Ma8	1.5	08Mu05	
^{39}Ca ^{19}F−^{39}K$_{1.487}$	23082.43	0.64	23082.4	0.6	−0.0	1	100	100 ^{39}Ca	MA8	1.0	08Ge08	
^{39}K−^{40}Ar	1323.3631	0.0043	1323.363	0.004	−0.1	1	100	100 ^{39}K	FS1	1.0	10Mo30	
^{39}K(p,α)^{36}Ar	1287	7	1288.398	0.027	0.2	U			MIT		64Sp12	
^{37}Cl(t,p)^{39}Cl	5701.9	2.5	5699.5	1.7	−1.0	2			Str		84An03	
^{39}K(p,^{3}He)^{37}Ar^i	−15493.4	6.				2			MSU		73Be23	*
^{39}K(p,t)^{37}K^i	−21713.1	3.	−21718.1	0.8	−1.7	U			MSU		73Be23	*
^{39}Sc^i(2p)^{37}K	4969	120	5170	40	1.7	U			Lis		90De43	*
	4877	40			7.3	B			Brk		92Mo15	*
	5146	40			0.6	o			Bor		01Gi01	*
	5170	40				3			Bor		07Do17	*
^{38}Ar(p,γ)^{39}K	6380.9	1.1	6381.34	0.19	0.4	U					70Ma31	Z
	6382.2	0.8			−1.1	U					84Ha27	Z
^{39}K(p,d)^{38}K	−10851	2	−10853.19	0.20	−1.1	U			MSU		74Wi17	
^{39}K(^{3}He,α)^{38}K	7498	15	7499.87	0.20	0.1	U			Roc		66Bl04	
	7483	10			1.7	U			Roc		72Fe06	
^{39}Cl(β^-)^{39}Ar	3440	20	3442	5	0.1	U					56Pe38	
^{39}Ar(β^-)^{39}K	565	5				2					50Br66	
^{39}Ca(β^+)^{39}K	6512	25	6524.5	0.6	0.5	U					58Ki40	
^{39}K(p,n)^{39}Ca	−7302.5	6.	−7306.8	0.6	−0.7	U			Tal		70Ke08	
	−7314.9	1.8			4.5	B					78Ra15	Z
*^{39}Al−u	Trends in Mass Surface TMS suggest ^{39}Al 1320 keV less bound										HWJ266	**
*^{39}K(p,^{3}He)^{37}Ar^i	$M-A$=–25954(6); rebuilt Q =–15493.8(6.) with Ame1971; recalibration +0.35										MMC123	**
*^{39}K(p,t)^{37}K^i	$M-A$=–19753(3); Q rebuilt with Ame1971										MMC123	**
*^{39}Sc^i(2p)^{37}K	E_{2p}=3600(120) to 1/2$^+$ level at 1370.85 keV										Ens123	**
*^{39}Sc^i(2p)^{37}K	Other possibility ^{39}Sc^i(α)^{35}K=3600(120) keV										90De43	**
*^{39}Sc^i(2p)^{37}K	E_{2p}=4750(40) p+p at 90 degrees; deduced Q =E_{2p}[1 + Mp/M(^{37}K)]										MMC123	**
*^{39}Sc^i(2p)^{37}K	E_{2p}=4880(40) + recoil 266 keV; data re-analysed and included in 07Do17										MMC135	**
*^{39}Sc^i(2p)^{37}K	IAS identification not sure										MMC135	**
^{40}Si−u	5290	1010	6080	130	0.5	o			GA4	1.5	00Sa21	
	6180	740			−0.1	o			GA5	1.5	00Sa21	
	5829	247			0.7	2			GA7	1.5	07Ju03	
	6120	140			−0.3	2			RT1	1.0	18Mi08	
^{40}P−u	−8800	200	−8740	90	0.2	o			GA4	1.5	00Sa21	
	−8950	210			0.7	2			TO4	1.5	91Zh24	
	−8200	320			−1.1	o			GA5	1.5	00Sa21	
	−8621	129			−0.6	2			GA7	1.5	07Ju03	
	−8749	107			0.1	2			RT1	1.0	18Mi08	
^{40}S−u	−24440	190	−24517	4	−0.3	o			GA4	1.5	00Sa21	
	−24530	250			0.0	U			TO4	1.5	91Zh24	
	−24910	340			0.8	o			GA5	1.5	00Sa21	
	−24627	129			0.6	U			GA7	1.5	07Ju03	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$C_3\ H_4-{}^{40}Ar$		68917.0053	0.0035	68917.0101	0.0020	1.4	1	34	34 ^{40}Ar	MI1	1.0	95Di08	
$C_2\ D_8-{}^{40}Ar$		150431.1045	0.0040	150431.1053	0.0020	0.2	1	26	26 ^{40}Ar	MI1	1.0	95Di08	
$^{20}Ne_2-{}^{40}Ar$		22497.2245	0.0042	22497.2201	0.0020	−1.1	–			MI1	1.0	95Di08	
		22497.2280	0.0060			−1.3	–			MI1	1.0	95Di08	
	ave.	22497.2260	0.0030			−1.6	1	35	35 ^{40}Ar			average	
$^{40}Ar-u$		−37616.878	0.040	−37616.8825	0.0020	−0.1	U			ST2	1.0	02Bf02	
$^{40}Ca-H_{40}$		−350410.425	0.022	−350410.417	0.013	0.4	1	34	34 ^{40}Ca	ST2	1.0	06Na18	
$^{40}Ti-u$		−9689	81	−9650	70	0.4	1	82	82 ^{40}Ti	LZ1	1.0	20Fu05	
$^{40}S\ O-{}^{41}K_{1.366}$		22541	16	22544	4	0.2	U			MS1	1.0	09Ri12	
$^{40}S-{}^{41}K_{.976}$		12752.0	9.4	12741	4	−1.2	1	21	21 ^{40}S	MS1	1.0	09Ri12	
$^{40}S-{}^{40}Ar$		13096.6	4.8	13099	4	0.6	1	79	79 ^{40}S	MS1	1.0	09Ri12	
$^{40}Ca-{}^{40}Ar$		208.2	0.5	207.742	0.013	−0.4	U			J3	2.5	68Fu11	
$^{40}Ca(^3He,^8Li)^{35}K$		−29693	20	−29688.1	0.5	0.2	U			MSU		76Be08	
$^{40}Ca(\alpha,^8He)^{36}Ca$		−57580	40	−57548	6	0.8	U			Tex		77Tr03	
$^{40}Ar(n,\alpha)^{37}S$		−2500	50	−2497.08	0.20	0.1	U					55Be78	
		−2490	50			−0.1	U			Ric		64Da11	
$^{40}K(n,\alpha)^{37}Cl$		3866	7	3872.46	0.08	0.9	U			BNL		68Sc01	
$^{40}Ca(p,\alpha)^{37}K$		−5179	9	−5182.14	0.09	−0.3	U			CIT		66Mc13	
$^{40}Ca(^3He,^6He)^{37}Ca$		−24270	50	−24371.2	0.6	−2.0	U			Brk		68Bu02	
		−24368	25			−0.1	U			MSU		73Be23	*
$^{40}Ar(p,^3He)^{38}Cl^i$		−21092	24				2			Brk		70Ha10	*
$^{40}Ar(p,t)^{38}Ar^j$		−26765	31				2			Brk		70Ha10	*
$^{40}Ca(d,\alpha)^{38}K$		4655	10	4665.18	0.20	1.0	U			MIT		64Sp12	
$^{40}Ca(p,t)^{38}Ca$		−20459	25	−20448.73	0.19	0.4	U					66Ha32	
		−20428	11			−1.9	U			MSU		72Pa02	
		−20452	5			0.7	U			MSU		74Se05	
$^{40}Ar(^{18}O,^{19}Ne)^{39}S$		−14504	200	−14410	50	0.5	U			Can		84Ho.B	
$^{40}Ar(^{13}C,^{14}O)^{39}S$		−16760	50				2			Can		89Dr03	
$^{40}Ar(t,\alpha)^{39}Cl$		7256	40	7285.2	1.7	0.7	U			LAl		61Ja07	
$^{40}Ar(d,^3He)^{39}Cl-{}^{36}Ar()^{35}Cl$		−4024.13	2.42	−4021.7	1.7	1.0	R			Hei		93Ma50	
$^{40}Ar(^3He,\alpha)^{39}Ar^i$		1604	19	1627	7	1.2	2					67Gr01	*
		1631.3	8.0			−0.5	2					72Wi07	*
$^{39}K(n,\gamma)^{40}K$		7799.50	0.08	7799.62	0.06	1.5	–			ILn		84Vo01	Z
		7799.56	0.16			0.4	–			Bdn		06Fi.A	
$^{39}K(d,p)^{40}K$		5579	10	5575.05	0.06	−0.4	U			MIT		64Sp12	
$^{39}K(n,\gamma)^{40}K$	ave.	7799.51	0.07	7799.62	0.06	1.5	1	61	61 ^{40}K			average	
$^{39}K(p,\gamma)^{40}Ca$		8329.5	0.9	8328.165	0.013	−1.5	U					68Do12	
		8329.6	0.9			−1.6	U					68Li12	*
		8328.24	0.09			−0.8	U			Utr		90Ki07	Z
$^{39}K(^3He,d)^{40}Ca$		2845	8	2834.690	0.013	−1.3	U			Oak		67Se10	
$^{40}Ca(^3He,\alpha)^{39}Ca$		4950	20	4942.6	0.6	−0.4	U			Ald		66Hi06	
		4919	15			1.6	U			MIT		71Ra35	
$^{40}Ca(^7Li,^8He)^{39}Sc$		−37400	40	−37376	24	0.6	2			MSU		88Mo18	
$^{40}Ca(^{14}N,^{15}C)^{39}Sc$		−27670	30	−27683	24	−0.4	2			Can		88Wo07	
$^{40}Sc^i(p)^{39}Ca$		3840	120	3830	6	−0.1	o			Lis		90De43	
		3820	30			0.3	U					90Zh.A	*
		3827.7	10.			0.2	2			GSI		97Li25	*
		3830.8	7.			−0.1	2			Lis		98Bh12	
		3841	20			−0.6	U			Bor		07Do17	
$^{40}Cl(\beta^-)^{40}Ar$		7320	80	7480	30	2.0	2					89Mi03	
$^{40}Ar(^7Li,^7Be)^{40}Cl$		−8375	35	−8340	30	0.9	2					84Fi02	
$^{40}K(\varepsilon)^{40}Ar$		1504	7	1504.40	0.06	0.1	U					67Mc10	*
		1497	8			0.9	U					68Az01	*
$^{40}K(n,p)^{40}Ar$		2270	5	2286.75	0.06	3.4	B			BNL		68Sc01	
		2286.7	1.0			0.1	U			ILL		81We12	
$^{40}Ar(p,n)^{40}K$		−2286.3	1.0	−2286.75	0.06	−0.5	U			Duk		66Pa18	Z
		−2286.3	1.0			−0.5	U					01Wa50	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{40}K(β^-)^{40}Ca		1325	15	1310.89	0.06	−0.9	U					52Fe16	
		1350	20			−2.0	U					59Ke26	
^{40}Sc(β^+)^{40}Ca		14330	40	14323.0	2.8	−0.2	U					68Ar03	*
^{40}Ca(p,n)^{40}Sc		−15105.4	2.9				2			Yal		69Ov01	Z
^{40}Ca(^{3}He,t)^{40}Sc		−14490	60	−14341.6	2.8	2.5	U			Bld		65Ri06	
^{40}Ca(π^+,π^-)^{40}Ti		−24974	160	−24830	70	0.9	1	18	18 ^{40}Ti			82Mo12	*
*^{40}Ca(^{3}He,^{6}He)^{37}Ca	Average of 2 values with small calibration correction											AHW	**
*^{40}Ar(p,^{3}He)^{38}Cl^i	IT=8216(25); rebuilt Q =–21093.65(23.68); recalibrated for ^{10}B +1.5 keV											MMC123	**
*^{40}Ar(p,t)^{38}Ar^j	IT=18784(30); Q rebuilt with ^{10}C=15702.5(1.8) from reference											68Br23	**
*^{40}Ar(^{3}He,α)^{39}Ar^i	IT=9089(20); Q rebuilt with Ame1961											MMC123	**
*^{40}Ar(^{3}He,α)^{39}Ar^i	IT=9075(10); Q rebuilt with Ame1964											MMC123	**
*^{39}K(p,γ)^{40}Ca	E_{res}=1345.4(0.5) to 2$^-$ level at 9641.1(0.8) keV											Ens048	**
*^{40}Sc^i(p)^{39}Ca	Uncertainty not given, estimated from graph: stat(9keV), calib(11)											GAu	**
*^{40}Sc^i(p)^{39}Ca	E_p=3731(10); also E_p=1330(20) Q_p=1364.4(20) keV to 2468.5 level											Ens062	**
*^{40}Sc^i(p)^{39}Ca	IT=4370(10) in original paper											MMC123	**
*^{40}K(ε)^{40}Ar	LMK=0.34(0.08) 0.47(0.16) respectively, to 2$^+$ level at 1460.851, recalculated Q											Ens048	**
*^{40}Sc(β^+)^{40}Ca	E_{β^+}=9580(40) to 3$^-$ level at 3736.69 keV, and other E_{β^+}											Ens048	**
*^{40}Ca(π^+,π^-)^{40}Ti	Recalibrated to ^{16}O(π^+,π^-) Q =–27704(20) keV											GAu	**
^{41}Si−u		14560	1980	14170#	320#	−0.1	o			GA5	1.5	00Sa21	
		13011	397			1.9	D			GA7	1.5	07Ju03	*
^{41}P−u		−5930	300	−5350	130	1.3	o			GA4	1.5	00Sa21	
		−5200	500			−0.2	U			TO4	1.5	91Zh24	
		−5290	420			−0.1	o			GA5	1.5	00Sa21	
		−5346	86				2			GA7	1.5	07Ju03	
^{41}S−u		−20500	150	−20407	4	0.4	U			GA4	1.5	00Sa21	
		−19970	230			−1.3	U			TO4	1.5	91Zh24	
		−20430	330			0.0	U			GA5	1.5	00Sa21	
		−20494	75			0.8	U			GT1	1.5	04Ma.A	
^{41}S−C$_2$ H O		−23146.2	4.4				2			MS1	1.0	09Ri12	*
^{41}Cl−u		−29620	190	−29320	70	1.1	2			TO3	1.5	90Tu01	
		−29500	270			0.5	2			TO4	1.5	91Zh24	
^{41}Sc−u		−30741	12	−30748.97	0.08	−0.7	U			LZ1	1.0	11Tu09	
^{41}Ti−u		−16200	390	−16872	17	−1.1	U			GT1	1.5	04St05	
		−16852	30			−0.7	2			LZ1	1.0	12Zh34	*
		−16880.7	20.1			0.4	2			LZ2	1.0	23Zh10	
^{41}K−^{39}K$_{1.051}$		−30.05	0.32	−30.259	0.006	−0.7	U			MA8	1.0	02He23	
		−29.5	2.4			−0.3	U			MA8	1.0	09Na.A	
^{41}K−^{40}Ar H		−8382.9005	0.0061	−8382.8980	0.0030	0.4	–			FS1	1.0	10Mo30	
^{41}K−^{40}Ar		−557.8652	0.0039	−557.8660	0.0030	−0.2	–			FS1	1.0	10Mo30	
^{41}K−^{40}Ar H	ave.	−8382.8980	0.0030	−8382.8980	0.0030	0.1	1	100	100 ^{41}K			average	
^{41}K(p,α)^{38}Ar		4002	20	4019.33	0.19	0.9	U			ChR		60Cl02	
		4018	10			0.1	U			MIT		64Sp12	
^{41}K(d,α)^{39}Ar		8397	15	8393	5	−0.2	U			MIT		67Sp09	
^{39}K(^{3}He,p)^{41}Ca		8920	20	8972.97	0.05	2.6	U			MIT		67Sp09	
^{40}Ar(^{18}O,^{17}F)^{41}Cl		−10530	83	−10470	70	0.8	R			Can		84Ho.B	
^{40}Ar(n,γ)^{41}Ar		6098.4	0.7	6098.9	0.3	0.8	2					70Ha56	Z
		6099.1	0.4			−0.4	2			Bdn		06Fi.A	
^{40}Ar(d,p)^{41}Ar		3878	6	3874.4	0.3	−0.6	U			MIT		64Sp12	
^{40}Ar(p,γ)^{41}K		7807.8	0.3	7808.6200	0.0030	2.7	U					89Sm06	Z
^{40}Ar(^{3}He,d)^{41}K^i		−6034	15				2					75Me10	*
^{40}K(n,γ)^{41}K		10095.19	0.10	10095.37	0.06	1.8	–			ILn		84Kr05	Z
		10095.25	0.20			0.6	–			Bdn		06Fi.A	
	ave.	10095.20	0.09			1.9	1	39	39 ^{40}K			average	
^{40}Ca(n,γ)^{41}Ca		8363.0	0.5	8362.85	0.05	−0.3	U					69Ar.A	Z
		8362.5	0.5			0.7	U					70Cr04	Z
		8362.72	0.3			0.4	U			MMn		80Is02	Z
		8362.86	0.17			−0.1	–			Bdn		06Fi.A	
		8362.86	0.06			−0.2	–			ILn		21Bo02	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{40}Ca(d,p)^{41}Ca		6134	4	6138.28	0.05	1.1	U			MIT		68Be36	
^{40}Ca(n,γ)^{41}Ca	ave.	8362.86	0.06	8362.85	0.05	−0.2	1	87	87 ^{41}Ca			average	
^{40}Ca(p,γ)^{41}Sc		1085.7	1.4	1085.06	0.07	−0.5	U					73Al11	
		1085.09	0.09			−0.3	1	68	68 ^{41}Sc	Utr		87Zi02	*
^{40}Ca(d,n)^{41}Sc		−1145	15	−1139.50	0.07	0.4	U					61Ma08	Y
^{40}Ca(p,γ)^{41}Sc^{r}		−1796.4	1.5	−1797.24	0.08	−0.6	U					77Ko10	
^{40}Ca(p,γ)^{41}Sc^{i}		−4851.4	4.9	−4854	3	−0.5	2					76Fo01	
^{41}Sc^{i}(p)^{40}Ca		4855.6	5.	4854	3	−0.4	2			Jyp		97Ho12	
		4855.6	8.			−0.2	2			Lis		98Bh12	
		4857	16			−0.2	U			Bor		07Do17	
^{41}Cl(β^-)^{41}Ar		5670	150	5760	70	0.6	R					74Gu10	
^{41}Ar(β^-)^{41}K		2492.0	1.1	2492.0	0.3	0.0	U					64Pa03	*
^{41}K(p,n)^{41}Ca		−1209.6	1.5	−1203.97	0.05	3.8	B			Oak		64Jo11	Z
		−1203.8	0.5			−0.3	U			Can		70Kn03	Z
^{41}Sc(β^+)^{41}Ca		6630	100	6495.44	0.10	−1.3	U					62Cr04	
^{41}Sc^{r}(IT)^{41}Sc		2882.6	0.3	2882.30	0.05	−1.0	U					77Ko10	
		2882.39	0.10			−0.9	–			Utr		87Zi02	Z
		2882.26	0.06			0.7	–			Utr		89Ki11	Z
	ave.	2882.29	0.05			0.2	1	90	58 ^{41}Sc^{r}			average	
*^{41}Si−u	Trends in Mass Surface TMS suggest ^{41}Si 1080 keV less bound											HWJ266	**
*^{41}S−C_2 H O	For original doublet ^{41}S C H−C_3 H_2 O											09Ri12	**
*^{41}Ti−u	Same result in reference											13Ya03	**
*^{40}Ar(^{3}He,d)^{41}K^{i}	IT=8349(15); Q rebuilt with Ame1971											MMC123	**
*^{40}Ca(p,γ)^{41}Sc	E_p=647.25(0.05,Z) to 1716.43(0.08,Z) level											87Zi02	**
*^{41}Ar(β^-)^{41}K	E_{β^-}=1198.3(1.1) to 7/2$^-$ level at 1293.609 keV											Ens163	**
^{42}Si−u		20860	3990	18790#	320#	−0.3	o			GA5	1.5	99Sa.A	
		16275	623			2.7	D			GA7	1.5	07Ju03	*
^{42}P−u		260	740	1170	100	0.8	o			GA4	1.5	00Sa21	
		1550	630			−0.4	o			GA5	1.5	00Sa21	
		1084	225			0.3	2			GA7	1.5	07Ju03	
		1181	107			−0.1	2			RT1	1.0	18Mi08	
^{42}S−u		−18940	150	−18935	3	0.0	U			GA4	1.5	00Sa21	
		−18510	350			−0.8	U			TO4	1.5	91Zh24	
		−19390	350			0.9	U			GA5	1.5	00Sa21	
		−18934.9	3.0				2			MS1	1.0	09Ri12	*
^{42}Cl−u		−27000	190	−26660	60	1.2	U			TO3	1.5	90Tu01	
		−26870	190			0.7	U			TO4	1.5	91Zh24	
		−26658	64				2			LZ1	1.0	15Xu14	
^{42}Ti−u		−26973	25	−26951.02	0.24	0.9	U			LZ1	1.0	17Zh12	
^{42}Ar−^{36}Ar$_{1.167}$		920.6	6.2				2			MA6	1.0	01He29	
^{42}Ca−^{40}Ca$_{1.050}$		−2103.04358	0.00068	−2103.0436	0.0007	−0.0	1	100	100 ^{42}Ca	Hep	1.0	25Wi05	
^{42}S−^{41}K$_{1.024}$		20151.8	9.5	20156	3	0.4	U			MS1	1.0	09Ri12	
^{42}Sc−^{42}Ca		6898.74	0.22	6898.95	0.05	1.0	o			JY1	1.0	06Er08	
		6898.960	0.064			−0.1	1	66	66 ^{42}Sc	JY1	1.0	17Er01	*
^{42}Sc^{m}−^{42}Ca		7560.68	0.23	7561.10	0.06	1.8	o			JY1	1.0	06Er08	
		7561.120	0.081			−0.2	1	62	62 ^{42}Sc^{m}	JY1	1.0	17Er01	*
^{42}Ti−^{42}Ca		14431.69	0.71	14431.62	0.24	−0.1	1	12	12 ^{42}Ti	JY1	1.0	09Ku19	
^{42}Sc^{m}−^{42}Sc		661.97	0.24	662.15	0.07	0.7	U			JY1	1.0	06Er08	
		662.50	0.42			−0.8	U			JY1	1.0	09Ku19	
		662.109	0.088			0.4	1	55	37 ^{42}Sc^{m}	JY1	1.0	17Er01	*
^{42}Ti−^{42}Sc		7532.92	0.34	7532.67	0.24	−0.7	1	50	49 ^{42}Ti	JY1	1.0	09Ku19	
^{42}Ti−^{42}Sc^{m}		6870.19	0.38	6870.52	0.24	0.9	1	40	39 ^{42}Ti	JY1	1.0	09Ku19	
^{28}Si(^{16}O,2n)^{42}Ti		−17250	13	−17267.72	0.23	−1.4	U					72Zi02	
^{42}Ca(p,α)^{39}K		118	7	123.570	0.013	0.8	U			MIT		64Sp12	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{39}K(α,n)^{42}Sc	−7160	60	−7332.25	0.05	−2.9	U			Yal		61Sm05	
	−7455	30			4.1	B			Tal		65Ne02	
^{40}Ar(t,p)^{42}Ar	7043	40	7044	6	0.0	U			LAl		61Ja07	
^{40}Ca(^{3}He,p)^{42}Sc	4966	20	4917.20	0.05	−2.4	U			MIT		64Sp12	
	4905	5			2.4	U			ANL		74Ha55	
^{40}Ca(^{3}He,n)^{42}Ti	−2865	6	−2881.78	0.23	−2.8	U			CIT		67Mi02	
^{41}K(n,γ)^{42}K	7533.78	0.15	7533.80	0.11	0.1	2			ILn		85Kr06	Z
	7533.82	0.15			−0.1	2			Bdn		06Fi.A	
^{41}K(d,p)^{42}K	5314	12	5309.23	0.11	−0.4	U			MIT		64Sp12	
^{41}K(p,γ)^{42}Ca	10275.5	3.4	10277.102	0.013	0.5	U					71Vi14	
^{41}Ca(n,γ)^{42}Ca	11480.63	0.06	11481.08	0.05	7.4	B			ORn		89Ki11	Z
^{42}Ca(^{3}He,α)^{41}Ca	9102	15	9096.54	0.05	−0.4	U			MIT		71Ra35	
^{41}Ca(p,γ)^{42}Sc^{r}−^{40}Ca()^{41}Sc^{r}	−6.67	0.05	−6.68	0.05	−0.2	1	90	42 ^{41}Sc^{r}	Utr		89Ki11	*
^{42}Cl(β^-)^{42}Ar	9760	220	9590	60	−0.8	U					89Mi03	
^{42}K(β^-)^{42}Ca	3519.	3.5	3525.65	0.11	1.9	U					68Va06	
	3524	6			0.3	U					75Ra09	
^{42}Sc(β^+)^{42}Ca	6342	100	6426.33	0.05	0.8	U					61Ja22	
	6486	100			−0.6	U					63Ro10	*
^{42}Ca(p,n)^{42}Sc	−7213.7	2.3	−7208.68	0.05	2.2	U			Har		75Fr.A	
^{42}Ca(^{3}He,t)^{42}Sc	−6442.3	0.4	−6444.93	0.05	−6.6	F			Mun		77Vo02	*
^{42}Ca(^{3}He,t)^{42}Sc−^{27}Al()^{27}Si	−1611.7	2.6	−1613.98	0.11	−0.9	U			ChR		74Ha35	
^{42}Ca(^{3}He,t)^{42}Sc−^{26}Mg()^{26}Al	−2417.8	3.5	−2421.93	0.08	−1.2	U			ChR		74Ha35	
	−2421.83	0.23			−0.4	1	11	7 ^{26}Al	ChR		87Ko34	*
^{42}Sc^{m}(IT)^{42}Sc	616.28	0.06	616.79	0.06	8.5	B					Ens167	
^{42}Sc^{r}(IT)^{42}Sc	6076.33	0.08	6076.31	0.07	−0.2	1	75	65 ^{42}Sc^{r}	Utr		89Ki11	Z
*^{42}Si−u	Trends in Mass Surface TMS suggest ^{42}Si 2340 keV less bound										GAu	**
*^{42}S−u	For original doublet ^{42}S−(C_2 H O)$_{1.024}$, D_M=–21740.2(3.1) μu										09Ri12	**
*^{42}Sc−^{42}Ca	Frequency ratio not correct for unc., values taken from paper										HWJ17b	**
*^{42}Sc^{m}−^{42}Ca	Frequency ratio not correct for unc., values taken from paper										HWJ17b	**
*^{42}Sc^{m}−^{42}Sc	Frequency ratio not correct for unc., values taken from paper										HWJ17b	**
*^{41}Ca(p,γ)^{42}Sc^{r}−^{40}Ca()^{41}Sc^{r}	Calculated from resonance energy difference = 5.73(0.05) keV										GAu	**
*^{42}Sc(β^+)^{42}Ca	E_{β^+}=2870(100) from ^{42}Sc^{m} at 616.28 to 6$^+$ level at 3189.26 keV										Ens167	**
*^{42}Ca(^{3}He,t)^{42}Sc	F : rejected in reference of same group										09Fa15	**
*^{42}Ca(^{3}He,t)^{42}Sc−^{26}Mg()^{26}Al	Q =–2193.52(0.23) to ^{26}Al^{m} at 228.306 keV										Nub269	**
^{43}P−u	4220	1620	5900#	320#	0.7	U			GA4	1.5	00Sa21	
	6190	1040			−0.2	o			GA5	1.5	00Sa21	
	5024	397			1.5	D			GA7	1.5	07Ju03	*
^{43}S−u	−12810	250	−13092	5	−0.8	o			GA4	1.5	00Sa21	
	−13400	900			0.2	U			TO4	1.5	91Zh24	
	−12900	460			−0.3	o			GA5	1.5	00Sa21	
	−12958	107			−0.8	U			GA7	1.5	07Ju03	
	−13087	22			−0.2	2			MS1	1.0	09Ri12	*
	−13092.7	5.5			0.1	2			MS1	1.0	09Ri12	*
	−13022	236			−0.2	U			GA8	1.5	12Ga45	
^{43}Cl−u	−26090	300	−25888.0	1.9	0.4	o			GA4	1.5	00Sa21	
	−25740	200			−0.5	o			TO3	1.5	90Tu01	
	−25970	350			0.2	U			TO4	1.5	91Zh24	
	−26010	330			0.2	o			GA5	1.5	00Sa21	
	−25905	86			0.1	o			GT1	1.5	04Ma.A	
	−25894	140			0.0	U			GA7	1.5	07Ju03	
	−26361	100			1.9	C			GT2	2.5	08Kn.A	
	−25941	70			0.8	U			LZ1	1.0	15Xu14	
^{43}Cl−^{39}K$_{1.103}$	14143.8	1.9				2			MS1	1.0	25Er02	
^{43}Ti−u	−31461	10	−31461	4	−0.0	–			LZ1	1.0	18Zh29	*
	−31457.2	4.5			−0.9	–			LZ2	1.0	22Wa39	*
	−31457.2	4.5			−0.9	o			LZ2	1.0	23Zh10	
ave.	−31458	4			−0.8	1	85	85 ^{43}Ti			average	
^{43}V−u	−19234	46	−19222	27	0.3	2			LZ1	1.0	13Ya03	
	−19215.7	33.9			−0.2	2			LZ2	1.0	23Zh10	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{43}Ar−^{36}Ar$_{1.194}$		4387.2	5.7				2			MA6	1.0	01He29	
^{43}K−^{39}K$_{1.103}$		766.45	0.44				2			MA8	1.0	07Ya08	
^{43}Ca(p,α)^{40}K		−14	8	−9.72	0.17	0.5	U			MIT		64Sp12	
^{40}Ca(α,p)^{43}Sc		−3470	30	−3521.9	1.9	−1.7	U					61Ma03	
^{40}Ca(α,n)^{43}Ti		−11169.9	10.	−11187	4	−1.9	1	15	15 ^{43}Ti	Tal		67Al08	
^{41}K(^{3}He,p)^{43}Ca^i		2452	30	2497	14	1.5	1	23	23 ^{43}Ca^i	MIT		68Do02	
^{43}V^i(2p)^{41}Sc		4320	50	4359	15	0.8	U			Lis		92Bo37	
		4503	22			−6.5	B			Bor		01Gi01	*
		4348	16			0.7	1	89	89 ^{43}V^i	Bor		07Do17	
^{42}Ca(n,γ)^{43}Ca		7933.1	0.5	7932.91	0.16	−0.4	–					69Ar.A	Z
		7933.1	0.5			−0.4	–			Ptn		69Gr08	Z
		7933.1	0.4			−0.5	–					71Bi.A	
		7932.73	0.23			0.8	–			Bdn		06Fi.A	
^{42}Ca(d,p)^{43}Ca		5716	10	5708.34	0.16	−0.8	U			MIT		64Sp12	
		5707	12			0.1	U			MIT		66Do02	
^{43}Ca(d,t)^{42}Ca		−1672	10	−1675.68	0.16	−0.4	U			Ald		64Bj02	
^{42}Ca(n,γ)^{43}Ca	ave.	7932.89	0.17	7932.91	0.16	0.1	1	84	84 ^{43}Ca			average	
^{42}Ca(p,γ)^{43}Sc		4935	5	4929.8	1.9	−1.0	2					65Br31	
		4929	2			0.4	2					69Wa19	
^{42}Ca(^{3}He,d)^{43}Sc^i		−4808	8	−4795	3	1.6	1	17	17 ^{43}Sc^i			66Sc17	*
^{43}V^i(p)^{42}Ti		8200	45	8111	15	−2.0	1	11	11 ^{43}V^i	Bor		01Gi01	*
^{43}K(β^-)^{43}Ca		1817	20	1833.9	0.4	0.8	U					54Li24	*
		1815	10			1.9	U					59Be72	*
^{43}Sc(β^+)^{43}Ca		2200	20	2220.7	1.9	1.0	U					52Ha44	
		2220	10			0.1	U					54Li42	
^{43}Ca(p,n)^{43}Sc		−3005	10	−3003.1	1.9	0.2	U			Har		60Mc12	Y
		−2998	10			−0.5	U					67Mc07	
^{43}Ca(^{3}He,t)^{43}Sc^i		−6467	8	−6471	3	−0.5	–					71Al19	*
		−6469	4			−0.5	–					71Be29	*
	ave.	−6469	4			−0.7	1	83	83 ^{43}Sc^i			average	
*^{43}P−u	Trends in Mass Surface TMS suggest ^{43}P 820 keV less bound											HWJ266	**
*^{43}S−u	For original doublet ^{43}S−(C$_3$ H$_5$ O)$_{0.754}$, D_M=–38753(22) μu											09Ri12	**
*^{43}S−u	For original doublet ^{43}S C H−(C$_3$ H$_5$ O)$_{0.982}$, D_M=–38694.8(5.5) μu											09Ri12	**
*^{43}Ti−u	could be mixed with 313(1) state, without correction											18Zh29	**
*^{43}Ti−u	could be mixed with 313(1) state, without correction											22Wa39	**
*^{43}V^i(2p)^{41}Sc	E_{2p}=4292(22) + recoil=211; data re-analysed and included in 07Do17											MMC135	**
*^{42}Ca(^{3}He,d)^{43}Sc^i	IT=4238(8); Q rebuilt with Ame1961											MMC123	**
*^{43}V^i(p)^{42}Ti	Q_p=4590(45) followed by γ's 1938+1554 keV + recoil=118 keV											MMC135	**
*^{43}K(β^-)^{43}Ca	E_{β^-}=827(20) 825(10) respectively, to 3/2$^+$ level at 990.257 keV											Ens156	**
*^{43}Ca(^{3}He,t)^{43}Sc^i	IT=4226(8); Q rebuilt with Ame1965											MMC123	**
*^{43}Ca(^{3}He,t)^{43}Sc^i	CDE=7238(4) Q =–6474(4); recalibration +6 keV for ^{42}Ca(p,n)^{42}Sc from Ame1961											MMC123	**
^{44}P−u		10070	966	12400#	430#	1.6	D			GA7	1.5	07Ju03	*
^{44}S−u		−10510	580	−9881	6	0.7	o			GA4	1.5	00Sa21	
		−8960	620			−1.0	o			GA5	1.5	00Sa21	
		−9769	150			−0.5	U			GA7	1.5	07Ju03	
		−10027	301			0.3	U			GA8	1.5	12Ga45	
^{44}S−C$_2$ H$_4$ O		−36095.9	5.6				2			MS1	1.0	09Ri12	*
^{44}Cl−u		−21700	130	−21954	11	−1.3	U			GA4	1.5	00Sa21	
		−21500	500			−0.6	U			TO3	1.5	90Tu01	
		−21450	270			−1.2	U			TO4	1.5	91Zh24	
		−22150	370			0.4	U			GA5	1.5	00Sa21	
		−22115	161			0.7	U			GT1	1.5	04Ma.A	
		−22051	118			0.8	U			RT1	1.0	18Mi08	
^{44}Cl−^{12}C ^{14}N ^{1}H$_2$ ^{16}O		−35596	17	−35593	11	0.2	1	44	44 ^{44}Cl	MS1	1.0	25Er02	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{44}Ca−u		−44515.3	10.0	−44518.245	0.014	−0.3	U			MR1	1.0	15Wi.A	
^{44}Sc−u		−40480	410	−40597.6	1.9	−0.2	U			TO6	1.5	98Ba.A	*
^{44}V−u		−25890	120	−25555	5	1.9	B			GT1	1.5	04St05	*
		−25579	21			1.2	–			LZ1	1.0	18Zh29	
		−25550.8	7.6			−0.5	–			LZ2	1.0	22Wa39	
		−25550.8	7.6			−0.5	o			LZ2	1.0	23Zh10	
	ave.	−25554	7			−0.1	1	58	58 ^{44}V			average	
^{44}V^m−u		−25272	20	−25267	5	0.3	U			LZ1	1.0	18Zh29	
		−25265.1	7.8			−0.2	1	36	36 ^{44}V^m	LZ2	1.0	22Wa39	
		−25265.1	7.8			−0.2	o			LZ2	1.0	23Zh10	
^{44}Cr−u		−14409	55				2			LZ1	1.0	20Fu05	
^{44}Cl−^{39}K$_{1.128}$		18987	15	18985	11	−0.2	1	56	56 ^{44}Cl	MS1	1.0	25Er02	
^{44}Ar−^{39}K$_{1.128}$		5862.9	1.7				2			MA8	1.0	03Bl17	
^{44}K−^{39}K$_{1.128}$		2526.07	0.45				2			MA8	1.0	07Ya08	
		2529.2	2.2	2526.1	0.5	−1.4	o			TT1	1.0	10La.A	
		2529.1	1.7			−1.8	U			TT1	1.0	12La05	
^{44}Ca−^{40}Ca$_{1.100}$		−3368.18938	0.00069	−3368.1894	0.0007	0.0	1	100	100 ^{44}Ca	Hep	1.0	25Wi05	
^{44}V−^{32}S		2373.0	8.4	2374	5	0.1	1	42	42 ^{44}V	MS1	1.0	20Pu02	*
^{44}V^m−^{32}S		2660.8	5.9	2662	5	0.2	1	64	64 ^{44}V^m	MS1	1.0	20Pu02	*
^{40}Ca(α,γ)^{44}Ti		5127.1	0.7				2					82Di05	
^{44}Ca(p,α)^{41}K		−1058	10	−1044.875	0.014	1.3	U			MIT		64Sp12	
^{41}K(α,n)^{44}Sc		−3420	60	−3389.5	1.7	0.5	U			Yal		61Sm05	
^{44}Ca(d,α)^{42}K		4273	20	4264.36	0.11	−0.4	U					77Pa24	
^{42}Ca(t,p)^{44}Ca		10593	15	10581.6379	0.0011	−0.8	U			Ald		67Bj06	
^{42}Ca(^{3}He,p)^{44}Sc		6920	20	6911.0	1.7	−0.4	U			Hei		70Sc22	
^{43}Ca(n,γ)^{44}Ca		11130.6	0.5	11130.52	0.16	−0.2	–					69Ar.A	Z
		11130.1	0.7			0.6	–					72Wh02	Z
		11131.54	0.29			−3.5	B			Bdn		06Fi.A	
^{43}Ca(d,p)^{44}Ca		8922	14	8905.96	0.16	−1.1	U			MIT		64Sp12	
		8920	10			−1.4	U			Kop		67Bj02	
^{44}Ca(^{3}He,α)^{43}Ca		9452	15	9447.10	0.16	−0.3	U			MIT		71Ra35	
^{43}Ca(n,γ)^{44}Ca	ave.	11130.4	0.4	11130.52	0.16	0.2	1	15	15 ^{43}Ca			average	
^{44}Ca(p,d)^{43}Ca^i		−16880	30	−16901	14	−0.7	–					72Ma23	*
^{44}Ca(d,t)^{43}Ca^i		−12858.7	19.7	−12869	14	−0.5	–					76Do05	*
^{44}Ca(p,d)^{43}Ca^i	ave.	−16888	16	−16901	14	−0.8	1	77	77 ^{43}Ca^i			average	
^{43}Ca(p,γ)^{44}Sc		6694	2	6696.1	1.7	1.1	2					71Po.A	
^{43}Ca(^{3}He,d)^{44}Sc^i		−1583	5	−1575.6	2.5	1.5	1	24	24 ^{44}Sc^i			68Sc15	
^{44}V^i(p)^{43}Ti		950	50	908	11	−0.8	U			Lis		92Bo37	
		908	11				2			Bor		07Do17	
		917	53			−0.2	U					14Po05	
^{44}K(β^-)^{44}Ca		5580	80	5687.0	0.4	1.3	U					70Le05	
^{44}Ca(t,^{3}He)^{44}K		−5660	40	−5668.4	0.4	−0.2	U			LAl		70Aj01	
^{44}Sc(β^+)^{44}Ca		3642	5	3652.0	1.7	2.0	R					50Br52	*
		3650	5			0.4	R					55Bl23	*
^{44}Ca(p,n)^{44}Sc		−4410	15	−4434.4	1.7	−1.6	U			Har		60Mc12	Y
		−4447	10			1.3	U					67Mc07	
^{44}Ca(^{3}He,t)^{44}Sc^i		−6444	4	−6448.9	2.5	−1.2	–					71Be29	*
		−6449	4			0.0	–					72Ma50	*
	ave.	−6446.5	2.8			−0.8	1	76	76 ^{44}Sc^i			average	
*^{44}P−u	Trends in Mass Surface TMS suggest ^{44}P 2170 keV less bound											HWJ266	**
*^{44}S−C$_2$ H$_4$ O	For original doublet ^{44}S C H−C$_3$ H$_5$ O											09Ri12	**
*^{44}Sc−u	$M-A$=–37570(370) keV for mixture gs+p at 271.241 keV											Nub269	**
*^{44}V−u	$M-A$=–23980(80) keV for mixture gs+m at 268(7) keV											Nub269	**
*^{44}V−u	Authors have unduly increased the lower error to 380 keV											GAu	**
*^{44}V−^{32}S	From freq. ratio ^{44}VO$^+$/(^{32}SCO)$^+$ =0.99996043(14)											HWJ207	**
*^{44}V^m−^{32}S	From freq. ratio ^{44}VmO$^+$/(^{32}SCO)$^+$ =0.999955631(98)											HWJ207	**
*^{44}Ca(p,d)^{43}Ca^i	IT=7970(30); Q rebuilt with Ame1965											MMC123	**
*^{44}Ca(d,t)^{43}Ca^i	IT=7980(20); Q rebuilt with Ame1971											MMC123	**
*^{43}Ca(^{3}He,d)^{44}Sc^i	IT=2796(5); Q rebuilt with Ame1965											MMC123	**
*^{44}Sc(β^+)^{44}Ca	E_{β^+}=1463(5) 1471(5) respectively, to 2$^+$ level at 1157.019 keV											Ens119	**
*^{44}Ca(^{3}He,t)^{44}Sc^i	CDE=7214(4) Q =–6450(4); recalibration +6 keV for ^{42}Ca(p,n)^{42}Sc from Ame1961											MMC123	**
*^{44}Ca(^{3}He,t)^{44}Sc^i	IT=2781(5); Q rebuilt with Ame1971											MMC123	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{45}S−u		−3610	2460	−3590#	320#	0.0	o			GA4	1.5	00Sa21	
		−3330	2880			−0.1	o			GA5	1.5	00Sa21	
		−4283	741			0.6	D			GA7	1.5	07Ju03	*
^{45}Cl−u		−19690	140	−19582	4	0.5	o			GA4	1.5	00Sa21	
		−20300	700			0.7	U			TO3	1.5	90Tu01	
		−19850	460			0.4	o			GA5	1.5	00Sa21	
		−19710	107			0.8	U			GA7	1.5	07Ju03	
		−19098	236			−1.4	U			GA8	1.5	12Ga45	
^{45}Cl−^{28}Si ^{16}O ^{1}H		752.2	4.0	752	4	−0.0	1	100	100 ^{45}Cl	MS1	1.0	25Er02	
^{45}V−u		−34225.7	9.7	−34231.5	0.9	−0.6	U			LZ1	1.0	11Tu09	
		−34230	11			−0.1	U			LZ1	1.0	18Zh29	
^{45}Cr−u		−20390	540	−20911	11	−0.6	U			GT1	1.5	04St05	*
		−20950	38			1.0	2			LZ1	1.0	12Zh34	*
		−20906.8	11.8			−0.3	2			LZ2	1.0	23Zh10	
^{45}Ar−^{39}K$_{1.154}$		9922.45	0.55				2			MA8	1.0	03Bl17	
^{45}K−^{39}K$_{1.154}$		2574.21	0.56				2			MA8	1.0	07Ya08	
^{45}V−^{45}Ti		7647.74	0.23	7647.74	0.23	0.0	1	100	100 ^{45}V	JY1	1.0	14Ka22	
^{45}Sc(p,α)^{42}Ca		2343	8	2339.4	0.7	−0.4	U			MIT		64Sp12	
^{45}Sc(d,α)^{43}Ca		8028	12	8047.8	0.7	1.6	U			MIT		64Sp12	
		8059	12			−0.9	U			Kop		67Ha.A	
^{43}Ca(^{3}He,p)^{45}Sc		10310	20	10305.3	0.7	−0.2	U			Hei		70Sc22	
^{45}Fe(2p)^{43}Cr		1140	40	1143	15	0.1	o					02Gi09	
		1100	100			0.4	U					02Pf02	
		1154	16			−0.7	3					05Do20	
		1077	39			1.7	3					25Or05	
^{44}Ca(n,γ)^{45}Ca		7414.8	1.0	7414.82	0.17	0.0	U					69Ar.A	Z
		7414.83	0.3			−0.0	–			MMn		80Is02	Z
		7414.79	0.21			0.1	–			Bdn		06Fi.A	
^{44}Ca(d,p)^{45}Ca		5184	4	5190.25	0.17	1.6	U			MIT		68Be36	
^{44}Ca(n,γ)^{45}Ca	ave.	7414.80	0.17	7414.82	0.17	0.1	1	99	99 ^{45}Ca			average	
^{44}Ca(p,γ)^{45}Sc		6887.8	1.2	6892.8	0.7	4.2	B					74Sc02	Z
^{45}Sc(^{3}He,α)^{44}Sc		9249	15	9250.4	1.9	0.1	U			MIT		71Ra09	
^{45}Sc(d,t)^{44}Sc^{i}		−7846	10	−7848.2	2.5	−0.2	U					71Oh01	*
^{45}V^{i}(p)^{44}Ti		3190	50	3170	9	−0.4	U					74Ja10	*
		3170	9				3			Bor		07Do17	*
^{45}K(β^-)^{45}Ca		4180	200	4196.3	0.5	0.1	U					64Mo18	
^{45}Ca(β^-)^{45}Sc		258	2	260.3	0.7	1.2	1	12	11 ^{45}Sc			65Fr12	
^{45}Ti(β^+)^{45}Sc		2066	5	2062.1	0.5	−0.8	U					66Po04	
^{45}Sc(p,n)^{45}Ti		−2844.2	4.	−2844.4	0.5	−0.1	U			Ric		55Br16	Y
		−2843.6	4.0			−0.2	U			Can		70Kn03	
		−2844.4	0.5			−0.0	1	100	100 ^{45}Ti	PTB		85Sc16	Z
^{45}Sc(^{3}He,t)^{45}Ti^{i}		−6801	4	−6800	3	0.3	1	61	60 ^{45}Ti^{i}			71Be29	*
*^{45}S−u	Trends in Mass Surface TMS suggest ^{45}S 650 keV less bound											HWJ266	**
*^{45}Cr−u	$M-A$=–18940(500) keV for mixture gs+m at 107(1) keV											Nub269	**
*^{45}Cr−u	Same result in reference											13Ya03	**
*^{45}Sc(d,t)^{44}Sc^{i}	IT=2784(10) combined with Q =–5062; Q rebuilt											MMC123	**
*^{45}V^{i}(p)^{44}Ti	Q_p=2060(50) 2087(9) respectively, to 2$^+$ level at 1083.06 keV											Ens119	**
*^{45}Sc(^{3}He,t)^{45}Ti^{i}	CDE=7571(4) Q =–6807(4); recalibration +6 keV for ^{42}Ca(p,n)^{42}Sc from Ame1961											MMC123	**
^{46}Cl−u		−16000	860	−14750	100	1.0	o			GA4	1.5	00Sa21	
		−14940	1730			0.1	o			GA5	1.5	00Sa21	
		−14826	172			0.3	2			GA7	1.5	07Ju03	
		−15040	301			0.7	2			GA8	1.5	12Ga45	
		−14708	118			−0.3	2			RT1	1.0	18Mi08	
^{46}Ar−u		−32013	107	−31960.8	2.5	0.3	U			GT1	1.5	04Ma.A	
^{46}Sc−u		−44650	230	−44833.0	0.7	−0.5	U			TO6	1.5	98Ba.A	*
C_2 H_8 N−^{46}Ti		113071	7	113047.92	0.09	−0.8	U			R09	4.0	72De11	
C ^{13}C H_5 O−^{46}Ti		84799	13	84768.28	0.09	−0.6	U			R09	4.0	72De11	
C H_4 N O−^{46}Ti		76672	8	76662.41	0.09	−0.3	U			R09	4.0	72De11	
C_5 H_2−^{46}Ti O		68145	15	68109.11	0.09	−0.6	U			R09	4.0	72De11	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
C H_2 O_2 − ^{46}Ti	52881	14	52852.97	0.09	−0.5	U			R09	4.0	72De11	
^{13}C H O_2 − ^{46}Ti	48423	9	48382.77	0.09	−1.1	U			R09	4.0	72De11	
^{46}Ti − ^{22}Ne$_{2.091}$	−29358.74	0.48	−29359.90	0.09	−2.4	U			CP1	1.0	05Sa44	
^{46}V − ^{22}Ne$_{2.091}$	−21787.10	0.58	−21788.87	0.14	−3.1	B			CP1	1.0	05Sa44	
^{46}Cr − u	−31638	15	−31645.1	2.8	−0.5	U			LZ1	1.0	17Zh12	
	−31645.1	2.8				2			LZ2	1.0	22Wa39	
	−31645.1	2.8			0.0	o			LZ2	1.0	23Zh10	
^{46}Mn − u	−13331	93				2			LZ1	1.0	20Fu05	*
^{46}Ar − ^{39}K$_{1.179}$	10827.5	1.2	10829.3	2.5	1.5	o			MA8	1.0	15Mo.A	
	10829.3	2.5				2			MA8	1.0	20Mo25	
^{46}K − ^{39}K$_{1.179}$	4771.64	0.78				2			MA8	1.0	07Ya08	
^{46}Ca − ^{40}Ca$_{1.150}$	−3289.8095	0.0012	−3289.8095	0.0012	−0.0	1	100	100 ^{46}Ca	Hep	1.0	25Wi05	
^{46}Ti − ^{48}Ti$_{.958}$	2499.165	0.062	2499.19	0.06	0.4	1	94	94 ^{46}Ti	MS1	1.0	17Ka53	
^{46}V − ^{46}Ti	7571.67	0.41	7571.03	0.10	−1.6	U			CP1	1.0	05Sa44	
	7571.41	0.33			−1.1	o			JY1	1.0	06Er08	
	7571.10	0.11			−0.6	1	81	87 ^{46}V	JY1	1.0	11Er02	
^{32}S(^{16}O,2n)^{46}Cr	−17421.6	20.	−17418.0	2.6	0.2	U					72Zi02	
^{46}Ti(p,α)^{43}Sc	−3065	14	−3075.7	1.9	−0.8	U			MIT		64Sp12	*
	−3083	10			0.7	U			Tal		65Pl01	
^{46}Ti(^{3}He,^{6}He)^{43}Ti	−17470	12	−17483	4	−1.1	U			MSU		77Mu03	*
^{44}Ca(t,p)^{46}Ca	9339	20	9330.1475	0.0014	−0.4	U			Kop		67Bj06	
^{44}Ca(^{3}He,p)^{46}Sc	7940	20	7935.4	0.7	−0.2	U			Hei		70Sc22	
^{46}Ti(d,α)^{44}Sc	4400	12	4399.0	1.8	−0.1	U			Kop		67Ha.A	
^{46}Ti(p,t)^{44}Ti	−14235	10	−14240.6	0.7	−0.6	U			Oak		72Ra05	
^{46}Ca(t,α)^{45}K	5998	10	6002.8	0.5	0.5	U			Ald		68Sa09	
^{46}Ca(d,t)^{45}Ca	−4144	10	−4139.89	0.17	0.4	U			Ald		67Bj05	
^{46}Ca(^{3}He,α)^{45}Ca	10194	10	10180.50	0.17	−1.4	U			MIT		71Ra35	
^{45}Sc(n,γ)^{46}Sc	8760.61	0.3	8760.64	0.10	0.1	2			BNn		80Li07	Z
	8760.58	0.14			0.4	2			Utr		82Ti02	Z
	8760.75	0.18			−0.6	2			Bdn		06Fi.A	
^{45}Sc(d,p)^{46}Sc	6541	8	6536.07	0.10	−0.6	U			MIT		64Sp12	
	6543	8			−0.9	U			Kop		67Ha.A	
^{45}Sc(p,γ)^{46}Ti	10344.7	0.7	10344.9	0.7	0.4	1	88	88 ^{45}Sc			71Gu.A	
^{46}Ti(p,d)^{45}Ti^{i}	−15682	5	−15684	3	−0.4	1	40	40 ^{45}Ti^{i}			78Ko27	
^{46}Cr^{i}(p)^{45}V	4350	50	4269	13	−1.6	U			Lis		92Bo37	
	4269	13				2			Bor		07Do17	*
^{46}Mn^{i}(p)^{45}Cr	3520	100	4840	30	13.2	B			Lis		92Bo37	
	4840	33				3			Bor		07Do17	*
^{46}K(β^-)^{46}Ca	7650	300	7723.9	0.7	0.2	U					66Pa20	
^{46}Ca(^{3}He,t)^{46}Sc^{i}	−6407	4	−6410	3	−0.6	1	69	69 ^{46}Sc^{i}			71Be29	*
^{46}Sc(β^-)^{46}Ti	2367	3	2366.7	0.7	−0.1	U					53Yo03	*
	2364	6			0.4	U					56Wo09	*
^{46}Ti(p,n)^{46}V	−7844	9	−7834.72	0.09	1.0	U			Tal		63Ja12	
	−7835.8	1.8			0.6	U			Har		76Sq01	Z
^{46}Ti(^{3}He,t)^{46}V	−7069.0	0.6	−7070.96	0.09	−3.3	F			Mun		77Vo02	*
^{46}Ti(^{3}He,t)^{46}V − ^{27}Al()^{27}Si	−2230.8	2.7	−2240.01	0.13	−3.4	B			ChR		74Ha35	
^{46}Ti(^{3}He,t)^{46}V − ^{47}Ti()^{47}V	−4121.62	0.19	−4121.83	0.11	−1.1	1	35	14 ^{47}V	Mun		09Fa15	*
^{46}Ti(^{3}He,t)^{46}V − ^{48}Ti()^{48}V^{i}	−18.57	0.20	−18.58	0.20	−0.0	1	100	100 ^{48}V^{i}	Mun		09Fa15	
^{46}Ti(^{3}He,t)^{46}V − ^{50}Ti()^{50}V^{i}	−31.21	0.25	−31.21	0.25	0.0	1	100	100 ^{50}V^{i}	Mun		09Fa.A	
*^{46}Sc − u	$M - A$=–41520(210) keV for mixture gs+n at 142.528 keV										Nub269	**
*^{46}Ti(p,α)^{43}Sc	Q =–3217 probably to ^{43}Sc^{m} at 151.79 keV										Nub269	**
*^{46}Ti(^{3}He,^{6}He)^{43}Ti	Averaged with reference ; Q reduced by 3 for recalibration ^{27}Al(^{3}He,^{6}He)										75Mu09	**
*^{46}Cr^{i}(p)^{45}V	Q_p=4254(15) 3494(25) 3003(13) to ground state, (5/2$^+$) level at 797.2, (7/2$^+$) at 1272.2 keV										MMC128	**
*											Ens082	**
*^{46}Mn^{i}(p)^{45}Cr	Q_p=4239(33) to (5/2$^+$) state at 493.6 + 107(1)										11Ho02	**
*^{46}Ca(^{3}He,t)^{46}Sc^{i}	CDE=7177(4) Q =–6413(4); recalibration +6 keV for ^{42}Ca(p,n)^{42}Sc from Ame1961										MMC123	**
*^{46}Sc(β^-)^{46}Ti	E_{β^-}=357(3) to 4$^+$ level at 2009.846 keV										Ens00b	**
*^{46}Sc(β^-)^{46}Ti	E_{β^-}=1475(6) to 2$^+$ level at 889.286 keV										Ens00b	**
*^{46}Ti(^{3}He,t)^{46}V	F : rejected in reference of same group										09Fa15	**
*^{46}Ti(^{3}He,t)^{46}V − ^{47}Ti()^{47}V	$Q-Q$ =28.73(0.16) keV to ^{47}V^{i} IAS at 4150.35(0.11) keV										Ens075	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{47}Cl−u		−9576	1074	−10290#	220#	−0.4	D			GA7	1.5	07Ju03	*
^{47}Ar−u		−25400	600	−27232.9	1.3	−2.0	U			TO3	1.5	90Tu01	
		−26570	1360			−0.3	U			GA5	1.5	00Sa21	
		−26903	204			−1.1	U			GA8	1.5	12Ga45	
^{47}Sc−u		−47630	230	−47595.5	1.1	0.1	U			TO6	1.5	98Ba.A	*
C ^{35}Cl−^{47}Ti		17085.94	0.82	17095.23	0.09	4.5	B			H32	2.5	79Ko10	
C ^{13}C H_8 N−^{47}Ti		117329	14	117271.62	0.08	−1.0	U			R09	4.0	72De11	
C_2 H_7 O−^{47}Ti		98012	7	97932.37	0.08	−2.8	B			R09	4.0	72De11	
C_5 H_3−^{47}Ti O		76869	10	76803.01	0.08	−1.6	U			R09	4.0	72De11	
C H_3 O_2−^{47}Ti		61608	10	61546.86	0.08	−1.5	U			R09	4.0	72De11	
^{47}Cr−u		−37103.8	8.6	−37103	5	0.1	–			LZ1	1.0	11Tu09	
		−37107	11			0.4	–			LZ1	1.0	18Zh29	
	ave.	−37105	7			0.3	1	56	56 ^{47}Cr			average	
^{47}Mn−u		−24226	34	−24222	18	0.1	2			LZ1	1.0	13Ya03	
		−24220.0	20.7			−0.1	2			LZ2	1.0	23Zh10	
^{47}Ar−$^{39}K_{1.205}$		16501.8	1.2	16500.8	1.3	−0.8	o			MA8	1.0	15Mo.A	
		16500.8	1.3				2			MA8	1.0	20Mo25	
^{47}K−$^{39}K_{1.205}$		5398.5	2.7	5395.3	1.5	−1.2	o			TT1	1.0	10La.A	
		5395.3	1.5				2			TT1	1.0	12La05	
^{46}Ti ^{13}C−^{47}Ti C		4218.03	0.94	4223.70	0.07	2.4	U			H32	2.5	79Ko10	
^{47}Ti−$^{48}Ti_{.979}$		2723.600	0.058	2723.57	0.03	−0.6	1	35	37 ^{47}Ti	MS1	1.0	17Ka53	
^{47}Ti−^{46}Ti		−929	41	−868.87	0.07	0.4	U			R09	4.0	72De11	
^{47}Ti(d,α)^{45}Sc		6830	12	6845.5	0.7	1.3	U			Kop		67Ha.A	
^{46}Ar(d,p)^{47}Ar		1327	80	1442.8	2.6	1.4	U					06Ga28	
^{46}Ca(n,γ)^{47}Ca		7277.4	0.6	7275.58	0.10	−3.0	B					70Cr04	Z
		7276.1	0.3			−1.7	–			Bdn		06Fi.A	
		7275.50	0.11			0.7	–			ILn		21Bo02	*
^{46}Ca(d,p)^{47}Ca		5055	8	5051.01	0.10	−0.5	U			Kop		67Ha.A	
		5044	4			1.8	U			MIT		68Be36	
^{46}Ca(n,γ)^{47}Ca	ave.	7275.57	0.10	7275.58	0.10	0.1	1	100	100 ^{47}Ca			average	
^{46}Ti(n,γ)^{47}Ti		8875.1	3.0	8880.66	0.06	1.9	U					69Te01	Z
		8880.5	0.3			0.5	U			Bdn		06Fi.A	
^{46}Ti(d,p)^{47}Ti		6658	6	6656.09	0.06	−0.3	U			MIT		67Ba32	*
		6659	8			−0.4	U			Kop		67Ba32	*
		6654.3	1.7			1.1	U			NDm		76Jo01	
^{46}Ti(d,p)^{47}Ti−^{48}Ti()^{49}Ti		738.15	0.25	738.28	0.07	0.5	U			Mun		09Fa15	
^{46}Ti(p,γ)^{47}V		5167.80	0.07	5167.77	0.07	−0.4	1	91	86 ^{47}V	Utr		86De13	*
^{46}Ti(^{3}He,d)^{47}V		−317	15	−325.70	0.07	−0.6	U			MIT		67Do03	
^{47}Mn^{i}(p)^{46}Cr		6867	20	6992	13	6.2	B			Bor		01Gi01	*
		6992	13				3			Bor		07Do17	*
^{47}K(β^-)^{47}Ca		6700	300	6630.1	1.4	−0.2	U					64Ku02	*
^{47}Ca(β^-)^{47}Sc		1984.6	5.	1992.9	1.0	1.7	U					67Hs03	*
		1992.3	5.			0.1	U					68Fi04	*
		1991.9	1.2			0.8	1	74	73 ^{47}Sc			87Ju04	
^{47}Sc(β^-)^{47}Ti		600	2	602.7	1.0	1.4	1	27	27 ^{47}Sc			56Gr12	
^{47}V(β^+)^{47}Ti		2912	10	2930.54	0.09	1.9	U					54Da31	
^{47}Ti(p,n)^{47}V		−3706	13	−3712.89	0.09	−0.5	U			Har		60Mc12	Y
*^{47}Cl−u	Trends in Mass Surface TMS suggest ^{47}Cl 660 keV more bound											HWJ266	**
*^{47}Sc−u	$M-A$=–44320(210) keV for mixture gs+m at 766.83 keV and											Nub269	**
*	assuming ratio R=0.07(3), from half-life=272 ns and TOF=1 μs											GAu	**
*^{47}Cr−u	independent from 11Tu09											WgM194	*W
*^{46}Ca(n,γ)^{47}Ca	From least squares fit to E—g, origianal 7275.4(0.2)											FGK266	**
*^{46}Ti(d,p)^{47}Ti	All 67Ba32 results decreased 0.2% for recalibration											AHW	**
*^{46}Ti(p,γ)^{47}V	E_p=985.94(0.05,Z) to 1/2$^+$ level at 6132.60(0.09) keV											Ens075	**
*^{47}Mn^{i}(p)^{46}Cr	Q_p=5975(25) 4880(20) to 2$^+$ level at 892.16 and (4$^+$) at 1987.1 keV;											Ens102	**
*	data re-analysed and included in 07Do17											07Do17	**
*^{47}Mn^{i}(p)^{46}Cr	Q_p=6104(24) 5000(15) to 2$^+$ level at 892.16 and (4$^+$) at 1987.1 keV											Ens102	**
*	also tentatively Q_p=3973(20) to (3$^-$) at 3196.5 keV, not used											MMC128	**
*^{47}K(β^-)^{47}Ca	E_{β^-}=4100(300) to 2578.33 3/2$^+$ and 2599.53 1/2$^+$ levels											Ens075	**
*^{47}Ca(β^-)^{47}Sc	Original values increased by 4(4) for shape factor											87Ju04	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{48}Ar−u	−23920	330	−23966	4	−0.1	o			MT1	1.0	15Me01	
	−23972	129			0.0	U			RT1	1.0	18Mi08	
	−24037	279			0.3	U			MT1	1.0	20Me06	
	−23999	18			1.8	U			MR1	1.0	20Mo25	
^{48}K−^{39}K$_{1.231}$	10017.7	2.5	10018.5	0.8	0.3	o			TT1	1.0	10La.A	
	10018.50	0.83				2			TT1	1.0	12La05	
^{48}Ca−^{39}K$_{1.231}$	−2799.93	0.22	−2800.032	0.016	−0.5	U			MS1	1.0	12Re17	
^{48}Ca−N ^{18}O O	−44625.15	0.29	−44625.588	0.015	−1.5	U			TT1	1.0	14Kw04	
^{48}Ca−C$_4$	−47477.348	0.019	−47477.352	0.015	−0.2	1	65	65 ^{48}Ca	SH2	1.0	16Ko45	
^{13}C ^{35}Cl−^{48}Ti	24261.73	0.75	24266.88	0.09	2.7	U			H32	2.5	79Ko10	
C$_5$ H$_4$−^{48}Ti O	88492	24	88444.85	0.08	−0.5	U			R09	4.0	72De11	
	88494	27			−0.5	U			R09	4.0	72De11	
C$_4$ H$_2$ N−^{48}Ti O	75935	17	75868.79	0.08	−1.0	U			R09	4.0	72De11	
C$_4$−^{48}Ti	52109	19	52059.34	0.08	−0.7	U			R09	4.0	72De11	
^{48}Ti O−^{85}Rb$_{.753}$	9277.7	1.2	9277.61	0.08	−0.1	U			MA8	1.0	12Na15	
^{48}Ti−N ^{18}O O	−49207.30	0.23	−49207.58	0.08	−1.2	1	11	11 ^{48}Ti	TT1	1.0	14Kw04	
^{48}Mn−u	−31480	120	−31444.8	3.0	0.2	U			GT1	1.5	04St05	
	−31454	10			0.9	U			LZ1	1.0	17Zh12	
	−31444.5	3.1			−0.1	1	92	92 ^{48}Mn	LZ2	1.0	22Wa39	
	−31444.5	3.1			−0.1	o			LZ2	1.0	23Zh10	
^{48}Fe−u	−19333	99				2			LZ1	1.0	20Fu05	
^{48}Ca−^{40}Ca$_{1.200}$	−2586.23	0.23	−2586.3824	0.0016	−0.7	U			MS1	1.0	12Re17	
	−2586.3824	0.0016			0.0	1	100	66 ^{40}Ca	Hep	1.0	25Wi05	
^{48}Ca−^{41}K$_{1.171}$	−2774.65	0.22	−2774.722	0.016	−0.3	U			MS1	1.0	12Re17	
^{48}Ar−^{48}K	10692.8	3.9				3			RKM	1.0	26Fu08	
^{48}Ca−^{48}Sc	298.69	0.82	299.2	0.5	0.6	1	44	44 ^{48}Sc	TT1	1.0	26Ka.A	*
^{48}Ti O−^{55}Mn$_{1.164}$	14972.6	1.2	14972.3	0.4	−0.3	1	12	11 ^{55}Mn	MA8	1.0	12Na15	
^{46}Ti ^{37}Cl−^{48}Ti ^{35}Cl	1726.8	1.1	1735.55	0.09	2.0	U			H18	4.0	64Ba03	
	1730.29	0.87			2.4	U			H32	2.5	79Ko10	
^{48}Ti−^{48}Ca	−4582.018	0.086	−4581.99	0.08	0.3	1	77	76 ^{48}Ti	MS1	1.0	13Bu12	
	−4581.86	0.22			−0.6	U			TT1	1.0	14Kw04	
^{48}Ca−^{48}Ti	4583.20	0.91	4581.99	0.08	−1.3	U			TT1	1.0	26Ka.A	
^{48}Sc−^{48}Ti	4282.47	0.72	4282.8	0.5	0.5	1	57	56 ^{48}Sc	TT1	1.0	26Ka.A	
^{48}Ti−^{47}Ti	−3791	48	−3816.81	0.03	−0.1	U			R09	4.0	72De11	
^{48}Ca(^{3}He,^{11}C)^{40}S	−17416	35	−17105	4	8.9	F			Pri		79Ko.B	*
^{48}Ca(^{3}He,^{8}B)^{43}Cl	−29070	60	−28100.8	2.0	16.2	F			MSU		76Ka24	*
^{48}Ca(α,^{9}Be)^{43}Ar	−21165.5	70.2	−21139	5	0.4	U			Brk		74Je01	
^{48}Ca(^{3}He,^{7}Be)^{44}Ar	−12362	20	−12389.5	1.6	−1.4	U			MSU		76Cr03	*
^{48}Ca(α,^{7}Be)^{45}Ar	−27840.4	60.2	−27798.2	0.5	0.7	U			Brk		74Je01	
^{48}Ti(p,α)^{45}Sc	−2560	5	−2556.6	0.7	0.7	U			ANL		64Yn03	
	−2545	15			−0.8	U			Tal		65Pl01	
^{48}Ca(^{6}Li,^{8}B)^{46}Ar	−23324.8	70.2	−23288.3	2.5	0.5	U			Brk		74Je01	
^{48}Ca(^{14}C,^{16}O)^{46}Ar	−6739	50	−6696.7	2.3	0.8	U			Mun		80Ma40	
^{48}Ca(d,α)^{46}K	1915	15	1899.9	0.7	−1.0	U			ANL		65Ma07	
^{46}Ca(t,p)^{48}Ca	8752	20	8747.9238	0.0019	−0.2	U			Ald		67Bj06	
^{48}Ti(d,α)^{46}Sc	3967	12	3979.5	0.7	1.0	U			Kop		67Ha.A	
^{48}Ti(p,^{3}He)^{46}Sc^{i}	−19394	6	−19388	3	0.9	1	31	31 ^{46}Sc^{i}			78Ko27	
^{48}Ti(p,t)^{46}Ti^{i}	−21192	7				2					78Ko27	
^{48}Ti(p,t)^{46}Ti^{j}	−26177	6				2					78Ko27	
^{46}Ti(^{3}He,n)^{48}Cr	5550	18	5553	7	0.2	R			CIT		67Mi02	
^{48}Ni(2p)^{46}Fe	1400	100	1283	23	−1.2	U					05Gi15	*
	1280	120			0.0	o					11Po09	
	1280	60			0.1	o					12Po03	*
	1290	40			−0.2	3					14Po05	
	1280	28			0.1	3					25Or05	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{48}Ca(^{14}C,^{15}O)^{47}Ar	−18142	100	−18693.3	1.3	−5.5	B			MSU		85Be50	
^{48}Ca(d,^{3}He)^{47}K	−10304	12	−10308.4	1.4	−0.4	U			ANL		66Ne01	
^{48}Ca(t,α)^{47}K	4006	15	4012.0	1.4	0.4	U			LAl		66Wi11	
	4001	10			1.1	U			Ald		68Sa09	
^{48}Ca(d,t)^{47}Ca	−3699	10	−3696.91	0.10	0.2	U			ANL		66Er02	
^{48}Ca(^{3}He,α)^{47}Ca	10630	12	10623.48	0.10	−0.5	U			ANL		66Er02	
	10642	10			−1.9	U			MIT		71Ra35	
^{47}Ti(n,γ)^{48}Ti	11626.39	0.3	11626.66	0.03	0.9	U			MMn		80Is02	Z
	11626.65	0.04			0.2	1	63	61 ^{47}Ti	Ptn		84Ru06	Z
	11626.66	0.23			−0.0	U			Bdn		06Fi.A	
^{47}Ti(d,p)^{48}Ti	9401	8	9402.09	0.03	0.1	U			Kop		67Ba32	
	9403	6			−0.2	U			MIT		67Ba32	
^{47}Ti(^{3}He,d)^{48}V	1337	15	1335.9	1.0	−0.1	U			MIT		68Do06	
^{47}Ti(^{3}He,d)^{48}V^{i}	−1706	20	−1682.96	0.22	1.2	U					68Do06	
^{48}Mn^{i}(p)^{47}Cr	979.6	32.	1018	5	1.2	o			Bor		95Bl05	
	979.6	33.			1.2	o			Bor		96Fa09	*
	1013	12			0.4	–			Bor		07Do17	*
	1018	10			0.0	–					16Or03	
ave.	1016	8			0.3	1	42	33 ^{47}Cr			average	
^{48}K(β^-)^{48}Ca	12000	500	11940.4	0.8	−0.1	U					75Mu08	
^{48}Ca(^{7}Li,^{7}Be)^{48}K	−12959	27	−12802.3	0.8	5.8	B			Can		78We14	
^{48}Ca(^{14}C,^{14}N)^{48}K	−11910	50	−11783.9	0.8	2.5	U			Mun		80Ma40	
^{48}Ca(p,n)^{48}Sc	−534	15	−503.7	0.5	2.0	U					67Mc07	Z
	−506	7			0.3	U					68Mc10	
^{48}Sc(β^-)^{48}Ti	3986	7	3989.4	0.5	0.5	U					57Va08	*
^{48}V(β^+)^{48}Ti	4008	5	4014.9	1.0	1.4	U					53Ma64	*
	4013.6	3.			0.4	1	10	10 ^{48}V			67Ko01	*
	4014	7			0.1	U					74Me15	*
^{48}Ti(p,n)^{48}V	−4803	10	−4797.3	1.0	0.6	U			Tal		62Ne08	Y
^{48}Ti(^{3}He,t)^{48}V^{i}	−7048	4	−7052.39	0.22	−1.1	U					71Be29	*
^{48}V^{i}(IT)^{48}V	3018.7	1.0	3018.8	0.9	0.1	1	90	90 ^{48}V			Ens067	
^{48}Mn^{i}(IT)^{48}Mn	3036.7	0.9	3036.7	0.9	−0.0	1	99	91 ^{48}Mn^{i}			07Do17	*
*^{48}Ca−^{48}Sc	average frequency ratio ^{48}Ca^{+}/(^{48}Sc)$^{+}$ =0.999993737(18),										26Ka.A	**
*^{48}Ca−^{48}Sc	0.999993805(45), additional sys. uncertainty 3.6e−9										26Ka.A	**
*^{48}Ca(^{3}He,^{11}C)^{40}S	F : possible ^{40}Ca contamination; mismatch in cross-sections										GAu	**
*^{48}Ca(^{3}He,^{8}B)^{43}Cl	F : poor spectrum. Authors say: possibly not to ground state										76Ka24	**
*^{48}Ca(^{3}He,^{7}Be)^{44}Ar	$M-A$=–32270(20) Q =–12791(20) for ^{7}Be 429 keV level										GAu	**
*^{48}Ni(2p)^{46}Fe	From only 1 event, Si detector										05Gi15	**
*^{48}Ni(2p)^{46}Fe	From 4 events, gaseous detector										12Po03	**
*^{48}Mn^{i}(p)^{47}Cr	Unexpectedly low intensity 3.6(1.1)%										96Fa09	**
*^{48}Mn^{i}(p)^{47}Cr	Measured intensity 1.8(0.3)%										07Do17	**
*^{48}Sc(β^-)^{48}Ti	E_{β^-}=654(7) to 6^{+} level at 3333.196 keV										Ens067	**
*^{48}V(β^+)^{48}Ti	E_{β^+}=692(5) 698(3) 698(7) respectively, to 4^{+} level at 2295.654 keV										Ens067	**
*^{48}Ti(^{3}He,t)^{48}V^{i}	CDE=7818(4) Q =–7054(4); recalibration +6 keV for ^{42}Ca(p,n)^{42}Sc from Ame1961										MMC123	**
*^{48}Mn^{i}(IT)^{48}Mn	γ cascade 2633.5(0.5)+89.9(0.6)+313.3(0.4)										GAu	**
^{49}Ar−u	−19110	1180	−18223	4	0.8	o			MT1	1.0	15Me01	
	−17499	1180			−0.6	B			MT1	1.0	20Me06	
^{49}K−u	−31981	225	−31789.3	0.9	0.6	U			GT1	1.5	04Ma.A	
^{49}K−^{39}K$_{1.256}$	13794.9	2.8	13795.4	0.9	0.2	o			TT1	1.0	10La.A	
	13795.41	0.86				2			TT1	1.0	12La05	
^{49}Ca−^{39}K$_{1.256}$	1247.1	2.9	1247.34	0.07	0.1	o			TT1	1.0	10La.A	
	1247.1	1.2			0.2	U			TT1	1.0	12La05	
^{49}Sc−u	−49989.3	4.5	−49986.8	2.4	0.5	1	29	29 ^{49}Sc	MR1	1.0	20Ku19	
C H$_2$ ^{35}Cl−^{49}Ti	36637	13	36638.39	0.09	0.0	U			R09	4.0	72De11	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
C_4 H−^{49}Ti		59967	10	59960.66	0.08	−0.2	U			R09	4.0	72De11	
C_5 H_5−^{49}Ti O		96348	19	96346.17	0.08	−0.0	U			R09	4.0	72De11	
C H_5 ^{32}S−^{49}Ti		63365	14	63331.96	0.08	−0.6	U			R09	4.0	72De11	
^{49}Mn−u		−40410	12	−40386.7	2.4	1.9	U			LZ1	1.0	11Tu09	
		−40373	15			−0.9	U			LZ1	1.0	18Zh29	
^{49}Fe−u		−26571	26	−26486	9	3.3	B			LZ1	1.0	12Zh34	*
		−26486.1	9.0				2			LZ2	1.0	23Zh10	
^{47}Ti ^{37}Cl−^{49}Ti ^{35}Cl		946.4	1.1	942.97	0.08	−0.8	U			H18	4.0	64Ba03	
		944.46	0.35			−1.7	U			H32	2.5	79Ko10	
^{48}Ti ^{13}C−^{49}Ti C		3432.64	0.80	3431.122	0.025	−0.8	U			H32	2.5	79Ko10	
^{48}Ti H−^{49}Ti		7876	7	7901.319	0.025	0.9	U			R09	4.0	72De11	
		7874	27			0.3	U			R09	4.0	72De11	
^{49}Ti−^{48}$Ti_{1.021}$		1016.967	0.053	1016.961	0.025	−0.1	1	23	20 ^{49}Ti	MS1	1.0	17Ka53	
^{49}Ar−^{49}K		13566.4	4.2				3			RKM	1.0	26Fu08	
^{49}Mn−^{49}Cr		8279.63	0.25	8279.63	0.25	−0.0	1	100	100 ^{49}Mn	JY1	1.0	14Ka22	
^{49}Ti−^{48}Ti		−43	36	−76.287	0.025	−0.2	U			R09	4.0	72De11	
^{49}Ti(d,α)^{47}Sc		6476	12	6481.7	1.0	0.5	U			Kop		67Ha.A	
^{48}Ca(n,γ)^{49}Ca		5146.6	0.7	5146.40	0.07	−0.3	U					69Ar.A	Z
		5146.38	0.30			0.1	U					70Cr04	Z
		5146.48	0.23			−0.4	2			Bdn		06Fi.A	
		5146.39	0.07			0.1	2			ILn		21Bo02	*
^{48}Ca(d,p)^{49}Ca		2917	7	2921.83	0.07	0.7	U			ANL		66Er02	
		2917	4			1.2	U			MIT		68Be36	
^{48}Ca(p,γ)^{49}Sc		9628.7	3.6	9626.6	2.3	−0.6	–					68Vi01	Z
^{48}Ca(d,n)^{49}Sc		7404	7	7402.0	2.3	−0.3	–					68Gr09	
^{48}Ca(^{3}He,d)^{49}Sc		4150	12	4133.1	2.3	−1.4	U			ANL		66Er02	
^{48}Ca(p,γ)^{49}Sc	ave.	9629	3	9626.6	2.3	−0.7	1	50	50 ^{49}Sc			average	
^{48}Ti(n,γ)^{49}Ti		8142.22	0.3	8142.379	0.023	0.5	U			MMn		80Is02	Z
		8142.39	0.03			−0.4	1	61	67 ^{49}Ti	Ptn		83Ru08	Z
		8142.35	0.16			0.2	U			Bdn		06Fi.A	
^{48}Ti(d,p)^{49}Ti		5907	8	5917.813	0.023	1.4	U			Kop		67Ba32	
		5918	6			−0.0	U			MIT		67Ba32	
		5918.6	1.7			−0.5	U			NDm		76Jo01	
^{48}Ti(p,γ)^{49}V		6756.8	1.5	6758.2	0.8	0.9	R					72Ki06	
^{49}Mn^i(p)^{48}Cr		2712.2	50.	2729	16	0.3	U					70Ce02	*
		2730	29			−0.0	o			Bor		96Fa09	*
		2729	16				3			Bor		07Do17	*
^{49}K(β^-)^{49}Ca		10970	70	11688.5	0.8	10.3	B					86Mi08	
^{49}Ca(β^-)^{49}Sc		5200	100	5262.5	2.3	0.6	U					56Ma27	
		4970	50			5.9	B					56Ok02	
^{49}Sc(β^-)^{49}Ti		2010	5	2001.6	2.3	−1.7	1	21	21 ^{49}Sc			61Re06	
		1983	7			2.7	B					69Fl02	
^{49}V(ε)^{49}Ti		626	10	601.9	0.8	−2.4	U					56Ha59	
^{49}Ti(p,n)^{49}V		−1383	9	−1384.2	0.8	−0.1	U			Har		60Mc12	Z
		−1383.6	1.0			−0.6	2			Oak		64Jo11	Z
^{49}Ti(^{3}He,t)^{49}V^i		−7052	4				2					71Be29	*
^{49}Cr(β^+)^{49}V		2590	20	2629.8	2.3	2.0	U					53Cr18	*
*^{49}Fe−u	Same result in reference											13Ya03	**
*^{48}Ca(n,γ)^{49}Ca	From least squares fit to Eg, original 5146.3(0.3)											FGK266	**
*^{49}Mn^i(p)^{48}Cr	Q_p=1960(50) 1978(29) 1977(16) respectively, to 2^+ level at 752.19(0.11) keV											Ens067	**
*^{49}Ti(^{3}He,t)^{49}V^i	CDE=7822(4) Q =–7058(4); recalibration +6 keV for ^{42}Ca(p,n)^{42}Sc from Ame1961											MMC123	**
*^{49}Cr(β^+)^{49}V	E_{β^+}=1540(10) 1390(20) to (7/2$^-$)ground state + (5/2$^-$)90.6392 and 3/2$^-$ at 152.9282											Ens089	**
^{50}Ar−C F_2		−10943.0	9.1				2			RKM	1.0	26Fu08	
^{50}K−u		−26100	800	−27620	8	−1.3	U			TO3	1.5	90Tu01	
^{50}K−^{39}$K_{1.282}$		18899	11	18908	8	0.8	o			TT1	1.0	10La.A	
		18908.3	8.3				2			TT1	1.0	12La05	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{50}Ca−^{39}K$_{1.282}$	4027.0	4.0	4027.5	1.7	0.1	o			TT1	1.0	10La.A	
	4027.5	1.7				2			TT1	1.0	12La05	
^{50}Sc−u	−47940	250	−47812.6	2.7	0.3	U			TO6	1.5	98Ba.A	*
	−47822.6	9.8			1.0	U			MR1	1.0	20Ku19	
C H$_3$ ^{35}Cl−^{50}Ti	47550	23	47542.19	0.09	−0.1	U			R09	4.0	72De11	
C$_4$ H$_2$−^{50}Ti	70860	8	70864.46	0.09	0.1	U			R09	4.0	72De11	
C$_5$ H$_6$−^{50}Ti O	107253	18	107249.97	0.09	−0.0	U			R09	4.0	72De11	
C$_3$ ^{13}C H−^{50}Ti	66401	21	66394.27	0.09	−0.1	U			R09	4.0	72De11	
C$_3$ N−^{50}Ti	58279	43	58288.40	0.09	0.1	U			R09	4.0	72De11	
C$_4$ H$_2$−^{50}V	68485	14	68493.40	0.10	0.2	U			R09	4.0	72De11	
C$_3$ N−^{50}V	55903	23	55917.34	0.10	0.2	U			R09	4.0	72De11	
C H$_3$ ^{35}Cl−^{50}V	45158	17	45171.13	0.10	0.2	U			R09	4.0	72De11	
C$_4$ H$_2$−^{50}Cr	69608	8	69607.87	0.10	−0.0	U			R09	4.0	72De11	
C$_3$ N−^{50}Cr	57051	7	57031.81	0.10	−0.7	U			R09	4.0	72De11	
C H$_3$ ^{35}Cl−^{50}Cr	46290	14	46285.60	0.10	−0.1	U			R09	4.0	72De11	
^{50}Fe−u	−37012	9	−37011.4	2.9	0.1	2			LZ1	1.0	17Zh12	
	−37011.3	3.1			−0.0	2			LZ2	1.0	22Wa39	
	−37011.3	3.1			−0.0	o			LZ2	1.0	23Zh10	
^{50}Co−u	−18883	135				2			LZ1	1.0	20Fu05	*
^{50}Ti−^{48}Ti$_{1.042}$	−968.526	0.048	−968.56	0.03	−0.7	1	50	45 ^{50}Ti	MS1	1.0	17Ka53	
^{50}V−^{48}Ti$_{1.042}$	1402.62	0.11	1402.50	0.06	−1.1	1	26	25 ^{50}V	MS1	1.0	17Ka53	
^{50}Cr−^{48}Ti$_{1.042}$	287.979	0.072	288.03	0.06	0.7	1	65	63 ^{50}Cr	MS1	1.0	17Ka53	
^{49}Ti ^{13}C−^{50}Ti C	6440.47	0.88	6433.60	0.03	−3.1	B			H32	2.5	79Ko10	
^{50}V−^{50}Ti	2371.14	0.12	2371.06	0.06	−0.7	1	27	20 ^{50}V	MS1	1.0	17Ka53	
^{50}V−^{50}Cr	1114.414	0.075	1114.47	0.06	0.8	1	62	33 ^{50}Cr	MS1	1.0	17Ka53	
^{50}Mn−^{50}Cr	8195.91	0.10	8195.95	0.07	0.4	1	52	52 ^{50}Mn	JY1	1.0	08Er04	
^{50}Mn^m−^{50}Cr	8437.852	0.065	8437.83	0.06	−0.3	1	81	81 ^{50}Mn^m	JY1	1.0	08Er04	
^{50}Mn^m−^{50}Mn	241.840	0.100	241.88	0.07	0.4	1	55	37 ^{50}Mn	JY1	1.0	08Er04	
^{50}Ti−^{49}Ti	−3075	38	−3078.77	0.03	−0.0	U			R09	4.0	72De11	
^{50}Sc O−^{19}F ^{35}Cl	−20153.8	2.7	−20153.8	2.7	0.0	1	100	100 ^{50}Sc	MS1	1.0	21Le02	
^{50}Cr(p,^{6}He)^{45}V	−28686	17	−28678.1	0.9	0.5	U			MSU		75Mu09	*
^{50}Ti(p,α)^{47}Sc	−2231	15	−2232.9	1.0	−0.1	U			Tal		65Pl01	
^{50}V(p,α)^{47}Ti	572	23	578.43	0.06	0.3	U			MIT		67Sp09	
^{50}Cr(p,α)^{47}V	−3387	10	−3390.24	0.10	−0.3	U			Ald		66Br06	
^{50}Cr(^{3}He,^{6}He)^{47}Cr	−18365	14	−18361	5	0.3	1	11	11 ^{47}Cr	MSU		77Mu03	*
^{48}Ca(t,p)^{50}Ca	3012	15	3025.2	1.6	0.9	U			Ald		66Hi01	
	3020	10			0.5	U			LAl		66Wi11	
^{48}Ca(^{3}He,p)^{50}Sc	7965	15	7954.5	2.5	−0.7	U			ANL		69Oh01	
^{50}V(d,α)^{48}Ti	9982	15	9980.52	0.05	−0.1	U			MIT		66Do06	
	9988	20			−0.4	U			Kop		67Ha.A	
^{50}Cr(d,α)^{48}V	4928	12	4927.4	1.0	−0.0	U			Kop		67Ha.A	
	4923	15			0.3	U			MIT		68Do03	
^{50}Cr(d,α)^{48}V^i	1880	11	1908.60	0.23	2.6	U			MIT		68Do03	*
^{50}Cr(p,t)^{48}Cr	−15100	8	−15101	7	−0.1	2			Oak		71Do18	
	−15100	30			−0.0	U			Bld		72Sh27	
^{50}Cr(p,t)^{48}Cr^j	−23861	15				2			MSU		75Mo26	*
^{50}Co^i(2p)^{48}Mn	1972	13				2			Bor		07Do17	
^{50}Ti(t,α)^{49}Sc	7644	25	7655.5	2.3	0.5	U			LAl		66Wi11	
^{49}Ti(n,γ)^{50}Ti	10939.6	0.3	10939.17	0.03	−1.4	U			MMn		80Is02	Z
	10939.19	0.04			−0.4	1	60	48 ^{50}Ti	Ptn		84Ru06	Z
	10939.20	0.22			−0.1	U			Bdn		06Fi.A	
^{49}Ti(d,p)^{50}Ti	8723	8	8714.61	0.03	−1.0	U			Kop		67Ba32	
	8721	6			−1.1	U			MIT		67Ba32	
^{50}Cr(p,d)^{49}Cr	−10790	30	−10775.8	2.2	0.5	U			Pri		67Wh03	
^{50}Cr(d,t)^{49}Cr	−6743.1	2.2	−6743.1	2.2	0.0	1	100	100 ^{49}Cr	NDm		76Jo01	
^{50}Fe^i(p)^{49}Mn	4389	41	4332	10	−1.4	o			Bor		96Fa09	*
	4332	10				2			Bor		07Do17	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{50}K(β^-)^{50}Ca	14050	300	13861	8	−0.6	U					86Mi08	
^{50}Sc(β^-)^{50}Ti	6500	200	6894.8	2.5	2.0	U					63Ch03	
	6260	100			6.3	B					69Wa24	
^{50}V(n,p)^{50}Ti	2979	15	2990.98	0.06	0.8	U			ILL		81Wa31	
	2984	10			0.7	U			ILL		94Wa17	
^{50}Ti(p,n)^{50}V	−2991	10	−2990.98	0.06	0.0	U			Har		60Mc12	Y
^{50}Ti(^{3}He,t)^{50}V^i	−7032	4	−7039.75	0.27	−1.9	U					71Be29	*
^{50}Cr(p,n)^{50}Mn	−8416.1	1.9	−8416.82	0.07	−0.4	U			Har		75Fr.A	
^{50}Cr(^{3}He,t)^{50}Mn	−7650.5	0.4	−7653.07	0.07	−6.4	F			Mun		77Vo02	*
^{50}Cr(^{3}He,t)^{50}Mn−^{27}Al()^{27}Si	−2820.0	2.8	−2822.12	0.12	−0.8	U			ChR		74Ha35	
^{50}Cr(^{3}He,t)^{50}Mn−^{42}Ca()^{42}Sc	−1207.6	2.3	−1208.14	0.08	−0.2	U			ChR		74Ha35	
^{50}Cr(^{3}He,t)^{50}Mn−^{54}Fe()^{54}Co	610.09	0.17	610.07	0.10	−0.1	1	35	23 ^{54}Co	ChR		87Ko34	*
*^{50}Sc−u	$M-A$=–44530(220) keV for mixture gs+m at 256.894 keV										Nub269	**
*^{50}Co−u	Authors suggested ground state										20Fu05	**
*^{50}Cr(p,^{6}He)^{45}V	Original Q increase by 1 for recalibration										AHW	**
*^{50}Cr(^{3}He,^{6}He)^{47}Cr	Original Q reduced by 3, see ^{46}Ti(^{3}He,^{6}He)										AHW	**
*^{50}Cr(d,α)^{48}V^i	IT=3043(9); rebuilt from their Q_{gs}=4923(15) keV										MMC124	**
*^{50}Cr(p,t)^{48}Cr^j	Strongest of two fragments given as IT=8760(15); Q rebuilt with Ame1971										75Mo26	**
*^{50}Fe^i(p)^{49}Mn	E_p=2790(41) to 11/2$^{(-)}$ level at 1541.3125 keV										Ens089	**
*^{50}Fe^i(p)^{49}Mn	Q_p=2770(12) 41.1%, 1874(16) 1.0% to 11/2$^{(-)}$ level at 1541.3125, and										Ens089	**
*	13/2$^{(-)}$ at 2481.3 keV										Ens089	**
*^{50}Ti(^{3}He,t)^{50}V^i	CDE=7802(4) Q =–7038(4); recalibration +6 keV for ^{42}Ca(p,n)^{42}Sc from Ame1961										MMC123	**
*^{50}Cr(^{3}He,t)^{50}Mn	F : rejected in reference of same group										09Fa15	**
*^{50}Cr(^{3}He,t)^{50}Mn−^{54}Fe()^{54}Co	$Q-Q$ =40.90(0.16) to 650.99(0.06) level in ^{50}Mn										Ens10c	**
^{51}Ca−^{39}K$_{1.308}$	8467.58	0.56				2			MA8	1.0	13Wi06	
^{51}Ca−u	−38800	350	−39004.3	0.6	−0.4	U			TO3	1.5	90Tu01	
	−38900	400			−0.2	U			TO5	1.5	94Se12	
	−39249	183			0.9	U			GT1	1.5	04Ma.A	
^{51}Sc H−^{19}F ^{33}S	−8468.2	2.7	−8468.2	2.7	−0.0	1	100	100 ^{51}Sc	MS1	1.0	21Le02	
^{51}Ti−^{39}K$_{1.308}$	−5917.0	2.3	−5918.7	0.5	−0.7	U			TT1	1.0	18Re11	
^{51}Ti−u	−53378.8	16.1	−53390.6	0.5	−0.7	U			TR1	1.0	18Le03	
	−53389.0	2.3			−0.7	o			TT1	1.0	18Le03	*
^{51}V−^{39}K$_{1.308}$	−8571.23	0.73	−8570.43	0.10	1.1	U			TT1	1.0	14Ma21	
	−8570.68	0.99			0.3	U			TT1	1.0	14Ma21	
	−8570.9	2.0			0.2	U			TT1	1.0	18Re11	
C$_4$ H$_3$−^{51}V	79526	9	79517.45	0.10	−0.2	U			R09	4.0	72De11	
C$_5$ H$_7$−^{51}V O	115921	13	115902.95	0.10	−0.3	U			R09	4.0	72De11	
C$_4$ H$_5$ N−^{51}V O	103334	13	103326.89	0.10	−0.1	U			R09	4.0	72De11	
C$_3$ H N−^{51}V	66943	7	66941.39	0.10	−0.1	U			R09	4.0	72De11	
^{51}V−u	−56042.8	1.9	−56042.35	0.10	0.2	o			TT1	1.0	18Le03	
	−56042.34	0.26			−0.0	1	14	14 ^{51}V	MS2	1.0	26Ir.1	
^{51}Cr−^{39}K$_{1.308}$	−7763.66	0.78	−7762.70	0.18	1.2	–			TT1	1.0	14Ma21	
	−7764.0	1.2			1.1	–			TT1	1.0	14Ma21	
ave.	−7763.8	0.7			1.6	1	7	7 ^{51}Cr			average	
^{51}Fe−u	−43148	12	−43144.9	1.5	0.3	U			LZ1	1.0	11Tu09	
	−43154	15			0.6	U			LZ1	1.0	18Zh29	
	−43158.5	17.1			0.8	U			LZ2	1.0	22Wa39	
	−43158.5	17.1			0.8	o			LZ2	1.0	23Zh10	
^{51}Fe−C H$_2$ ^{37}Cl	−24697.5	1.5				2			MS1	1.0	18On01	
^{51}Co−u	−29353	52	−29391	6	−0.7	U			LZ1	1.0	14Sh14	
	−29342	44			−1.1	U			LZ1	1.0	21Zh.B	
	−29400.1	12.3			0.8	1	22	22 ^{51}Co	LZ2	1.0	23Zh10	
^{51}Co−^{14}N$_2$	−35536.2	6.5	−35539	6	−0.4	1	78	78 ^{51}Co	MS2	1.0	26Ma.A	
^{51}Ca−^{58}Ni$_{.879}$	17823	24	17829.8	0.7	0.3	U			TT1	1.0	12Ga29	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{47}Ti ^{37}Cl$_2$−^{51}V ^{35}Cl$_2$		1906.	1.8	1899.56	0.15	−0.9	U			H18	4.0	64Ba03	
^{51}V−^{48}Ti$_{1.063}$		−703.292	0.082	−703.27	0.06	0.3	1	58	51 ^{51}V	MS1	1.0	17Ka53	
^{49}Ti ^{37}Cl−^{51}V ^{35}Cl		956.7	0.7	956.59	0.10	−0.0	U			H18	4.0	64Ba03	
^{51}K−^{51}V		31871	14	31881	6	0.7	2			TT1	1.0	12Ga29	
		31882.9	6.4			−0.3	2			TR1	1.0	26Wa.A	
^{51}Cr−^{51}V		807.46	0.71	807.72	0.16	0.4	–			TT1	1.0	14Ma21	
		806.3	1.2			1.2	–			TT1	1.0	14Ma21	
	ave.	807.2	0.6			0.9	1	7	6 ^{51}Cr			average	
^{48}Ca(^{14}C,^{11}C)^{51}Ca		−15900	150	−15522.1	0.5	2.5	U			Mun		80Ma40	*
		−16886	100			13.6	B			MSU		85Be50	
^{48}Ca(^{18}O,^{15}O)^{51}Ca		−12040	120	−11531.0	0.7	4.2	B			Hei		85Br03	*
		−13900	40			59.2	B			Can		88Ca21	
^{48}Ca(α,p)^{51}Sc		−5860	20	−5838.6	2.5	1.1	U			ANL		66Er02	
^{51}V(p,α)^{48}Ti		1162	10	1153.91	0.06	−0.8	U			MIT		64Sp12	
^{51}V(d,α)^{49}Ti		7066	12	7071.72	0.06	0.5	U			Kop		67Ha.A	
^{50}Ti(n,γ)^{51}Ti		6372.3	1.2	6372.5	0.5	0.2	2					71Ar39	Z
		6372.6	0.6			−0.2	2			Bdn		06Fi.A	
^{50}Ti(d,p)^{51}Ti		4143	6	4147.9	0.5	0.8	U			MIT		67Ba32	
		4148	8			−0.0	U			Kop		67Ba32	
		4147.7	1.2			0.2	2			NDm		76Jo01	
^{50}Ti(p,γ)^{51}V		8063.3	2.0	8060.20	0.06	−1.5	U					70Kl05	Z
		8063.6	2.0			−1.7	U					70Ma36	Z
^{50}Ti(^{3}He,d)^{51}V		2555	15	2566.73	0.06	0.8	U			MIT		67Ob04	
^{50}V(n,γ)^{51}V		11051.18	0.10	11051.18	0.06	−0.0	–			MMn		78Ro03	Z
		11051.05	0.17			0.8	–			ILn		91Mi08	Z
		11051.14	0.22			0.2	–			Bdn		06Fi.A	
^{51}V(γ,n)^{50}V		−11040	60	−11051.18	0.06	−0.2	U			Phi		60Ge01	
^{50}V(d,p)^{51}V		8840	15	8826.61	0.06	−0.9	U			MIT		67De02	
		8828	20			−0.1	U			Kop		67Ha.A	
^{51}V(p,d)^{50}V		−8815	20	−8826.61	0.06	−0.6	U			Oak		65Ba29	
^{50}V(n,γ)^{51}V	ave.	11051.15	0.08	11051.18	0.06	0.4	1	57	31 ^{51}V			average	
^{50}V(^{3}He,d)^{51}Cr		4031	12	4022.97	0.15	−0.7	U			MIT		69Do01	
^{50}Cr(n,γ)^{51}Cr		9261.71	0.30	9260.67	0.15	−3.5	B			MMn		80Is02	Z
		9260.63	0.20			0.2	1	55	51 ^{51}Cr	Bdn		06Fi.A	
^{50}Cr(d,p)^{51}Cr		7049	8	7036.10	0.15	−1.6	U			Kop		67Ha.A	
		7041	6			−0.8	U			MIT		68Ro09	
^{50}Cr(p,γ)^{51}Mn		5270.8	0.3	5270.79	0.29	−0.0	1	94	93 ^{51}Mn			72Fo25	Z
^{50}Cr(^{3}He,d)^{51}Mn		−206	15	−222.69	0.29	−1.1	U			MIT		67Sp09	
^{50}Cr(p,γ)^{51}Mn^{i}		819	2	820.2	1.5	0.6	2					72Fo25	
^{50}Cr(^{3}He,d)^{51}Mn^{i}		−4652	20	−4673.3	1.5	−1.1	U			MIT		67Ra14	
		−4671.7	2.3			−0.7	2					79Pa14	*
^{51}Co^{i}(p)^{50}Fe		6513	16				3			Bor		07Do17	*
^{51}Ti(β^-)^{51}V		2440	30	2470.1	0.5	1.0	U					55Bu01	
		2450	30			0.7	U					55Ma01	
^{51}Cr(ε)^{51}V		756	5	752.39	0.15	−0.7	U					55Bi29	
^{51}V(p,n)^{51}Cr		−1533.5	2.0	−1534.74	0.15	−0.6	U			Nvl		59Go68	Z
		−1533.3	1.8			−0.8	U			Oak		64Jo11	Z
		−1533.7	1.5			−0.7	U			Can		70Kn03	Z
		−1534.93	0.24			0.8	1	39	36 ^{51}Cr	PTB		89Sc24	Z
^{51}V(^{3}He,t)^{51}Cr^{i}		−7384	5				2					71Be29	
^{51}Mn(β^+)^{51}Cr		3232	20	3207.5	0.3	−1.2	U					66Gl02	
*^{51}Ti−u	Frequency ratios given in 18Re11											HWJ192	**
*^{48}Ca(^{14}C,^{11}C)^{51}Ca	May be a ^{40}Ca contamination. There is a –16900(150) peak											85Be50	**
*^{48}Ca(^{18}O,^{15}O)^{51}Ca	Proposed 970(90) level reinterpreted as ground state in reference											85Be50	**
*^{48}Ca(^{18}O,^{15}O)^{51}Ca	Weak $M-A$=–36120(120) level disregarded											AHW	**
*^{50}Cr(^{3}He,d)^{51}Mn^{i}	IT=4449(3); Q rebuilt with Ame1977											MMC124	**
*^{51}Co^{i}(p)^{50}Fe	Q_p=4662(16) to (4^+) level at 1851.5 keV											Ens10c	**
*^{51}Co^{i}(p)^{50}Fe	Q_p=6153 was a misprint in Ame2012											XuX16b	**
*^{51}V(^{3}He,t)^{51}Cr^{i}	CDE=8145(5) Q =–7881(5); recalibration –3 keV for ^{50}Cr(p,n)^{50}Mn from Ame1961											MMC123	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{52}K−u		−18398	36	−18462	6	−1.8	U			MR1	1.0	15Ro10	
^{52}Ca−^{39}K$_{1.333}$		11592.90	0.72				2			MA8	1.0	13Wi06	
^{52}Ca−u		−34900	500	−36786.4	0.7	−2.5	U			TO3	1.5	90Tu01	
		−36792	11			0.5	U			MR1	1.0	13Wi06	
^{52}Sc−^{33}S ^{19}F		−13365.9	3.3	−13366	3	0.0	1	100	100 ^{52}Sc	MS1	1.0	21Le02	
^{52}Sc−u		−43500	230	−43504	3	−0.0	U			TO3	1.5	90Tu01	
		−43350	250			−0.4	U			TO5	1.5	94Se12	
		−43110	240			−1.1	U			TO6	1.5	98Ba.A	
		−43438	97			−0.7	o			LZ1	1.0	15Xu14	
		−43260	560			−0.4	o			MT1	1.0	15Me08	
		−43505	70			0.0	U			LZ1	1.0	19Xu09	
		−43607	247			0.4	U			MT1	1.0	20Me06	
^{52}Ti−^{39}K$_{1.333}$		−4738.7	3.2	−4738.4	1.9	0.1	1	36	36 ^{52}Ti	TT1	1.0	18Re11	
^{52}Ti−u		−53104	17	−53117.7	1.9	−0.8	U			TR1	1.0	18Le03	
		−53118.0	3.2			0.1	o			TT1	1.0	18Le03	
		−53117.5	2.4			−0.1	1	64	64 ^{52}Ti	MS1	1.0	22Po06	*
^{52}V−u		−55198	28	−55226.38	0.17	−1.0	U			TR1	1.0	18Re11	
^{52}Cr−^{39}K$_{1.333}$		−11116.29	0.48	−11116.06	0.12	0.5	U			MA8	1.0	19Hu15	
		−11118.0	2.1			0.9	U			TT1	1.0	18Re11	
C$_4$ H$_4$−^{52}Cr		90826	9	90795.44	0.12	−0.8	U			R09	4.0	72De11	
C$_3$ ^{13}C H$_3$−^{52}Cr		86373	18	86325.25	0.12	−0.7	U			R09	4.0	72De11	
C$_3$ H$_2$ N−^{52}Cr		78253	6	78219.38	0.12	−1.4	U			R09	4.0	72De11	
^{52}Cr−^{85}Rb$_{.612}$		−5510.46	0.57	−5510.63	0.12	−0.3	U			MA8	1.0	19Hu15	
^{52}Cr−u		−59497.2	2.1	−59495.32	0.12	0.9	o			TT1	1.0	18Le03	
^{52}Co−u		−36888	9	−36874	4	1.5	–			LZ1	1.0	16Xu10	
		−36879.0	5.9			0.8	–			LZ2	1.0	22Wa39	
		−36879.0	5.9			0.8	o			LZ2	1.0	23Zh10	
	ave.	−36882	5			1.5	1	69	69 ^{52}Co			average	
^{52}Co^m−u		−36473	11	−36468	7	0.5	–			LZ1	1.0	16Xu10	
		−36471.5	11.4			0.3	–			LZ2	1.0	22Wa39	
		−36471.5	11.4			0.3	o			LZ2	1.0	23Zh10	
	ave.	−36472	8			0.6	1	70	70 ^{52}Co^m			average	
^{52}Ni−u		−24219	89	−24127.6	1.8	1.0	U			LZ1	1.0	20Fu05	
^{52}Ni−^{12}C$_2$ ^{1}H$_2$		−39777.7	1.8				2			MS2	1.0	26Ma.A	
^{52}Ca−^{58}Ni$_{.897}$		21220	110	21211.6	0.8	−0.1	U			TT1	1.0	12Ga29	
^{52}Cr−^{48}Ti$_{1.083}$		−3114.916	0.096	−3115.05	0.08	−1.4	1	78	77 ^{52}Cr	MS1	1.0	17Ka53	
^{52}K−^{52}Cr		41032.9	6.0				2			TR1	1.0	26Wa.A	
^{52}Ca−^{52}Cr		22740	82	22709.0	0.7	−0.4	U			TT1	1.0	12Ga29	
^{52}Mn−^{52}Cr		5054.376	0.068				2			JY1	1.0	17Ne05	
^{52}Fe−^{52}Cr		7608.65	0.15	7608.65	0.15	−0.0	1	100	100 ^{52}Fe	JY1	1.0	17Ne05	
^{52}Fe^m−^{52}Cr		15081.23	0.29	15077.8	0.6	−11.7	B			JY1	1.0	17Ne05	
^{52}Co−^{52}Cr		22637.5	7.3	22621	4	−2.2	1	31	31 ^{52}Co	JY1	1.0	17Ne05	
^{52}Co^m−^{52}Cr		23038	12	23028	7	−0.9	1	30	30 ^{52}Co^m	JY1	1.0	17Ne05	
^{52}Cr−^{50}Cr		−5566	41	−5537.51	0.10	0.2	U			R09	4.0	72De11	
^{52}Cr(p,α)^{49}V		−2596	10	−2593.3	0.8	0.3	U			Ald		66Br06	
^{50}Ti(t,p)^{52}Ti		5698	10	5707.7	1.8	1.0	U			LAl		66Wi11	
		5700	10			0.8	U			LAl		71Ca19	
^{50}Ti(^{3}He,p)^{52}V		7653	15	7653.41	0.14	0.0	U			Phi		75Ca07	
^{52}Cr(d,α)^{50}V		4517	12	4514.53	0.09	−0.2	U			Kop		67Ha.A	
^{52}Cr(p,^{3}He)^{50}V^i		−18645	6	−18651.06	0.28	−1.0	U					78Ko27	
^{52}Cr(p,t)^{50}Cr^i		−21244	7				2					78Ko27	
^{52}Cr(p,t)^{50}Cr^j		−26041	6				2					78Ko27	
^{51}V(n,γ)^{52}V		7311.2	0.5	7311.24	0.13	0.1	2					84De15	
		7311.18	0.26			0.2	2			ILn		91Mi08	Z
		7311.27	0.15			−0.2	2			Bdn		06Fi.A	
^{51}V(d,p)^{52}V		5098	9	5086.68	0.13	−1.3	U			MIT		64Sp12	
		5086	8			0.1	U			Kop		67Ha.A	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{51}V(p,γ)^{52}Cr	10500.7	2.8	10505.39	0.10	1.7	U					74Ro44	Z
^{52}Co^{i}(p)^{51}Fe	1367	60	1479	7	1.9	o			Bor		94Fa06	
	1349	10			13.0	B			Bor		07Do17	
	1352	10			12.7	B					16Or03	
^{52}Ca(β^-)^{52}Sc	5700	200	6257	3	2.8	B					85Hu03	
^{52}Sc(β^-)^{52}Ti	8020	250	8955	4	3.7	B					85Hu03	
^{52}Ti(β^-)^{52}V	1940	200	1964.3	1.8	0.1	U					67Mo11	*
^{52}V(β^-)^{52}Cr	3904	30	3976.49	0.16	2.4	U					65Ko09	*
	3854	30			4.1	B					67Va27	*
^{52}Mn(β^+)^{52}Cr	4710.9	4.	4708.12	0.06	−0.7	U					58Ko57	*
	4707.9	6.			0.0	U					60Ka20	*
^{52}Cr(p,n)^{52}Mn	−5479	10	−5490.47	0.06	−1.1	U			Ric		66Ri09	
^{52}Cr(^{3}He,t)^{52}Mn^{i}	−7653	5				2					71Be29	*
^{52}Fe(β^+)^{52}Mn	2372	10	2379.29	0.15	0.7	U					56Ar33	*
	2229	130			1.2	U					79Ge02	*
	2510	100			−1.3	U					95Ir01	
^{52}Fe^{m}(IT)^{52}Fe	6957.5	0.5				2					03Ax01	*
^{52}Co^{i}(IT)^{52}Co^{m}	2548	2				2					16Or03	
*^{52}Ti(β^-)^{52}V	E_{β^-}=1800(200) to 1^+ level at 141.610 keV										Ens159	**
*^{52}V(β^-)^{52}Cr	E_{β^-}=2470(30) 2420(30) respectively, to 2^+ level at 1434.091 keV										Ens159	**
*^{52}Mn(β^+)^{52}Cr	E_{β^+}=575(4) and 572(6) respectively, to 6^+ level at 3113.858 keV										Ens159	**
*^{52}Cr(^{3}He,t)^{52}Mn^{i}	CDE=8414(5) Q =–7650(5); recalibration –3 keV for ^{50}Cr(p,n)^{50}Mn from Ame1961										MMC123	**
*^{52}Fe(β^+)^{52}Mn	E_{β^+}=804(10) to 1^+ level at 546.438 keV										Ens159	**
*^{52}Fe(β^+)^{52}Mn	E_{β^+}=5350(130) from ^{52}Fe^{m} 12^+ at 6958.0 to 11^+ level at 3837.2 keV										Ens159	**
*^{52}Fe^{m}(IT)^{52}Fe	E=6958.0(0.4) but 0.3 keV unc assumed for 465.0 and 597.1 γs										Ens157	**
^{53}K−u	−13200	120	−13228	7	−0.2	U			MR1	1.0	15Ro10	
^{53}Ca−u	−31549	47				2			MR1	1.0	13Wi06	
^{53}Sc−^{34}S ^{19}F	−7891	19	−7891	19	−0.0	1	100	100 ^{53}Sc	MS1	1.0	21Le02	
^{53}Sc−u	−41440	260	−41621	19	−0.5	o			TO3	1.5	90Tu01	
	−41830	280			0.5	U			TO5	1.5	94Se12	
	−41100	400			−0.9	U			TO6	1.5	98Ba.A	
	−41694	118			0.4	U			GT1	1.5	04Ma.A	
	−40910	290			−2.5	B			MT1	1.0	11Es06	
	−41804	123			1.5	o			LZ1	1.0	15Xu14	
	−40980	610			−1.1	o			MT1	1.0	15Me08	
	−41772	86			1.8	U			LZ1	1.0	19Xu09	
	−41224	225			−1.8	U			MT1	1.0	20Me06	
^{53}Ti−^{39}K$_{1.359}$	−1006.4	3.1				2			TT1	1.0	18Re11	
^{53}Ti−u	−50325	19	−50329	3	−0.2	U			TR1	1.0	18Le03	
	−50329.3	3.1			0.0	o			TT1	1.0	18Le03	
^{53}V−u	−55664	20	−55665	3	−0.1	U			TR1	1.0	18Re11	
^{53}Cr−^{39}K$_{1.359}$	−10031.7	2.0	−10030.86	0.12	0.4	U			TT1	1.0	18Re11	
C$_4$ H$_5$−^{53}Cr	98529	8	98478.91	0.12	−1.6	U			R09	4.0	72De11	
C$_3$ H$_3$ N−^{53}Cr	85958	10	85902.85	0.12	−1.4	U			R09	4.0	72De11	
C$_2$ ^{13}C H$_2$ N−^{53}Cr	81507	27	81432.65	0.12	−0.7	U			R09	4.0	72De11	
C$_3$ H O−^{53}Cr	62152	14	62093.40	0.12	−1.0	U			R09	4.0	72De11	
^{53}Cr−^{85}Rb$_{.624}$	−4311.80	0.58	−4310.54	0.12	2.2	U			MA8	1.0	19Hu15	
^{53}Cr−u	−59354.5	2.0	−59353.75	0.12	0.4	o			TT1	1.0	18Le03	
^{53}Co−u	−45783	18	−45795.9	1.4	−0.7	U			LZ1	1.0	11Tu09	
^{53}Ni−u	−31810	27	−31792	6	0.7	U			LZ1	1.0	12Zh34	*
	−31791.6	6.2				2			LZ2	1.0	23Zh10	
^{53}Cr−^{48}Ti$_{1.104}$	−1880.44	0.12	−1880.23	0.09	1.7	1	57	56 ^{53}Cr	MS1	1.0	17Ka53	
^{53}K−^{53}Cr	46126.0	6.5				2			TR1	1.0	26Wa.A	
^{53}Co−^{53}Fe	8897.67	0.49	8897.7	0.5	−0.0	1	94	76 ^{53}Co	JY1	1.0	10Ka26	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{53}Co^{m}−^{53}Fe		12305.2	1.3	12305.4	0.9	0.1	1	51	27 ^{53}Fe	JY1	1.0	10Ka26	
^{53}Co^{m}−^{53}Co		3407.9	1.5	3407.7	1.0	−0.1	1	41	24 ^{53}Co	JY1	1.0	10Ka26	
^{53}Cr−^{52}Cr		115	46	141.57	0.10	0.1	U			R09	4.0	72De11	
^{51}V(t,p)^{53}V		7325	25	7309	3	−0.6	U			Ald		67Hi02	
^{53}Cr(d,α)^{51}V		7635	12	7626.26	0.10	−0.7	U			Kop		67Ha.A	
^{52}Cr(n,γ)^{53}Cr		7939.52	0.3	7939.45	0.10	−0.2	–			MMn		80Is02	Z
		7939.01	0.2			2.2	–			BNn		80Ko01	Z
		7939.10	0.28			1.2	–			Bdn		06Fi.A	
^{52}Cr(d,p)^{53}Cr		5725	6	5714.88	0.10	−1.7	U			MIT		64Sp12	
		5719	8			−0.5	U			Kop		67Ha.A	
^{52}Cr(n,γ)^{53}Cr	ave.	7939.15	0.14	7939.45	0.10	2.1	1	46	26 ^{53}Cr			average	
^{52}Cr(p,γ)^{53}Mn		6559.1	1.1	6559.7	0.3	0.6	U					70Ma25	Z
		6559.72	0.36			−0.0	1	85	84 ^{53}Mn			79Sw01	Z
^{52}Cr(^{3}He,d)^{53}Mn		1070	15	1066.2	0.3	−0.3	U			MIT		67Ob04	
^{53}Co^{m}(p)^{52}Fe		1559.7	40.	1558.8	1.2	−0.0	o					70Ja22	
		1600.5	30.			−1.4	U					70Ce04	
		1590	30			−1.0	U					72Ce01	
		1590	30			−1.0	U					76Vi02	
		1552.3	8.0			0.8	U					15Sh16	*
		1558.9	1.6			−0.1	1	59	59 ^{53}Co^{m}			23Sa59	*
^{53}Co^{i}(p)^{52}Fe		2789.5	50.	2709.5	2.4	−1.6	U					76Vi02	*
		2778.5	18.			−3.8	B			Bor		07Do17	*
^{53}Ti(β^{-})^{53}V		5020	100	4970	4	−0.5	U			ANB		77Pa01	
^{53}V(β^{-})^{53}Cr		3536	50	3436	3	−2.0	U					56Sc.A	*
^{53}Cr(p,n)^{53}Mn		−1379	8	−1379.7	0.3	−0.1	U			MIT		52Lo06	Y
		−1381.1	1.6			0.8	U			Oak		64Jo11	Z
^{53}Cr(^{3}He,t)^{53}Mn^{i}		−7589	4				2					71Be29	*
^{53}Fe(β^{+})^{53}Mn		3860	100	3743.5	1.3	−1.2	U					59Ju40	
		3820	100			−0.8	U					75Bl01	
^{53}Co^{i}(IT)^{53}Co		4325	2				2					16Su10	
*^{53}Ni−u	Same result in reference											13Ya03	**
*^{53}Co^{m}(p)^{52}Fe	Original Q =1558(8) corrected for recoil											16Hu.A	**
*^{53}Co^{m}(p)^{52}Fe	also Q_p=709.5(1.6) keV to 2^{+} level at 849.45(0.10) keV											23Sa59	**
*^{53}Co^{i}(p)^{52}Fe	Q_p=1940(50) 1929(18) respectively, to 2^{+} level at 849.45 keV											Ens159	**
*^{53}V(β^{-})^{53}Cr	$E_{\beta^{-}}$=2530(50) to 5/2^{-} level at 1006.27 keV											Ens09a	**
*^{53}Cr(^{3}He,t)^{53}Mn^{i}	CDE=8350(4) Q =–7586(4); recalibration –3 keV for ^{50}Cr(p,n)^{50}Mn from Ame1961											MMC123	**
^{54}Ca−u		−27011	52	−26967	13	0.9	o			MR1	1.0	13Wi06	
^{54}Sc−u		−36060	500	−36972	19	−1.2	o			TO3	1.5	90Tu01	*
		−37060	500			0.1	o			TO5	1.5	94Se12	*
		−36960	400			−0.0	U			TO6	1.5	98Ba.A	*
		−37059	225			0.3	U			GT1	1.5	04Ma.A	
		−36070	390			−2.3	U			MT1	1.0	11Es06	*
		−37202	585			0.4	o			LZ1	1.0	15Xu14	*
		−36230	680			−1.1	o			MT1	1.0	15Me08	
		−37021	386			0.1	U			LZ1	1.0	19Xu09	
		−36554	258			−1.6	U			MT1	1.0	20Me06	
^{54}Ti−u		−48820	230	−49106	15	−0.8	U			TO3	1.5	90Tu01	
		−49130	250			0.1	U			TO5	1.5	94Se12	
		−48820	280			−0.7	U			TO6	1.5	98Ba.A	
		−48998	118			−0.9	o			LZ1	1.0	15Xu14	
		−49108	17			0.1	1	74	74 ^{54}Ti	TR1	1.0	18Le03	
		−49050	107			−0.5	U			LZ1	1.0	19Xu09	
^{54}V−u		−53574	18	−53568	8	0.3	1	20	20 ^{54}V	TR1	1.0	18Re11	
^{54}Cr−^{39}K$_{1.385}$		−10849.6	5.1	−10856.22	0.14	−1.3	U			TT1	1.0	18Re11	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$C_4\ H_6-{}^{54}Cr$		108018	17	108072.94	0.14	0.8	U			R09	4.0	72De11	
$C_3\ {}^{13}C\ H_5-{}^{54}Cr$		103569	15	103602.74	0.14	0.6	U			R09	4.0	72De11	
$C_3\ H_4\ N-{}^{54}Cr$		95445	13	95496.88	0.14	1.0	U			R09	4.0	72De11	
$C_2\ {}^{13}C\ H_3\ N-{}^{54}Cr$		90960	24	91026.68	0.14	0.7	U			R09	4.0	72De11	
$C_2\ N\ O-{}^{54}Cr$		59057	26	59111.37	0.14	0.5	U			R09	4.0	72De11	
${}^{54}Cr-{}^{85}Rb_{.635}$		−5110.22	0.69	−5109.23	0.14	1.4	U			MA8	1.0	19Hu15	
${}^{54}Cr-u$		−61116.1	4.9	−61122.75	0.14	−1.4	o			TT1	1.0	18Le03	
${}^{13}C\ {}^{37}Cl_3-{}^{54}Fe\ {}^{35}Cl_2$		23744.46	1.26	23748.3	0.5	0.8	U			H39	4.0	84Ha20	
$C_4\ H_6-{}^{54}Fe$		107368	11	107341.3	0.4	−0.6	U			R09	4.0	72De11	
$C_3\ H_4\ N-{}^{54}Fe$		94791	8	94765.2	0.4	−0.8	U			R09	4.0	72De11	
$C_2\ N\ O-{}^{54}Fe$		58411	8	58379.7	0.4	−1.0	U			R09	4.0	72De11	
$C_3\ {}^{13}C\ H_5-{}^{54}Fe$		102908	48	102871.1	0.4	−0.2	U			R09	4.0	72De11	
${}^{54}Ni-u$		−42167	5	−42172.7	2.5	−1.1	2			LZ1	1.0	17Zh12	
		−42174.6	2.9			0.7	2			LZ2	1.0	22Wa39	
		−42174.6	2.9			0.7	o			LZ2	1.0	23Zh10	
${}^{54}Cr-{}^{48}Ti_{1.125}$		−2555.97	0.16	−2555.99	0.12	−0.1	1	52	52 ${}^{54}Cr$	MS1	1.0	17Ka53	
${}^{54}K-{}^{54}Cr$		55311	14				2			TR1	1.0	26Wa.A	
${}^{54}Ca-{}^{54}Cr$		34156	13				2			TR1	1.0	22Po06	
${}^{54}Sc-{}^{54}Cr$		24151	19				2			TR1	1.0	21Le02	
${}^{54}Ti-{}^{54}Cr$		12021	29	12016	15	−0.2	1	26	26 ${}^{54}Ti$	TR1	1.0	22Po06	
${}^{54}V-{}^{54}Cr$		7554	11	7554	8	0.0	1	54	54 ${}^{54}V$	TR1	1.0	22Po06	
${}^{54}Co-{}^{54}Fe$		8850.94	0.14	8850.89	0.10	−0.4	1	47	47 ${}^{54}Co$	JY1	1.0	08Er04	
${}^{54}Co^m-{}^{54}Fe$		9062.960	0.092	9062.99	0.08	0.3	1	81	81 ${}^{54}Co^m$	JY1	1.0	08Er04	
${}^{54}Co^m-{}^{54}Co$		212.18	0.15	212.10	0.10	−0.5	1	49	30 ${}^{54}Co$	JY1	1.0	08Er04	
${}^{54}Cr-{}^{53}Cr$		−1662	48	−1769.00	0.11	−0.6	U			R09	4.0	72De11	
${}^{54}Fe(p,{}^6He){}^{49}Mn$		−28943	24	−28937.1	2.2	0.2	U			MSU		75Mu09	*
${}^{54}Fe(\alpha,{}^8He){}^{50}Fe$		−50950	60	−50962.9	2.8	−0.2	U			Tex		77Tr05	
${}^{54}Cr(p,\alpha){}^{51}V$		130	30	131.69	0.12	0.1	U			Kop		64Ve02	
${}^{54}Fe(p,\alpha){}^{51}Mn$		−3145	9	−3146.7	0.5	−0.2	U			Ald		66Br05	
		−3146.9	1.1			0.2	1	18	11 ${}^{54}Fe$	NDm		74Jo14	
${}^{54}Fe(p,\alpha){}^{51}Mn^i$		−7606.6	5.0	−7597.3	1.6	1.9	U					79Ta22	*
${}^{54}Fe({}^3He,{}^6He){}^{51}Fe$		−18694	15	−18725.7	1.5	−2.1	U			MSU		77Mu03	*
${}^{54}Cr(d,\alpha){}^{52}V$		5225	12	5218.37	0.18	−0.6	U			Kop		67Ha.A	
${}^{52}Cr(t,p){}^{54}Cr$		9171	10	9176.78	0.12	0.6	U			LAl		71Ca19	
${}^{52}Cr({}^3He,p){}^{54}Mn$		7785	15	7781.0	1.0	−0.3	U			MIT		69Ly06	
		7788	9			−0.8	U			Phi		72Be07	
${}^{52}Cr({}^3He,p){}^{54}Mn^i$		1633.6	3.9	1634.7	2.8	0.3	1	51	51 ${}^{54}Mn^i$			72Be07	*
${}^{52}Cr({}^3He,n){}^{54}Fe^j$		−7173	20				2					75Bo14	
${}^{54}Fe(d,\alpha){}^{52}Mn$		5169	12	5168.3	0.4	−0.1	U			Kop		67Ha.A	
		5159	15			0.6	U			MIT		67Sp09	
		5163.3	2.2			2.3	U			NDm		76Jo01	
${}^{54}Fe(p,t){}^{52}Fe$		−15584	8	−15582.7	0.4	0.2	U					78Ko27	*
${}^{54}Fe(p,t){}^{52}Fe^j$		−24139	7	−24140	6	−0.1	2					78Ko27	
		−24141.3	11.0			0.1	2					78De18	*
${}^{54}Zn(2p){}^{52}Ni$		1480	20	1480	20	−0.0	o					05Gi15	
		1480	20				3					05Bl15	
		1280	210			1.0	U					11As08	
		1363	25			4.7	B					25Ku.B	
${}^{54}Cr(d,{}^3He){}^{53}V$		−6879.2	3.1				2			NDm		79Br.B	
${}^{53}Cr(n,\gamma){}^{54}Cr$		9719.30	0.16	9719.13	0.10	−1.1	–					68Wh03	Z
		9718.3	0.4			2.1	–					72Lo26	Z
		9718.91	0.27			0.8	–			MMn		80Is02	Z
		9719.7	0.5			−1.1	–			SAn		89Ho15	Z
		9720.00	0.20			−4.3	C			Bdn		06Fi.A	
${}^{53}Cr(d,p){}^{54}Cr$		7480	12	7494.56	0.10	1.2	U			MIT		64Sp12	
		7514	10			−1.9	U			Kop		67Ha.A	
${}^{53}Cr(n,\gamma){}^{54}Cr$	ave.	9719.14	0.13	9719.13	0.10	−0.1	1	66	48 ${}^{54}Cr$			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{53}Cr(p,γ)^{54}Mn		7559.6	1.0				2					75We10	Z
^{53}Cr(^{3}He,d)^{54}Mn		2080	12	2066.1	1.0	−1.2	U			MIT		69Ly06	
^{54}Fe(d,t)^{53}Fe		−7121.5	2.1	−7121.3	1.2	0.1	–			NDm		74Jo14	
^{54}Fe(^{3}He,α)^{53}Fe		7197	20	7199.1	1.2	0.1	U			MIT		68Tr01	
		7199.6	2.6			−0.2	–			NDm		74Jo14	
^{54}Fe(d,t)^{53}Fe	ave.	−7121.2	1.6	−7121.3	1.2	−0.1	1	58	55 ^{53}Fe			average	
^{54}Ti(β^-)^{54}V		4280	160	4156	16	−0.8	U					96Do23	
^{54}V(β^-)^{54}Cr		7000	100	7037	8	0.4	U					70Wa14	
^{54}Cr(t,^{3}He)^{54}V		−7023	15	−7018	8	0.3	1	25	25 ^{54}V	LAl		77Fl03	
^{54}Mn(ε)^{54}Cr		1359	8	1377.2	1.0	2.3	U					72Ko47	*
		1379	8			−0.2	U					00Hi08	*
^{54}Cr(p,n)^{54}Mn		−2160	5	−2159.5	1.0	0.1	U			MIT		52Lo06	Z
^{54}Cr(^{3}He,t)^{54}Mn^{i}		−7541	4	−7542.1	2.8	−0.3	1	49	49 ^{54}Mn^{i}			71Be29	*
^{54}Co(β^+)^{54}Fe		8023	110	8244.55	0.09	2.0	U					59Su.A	*
		8459	41			−5.2	C					60Mi.A	
^{54}Fe(p,n)^{54}Co		−9031.1	2.5	−9026.89	0.09	1.7	U			Yal		69Ov01	*
		−9023.7	1.8			−1.8	U			Har		74Ho21	Z
^{54}Fe(^{3}He,t)^{54}Co		−8261.2	1.0	−8263.14	0.09	−1.9	F			Mun		77Vo02	*
^{54}Fe(^{3}He,t)^{54}Co−^{27}Al()^{27}Si		−3432.5	3.0	−3432.19	0.13	0.1	U			ChR		74Ha35	
^{54}Fe(^{3}He,t)^{54}Co−^{42}Ca()^{42}Sc		−1817.24	0.18	−1818.21	0.10	−5.4	B			ChR		87Ko34	
^{54}Co^{m}(IT)^{54}Co		197.1	0.4	197.57	0.10	1.2	U					Ens146	
*^{54}Sc−u	Original –36000(500) μu or $M-A$=–33500(470) keV											GAu	**
*^{54}Sc−u	Original –37000(500) μu or $M-A$=–34470(470) keV											GAu	**
*^{54}Sc−u	$M-A$=–34370(370) keV for mixture gs+m at 110.6 keV											Nub269	**
*^{54}Sc−u	$M-A$=–33540(360) keV for mixture gs+m at 110.6 keV											Nub269	**
*^{54}Sc−u	No isomeric correction needed											HWJ151	**
*^{54}Fe(p,^{6}He)^{49}Mn	Q increased 1 for recalibration											AHW	**
*^{54}Fe(p,α)^{51}Mn^{i}	IT=4459(5); Q rebuilt with Ame1977											MMC124	**
*^{54}Fe(^{3}He,^{6}He)^{51}Fe	Averaged with reference See ^{46}Ti(^{3}He,^{6}He)											75Mu09	**
*^{52}Cr(^{3}He,p)^{54}Mn^{i}	IT=6151(5); Q rebuilt with Ame1971											MMC124	**
*^{54}Fe(p,t)^{52}Fe	Q =–21239(8) to ^{52}Fe^{i} at 5654.5											Nub269	**
*^{54}Fe(p,t)^{52}Fe^{j}	IT=8561(5); Q rebuilt with Ame1977											MMC124	**
*^{54}Mn(ε)^{54}Cr	IBE=518(8) to 2$^+$ level at 834.855 keV, B(K)=5.99											Ens148	**
*^{54}Mn(ε)^{54}Cr	IBE=544(8) to 2$^+$ level at 834.855 keV											Ens148	**
*^{54}Cr(^{3}He,t)^{54}Mn^{i}	CDE=8302(4) Q =–7538(4); recalibration –3 keV for ^{50}Cr(p,n)^{50}Mn from Ame1961											MMC123	**
*^{54}Co(β^+)^{54}Fe	E_{β^+}=4250(110) from ^{54}Co^{m} at 197.57 to 2949.2 6$^+$ level											Nub269	**
*^{54}Fe(p,n)^{54}Co	Uncorrected for resonance. Orig T=9204.1(1.8) corrected in reference											76Fr13	**
*^{54}Fe(^{3}He,t)^{54}Co	Original value –8260.2(0.6) recalibrated											AHW	**
*^{54}Fe(^{3}He,t)^{54}Co	F : rejected in reference of same group											09Fa15	**
^{55}Ca−u		−20022	172				2			RT1	1.0	18Mi08	
^{55}Sc−u		−30600	1100	−33101	23	−1.5	U			TO3	1.5	90Tu01	
		−32100	600			−1.1	U			TO6	1.5	98Ba.A	
		−32460	640			−1.0	o			MT1	1.0	11Es06	
		−32760	620			−0.5	o			MT1	1.0	15Me08	
		−33376	236			1.2	U			MT1	1.0	20Me06	
		−32861	204			−1.2	U			RT1	1.0	20Mi13	
^{55}Ti−u		−44650	280	−44902	6	−0.6	U			TO3	1.5	90Tu01	
		−44880	260			−0.1	U			TO5	1.5	94Se12	
		−44360	350			−1.0	U			TO6	1.5	98Ba.A	
		−44909	31			0.2	U			TR1	1.0	18Le03	
		−44903.1	6.1			0.1	1	93	93 ^{55}Ti	MS1	1.0	22Po06	
^{55}V−u		−52738	29	−52752	7	−0.5	U			TR1	1.0	18Re11	
^{55}Cr−^{85}Rb$_{.647}$		−2093.4	1.9	−2091.43	0.25	1.0	U			MA8	1.0	19Hu15	
C_4 H_7−^{55}Mn		116757	8	116731.4	0.3	−0.8	U			R09	4.0	72De11	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
C_3 ^{13}C $H_6-^{55}Mn$		112281	25	112261.2	0.3	−0.2	U				R09	4.0	72De11	
C_3 H_5 $N-^{55}Mn$		104202	10	104155.4	0.3	−1.2	U				R09	4.0	72De11	
C_2 H_3 $N_2-^{55}Mn$		91618	28	91579.3	0.3	−0.3	U				R09	4.0	72De11	
C_3 H_3 $O-^{55}Mn$		80372	10	80345.9	0.3	−0.7	U				R09	4.0	72De11	
$^{55}Mn-^{85}Rb_{.647}$		−4884.41	0.84	−4884.2	0.3	0.3	o				MA8	1.0	12Na15	
		−4883.81	0.80			−0.4	1	19	19	^{55}Mn	MA8	1.0	19Hu15	
$^{55}Ni-u$		−48678	18	−48669.5	0.8	0.5	U				LZ1	1.0	11Tu09	
$^{55}Cu-u$		−33962	167	−34146	27	−1.1	U				LZ1	1.0	13Ya03	
		−34146.2	26.6				2				LZ2	1.0	23Zh10	
$^{55}Sc-^{55}Ti$		11803	24	11802	23	−0.1	1	88	88	^{55}Sc	RI1	1.0	23Ii01	
$^{55}Sc-^{55}Cr$		26053	66	26063	23	0.1	1	12	12	^{55}Sc	TR1	1.0	21Le02	
$^{55}Ti-^{55}Cr$		14273	24	14261	6	−0.5	1	6	6	^{55}Ti	TR1	1.0	22Po06	
$^{55}V-^{55}Ti$		−7850	98	−7850	9	0.0	U				RI1	1.0	23Ii01	
$^{55}V-^{55}Cr$		6411.4	7.1				2				TR1	1.0	22Po06	
$^{55}Ni-^{55}Co$		9333.43	0.62				2				JY1	1.0	10Ka26	
$^{55}Mn(p,\alpha)^{52}Cr$		2570	8	2571.8	0.3	0.2	U				MIT		64Sp12	
		2600	10			−2.8	U				ANL		67Ka11	
$^{55}Mn(d,\alpha)^{53}Cr$		8283	8	8286.6	0.3	0.5	U				MIT		64Sp12	
		8277	15			0.6	U				Kop		67Ha.A	
$^{54}Cr(n,\gamma)^{55}Cr$		6246.2	0.4	6246.26	0.19	0.2	–						72Wh05	Z
		6246.28	0.21			−0.1	–				Bdn		06Fi.A	
$^{54}Cr(d,p)^{55}Cr$		4027	8	4021.70	0.19	−0.7	U				MIT		64Sp12	
		4035	8			−1.7	U				Kop		67Ha.A	
		4022.1	1.2			−0.3	U				NDm		74Jo14	
$^{54}Cr(n,\gamma)^{55}Cr$	ave.	6246.26	0.19	6246.26	0.19	0.0	1	100	100	^{55}Cr			average	
$^{54}Cr(p,\gamma)^{55}Mn$		8067.2	0.4	8065.3	0.3	−4.7	B						78We12	
$^{54}Cr(^3He,d)^{55}Mn$		2568	18	2571.9	0.3	0.2	U				MIT		69Ra02	
$^{55}Mn(\gamma,n)^{54}Mn$		−10192	20	−10224.9	1.1	−1.6	U				Phi		60Ge01	
$^{54}Fe(n,\gamma)^{55}Fe$		9297.91	0.3	9298.09	0.19	0.6	–				MMn		80Is02	Z
		9298.53	0.27			−1.6	–				Bdn		06Fi.A	
$^{54}Fe(d,p)^{55}Fe$		7084	8	7073.53	0.19	−1.3	U				MIT		64Sp12	
		7083	10			−0.9	U				Kop		67Ha.A	
		7072.3	1.7			0.7	U				NDm		74Jo14	
$^{54}Fe(n,\gamma)^{55}Fe$	ave.	9298.25	0.20	9298.09	0.19	−0.8	1	89	68	^{54}Fe			average	
$^{54}Fe(p,\gamma)^{55}Co$		5064.0	0.7	5064.38	0.30	0.5	–						77Er02	Z
		5063.9	0.4			1.2	–						80Ha36	Z
$^{54}Fe(^3He,d)^{55}Co$		−428	15	−429.09	0.30	−0.1	U				MIT		67Ob04	
		−426.9	2.2			−1.0	U				NDm		74Jo14	
$^{54}Fe(p,\gamma)^{55}Co$	ave.	5063.9	0.3	5064.38	0.30	1.3	1	74	55	^{55}Co			average	
$^{55}Ti(\beta^-)^{55}V$		7440	200	7312	9	−0.6	U						96Do23	
$^{55}V(\beta^-)^{55}Cr$		5956	100	5972	7	0.2	U				ANB		77Na17	
$^{55}Cr(\beta^-)^{55}Mn$		2500	40	2601.4	0.4	2.5	U						63Me06	
		2494	25			4.3	B						65Ko09	
$^{55}Fe(\varepsilon)^{55}Mn$		224.5	4.	231.09	0.18	1.6	U						65Be19	
		224.5	3.			2.2	U						69Ka13	
		231.4	0.4			−0.8	–						89Zl.A	
		230.7	1.9			0.2	U						90Is06	
		231.0	1.0			0.1	U						93Wi05	*
		231.37	0.30			−0.9	–						95Da14	*
		231.0	0.3			0.3	–						95Sy01	*
		232.36	0.64			−2.0	U						01Ke14	
$^{55}Mn(p,n)^{55}Fe$		−1015.7	2.	−1013.44	0.18	1.1	U				Nvl		59Go68	Z
		−1014.6	0.8			1.4	U				Oak		64Jo11	Z
$^{55}Fe(\varepsilon)^{55}Mn$	ave.	231.23	0.19	231.09	0.18	−0.7	1	91	78	^{55}Fe			average	
$^{55}Mn(^3He,t)^{55}Fe^i$		−7883	6				2						71Be29	*
$^{55}Co(\beta^+)^{55}Fe$		3466	2	3451.4	0.3	−7.3	B						66Fi06	*
$*^{55}Fe(\varepsilon)^{55}Mn$	Error estimated by evaluator												AHW	**
$*^{55}Fe(\varepsilon)^{55}Mn$	Original error 0.10 increased by evaluator												GAu	**
$*^{55}Fe(\varepsilon)^{55}Mn$	Original statistical error 0.10 increased by evaluator												GAu	**
$*^{55}Mn(^3He,t)^{55}Fe^i$	CDE=8654(6) Q =–7890(6); recalibration +7 keV for $^{54}Fe(p,n)^{54}Co$ from Ame1961												MMC123	**
$*^{55}Co(\beta^+)^{55}Fe$	E_{β^+}=1513(2) to $5/2^-$ level at 931.29 keV												Ens097	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{56}Ca−u	−14504	268				2			RT1	1.0	18Mi08	
^{56}Sc−u	−26680	630	−27390	280	−1.1	o			MT1	1.0	15Me08	*
	−27520	350			0.4	2			MT1	1.0	20Me06	*
	−27171	461			−0.5	2			RT1	1.0	20Mi13	*
^{56}Ti−u	−41300	350	−42304	7	−1.9	U			TO3	1.5	90Tu01	
	−42010	300			−0.7	U			TO5	1.5	94Se12	
	−41770	270			−1.3	U			TO6	1.5	98Ba.A	
	−42319	129			0.1	U			GT1	1.5	04Ma.A	
	−42700	290			1.4	o			LZ1	1.0	15Xu14	
	−42738	203			2.1	U			LZ1	1.0	19Xu09	
	−42384	258			0.3	U			MT1	1.0	20Me06	
^{56}Ti−N_4	−54602.5	7.9	−54600	7	0.3	1	89	89 ^{56}Ti	RI1	1.0	23Ii01	
^{56}V−u	−49470	250	−49663	6	−0.5	U			TO3	1.5	90Tu01	
	−49640	260			−0.1	U			TO5	1.5	94Se12	
	−49310	250			−0.9	U			TO6	1.5	98Ba.A	
^{56}V−N_4	−61957.7	6.6	−61959	6	−0.2	1	84	84 ^{56}V	RI1	1.0	23Ii01	
^{56}V−u	−49596	39	−49663	6	−1.7	U			MS1	1.0	22Po06	
^{56}Cr−^{85}Rb$_{.659}$	−1216.3	2.0	−1219.8	0.6	−1.8	U			MA8	1.0	05Gu27	
^{56}Cr−N_4	−71657.7	5.7	−71646.4	0.6	2.0	U			RI1	1.0	23Ii01	
^{56}Mn−^{85}Rb$_{.659}$	−2965.1	1.5	−2965.9	0.4	−0.5	U			MA8	1.0	05Gu37	
^{56}Mn−^{39}K$_{1.436}$	−8979.0	2.7	−8978.9	0.4	0.0	U			MA8	1.0	09Na.A	
C_4 H_8−^{56}Fe	127754	10	127664.1	0.3	−2.2	U			R09	4.0	72De11	
C_3 ^{13}C H_7−^{56}Fe	123300	47	123193.9	0.3	−0.6	U			R09	4.0	72De11	
C_3 H_6 N−^{56}Fe	115171	13	115088.0	0.3	−1.6	U			R09	4.0	72De11	
C_3 H_4 O−^{56}Fe	91381	15	91278.6	0.3	−1.7	U			R09	4.0	72De11	
C_2 H_2 N O−^{56}Fe	78790	24	78702.5	0.3	−0.9	U			R09	4.0	72De11	
C_2 O_2−^{56}Fe	54990	9	54893.1	0.3	−2.7	B			R09	4.0	72De11	
^{56}Fe−^{85}Rb$_{.659}$	−6933.40	0.60	−6933.3	0.3	0.2	1	31	31 ^{56}Fe	MA8	1.0	19Hu15	
^{56}Cu−u	−41485	16	−41467	5	1.1	2			LZ1	1.0	18Zh29	
	−41467.5	7.6			0.1	2			MS1	1.0	18Va01	*
	−41463.1	6.4			−0.6	2			LZ2	1.0	22Wa39	
	−41463.1	6.4			−0.6	o			LZ2	1.0	23Zh10	
^{56}Fe−^{58}Ni$_{.966}$	−2604.70	0.47	−2604.44	0.26	0.6	1	31	25 ^{58}Ni	JY1	1.0	10Ka26	
^{56}Co−^{58}Ni$_{.966}$	2297.85	0.55	2298.0	0.4	0.3	1	58	51 ^{56}Co	JY1	1.0	10Ka26	
^{56}Ni−^{55}Co$_{1.018}$	1176.23	0.48	1175.4	0.4	−1.8	1	60	33 ^{55}Co	JY1	1.0	10Ka26	
^{56}Ti−^{56}Cr	17064	22	17046	7	−0.8	1	11	11 ^{56}Ti	TR1	1.0	22Po06	
^{56}V−^{56}Cr	9680	15	9687	6	0.5	1	16	16 ^{56}V	TR1	1.0	22Po06	
^{56}Cr−^{56}Fe	5713.44	0.55	5713.4	0.5	−0.0	1	100	100 ^{56}Cr	MA8	1.0	19Hu15	
^{56}Ni−^{56}Fe	7192.00	0.52	7192.2	0.3	0.4	1	42	40 ^{56}Ni	JY1	1.0	10Ka26	
^{56}Ni−^{56}Co	2289.61	0.49	2289.8	0.4	0.3	1	67	49 ^{56}Co	JY1	1.0	10Ka26	
^{56}Fe−^{54}Fe	−4755	47	−4672.71	0.30	0.4	U			R09	4.0	72De11	
^{56}Fe(p,α)^{53}Mn	−1060	9	−1052.2	0.4	0.9	U			MIT		64Sp12	
	−1056	9			0.4	U			Ald		66Br05	
	−1052.3	0.8			0.1	1	28	16 ^{53}Mn	NDm		74Jo14	
^{54}Cr(t,p)^{56}Cr	5995	30	6009.9	0.6	0.5	U			Ald		68Ch20	
	6024	10			−1.4	U			LAl		71Ca19	
^{56}Fe(d,α)^{54}Mn	5662	12	5662.5	1.1	0.0	U			Kop		67Ha.A	
	5673	30			−0.3	U					67Hj01	
^{54}Fe(^{3}He,p)^{56}Co	7410	10	7428.2	0.5	1.8	U			CIT		67Mi02	
	7408	15			1.3	U			MIT		68Be10	
^{54}Fe(^{3}He,n)^{56}Ni	4513	14	4513.0	0.4	−0.0	U			CIT		67Mi02	
^{55}Mn(n,γ)^{56}Mn	7270.53	0.3	7270.44	0.13	−0.3	2			MMn		80Is02	Z
	7270.42	0.15			0.1	2			Bdn		06Fi.A	
^{55}Mn(d,p)^{56}Mn	5052	5	5045.88	0.13	−1.2	U			MIT		64Sp12	
	5053	8			−0.9	U			Kop		67Ha.A	
^{55}Mn(p,γ)^{56}Fe	10189	7	10183.70	0.16	−0.8	U					69Fr22	
	10193.7	4.5			−2.2	U					70Sa19	*
	10195.7	3.6			−3.3	B					74Pe15	*
	10183.80	0.17			−0.6	1	89	57 ^{55}Mn	Utr		92Gu03	Z

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{56}Fe(d,t)^{55}Fe	−4938.3	1.3	−4939.91	0.23	−1.2	U			NDm		74Jo14	
^{56}Ni(p)^{55}Co	−7148.5	30.	−7166.6	0.3	−0.6	U					08Jo04	*
^{56}Cu^{i}(p)^{55}Ni	2929	31	2948	10	0.6	U			Bor		07Do17	*
	2948	10				3					14Or04	*
^{56}Ti(β^-)^{56}V	7030	330	6855	9	−0.5	U					96Do23	
^{56}Cr(β^-)^{56}Mn	1610	150	1626.4	0.6	0.1	U					60Dr03	
^{56}Mn(β^-)^{56}Fe	3685	5	3695.61	0.21	2.1	U					62Ho14	*
^{56}Co(β^+)^{56}Fe	4566.0	2.0	4566.6	0.4	0.3	U					65Pe18	*
^{56}Fe(p,n)^{56}Co	−5351	10	−5349.0	0.4	0.2	U			Tal		62Ne08	Y
^{56}Fe(^{3}He,t)^{56}Co^{i}	−8178	9				2					71Be29	*
*^{56}Sc−u	$M-A$=–24850(590) keV for mixture gs+m at 0#(100#) keV										15Me08	**
*^{56}Sc−u	$M-A$=–25380(260) keV for mixture gs+m at 0#(100#) keV and 775.0(0.1) keV										Nub269	**
*^{56}Sc−u	Isomer mixture corrected by 19(3) for 774.9(0.3) keV, 0(390) for										20Mi13	**
*^{56}Sc−u	0#(100#) keV										20Mi13	**
*^{56}Cu−u	represents frequency ratio $^{56}Cu^+/(C_4H_7)^+$ =1.01641577(12)										HWJ192	**
*^{55}Mn(p,γ)^{56}Fe	E_p=1537(2) to 11703(4) level										70Sa19	**
*^{55}Mn(p,γ)^{56}Fe	E_p=1537(2) to 11705(3) level										74Pe15	**
*^{56}Ni(p)^{55}Co	E_p=2540(30) from 9735 level										08Jo04	**
*^{56}Cu^{i}(p)^{55}Ni	Strongest fragment (strength 2.7); weaker fragment 85(70) keV lower (strength 1.3)										14Or04	**
*^{56}Mn(β^-)^{56}Fe	E_{β^-}=2838(5) to 2^+ level at 846.7778 keV										Ens115	**
*^{56}Co(β^+)^{56}Fe	E_{β^+}=1459(3) to 4^+ level at 2085.1045 keV										Ens115	**
*^{56}Fe(^{3}He,t)^{56}Co^{i}	Strongest fragment given as CDE=8950(6); Q =–8186(6) rebuilt with Ame1965										71Be29	**
*	recalibration +7 keV for ^{54}Fe(p,n)^{54}Co from Ame1961										MMC123	**
^{57}Ca−u	−7912	1063	−7040#	430#	0.8	D			RT1	1.0	18Mi08	*
^{57}Sc−u	−22540	1400	−22950	190	−0.3	o			MT1	1.0	15Me08	
	−21664	944			−1.4	U			MT1	1.0	20Me06	
	−22952	193				2			RT1	1.0	20Mi13	
^{57}Ti−u	−35700	1000	−36930	220	−0.8	o			TO3	1.5	90Tu01	
	−36200	400			−1.2	2			TO6	1.5	98Ba.A	
	−37102	408			0.3	2			GT1	1.5	04Ma.A	
	−36280	370			−1.8	o			MT1	1.0	11Es06	
	−37037	258			0.4	2			MT1	1.0	20Me06	
^{57}V−^{40}Ar O H	−12770	16	−12767	7	0.2	1	20	20 ^{57}V	RI1	1.0	23Ii01	
^{57}V−u	−47300	400	−47644	7	−0.6	U			TO3	1.5	90Tu01	
	−47640	270			−0.0	U			TO5	1.5	94Se12	
	−47320	250			−0.9	U			TO6	1.5	98Ba.A	
	−47703	91			0.6	U			LZ1	1.0	15Xu14	
	−47934	279			1.0	U			MT1	1.0	20Me06	
	−47634	16			−0.6	1	20	20 ^{57}V	MS1	1.0	22Po06	
^{57}Cr−u	−56240	250	−56387.8	2.0	−0.4	U			TO3	1.5	90Tu01	
	−56300	260			−0.2	U			TO5	1.5	94Se12	
	−56170	270			−0.5	U			TO6	1.5	98Ba.A	
^{57}Cr−^{85}Rb$_{.671}$	2802.1	2.0	2801.3	2.0	−0.4	o			MA8	1.0	05Gu27	
	2801.2	2.0			0.1	1	98	98 ^{57}Cr	MA8	1.0	19Hu15	
^{57}Cr−^{40}Ar O H	−21523	13	−21510.5	2.0	1.0	U			RI1	1.0	23Ii01	
^{57}Mn−^{85}Rb$_{.671}$	−2525.1	2.3	−2524.8	1.6	0.1	1	49	49 ^{57}Mn	MA8	1.0	05Gu37	
^{57}Mn−^{39}K$_{1.462}$	−8650.7	2.8	−8652.8	1.6	−0.7	1	33	33 ^{57}Mn	MA8	1.0	12Na15	
^{57}Mn−^{57}Ar O H	−26848	16	*			U			RI1	1.0	23Ii01	
C_7 H_8−^{57}Fe ^{35}Cl	158378.5	3.5	158355.0	0.3	−2.7	U			M18	2.5	68Hu05	
C_4 H_9−^{57}Fe	135085	11	135032.7	0.3	−1.2	U			R09	4.0	72De11	
C_3 H_7 N−^{57}Fe	122500	10	122456.6	0.3	−1.1	U			R09	4.0	72De11	
C_3 H_5 O−^{57}Fe	98684	8	98647.2	0.3	−1.2	U			R09	4.0	72De11	
C_2 H_3 N O−^{57}Fe	86104	17	86071.1	0.3	−0.5	U			R09	4.0	72De11	
^{57}Ni−^{85}Rb$_{.671}$	−1019.8	2.7	−1018.9	0.6	0.3	U			MA8	1.0	07Gu09	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{57}Cu−u		−50772	43	−50787.7	0.6	−0.4	U			LZ1	1.0	11Tu09	
^{56}Fe ^{13}C−^{57}Fe C		2897.67	0.47	2898.422	0.018	0.6	U			H30	2.5	77Ba10	
		2897.68	0.40			0.7	U			H30	2.5	77Ba10	
^{56}Fe H−^{57}Fe		7325	7	7368.619	0.018	1.6	U			R09	4.0	72De11	
^{57}Fe−^{56}Fe$_{1.018}$		1627.95	0.46	1627.563	0.019	−0.8	U			JY1	1.0	10Ka26	
^{57}Fe−^{58}Ni$_{.983}$		−1048.75	0.46	−1048.85	0.27	−0.2	1	33	28 ^{58}Ni	JY1	1.0	10Ka26	
^{57}Ni−^{58}Ni$_{.983}$		3350.77	0.72	3350.6	0.5	−0.3	1	55	50 ^{57}Ni	JY1	1.0	10Ka26	
^{57}Cu−^{56}Ni$_{1.018}$		8126.29	0.55	8125.6	0.4	−1.2	1	63	48 ^{57}Cu	JY1	1.0	10Ka26	
^{57}V−^{57}Cr		8741.4	9.2	8744	7	0.2	1	61	59 ^{57}V	TR1	1.0	22Po06	
^{57}Cu−^{57}Fe		13817.80	0.86	13819.7	0.5	2.2	1	29	29 ^{57}Cu	JY1	1.0	10Ka26	
^{57}Cu−^{57}Ni		9420.42	0.55	9420.3	0.5	−0.2	1	74	50 ^{57}Ni	JY1	1.0	10Ka26	
^{56}Fe ^{37}Cl−^{57}Fe ^{35}Cl		−3413.7	4.3	−3406.54	0.07	0.7	U			M18	2.5	68Hu05	
^{57}Fe−^{56}Fe		456.6	1.4	456.413	0.018	−0.1	U			M18	2.5	68Hu05	
		453.2	2.1			0.6	U			M18	2.5	68Hu05	
		491	39			−0.2	U			R09	4.0	72De11	
^{54}Cr(α,p)^{57}Mn		−4308	8	−4313.4	1.5	−0.7	U			NDm		76Ma03	
		−4302	8			−1.4	U			Can		78An10	
^{57}Fe(p,α)^{54}Mn		237	9	240.9	1.1	0.4	U			MIT		64Sp12	
^{54}Fe(α,p)^{57}Co		−1770.3	1.8	−1772.9	0.5	−1.5	U			NDm		74Jo14	
^{55}Mn(t,p)^{57}Mn		7438.2	3.6	7435.2	1.5	−0.8	1	18	17 ^{57}Mn	NDm		77Ma12	
^{57}Fe(d,α)^{55}Mn		8246	15	8241.22	0.16	−0.3	U			Kop		67Ha.A	
^{56}Fe(n,γ)^{57}Fe		7645.9	0.5	7646.172	0.017	0.5	U			Utr		68Sp01	
		7646.10	0.17			0.4	o			BNn		76Al16	Z
		7645.96	0.20			1.1	U			BNn		78St25	Z
		7646.13	0.21			0.2	U			MMn		80Is02	Z
		7645.93	0.15			1.6	U			Ptn		80Ve05	Z
		7646.0956	0.0500			1.5	–			PTc		97Ro26	*
		7646.08	0.09			1.0	U					02Bo11	
		7646.10	0.15			0.5	o			Bdn		06Fi.A	
		7646.183	0.018			−0.6	–			Bdn		17Fi03	
^{56}Fe(d,p)^{57}Fe		5425	8	5421.606	0.017	−0.4	U			MIT		64Sp12	
		5425	8			−0.4	U			Kop		67Ha.A	
		5419.8	1.3			1.4	U			NDm		74Jo14	
^{56}Fe(n,γ)^{57}Fe	ave.	7646.173	0.017	7646.172	0.017	−0.1	1	100	85 ^{57}Fe			average	
^{56}Fe(p,γ)^{57}Co		6027.7	1.0	6027.5	0.4	−0.2	–					70Ob02	Z
		6029.3	1.5			−1.2	–					71Le21	Z
^{56}Fe(^{3}He,d)^{57}Co		538	20	534.0	0.4	−0.2	U			LAl		65Bl13	
^{56}Fe(p,γ)^{57}Co	ave.	6028.2	0.8	6027.5	0.4	−0.9	1	29	28 ^{57}Co			average	
^{56}Fe(p,γ)^{57}Co^i		−1226.4	0.5	−1225.9	0.3	0.9	2					70Ob02	*
		−1225.6	0.4			−0.6	2					71Le21	*
^{57}Cu^i(p)^{56}Ni		4650	50	4609	25	−0.8	2					76Vi02	
		4568	10			4.1	C					98Jo.A	
		4595	29			0.5	2					02Jo09	
^{57}Ti(β^-)^{57}V		11020	950	9980	210	−1.1	U					96Do23	
^{57}Cr(β^-)^{57}Mn		5100	100	4961.3	2.4	−1.4	U			ANB		78Da04	
^{57}Mn(β^-)^{57}Fe		2690	50	2695.3	1.5	0.1	U					63Va37	
^{57}Co(ε)^{57}Fe		810	30	836.3	0.4	0.9	U					71La02	*
^{57}Fe(p,n)^{57}Co		−1619.4	2.0	−1618.7	0.4	0.3	–			Oak		64Jo11	Z
		−1618.2	2.0			−0.2	–			Can		70Kn03	
	ave.	−1618.8	1.4			0.1	1	10	10 ^{57}Co			average	
^{57}Fe(^{3}He,t)^{57}Co^i		−8122	7	−8108.3	0.3	2.0	U					71Be29	*
^{57}Ni(β^+)^{57}Co		3245	10	3261.7	0.6	1.7	U					50Fr10	*
		3235	10			2.7	U					51Ca28	*
		3246	10			1.6	U					58Ko60	*
^{57}Cu(β^+)^{57}Ni		8742	130	8775.0	0.4	0.3	U					84Sh28	
*^{57}Ca−u	Trends in Mass Surface TMS suggest ^{57}Ca 810 keV less bound											HWJ266	**
*^{56}Fe(n,γ)^{57}Fe	Original error 0.0005 increased for calibration											GAu	**
*^{56}Fe(p,γ)^{57}Co^i	T=1247.9 recalibrated to T=1248.5(0.6) keV											AHW	**
*^{56}Fe(p,γ)^{57}Co^i	T=1247.1 recalibrated to T=1247.7(0.4) keV											AHW	**
*^{57}Co(ε)^{57}Fe	IBE=674(30) to 5/2$^-$ level at 136.4743 keV											Ens98c	**
*^{57}Fe(^{3}He,t)^{57}Co^i	CDE=8893(7) Q =−8129(7); recalibration +7 keV for ^{54}Fe(p,n)^{54}Co from Ame1961											MMC123	**
*^{57}Ni(β^+)^{57}Co	E_{β^+}=845(10) 835(10) 849(10) respectively, to 13/2$^-$ level at 1377.663 keV											Ens98c	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{58}Sc−u		−16618	204				2				RT1	1.0	20Mi13	*
^{58}Ti−u		−33162	268	−33755	5	−2.2	U				MT1	1.0	20Me06	
		−33226	290			−1.8	U				RT1	1.0	20Mi13	
^{58}V−u		−43210	280	−43270	7	−0.1	U				TO3	1.5	90Tu01	
		−43350	280			0.2	U				TO5	1.5	94Se12	
		−42700	400			−1.0	U				TO6	1.5	98Ba.A	
		−43328	107			0.4	U				GT1	1.5	04Ma.A	
		−43457	134			1.4	U				LZ1	1.0	15Xu14	
		−42641	247			−2.5	B				MT1	1.0	20Me06	
^{58}Cr−u		−55680	230	−55819.2	2.6	−0.4	U				TO3	1.5	90Tu01	
		−55750	260			−0.2	U				TO5	1.5	94Se12	
		−55490	270			−0.8	U				TO6	1.5	98Ba.A	
^{58}Cr−^{85}Rb$_{.682}$		4343.9	3.2	4340.2	2.6	−1.2	1	65	65	^{58}Cr	MA8	1.0	18Mo14	
^{58}Mn−^{39}K$_{1.487}$		−5964.9	2.9	−5965.2	2.7	−0.1	1	88	88	^{58}Mn	MA8	1.0	12Na15	*
C_3 H_8 N−^{58}Fe		132382	12	132400.1	0.4	0.4	U				R09	4.0	72De11	
C_3 H_6 O−^{58}Fe		108576	13	108590.6	0.4	0.3	U				R09	4.0	72De11	
C_2 H_4 N O−^{58}Fe		95999	13	96014.6	0.4	0.3	U				R09	4.0	72De11	
C_3 H_6 O−^{58}Ni		106491	8	106522.6	0.4	1.0	U				R10	4.0	74De22	
C_3 ^{13}C H_9−^{58}Ni		138424	14	138437.9	0.4	0.2	U				R10	4.0	74De22	
C_3 H_8 N−^{58}Ni		130302	25	130332.0	0.4	0.3	U				R10	4.0	74De22	
C_2 H_4 O N−^{58}Ni		93926	10	93946.5	0.4	0.5	U				R10	4.0	74De22	
		93928	15			0.3	U				R10	4.0	74De22	
C_3 H_6 O−^{58}Ni		106504	14	106522.6	0.4	0.3	U				R10	4.0	74De22	
^{58}Ni O H_2−^{152}Pr$_{.500}$		−19850.3	2.2	−19861.4	1.5	−5.0	B				RI1	1.0	24Ki16	*
^{58}Zn−u		−45355.1	38.6	−45370	30	−0.5	1	66	66	^{58}Zn	LZ2	1.0	23Wa10	
^{58}V−^{58}Fe		23397	48	23455	7	1.2	U				TR1	1.0	22Po06	
^{58}Ni−^{58}Fe		2059	32	2068.1	0.3	0.1	U				R09	4.0	72De11	
^{58}Cu−^{58}Ni		9190.61	0.50	9190.6	0.5	0.0	1	90	90	^{58}Cu	JY1	1.0	10Ka26	
^{58}Ti H−^{19}F O		−19247.5	4.7	−19248	5	−0.0	1	100	100	^{58}Ti	RI1	1.0	23Ii01	
^{58}V H−^{19}F O		−28761.8	9.9	−28763	7	−0.1	1	48	48	^{58}V	RI1	1.0	23Ii01	
^{58}V−H_2 ^{32}S		−30992.7	9.5	−30992	7	0.1	1	52	52	^{58}V	RI1	1.0	23Ii01	
^{58}Cr H−^{19}F O		−41309.4	8.3	−41312.0	2.6	−0.3	1	10	10	^{58}Cr	RI1	1.0	23Ii01	
^{58}Cr−H_2 ^{32}S		−43551.3	5.2	−43540.5	2.6	2.1	1	25	25	^{58}Cr	RI1	1.0	23Ii01	
^{58}Mn−H_2 ^{32}S		−47657.	8.0	−47654.9	2.7	0.3	1	12	12	^{58}Mn	RI1	1.0	23Ii01	
^{58}Ni(p,^{6}He)^{53}Co		−27889	18	−27872.8	1.3	0.9	U				MSU		75Mu09	*
^{58}Ni(α,^{8}He)^{54}Ni		−50190	50	−50129.5	2.4	1.2	U				Tex		77Tr05	
^{58}Fe(p,α)^{55}Mn		420	9	421.20	0.24	0.1	U				MIT		64Sp12	
^{58}Ni(p,α)^{55}Co		−1341.0	2.9	−1334.9	0.4	2.1	U				BNL		73Go19	
		−1335.1	0.9			0.2	1	18	12	^{55}Co	NDm		74Jo14	
^{58}Ni(^{3}He,^{6}He)^{55}Ni		−17556	11	−17553.8	0.7	0.2	U				MSU		77Mu03	*
^{58}Fe(d,α)^{56}Mn		5470	12	5467.07	0.28	−0.2	U				Kop		67Ha.A	
^{56}Fe(^{3}He,p)^{58}Co		6853	15	6882.4	1.1	2.0	U				MIT		72Ly01	
^{58}Ni(d,α)^{56}Co		6522	12	6522.4	0.4	0.0	U				Kop		67Ha.A	
		6506	10			1.6	U				MIT		68Be10	
^{58}Ni(p,t)^{56}Ni		−13987	18	−13982.1	0.3	0.3	U				Bld		65Ho07	
^{58}Ni(p,t)^{56}Ni^{j}		−23926	4				2						84Ka07	*
^{57}Fe(n,γ)^{58}Fe		10044.60	0.3	10044.59	0.18	−0.0	–				MMn		80Is02	Z
		10044.65	0.24			−0.2	–				Bdn		06Fi.A	
^{57}Fe(d,p)^{58}Fe		7815	8	7820.03	0.18	0.6	U				MIT		64Sp12	
		7824	12			−0.3	U				Kop		67Ha.A	
^{57}Fe(n,γ)^{58}Fe	ave.	10044.63	0.19	10044.59	0.18	−0.2	1	93	84	^{58}Fe			average	
^{57}Fe(p,γ)^{58}Co		6952	3	6954.3	1.1	0.8	1	14	14	^{58}Co			70Er03	
^{58}Ni(p,d)^{57}Ni		−9971.2	7.	−9991.7	0.5	−2.9	U						79Ik04	*
^{58}Ni(^{3}He,α)^{57}Ni		8360.3	4.	8361.4	0.5	0.3	U				MSU		76Na23	
		8384.8	15.			−1.6	U						79Fo09	*
^{58}Ni(^{7}Li,^{8}He)^{57}Cu		−29564	50	−29622.5	0.5	−1.2	U				MSU		85Sh03	
		−29613	17			−0.6	U				Tex		86Ga19	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{58}Ni(^{14}N,^{15}C)^{57}Cu	−19900	40	−19929.6	0.9	−0.7	U			Ber		87St04	
^{58}Mn(β^-)^{58}Fe	5890	100	6326.9	2.6	4.4	B					69Wa10	*
	5958	100			3.7	B					71Dy01	*
^{58}Fe(t,^{3}He)^{58}Mn	−6318	15	−6308.3	2.6	0.6	U			LAl		77Fl03	*
^{58}Co(β^+)^{58}Fe	2305	6	2308.0	1.1	0.5	U					52Ch31	*
	2307	4			0.2	U					63Rh02	*
^{58}Fe(^{3}He,t)^{58}Co^i	−8079	8				2					71Be29	*
^{58}Ni(p,n)^{58}Cu	−9351	5	−9343.4	0.4	1.5	U			Mar		64Ma.A	
	−9352.6	3.4			2.7	U			Ric		66Bo20	Z
	−9346	10			0.3	U			Ric		66Ri09	
	−9346.6	1.7			1.9	U			Yal		69Ov01	Z
	−9347.8	4.0			1.1	U			Har		76Fr13	
^{58}Ni(π^+,π^-)^{58}Zn	−16908	50	−16941	29	−0.7	1	34	34 ^{58}Zn			86Se04	
*^{58}Sc−u	Isomer mixture was corrected.										20Mi13	**
*^{58}Mn−^{39}K$_{1.487}$	D_M=–5887.8(2.9) μu for ^{58}Mn^m at 71.77 keV; $M-A$=–55755.6(2.7) keV										Nub269	**
*^{58}Ni O H$_2$−^{152}Pr$_{.500}$	Reported $M-A$=–50382(10) keV, author confirmed uncertainty error										24Ki16	**
*^{58}Ni(p,^{6}He)^{53}Co	Q increased 1 for recalibration										AHW	**
*^{58}Ni(^{3}He,^{6}He)^{55}Ni	Averaged with reference See ^{46}Ti(^{3}He,^{6}He)										75Mu09	**
*^{58}Ni(p,t)^{56}Ni^j	Strongest of three fragments IT=9943(4); Q rebuilt with Ame1977										MMC129	**
*^{58}Ni(p,d)^{57}Ni	Q =–15210(7) for ^{57}Ni^i at 5238.8(0.7) keV, strongest fragment IT=5230(7);										Nub269	**
*	rebuilt with Q_{gs}=–9975 keV, average of 73Ed01 and 65Sh06										73Ed01	**
*^{58}Ni(^{3}He,α)^{57}Ni	IT=5235(15); Q =3146(15) for ^{57}Ni^i at 5238.8(0.7) rebuilt with Ame1977										MMC129	**
*^{58}Mn(β^-)^{58}Fe	Q_{β^-}=6100(300); and 5930(100) from ^{58}Mn^m at 71.77 keV										Nub269	**
*^{58}Mn(β^-)^{58}Fe	Q_{β^-}=6030(100) from ^{58}Mn^m at 71.77 keV										Nub269	**
*^{58}Fe(t,^{3}He)^{58}Mn	And Q =–6318(15)–77(8) to ^{58}Mn^m at 71.77 keV										Nub269	**
*^{58}Co(β^+)^{58}Fe	E_{β^+}=472(6) 474(4) respectively, to 2$^+$ level at 810.7662 keV										Ens104	**
*^{58}Fe(^{3}He,t)^{58}Co^i	Strongest of two fragments IT=5759(8); Q rebuilt with Ame1964										71Be29	**
*	recalibration +7 keV for ^{54}Fe(p,n)^{54}Co from Ame1961										MMC123	**
^{59}Sc−u	−11626	268				2			RT1	1.0	20Mi13	
^{59}Ti−u	−27075	290	−27780#	320#	−2.4	D			MT1	1.0	20Me06	
	−27053	258			−2.8	D			RT1	1.0	20Mi13	*
^{59}V−u	−38500	400	−40582	3	−3.5	B			TO3	1.5	90Tu01	
	−40700	350			0.2	U			TO5	1.5	94Se12	
	−39900	400			−1.1	U			TO6	1.5	98Ba.A	
	−40677	129			0.5	U			GT1	1.5	04Ma.A	
	−39764	279			−2.9	U			MT1	1.0	20Me06	
^{59}Cr−u	−51490	290	−51654.5	0.7	−0.4	U			TO3	1.5	90Tu01	*
	−51640	310			−0.0	U			TO5	1.5	94Se12	*
	−51100	310			−1.2	U			TO6	1.5	98Ba.A	*
	−52380	500			1.5	U			MT1	1.0	16Me07	*
	−51672	21			0.8	U			MR1	1.0	18Mo14	
	−51651.4	7.5			−0.4	U			TR1	1.0	22Si20	
^{59}Cr−^{85}Rb$_{.694}$	9563.35	0.72	9563.4	0.7	0.1	1	99	99 ^{59}Cr	MA8	1.0	18Mo14	
^{59}Mn−^{39}K$_{1.513}$	−4696.8	2.5	−4696.7	2.4	0.0	1	91	91 ^{59}Mn	MA8	1.0	12Na15	
^{59}Fe−^{85}Rb$_{.694}$	−3907.7	1.1	−3908.0	0.4	−0.2	1	13	13 ^{59}Fe	MA8	1.0	19Hu15	
C$_3$ H$_7$ O−^{59}Co	116467	12	116495.7	0.5	0.6	U			R10	4.0	74De22	
C$_2$ ^{13}C H$_6$ O−^{59}Co	112011	25	112025.5	0.5	0.1	U			R10	4.0	74De22	
C$_2$ H$_5$ O N−^{59}Co	103901	6	103919.7	0.5	0.8	U			R10	4.0	74De22	
^{59}Co−^{39}K$_{1.513}$	−11892.4	1.5	−11893.8	0.5	−0.9	U			MA8	1.0	19Hu15	
^{59}Zn−u	−50698	29	−50687.5	0.8	0.4	U			LZ1	1.0	11Tu09	
^{58}Ni H−^{59}Co	9970	15	9973.16	0.22	0.1	U			R10	4.0	74De22	
^{59}Zn−^{58}Cu$_{1.017}$	5722.4	1.3	5722.5	0.8	0.1	1	36	27 ^{59}Zn	JY1	1.0	10Ka26	
^{59}V−^{59}Cr	11057	12	11073	3	1.3	1	7	7 ^{59}V	RI1	1.0	23Ii01	
^{59}Mn−^{59}Cr	−7953.5	8.1	−7954.3	2.5	−0.1	1	9	9 ^{59}Mn	RI1	1.0	23Ii01	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{59}Zn−^{59}Cu		9815.22	0.72	9815.2	0.6	−0.1	1	80	73 ^{59}Zn	JY1	1.0	10Ka26	
^{59}Co−^{58}Ni		−2182	35	−2148.13	0.22	0.2	U			R10	4.0	74De22	
^{59}V−^{19}F O		−33911.0	7.1	−33900	3	1.6	1	19	19 ^{59}V	RI1	1.0	23Ii01	
^{59}V O H−^{32}S$_2$		18019.7	3.6	18015	3	−1.2	1	74	74 ^{59}V	RI1	1.0	23Ii01	
^{59}Cr O H−^{32}S$_2$		6954.8	5.0	6942.8	0.7	−2.4	U			RI1	1.0	23Ii01	
^{59}Co(p,α)^{56}Fe		3245	8	3241.3	0.3	−0.5	U			MIT		64Sp12	
		3243	9			−0.2	U			Ald		66Br05	
		3240.4	1.4			0.7	U			NDm		74Jo14	
^{59}Co(d,α)^{57}Fe		8667	15	8662.9	0.3	−0.3	U			Kop		67Ha.A	
		8659.3	3.2			1.1	U			NDm		74Jo14	
^{59}Ni(p,t)^{57}Ni		−12738.2	3.3	−12733.7	0.5	1.4	U			MSU		76Na23	
		−12738.4	5.0			0.9	U					78Na11	*
^{58}Fe(n,γ)^{59}Fe		6581.15	0.30	6581.01	0.11	−0.5	–			Ptn		73Sp06	Z
		6580.94	0.20			0.4	–			Ptn		80Ve05	Z
		6581.02	0.14			−0.1	–			Bdn		06Fi.A	
^{58}Fe(d,p)^{59}Fe		4357	8	4356.44	0.11	−0.1	U			MIT		64Sp12	
		4369	8			−1.6	U			Kop		67Ha.A	
^{58}Fe(n,γ)^{59}Fe	ave.	6581.01	0.11	6581.01	0.11	−0.0	1	99	87 ^{59}Fe			average	
^{58}Fe(p,γ)^{59}Co		7359.7	2.0	7363.6	0.4	1.9	U					74Ke14	Z
^{58}Fe(^{3}He,d)^{59}Co		1871	20	1870.1	0.4	−0.0	U			LAl		65Bl13	
^{58}Fe(p,γ)^{59}Co−^{56}Fe()^{57}Co		1336.5	0.7	1336.1	0.4	−0.6	1	41	28 ^{57}Co			75Br29	
^{59}Co(γ,n)^{58}Co		−10441	26	−10453.9	1.1	−0.5	U			Phi		60Ge01	
^{59}Co(d,t)^{58}Co		−4196.0	1.4	−4196.6	1.1	−0.5	1	62	61 ^{58}Co	NDm		74Jo14	
^{58}Ni(n,γ)^{59}Ni		8999.37	0.30	8999.28	0.05	−0.3	U					75Wi06	Z
		8999.38	0.20			−0.5	U			MMn		77Is01	Z
		8999.10	0.23			0.8	U			ILn		93Ha05	Z
		8999.28	0.05			0.0	1	99	71 ^{59}Ni	ORn		04Ra23	
		8999.15	0.18			0.7	U			Bdn		06Fi.A	
^{58}Ni(d,p)^{59}Ni		6797	10	6774.72	0.05	−2.2	U			Kop		67Ha.A	
		6785	5			−2.1	U			MIT		70An25	
		6773.5	1.7			0.7	U			NDm		74Jo14	
^{58}Ni(p,γ)^{59}Cu		3418.5	0.5	3418.6	0.4	0.1	1	62	62 ^{59}Cu			63Bo07	Z
		3419	2			−0.2	U					70Fo09	
		3416.7	2.0			0.9	U					75Kl06	Z
^{58}Ni(p,π^-)^{59}Zn		−144735	40	−144783.6	0.7	−1.2	U					83Sh31	
^{59}Mn(β^-)^{59}Fe		5200	100	5139.1	2.3	−0.6	U			ANB		77Pa18	
^{59}Fe(β^-)^{59}Co		1570	4	1564.9	0.4	−1.3	U					52Me53	*
		1563	3			0.6	U					63Wo01	*
^{59}Ni(ε)^{59}Co		1074.5	1.3	1073.01	0.19	−1.1	U					76Be02	*
^{59}Co(p,n)^{59}Ni		−1855.8	2.0	−1855.35	0.19	0.2	U			MIT		51Mc48	Z
		−1854.3	4.0			−0.3	U					57Bu37	Z
		−1861	5			1.1	U			Ric		57Ch30	Z
		−1855.8	1.6			0.3	U			Oak		64Jo11	Z
		−1855.33	0.20			−0.1	1	95	90 ^{59}Co	PTB		98Bo30	
^{59}Co(^{3}He,t)^{59}Ni^i		−8436	8	−8433.5	2.1	0.3	U					71Be29	*
^{59}Zn(β^+)^{59}Cu		9120	100	9142.8	0.6	0.2	U					81Ar13	
*^{59}Ti−u	Isomer mixture was corrected											20Mi13	**
*^{59}Ti−u	Trends in Mass Surface TMS suggest ^{59}Ti 670 keV more bound											GAu	**
*^{59}Cr−u	Original –51220(240) μu or $M-A$=–47710(230) keV											GAu	**
*^{59}Cr−u	Original –51370(270) μu or $M-A$=–47850(250) keV											GAu	**
*^{59}Cr−u	$M-A$=–47350(250) keV for mixture gs+m at 502.7 keV											Nub269	**
*^{59}Cr−u	$M-A$=–48540(440) keV for mixture gs+m at 502.7 keV											Nub269	**
*^{59}Ni(p,t)^{57}Ni	Strongest of three IAS fragments, Q =–17977.2(5.0) for ^{57}Ni^i at 5238.8											Nub269	**
*^{59}Fe(β^-)^{59}Co	E_{β^-}=475(3) to 3/2$^-$ level at 1099.256 keV											Ens024	**
*^{59}Fe(β^-)^{59}Co	E_{β^-}=462(3), 273(3) to 3/2$^-$ levels at 1099.256, 1291.605 keV											Ens024	**
*^{59}Ni(ε)^{59}Co	Authors add B(K)=8.3 of Ni, changed in 7.7 of Co											AHW	**
*^{59}Co(^{3}He,t)^{59}Ni^i	Strongest fragment Q =–8441(8); recalibration +5 keV for ^{58}Ni(p,n)^{58}Cu^i from Ame1961											MMC129	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{60}Sc−u	−3811	1116	−4890#	320#	−1.0	D			RT1	1.0	20Mi13	*
^{60}Ti−u	−23725	258				2			RT1	1.0	20Mi13	
^{60}V−u	−33860	700	−35490	190	−1.6	U			TO3	1.5	90Tu01	*
	−35560	600			0.1	2			TO5	1.5	94Se12	*
	−35160	510			−0.4	2			TO6	1.5	98Ba.A	*
	−35889	215			1.2	2			GT1	1.5	04Ma.A	
	−35500	420			0.0	o			MT1	1.0	11Es06	*
	−35280	250			−0.8	2			MT1	1.0	20Me06	*
^{60}Cr−u	−49680	240	−50358.3	1.2	−1.9	U			TO3	1.5	90Tu01	
	−50270	280			−0.2	U			TO5	1.5	94Se12	
	−49910	280			−1.1	U			TO6	1.5	98Ba.A	
	−50929	494			1.2	U			MT1	1.0	16Me07	
	−50368	20			0.5	U			MR1	1.0	18Mo14	
^{60}Cr−^{85}Rb$_{.706}$	11918.1	1.2				2			MA8	1.0	18Mo14	
^{60}Mn−u	−56550	240	−56863.4	2.5	−0.9	U			TO3	1.5	90Tu01	*
	−56810	290			−0.1	U			TO5	1.5	94Se12	*
	−56530	280			−0.8	U			TO6	1.5	98Ba.A	*
^{60}Mn−^{39}K$_{1.538}$	−1044.0	2.5				2			MA8	1.0	12Na15	*
^{60}Co−u	−66380	280	−66183.9	0.5	0.5	U			TO6	1.5	98Ba.A	*
C_3 H_8 O−^{60}Ni	126796	14	126729.1	0.4	−1.2	U			R10	4.0	74De22	
C_2 H_6 O N−^{60}Ni	114231	10	114153.1	0.4	−1.9	U			R10	4.0	74De22	
C_2 ^{13}C H_7 O−^{60}Ni	122315	10	122259.0	0.4	−1.4	U			R10	4.0	74De22	
C H_2 N O_2−^{60}Ni	77843	16	77767.6	0.4	−1.2	U			R10	4.0	74De22	
C_5−^{60}Ni	69275	14	69214.3	0.4	−1.1	U			R10	4.0	74De22	
^{60}Ni−^{85}Rb$_{.706}$	−6937.8	1.6	−6937.8	0.4	−0.0	U			MA8	1.0	07Gu09	
^{60}Ga−u	−42947.1	32.2	−42958	14	−0.3	–			TR1	1.0	21Pa44	
	−42978.3	49.4			0.4	–			LZ2	1.0	23Wa10	
ave.	−42956	27			−0.1	1	26	26 ^{60}Ga			average	
^{60}Zn−^{58}Ni$_{1.034}$	8698.02	0.55	8698.1	0.4	0.1	1	65	65 ^{60}Zn	JY1	1.0	10Ka26	
^{60}Zn−^{59}Cu$_{1.017}$	3373.19	0.55	3373.2	0.4	−0.1	1	65	35 ^{60}Zn	JY1	1.0	10Ka26	
^{60}Ni−^{58}Ni H	−12513	30	−12381.56	0.08	1.1	U			R10	4.0	74De22	
^{60}Ni−^{59}Co	−2503	40	−2408.40	0.22	0.6	U			R10	4.0	74De22	
^{60}Ni−^{58}Ni	−4624	25	−4556.52	0.08	0.7	U			R10	4.0	74De22	
	−4627	45			0.4	U			R10	4.0	74De22	
^{60}Ni H−^{59}Co	5310	40	5416.64	0.22	0.7	U			R10	4.0	74De22	
^{60}Ni(p,α)^{57}Co	−263.6	0.7	−263.6	0.4	0.1	1	37	33 ^{57}Co	NDm		74Jo14	
^{58}Fe(t,p)^{60}Fe	6907	15	6919	3	0.8	2			LAl		71Ca19	
	6947	10			−2.8	2			MSU		76St11	
	6913	4			1.4	2			LAl		78No05	
^{60}Ni(d,α)^{58}Co	6084.5	2.2	6084.8	1.1	0.2	1	25	25 ^{58}Co	NDm		74Jo14	
^{58}Ni(t,p)^{60}Ni	11905	10	11905.22	0.07	0.0	U			Ald		71Da16	
^{60}Ni(p,t)^{58}Ni^{i}	−20735	40				2					74Ko08	*
^{60}Ni(p,t)^{58}Ni^{j}	−26444	7				2					84Ka07	*
^{58}Ni(^{3}He,p)^{60}Cu	5770	12	5758.6	1.6	−0.9	U			CIT		67Mi02	
	5746	20			0.6	U			MIT		68Yo01	
^{58}Ni(^{3}He,p)^{60}Cu^{i}	3210	10	3218	5	0.8	1	26	26 ^{60}Cu^{i}	MIT		68Yo01	
^{58}Ni(^{3}He,n)^{60}Zn	818	18	805.5	0.4	−0.7	U			CIT		67Mi02	
	821	13			−1.2	U			Oak		72Gr39	
^{58}Ni(^{3}He,n)^{60}Zn^{j}	−6562	24				2					74Ev02	*
^{59}Co(n,γ)^{60}Co	7491.88	0.08	7491.92	0.07	0.5	2			BNn		84Ko29	Z
	7492.05	0.15			−0.9	2			Bdn		06Fi.A	
^{59}Co(d,p)^{60}Co	5267	11	5267.35	0.07	0.0	U			MIT		64Sp12	
	5272	8			−0.6	U			Kop		67Ha.A	
^{59}Co(p,γ)^{60}Ni^{i}	−1594	4				2					67Ar01	
^{59}Ni(n,γ)^{60}Ni	11387.6	0.4	11387.73	0.05	0.3	U					75Wi06	Z
	11387.73	0.05			0.0	1	99	74 ^{60}Ni	ORn		04Ra23	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{60}Ni(p,d)^{59}Ni	−9180	50	−9163.16	0.05	0.3	U			Pri		64Le10	
^{60}Ni(d,t)^{59}Ni	−5130.2	2.1	−5130.50	0.05	−0.1	U			NDm		74Jo14	
^{60}Ni(p,d)^{59}Ni^{i}	−16505.1	2.1				2					78Ik02	*
^{60}Ga^{i}(p)^{59}Zn	2522	15	2523	13	0.0	1	74	74 ^{60}Ga^{i}			21Or01	
^{60}Mn(β^-)^{60}Fe	8234	86	8445	4	2.4	U			ANB		78No03	*
^{60}Co(β^-)^{60}Ni	2823.6	1.0	2822.81	0.21	−0.8	U					68Wo02	*
^{60}Cu(β^+)^{60}Ni	6250	40	6128.0	1.6	−3.1	B					54Nu26	
^{60}Ni(p,n)^{60}Cu	−6912	20	−6910.3	1.6	0.1	U			ChR		58Go77	
	−6909	10			−0.1	U			Ric		66Ri09	
	−6910.3	1.6				2			Yal		69Ov01	Z
^{60}Ni(^{3}He,t)^{60}Cu^{i}	−8685	6	−8688	5	−0.5	1	74	74 ^{60}Cu^{i}			71Be29	*
^{60}Zn(β^+)^{60}Cu	4166	64	4170.8	1.6	0.1	U					86Ka38	
^{60}Ga^{i}(IT)^{60}Ga	2611.8	0.9	2611.8	0.9	−0.0	1	100	74 ^{60}Ga			21Or01	
*^{60}Sc−u	Trends in Mass Surface TMS suggest ^{60}Sc 1000 keV more bound										HWJ266	**
*^{60}V−u	Original –33800(700) μu or $M-A$=–31500(650) keV										GAu	**
*^{60}V−u	Original –35500(600) μu or $M-A$=–33070(560) keV										GAu	**
*^{60}V−u	$M-A$=–32700(470) keV for mixture gs+m+n at 60#50 and 103.2(0.1) keV										Nub269	**
*^{60}V−u	$M-A$=–33010(390) keV for mixture gs+m+n at 60#50 and 103.2(0.1) keV										Nub269	**
*^{60}V−u	$M-A$=–32810(230) keV for mixture gs+m+n at 60#50 and 103.2(0.1) keV										Nub269	**
*^{60}Mn−u	$M-A$=–52540(230) keV for mixture gs+m at 271.90 keV										Nub269	**
*^{60}Mn−u	$M-A$=–52780(260) keV for mixture gs+m at 271.90 keV										Nub269	**
*^{60}Mn−u	$M-A$=–52520(250) keV for mixture gs+m at 271.90 keV										Nub269	**
*^{60}Mn−^{39}K$_{1.538}$	D_M=–752.1(2.5) μu for ^{60}Mn^{m} at 271.90 keV; $M-A$=–52695.9(2.4) keV										Nub269	**
*^{60}Co−u	$M-A$=–61800(260) keV for mixture gs+m at 58.59 keV										Nub269	**
*^{60}Ni(p,t)^{58}Ni^{i}	IT=8830(40); Q rebuilt with Ame1971										MMC124	**
*^{60}Ni(p,t)^{58}Ni^{j}	IT=14537(7); Q rebuilt with Ame1977										MMC124	**
*^{58}Ni(^{3}He,n)^{60}Zn^{j}	IT=7380(30); Q rebuilt with Ame1971										MMC124	**
*^{60}Ni(p,d)^{59}Ni^{i}	Strongest fragment IT=7341; Q rebuilt with Ame1977										MMC129	**
*^{60}Mn(β^-)^{60}Fe	E_{β^-}=5714(86) from ^{60}Mn^{m} at 271.90 to (3^+,4^+) 2792.68 level										Ens13c	**
*^{60}Co(β^-)^{60}Ni	E_{β^-}=317.88(0.10) to 4^+ level at 2505.753 keV										Ens13c	**
*^{60}Ni(^{3}He,t)^{60}Cu^{i}	CDE=9454(6) Q =–8690(6); recalibration +5 keV for ^{58}Ni(p,n)^{58}Cu^{i} from Ame1961										MMC123	**
^{61}Ti−u	−17370	483				2			RT1	1.0	20Mi13	*
^{61}V−u	−32750	960	−32400	250	0.4	o			MT1	1.0	11Es06	
	−32614	301			0.7	2			MT1	1.0	20Me06	
	−31884	462			−1.1	2			RT1	1.0	20Mi13	
^{61}Cr−u	−44500	400	−45621.9	2.0	−1.9	U			TO3	1.5	90Tu01	
	−45910	300			0.6	U			TO5	1.5	94Se12	
	−45120	280			−1.2	U			TO6	1.5	98Ba.A	
	−45679	107			0.4	U			GT1	1.5	04Ma.A	
	−45634	176			0.1	U			LZ1	1.0	15Xu14	
	−46248	548			1.1	U			MT1	1.0	16Me07	
	−45629	22			0.3	U			MR1	1.0	18Mo14	
	−45622.4	15.0			0.0	U			TR1	1.0	22Si20	
^{61}Cr−^{85}Rb$_{.718}$	17713.1	2.0				2			MA8	1.0	18Mo14	
^{61}Mn−u	−55160	300	−55547.5	2.5	−0.9	U			TO3	1.5	90Tu01	
	−55540	280			−0.0	U			TO5	1.5	94Se12	
	−55320	270			−0.6	U			TO6	1.5	98Ba.A	
^{61}Mn−^{39}K$_{1.564}$	1215.6	2.5				2			MA8	1.0	12Na15	
^{61}Fe−^{39}K$_{1.564}$	−6490.7	2.8				2			MA8	1.0	12Na15	
C H$_3$ N O$_2$−^{61}Ni	85373	14	85322.9	0.4	−0.9	U			R10	4.0	74De22	
C$_5$ H−^{61}Ni	76810	10	76769.6	0.4	−1.0	U			R10	4.0	74De22	
^{61}Zn−^{42}Ca ^{19}F	−17523.3	3.6	−17521.1	1.2	0.6	–			MA8	1.0	26Gi02	
	−17520.5	1.5			−0.4	–			IS1	1.0	26Gi02	
ave.	−17520.9	1.4			−0.1	1	79	79 ^{61}Zn			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{61}Zn−^{85}Rb$_{.718}$		2833.2	3.1	2834.4	1.2	0.4	1	16	16 ^{61}Zn	MA8	1.0	26Gi02	
^{61}Zn−C$_5$ H		−68325.1	5.4	−68325.6	1.2	−0.1	1	5	5 ^{61}Zn	MS1	1.0	22Me04	
^{61}Zn−u		−60501	24	−60500.6	1.2	0.0	U			MR1	1.0	26Gi02	
^{61}Ga−u		−50654	59	−50592	11	1.0	U			LZ1	1.0	11Tu09	
		−50579.0	12.9			−1.0	2			TR1	1.0	21Pa44	
		−50636.9	23.5			1.9	2			LZ2	1.0	23Wa10	
^{60}Ni H−^{61}Ni		7539	14	7555.34	0.05	0.3	U			R10	4.0	74De22	
^{61}Ni−^{60}Ni		339	60	269.69	0.05	−0.3	U			R10	4.0	74De22	
^{61}Ni−^{58}Ni H		−12187	30	−12111.87	0.09	0.6	U			R10	4.0	74De22	
^{61}Ni−^{59}Co		−2220	30	−2138.71	0.22	0.7	U			R10	4.0	74De22	
^{58}Ni(α,n)^{61}Zn		−9810	30	−9518.8	1.2	9.7	B			Oak		64St01	
^{58}Ni(^{6}Li,t)^{61}Zn		−4736	23	−4735.3	1.2	0.0	R			LAl		78Wo01	
^{59}Co(^{3}He,p)^{61}Ni		9635	10	9634.44	0.21	−0.1	U			MIT		67Sp09	
^{60}Ni(n,γ)^{61}Ni		7820.22	0.40	7820.11	0.05	−0.3	U					75Wi06	Z
		7819.96	0.20			0.7	U			MMn		77Is01	Z
		7820.02	0.20			0.4	U			ILn		93Ha05	Z
		7820.11	0.05			−0.1	–			ORn		04Ra23	
		7820.06	0.16			0.3	–			Bdn		06Fi.A	
^{60}Ni(d,p)^{61}Ni		5604	8	5595.54	0.05	−1.1	U			MIT		70An25	
		5596.1	1.3			−0.4	U			NDm		74Jo14	
^{60}Ni(n,γ)^{61}Ni	ave.	7820.11	0.05	7820.11	0.05	0.0	1	100	78 ^{61}Ni			average	
^{61}Ga^{i}(p)^{60}Zn		3110	30				2					87Ho.A	
^{61}Fe(β^-)^{61}Co		3827	100	3977.2	2.7	1.5	U					67Eh02	*
		3887	100			0.9	U					67Gu06	*
^{61}Co(β^-)^{61}Ni		1290	40	1323.7	0.8	0.8	U					56Nu02	
^{61}Cu(β^+)^{61}Ni		2227	5	2237.7	1.0	2.1	U					50Ow03	
^{61}Ni(p,n)^{61}Cu		−3024.0	4.	−3020.0	1.0	1.0	U			Oak		64Jo11	Z
^{61}Ni(^{3}He,t)^{61}Cu^{i}		−8630	7				2					71Be29	*
^{61}Zn(β^+)^{61}Cu		5400	200	5627.9	1.5	1.1	U					59Cu86	
^{61}Ga(β^+)^{61}Zn		9255	50	9229	11	−0.5	U					02We07	
*^{61}Ti−u	Isomer mixture was corrected.											20Mi13	**
*^{61}Fe(β^-)^{61}Co	E_{β^-}=2800(100) 2860(100) respectively, to 3/2$^-$ level at 1027.48 keV											Ens153	**
*^{61}Ni(^{3}He,t)^{61}Cu^{i}	Strongest fragment IT=6380(7); Q rebuilt with Ame1964											MMC129	**
*	recalibration +5 keV for ^{58}Ni(p,n)^{58}Cu^{i} from Ame1961											MMC123	**
^{62}Ti−u		−14278	483	−13100#	430#	2.4	D			RT1	1.0	20Mi13	*
^{62}V−u		−27204	451	−27070	280	0.3	2			MT1	1.0	20Me06	
		−26978	365			−0.2	2			RT1	1.0	20Mi13	
^{62}Cr−u		−42400	600	−43857	4	−1.6	U			TO3	1.5	90Tu01	
		−44200	400			0.6	U			TO5	1.5	94Se12	
		−43100	350			−1.4	U			TO6	1.5	98Ba.A	
		−44026	118			1.0	U			GT1	1.5	04Ma.A	
		−43897	526			0.1	U			MT1	1.0	16Me07	
		−43845	19			−0.6	U			MR1	1.0	18Mo14	
		−43859.6	20.4			0.1	U			TR1	1.0	22Si20	
^{62}Cr−^{85}Rb$_{.729}$		20448.2	3.7				2			MA8	1.0	18Mo14	
^{62}Mn−u		−51510	270	−52093	7	−1.4	U			TO3	1.5	90Tu01	
		−52030	280			−0.1	U			TO5	1.5	94Se12	
		−51180	280			−2.2	U			TO6	1.5	98Ba.A	
^{62}Mn^{m}−^{39}K$_{1.590}$		5982.3	2.8				2			MA8	1.0	12Na15	
^{62}Fe−^{39}K$_{1.590}$		−5501.5	3.0				2			MA8	1.0	12Na15	
C$_5$ H$_2$−^{62}Ni		87299	10	87304.8	0.5	0.1	U			R10	4.0	74De22	
C H$_4$ N O$_2$−^{62}Ni		95859	12	95858.1	0.5	−0.0	U			R10	4.0	74De22	
^{62}Ga−u		−55815.7	15.0	−55809.9	0.7	0.4	U			TR1	1.0	21Pa44	
^{62}Ge−u		−45399.1	39.7				2			LZ2	1.0	23Wa10	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{62}Cu−^{62}Ni		4250.05	0.51				2			JY1	1.0	06Er03	
^{62}Zn−^{62}Ni		5988.49	0.58	5988.6	0.5	0.2	1	68	68 ^{62}Zn	JY1	1.0	06Er03	
^{62}Ga−^{62}Ni		15845.06	0.71	15844.9	0.5	−0.2	1	52	52 ^{62}Ga	JY1	1.0	06Er03	
^{62}Ga−^{62}Zn		9856.21	0.45	9856.3	0.4	0.2	1	81	48 ^{62}Ga	JY1	1.0	06Er03	
^{62}Ni−^{61}Ni		−2669	15	−2710.2	0.3	−0.7	U			R10	4.0	74De22	
^{62}Ni−^{60}Ni		−2333	30	−2440.5	0.3	−0.9	U			R10	4.0	74De22	
^{62}Ni(p,α)^{59}Co		342	10	347.4	0.4	0.5	U			MIT		64Sp12	
		343.3	0.7			5.8	B			NDm		74Jo14	
^{59}Co(α,p)^{62}Ni		−346.5	2.3	−347.4	0.4	−0.4	U			NDm		74Jo14	
^{62}Ni(^{18}O,^{20}Ne)^{60}Fe		911	20	926	3	0.7	U			Hei		84Ha31	
^{62}Ni(d,α)^{60}Co		5611.2	2.4	5614.7	0.4	1.5	U			NDm		74Jo14	
^{60}Ni(t,p)^{62}Ni		9937	10	9934.1	0.3	−0.3	U			Ald		71Da16	
^{60}Ni(^{3}He,p)^{62}Cu		5938	25	5956.6	0.6	0.7	U			MIT		67Sp09	
^{60}Ni(^{3}He,n)^{62}Zn		3580	30	3554.8	0.5	−0.8	U			Oak		72Gr39	
^{62}Ni(^{14}C,^{15}O)^{61}Fe		−7921	100	−7661.2	2.7	2.6	F			Ors		84De33	*
^{62}Ni(t,α)^{61}Co		8689	20	8676.6	0.7	−0.6	U			LAl		66Bl15	
^{61}Ni(n,γ)^{62}Ni		10596.2	1.5	10595.8	0.3	−0.3	–					70Fa06	
		10595.8	0.7			0.0	–					75Wi06	Z
		10595.6	0.4			0.5	–			Bdn		06Fi.A	
^{61}Ni(d,p)^{62}Ni		8379	8	8371.2	0.3	−1.0	U			MIT		64Sp12	
		8369	15			0.1	U			Ald		67Te02	
^{62}Ni(d,t)^{61}Ni		−4340.6	1.3	−4338.6	0.3	1.5	–			NDm		74Jo14	
^{61}Ni(n,γ)^{62}Ni	ave.	10595.8	0.3	10595.8	0.3	0.0	1	88	66 ^{62}Ni			average	
^{62}Mn^{m}(IT)^{62}Mn		343	6				3					15Ga38	*
^{62}Fe(β^-)^{62}Co		3000	200	2546	19	−2.3	U					75Fr16	
^{62}Co(β^-)^{62}Ni		5195	30	5322	19	4.2	C					57Ga15	*
^{62}Ni(t,^{3}He)^{62}Co		−5350	50	−5303	19	0.9	2					72Ba31	
		−5296	20			−0.4	2			LAl		76Aj03	
^{62}Cu(β^+)^{62}Ni		3932	10	3958.9	0.5	2.7	U					54Nu27	
		3942	10			1.7	U					64Sa32	
		3956	7			0.4	U					67An01	
^{62}Ni(p,n)^{62}Cu		−4733	10	−4741.2	0.5	−0.8	U			Bar		61Ri02	
		−4734.8	10.			−0.6	U			Ric		66Ri09	
^{62}Ni(^{3}He,t)^{62}Cu^{i}		−8591	6				2					71Be29	*
^{62}Zn(β^+)^{62}Cu		1682	10	1619.5	0.7	−6.3	B					50Ha65	
		1697	10			−7.8	B					54Nu27	
^{62}Ga(β^+)^{62}Zn		9171	26	9181.1	0.4	0.4	U			ANB		79Da04	
*^{62}Ti−u	Trends in Mass Surface TMS suggest ^{62}Ti 1200 keV less bound											GAu	**
*^{62}Ni(^{14}C,^{15}O)^{61}Fe	F : not unambiguously ground state transition											84De33	**
*^{62}Mn^{m}(IT)^{62}Mn	E_x =418(2) – 72(+8–3)											GAu	**
*^{62}Co(β^-)^{62}Ni	E_{β^-}=5217(30) from ^{62}Co^{m} at 22(5) keV											Nub269	**
*^{62}Ni(^{3}He,t)^{62}Cu^{i}	CDE=9360(6) Q =–8596(6); recalibration +5 keV for ^{58}Ni(p,n)^{58}Cu^{i} from Ame1961											MMC124	**
^{63}V−u		−23339	365				2			RT1	1.0	20Mi13	
^{63}Cr−u		−38819	462	−38867	19	−0.1	U			GT1	1.5	04Ma.A	
		−37870	700			−1.4	o			MT1	1.0	11Es06	
		−38583	462			−0.6	U			MT1	1.0	16Me07	
		−38839	78			−0.4	U			MR1	1.0	18Mo14	
		−38866.6	19.3				2			TR1	1.0	22Si20	
^{63}Mn−u		−49300	400	−50335	4	−1.7	U			TO3	1.5	90Tu01	
		−50190	300			−0.3	U			TO5	1.5	94Se12	
		−49600	290			−1.7	U			TO6	1.5	98Ba.A	
		−50500	107			1.0	o			GT1	1.5	04Ma.A	
		−50829	102			1.9	U			GT2	2.5	08Kn.A	
^{63}Mn−^{39}K$_{1.615}$		8278.7	4.0				2			MA8	1.0	12Na15	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{63}Fe−u		−59190	240	−59727	4	−1.5	U			TO3	1.5	90Tu01	
		−59570	290			−0.4	U			TO5	1.5	94Se12	
		−58990	300			−1.6	U			TO6	1.5	98Ba.A	
		−59727.1	5.8			−0.0	1	39	39 ^{63}Fe	TR1	1.0	22Po02	
^{63}Fe−^{39}K$_{1.615}$		−1114.5	6.1	−1113	4	0.2	1	35	35 ^{63}Fe	MA8	1.0	12Na15	
^{63}Fe−H C$_2$ F$_2$		−64354	10	−64359	4	−0.5	o			MS1	1.0	08Bl05	
		−64353	10			−0.6	1	13	13 ^{63}Fe	MS1	1.0	10Fe01	
^{63}Fe−C ^{32}S F		−30204	10	−30202	4	0.2	1	13	13 ^{63}Fe	MS1	1.0	10Fe01	
C$_5$ H$_3$−^{63}Cu		93930	4	93877.5	0.5	−3.3	B			R10	4.0	74De22	
C$_4$ H N−^{63}Cu		81347	10	81301.4	0.5	−1.1	U			R10	4.0	74De22	
C$_4$ ^{13}C H$_2$−^{63}Cu		89466	14	89407.3	0.5	−1.0	U			R10	4.0	74De22	
C$_2$ H$_7$ O$_2$−^{63}Cu		115064	16	115006.8	0.5	−0.9	U			R10	4.0	74De22	
^{13}C C H$_8$ O N−^{63}Cu		134404	18	134346.1	0.5	−0.8	U			R10	4.0	74De22	
^{47}Ti O−^{63}Cu		17036	23	17074.5	0.5	0.4	U			R09	4.0	72De11	
^{63}Cu H$_2$ O−^{81}Sr		16962	14	16951	3	−0.8	U			RI1	1.0	18Ki21	
^{63}Ga−^{85}Rb$_{.741}$		4658.0	1.4				2			MA8	1.0	07Gu09	
^{63}Ga−u		−60722.9	15.0	−60705.8	1.4	1.1	U			TR1	1.0	21Pa44	
C$_5$ H$_2$−^{63}Ga$_{.984}$		75382.6	6.7	75384.6	1.4	0.3	U			MS1	1.0	07Sc24	
^{63}Ge−u		−50372	40	−50426	15	−1.3	–			LZ1	1.0	11Tu02	
		−50433.0	16.1			0.4	–			LZ2	1.0	23Wa10	
	ave.	−50424	15			−0.1	1	99	99 ^{63}Ge			average	
^{63}Cu−^{62}Ni		1193	35	1252.36	0.06	0.4	U			R10	4.0	74De22	
^{63}Cu−^{61}Ni		−1449	30	−1457.8	0.3	−0.1	U			R10	4.0	74De22	
^{63}Cu(p,α)^{60}Ni		3757	8	3757.3	0.3	0.0	U			MIT		64Sp12	
		3780	10			−2.3	U			Min		67Jo03	
		3754.9	1.5			1.6	U			NDm		76Jo01	
^{60}Ni(α,n)^{63}Zn		−7970	40	−7906.0	1.6	1.6	U			Oak		64St01	
		−7910	20			0.2	U					67Bi04	
^{63}Cu(d,α)^{61}Ni		9376	30	9352.9	0.3	−0.8	U					67Hj01	
^{62}Ni(n,γ)^{63}Ni		6838.04	0.20	6837.77	0.06	−1.3	–			MMn		77Is01	Z
		6837.88	0.18			−0.6	–			ILn		92Ha21	Z
		6837.89	0.14			−0.8	–			Bdn		06Fi.A	
		6837.75	0.18			0.1	–			JAn		12Os04	
^{62}Ni(d,p)^{63}Ni		4620	6	4613.21	0.06	−1.1	U			MIT		70An25	
		4614.0	1.1			−0.7	U			NDm		74Jo14	
^{62}Ni(n,γ)^{63}Ni	ave.	6837.88	0.09	6837.77	0.06	−1.3	1	48	33 ^{63}Ni			average	
^{62}Ni(p,γ)^{63}Cu		6119.2	1.5	6122.40	0.06	2.1	U					72Ki15	
		6122.30	0.08			1.3	1	54	37 ^{63}Cu	Utr		86De14	Z
^{63}Cu(γ,n)^{62}Cu		−10833	17	−10863.6	0.5	−1.8	U			Phi		60Ge01	
^{63}Co(β^-)^{63}Ni		3590	50	3661	19	1.4	1	14	14 ^{63}Co			69Ki.A	
^{63}Ni(β^-)^{63}Cu		65.87	0.15	66.977	0.015	7.4	B					66Hs01	
		66.946	0.020			1.5	o					87He14	
		66.945	0.004			7.9	F					92Ka29	*
		66.9459	0.0054			5.7	F					93Oh02	*
		66.980	0.015			−0.2	1	98	55 ^{63}Ni			99Ho09	
^{63}Zn(β^+)^{63}Cu		3352	20	3366.3	1.5	0.7	U					61Cu02	
		3390	30			−0.8	U					61Va08	
^{63}Cu(p,n)^{63}Zn		−4146.5	4.	−4148.6	1.5	−0.5	–			Ric		55Br16	
		−4139.5	8.			−1.1	U			Oak		55Ki28	Z
		−4150.1	4.4			0.3	–			Tkm		63Ok01	
	ave.	−4148.1	2.9			−0.2	1	28	27 ^{63}Zn			average	
^{63}Cu(^{3}He,t)^{63}Zn^i		−8875	6				2					71Be29	*
^{63}Ga(β^+)^{63}Zn		5520	100	5666.0	2.0	1.5	U					72Fi.A	
*^{63}Ni(β^-)^{63}Cu	F : excitation of atomic electron not taken into account											99Ho09	**
*^{63}Cu(^{3}He,t)^{63}Zn^i	CDE=9644(6) Q =–8880(6); recalibration +5 keV for ^{58}Ni(p,n)^{58}Cu^i from Ame1961											MMC124	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{64}V−u	−17917	548	−17520#	430#	0.7	D			RT1	1.0	20Mi13	*
^{64}Cr−u	−35942	472	−36042	28	−0.2	o			MT1	1.0	16Me07	
	−36114	322			0.2	U			MT1	1.0	20Me06	
	−36042	28				2			TR1	1.0	22Si20	
^{64}Mn−u	−45340	350	−46151	4	−1.5	U			TO3	1.5	90Tu01	*
	−46340	350			0.4	U			TO5	1.5	94Se12	*
	−45664	306			−1.1	U			TO6	1.5	98Ba.A	*
	−46280	129			0.7	U			GT1	1.5	04Ma.A	
	−46122.7	20.4			−1.4	U			TR1	1.0	25Ch42	
^{64}Mn−^{85}Rb$_{.753}$	20271.7	3.8				2			MA8	1.0	12Na15	
^{64}Fe−u	−58600	400	−59012	4	−0.7	U			TO3	1.5	90Tu01	
	−59130	300			0.3	U			TO5	1.5	94Se12	
	−58500	350			−1.0	U			TO6	1.5	98Ba.A	
	−59012.2	5.3			−0.0	o			MS1	1.0	08Bl05	
	−59012.7	5.7			0.0	1	47	47 ^{64}Fe	TR1	1.0	22Po02	
H C$_2$ F$_2$−^{64}Fe$_{.984}$	62699.4	5.3	62700	4	0.0	1	53	53 ^{64}Fe	MS1	1.0	10Fe01	
^{64}Co^m−u	−64075.5	4.5	−64075	4	0.1	o			MS1	1.0	08Bl05	
H C$_2$ F$_2$−^{64}Co$^m_{.984}$	67681.6	4.6	67681	4	−0.1	1	87	87 ^{64}Co^m	MS1	1.0	10Fe01	
^{64}Co^m−^{32}S O$_2$	−25974	12	−25976	4	−0.1	1	13	13 ^{64}Co^m	MS1	1.0	10Fe01	
C$_5$ H$_4$−^{64}Ni	103278	10	103333.4	0.5	1.4	U			R10	4.0	74De22	
C$_4$ ^{13}C H$_3$−^{64}Ni	98809	12	98863.2	0.5	1.1	U			R10	4.0	74De22	
C$_4$ H$_2$ N−^{64}Ni	90703	16	90757.4	0.5	0.8	U			R10	4.0	74De22	
^{64}Ni−^{85}Rb$_{.753}$	−5609.2	1.4	−5611.0	0.5	−1.3	1	14	14 ^{64}Ni	MA8	1.0	07Gu09	
^{64}Cu−u	−70224	30	−70235.5	0.5	−0.4	U			TR1	1.0	25Ch42	
^{64}Zn−^{85}Rb$_{.753}$	−4430.1	8.4	−4435.6	0.7	−0.7	U			MA8	1.0	07Ke09	
^{64}Ga−^{85}Rb$_{.753}$	3261.3	2.5	3262.8	1.5	0.6	1	38	38 ^{64}Ga	MA8	1.0	07Gu09	
C$_5$ H$_2$−^{64}Ga$_{.969}$	76851.5	2.6	76851.7	1.5	0.1	1	33	33 ^{64}Ga	MS1	1.0	07Sc24	
^{64}Ge−u	−57090	690	−58310	4	−1.2	U			GA6	1.5	02Li24	
H ^{32}S O$_2$−^{64}Ge H	20210.5	4.0				2			MS1	1.0	07Sc24	
^{64}Ge−^{85}Rb$_{.753}$	8070	43	8112	4	1.0	U			MS1	1.0	19Sc11	
^{64}As−u	−42630	120	−42540	30	0.8	1	8	8 ^{64}As	LZ2	1.0	23Wa10	
^{64}Ga−^{64}Zn	7698.5	4.1	7698.4	1.6	−0.0	1	15	13 ^{64}Ga	CP1	1.0	07Cl01	
^{64}Ge−^{64}Zn	12517	33	12548	4	0.9	U			CP1	1.0	07Cl01	
^{64}Ni−^{63}Cu	−1523	30	−1630.90	0.22	−0.9	U			R10	4.0	74De22	
^{64}Ga−^{63}Ga	−2730	150	−2453.7	2.1	0.7	U			CR1	2.5	89Sh10	
^{64}Ni−^{62}Ni	−352	25	−378.54	0.23	−0.3	U			R10	4.0	74De22	
^{64}Ni(^{3}He,^{8}B)^{59}Mn	−19610	30	−19563.7	2.5	1.5	U			MSU		76Ka24	
^{64}Ni(^{3}He,^{7}Be)^{60}Fe	−6511	10	−6524	3	−1.3	R			MSU		76St11	
^{64}Ni(α,^{7}Be)^{61}Fe	−21523	20	−21522.2	2.7	0.0	U			Tex		77Co08	
^{64}Ni(^{14}C,^{17}O)^{61}Fe	−4609	100	−4349.4	2.7	2.6	U			Ors		84Be.A	*
^{64}Ni(p,α)^{61}Co	663.2	0.7				2			NDm		74Jo14	
^{64}Zn(p,α)^{61}Cu	830	15	844.1	0.7	0.9	U					67Br10	
	830	10			1.4	U			Min		67Jo03	
	844.1	0.7				2			NDm		76Jo01	
^{64}Zn(^{3}He,^{6}He)^{61}Zn	−12331	23	−12308.7	1.3	1.0	U			MSU		79We02	
	−12309.6	23.			0.0	U					58Ni+3	
^{64}Ni(^{11}B,^{13}N)^{62}Fe	−4930	70	−4898.3	2.8	0.5	U			Tex		77Co08	
^{64}Ni(^{14}C,^{16}O)^{62}Fe	−501	40	−463.6	2.8	0.9	U			Ors		81Be40	
^{64}Ni(^{18}O,^{20}Ne)^{62}Fe	−1915	50	−1961.4	2.8	−0.9	U			Can		76Hi14	
	−1920	21			−2.0	U			Hei		77Bh03	*
	−1947	26			−0.6	U			Hei		84Ha31	
^{64}Ni(d,α)^{62}Co	5190	20	5036	19	−7.7	B					72Ba31	
^{62}Ni(t,p)^{64}Ni	7999	20	8013.45	0.21	0.7	U			Ald		71Da16	
^{62}Ni(^{3}He,p)^{64}Cu	6299	25	6320.23	0.06	0.8	U			MIT		67Sp09	
^{62}Ni(^{3}He,n)^{64}Zn	6118	12	6117.7	0.6	−0.0	U			Oak		72Gr39	
^{64}Zn(d,α)^{62}Cu	7508	15	7494.1	0.8	−0.9	U			MIT		67Sp09	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{64}Zn(p,t)^{62}Zn		−12493	10	−12497.0	0.8	−0.4	U			Bld		72Fa08	
^{64}Ni(^{14}C,^{15}O)^{63}Fe		−11387	60	−11299	3	1.5	U			Ors		82De.A	*
^{64}Ni(^{34}S,^{35}Ar)^{63}Fe		−17931	260	−18347	3	−1.6	U			Hei		83Wi.B	
^{64}Ni(t,α)^{63}Co		7266	20	7277	19	0.6	1	86	86 ^{63}Co	LAl		66Bl15	
^{63}Ni(n,γ)^{64}Ni		9657.32	0.4	9657.47	0.20	0.4	–					75Wi06	Z
		9657.58	0.24			−0.5	–			ILn		92Ha21	
	ave.	9657.51	0.21			−0.2	1	98	86 ^{64}Ni			average	
^{63}Cu(n,γ)^{64}Cu		7916.07	0.12	7915.868	0.024	−1.7	U			BNn		83De28	Z
		7915.52	0.08			4.4	B					02Bo11	
		7916.14	0.16			−1.7	U			Bdn		06Fi.A	
		7915.867	0.024			0.0	1	100	90 ^{64}Cu			16Te05	
^{63}Cu(d,p)^{64}Cu		5697	8	5691.302	0.024	−0.7	U			MIT		64Sp12	
^{64}Zn(n,d)^{63}Cu		−5520	50	−5488.8	0.6	0.6	U					65Wa14	
^{64}Zn(d,t)^{63}Zn		−5604.9	1.7	−5604.7	1.5	0.1	1	76	73 ^{63}Zn	NDm		76Jo01	
^{64}As^{i}(p)^{63}Ge		2020	31	2014	30	−0.2	1	93	92 ^{64}As^{i}			25Ag12	*
^{64}Co(β^-)^{64}Ni		7000	500	7307	20	0.6	U					69Wa15	
		7000	400			0.8	U					74Ra31	
^{64}Ni(t,^{3}He)^{64}Co		−7288	20				2			LAl		72Fl17	
^{64}Cu(β^+)^{64}Ni		1673.4	1.0	1674.63	0.21	1.2	U					83Ch47	
^{64}Ni(p,n)^{64}Cu		−2458	6	−2456.97	0.21	0.2	U					61Va19	
		−2458.22	0.31			4.0	B			PTB		92Bo02	Z
^{64}Cu(β^-)^{64}Zn		577.8	1.0	579.8	0.6	2.0	1	42	32 ^{64}Zn			83Ch47	
^{64}Ga(β^+)^{64}Zn		7072	30	7171.0	1.5	3.3	B					60Ja07	
^{64}Zn(p,n)^{64}Ga		−7951	4	−7953.4	1.5	−0.6	1	14	12 ^{64}Ga	Tex		72Da.A	
^{64}Zn(^{3}He,t)^{64}Ga		−7206	8	−7189.6	1.5	2.0	U			MSU		74Ro16	*
^{64}Zn(^{3}He,t)^{64}Ga^{i}		−9141	17	−9096.7	2.5	2.6	U			MIT		70Hi06	
		−9110	6			2.2	1	17	17 ^{64}Ga^{i}			71Be29	*
^{64}Ga^{i}(IT)^{64}Ga		1905.1	2.3	1907.1	2.2	0.9	1	88	83 ^{64}Ga^{i}			74Ro16	
		1923	4			−4.0	B					19Di08	*
^{64}Ge(β^+)^{64}Ga		4410	250	4517	4	0.4	U					73Da01	*
^{64}As^{i}(IT)^{64}As		1956.5	0.2	1956.50	0.20	0.0	1	100	92 ^{64}As			25Ag12	
*^{64}V−u	Trends in Mass Surface TMS suggest ^{64}V 370 keV less bound											GAu	**
*^{64}Mn−u	Original –45270(350) μu or $M-A$=–42170(330) keV											GAu	**
*^{64}Mn−u	Original –46270(350) μu or $M-A$=–43100(330) keV											GAu	**
*^{64}Mn−u	$M-A$=–42430(280) keV for mixture gs+m at 174.5(0.4) (4^+) keV											Nub269	**
*^{64}Ni(^{14}C,^{17}O)^{61}Fe	Cited in reference and confirmed in PrvCom sep 88											84De33	**
*^{64}Ni(^{18}O,^{20}Ne)^{62}Fe	$Q-Q(^{62}\mathrm{Ni}(^{18}\mathrm{O},^{20}\mathrm{Ne}))$=–2843(20), Q(62)=923(4) keV											AHW	**
*^{64}Ni(^{14}C,^{15}O)^{63}Fe	Original –11743(60) reinterpreted as ($3/2^-$) 356.2 level in ^{63}Fe											GAu	**
*^{64}As^{i}(p)^{63}Ge	E_p=1982(30),1582(30) to the gs., 417.5(0.1) keV level											25Ag12	**
*^{64}Zn(^{3}He,t)^{64}Ga	$M-A$=–58819(8); Q rebuilt with Ame1971											GAu	**
*^{64}Zn(^{3}He,t)^{64}Ga^{i}	CDE=9879(6) Q =–9115(6); recalibration +5 keV for ^{58}Ni(p,n)^{58}Cu^{i} from Ame1961											MMC124	**
*^{64}Ga^{i}(IT)^{64}Ga	uncertainty assigned based on the deviation of known states in ^{64}Ga											19Di08	**
*^{64}Ge(β^+)^{64}Ga	E_{β^+}=2960(250) to (1^+) level at 427.03 keV											Ens073	**
^{65}Cr−u		−29286	837	−30280	50	−1.2	B			MT1	1.0	20Me06	
		−30283	48				2			TR1	1.0	22Si20	
^{65}Mn−u		−43900	600	−43980	4	−0.1	U			TO5	1.5	94Se12	
		−43500	500			−0.6	U			TO6	1.5	98Ba.A	
		−43790	330			−0.6	U			MT1	1.0	11Es06	
		−43979.9	20.4			−0.0	U			TR1	1.0	25Ch42	
^{65}Mn−^{85}Rb$_{.765}$		23500.6	4.0				2			MA8	1.0	12Na15	
^{65}Fe−u		−54680	300	−54984	4	−0.7	U			TO3	1.5	90Tu01	*
		−55270	320			0.6	U			TO5	1.5	94Se12	*
		−54290	380			−1.2	U			TO6	1.5	98Ba.A	*
		−54985.9	5.4			0.3	o			MS1	1.0	08Bl05	*
		−54985.5	9.0			0.2	–			TR1	1.0	22Po02	
		−54981.7	8.3			−0.3	–			TR1	1.0	22Po02	*
	ave.	−54983	6			−0.1	1	45	45 ^{65}Fe			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$O_2-{}^{65}Fe_{.492}$		16881.7	2.7	16881.4	2.0	−0.1	1	55	55 ^{65}Fe	MS1	1.0	10Fe01	*
^{65}Co−u		−63537.9	2.3	−63537.9	2.2	−0.0	o			MS1	1.0	08Bl05	
$O_2-{}^{65}Co_{.492}$		21089.9	1.1				2			MS1	1.0	10Fe01	
^{65}Ni−^{85}Rb$_{.765}$		−2438.0	2.4	−2434.1	0.5	1.6	U			MA8	1.0	07Gu09	
^{65}Ni−u		−69936	23	−69914.9	0.5	0.9	U			TR1	1.0	25Ch42	
$C_5\ H_5-{}^{65}Cu$		111384	4	111335.4	0.7	−3.0	B			R10	4.0	74De22	
$C_4\ H_3\ N-{}^{65}Cu$		98800	8	98759.4	0.7	−1.3	U			R10	4.0	74De22	
$C_4\ ^{13}C\ H_4-{}^{65}Cu$		106921	18	106865.3	0.7	−0.8	U			R10	4.0	74De22	
^{49}Ti O−^{65}Cu		15030	10	14989.3	0.7	−1.0	U			R09	4.0	72De11	
^{65}Cu−^{85}Rb$_{.765}$		−4730.6	1.2	−4729.4	0.7	1.0	1	33	33 ^{65}Cu	MA8	1.0	07Gu09	
^{65}Cu−u		−72224	20	−72210.3	0.7	0.7	U			TR1	1.0	25Ch42	
^{65}Ga−^{85}Rb$_{.765}$		215.4	1.5	215.5	0.9	0.1	1	32	32 ^{65}Ga	MA8	1.0	07Gu09	
^{65}Ge−u		−60080	270	−60631.9	2.3	−1.4	U			GA6	1.5	02Li24	
^{65}Ge ^{1}H−$^{12}C_5\ ^{1}H_2$		−68458.9	4.3	−68456.9	2.3	0.5	–			MS1	1.0	07Sc24	
^{65}Ge ^{1}H−$^{12}C_5\ ^{1}H_5$		−91931.6	4.3	−91932.0	2.3	−0.1	–			MS1	1.0	07Sc24	
^{65}Ge ^{1}H−$^{12}C_5\ ^{1}H_2$	ave.	−68458	3	−68456.9	2.3	0.3	1	57	57 ^{65}Ge			average	
^{65}Ge H−^{85}Rb$_{.776}$		15634.4	6.2	15644.3	2.3	1.6	1	14	14 ^{65}Ge	MS1	1.0	07Sc24	*
^{65}Ge O H−^{85}Rb$_{.965}$		27237.1	4.3	27230.7	2.3	−1.5	1	29	29 ^{65}Ge	MS1	1.0	19Sc11	
^{65}As−u		−50389	91	−50280	40	1.2	2			LZ1	1.0	11Tu02	
		−50248	45			−0.6	2			LZ2	1.0	23Wa10	
^{64}Zn H−^{65}Cu		9171	16	9177.4	0.4	0.4	U			RI1	1.0	18Ki21	
^{65}Zn−^{65}Cu		1459	13	1451.1	0.4	−0.6	U			RI1	1.0	18Ki21	
^{65}Ga−^{65}Cu		4945.2	2.1	4944.9	0.7	−0.1	1	12	9 ^{65}Ga	RI1	1.0	18Ki21	
^{65}Ge−^{65}Cu		11592	22	11578.4	2.4	−0.6	U			RI1	1.0	18Ki21	
^{65}Cu−^{64}Ni		−275	40	−177.0	0.7	0.6	U			R10	4.0	74De22	
^{65}Cu−^{63}Cu		−1784	10	−1807.9	0.7	−0.6	U			R10	4.0	74De22	
^{65}Cu(p,α)^{62}Ni		4345	8	4346.6	0.7	0.2	U			MIT		64Sp12	
		4340	10			0.7	U			Min		67Jo03	
		4344.6	1.8			1.1	1	14	10 ^{65}Cu	NDm		76Jo01	
^{62}Ni(α,n)^{65}Zn		−6510	40	−6480.6	0.7	0.7	U			Oak		64St01	
^{65}Cu(d,α)^{63}Ni		9012	40	8959.8	0.7	−1.3	U					67Hj01	
^{63}Cu(t,p)^{65}Cu		9351	25	9344.9	0.7	−0.2	U			Ald		66Bj02	
^{64}Ni(n,γ)^{65}Ni		6097.86	0.20	6098.08	0.14	1.1	2			MMn		77Is01	Z
		6098.28	0.19			−1.0	2			Bdn		06Fi.A	
^{64}Ni(d,p)^{65}Ni		3876	6	3873.51	0.14	−0.4	U			MIT		64Sp12	
		3867	15			0.4	U			Ald		67Te02	
		3870	5			0.7	U			MIT		70An25	
^{65}Cu(t,α)^{64}Ni		12352	11	12360.0	0.7	0.7	U					72He23	
^{65}Cu(γ,n)^{64}Cu		−9896	28	−9910.8	0.7	−0.5	U			Phi		60Ge01	
^{65}Cu(d,t)^{64}Cu		−3650	60	−3653.6	0.7	−0.1	U			ANL		60Ze02	
^{64}Zn(n,γ)^{65}Zn		7979.3	0.8	7979.33	0.17	0.0	U					71Ot01	Z
		7979.2	0.5			0.3	U					75De.A	Z
		7979.28	0.17			0.3	1	98	54 ^{65}Zn	Bdn		06Fi.A	
^{64}Zn(d,p)^{65}Zn		5758	10	5754.76	0.17	−0.3	U			ANL		67Vo05	
^{64}Zn(p,γ)^{65}Ga		3942.0	1.0	3942.5	0.6	0.5	–					75We24	Z
		3943.0	1.0			−0.5	–					87Vi01	
	ave.	3942.5	0.7			−0.0	1	76	59 ^{65}Ga			average	
^{65}Ge(εp)^{64}Zn		2300	100	2236.6	2.2	−0.6	U			ChR		81Ha44	
^{65}As^{i}(p)^{64}Ge		3603	30	3579	8	−0.8	U					93Ba12	
		3564	20			0.7	o					11Ro47	*
		3575	10			0.4	3					14Ro14	
		3588	16			−0.6	3					17GoZT	*
^{65}Ni(β^-)^{65}Cu		2140	10	2138.1	0.7	−0.2	U					64Fr04	
^{65}Zn(β^+)^{65}Cu		1347	2	1351.7	0.4	2.3	U					49Ma57	
		1347	2			2.3	U					53Ba82	
		1347	3			1.6	U					53Pe14	
		1349	3			0.9	U					53Sa26	
		1342	4			2.4	U					53Yu04	
		1346	4			1.4	U					56Av28	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{65}Cu(p,n)^{65}Zn		−2131.4	1.5	−2134.0	0.4	−1.7	U					56Ma14	Z
		−2135.8	2.5			0.7	U					57Be44	Z
		−2135.3	1.8			0.7	U			Tkm		63Ok01	
		−2135.6	1.7			0.9	U			Oak		64Jo11	Z
		−2134.6	0.8			0.7	–			Yal		69Ov01	Z
		−2133.55	0.43			−1.1	–			PTB		89Sc24	
	ave.	−2133.8	0.4			−0.6	1	89	46 ^{65}Zn			average	
^{65}Ga(β^+)^{65}Zn		3277	30	3254.5	0.6	−0.8	U					57Da07	
^{65}Ge(β^+)^{65}Ga		5220	400	6179.1	2.3	2.4	U					58Po79	
$*^{65}$Fe−u	$M-A$=–50740(250) keV for mixture gs+m at 393.64(0.18) keV											Nub269	**
$*^{65}$Fe−u	$M-A$=–51290(280) keV for mixture gs+m at 393.64(0.18) keV											Nub269	**
$*^{65}$Fe−u	$M-A$=–50370(330) keV for mixture gs+m at 393.64(0.18) keV and											Nub269	**
*	assuming ratio R=0.13(6), from half-life=430 ns and TOF=1 μs											GAu	**
$*^{65}$Fe−u	D_M=–54988.9(7.1) μu to ground state, D_M=–54557.0(8.4) to ^{65}Fe^m at 393.6 keV											Nub269	**
$*^{65}$Fe−u	D_M=–54559.0(8.3) μu for ^{65}Fe^m at 393.6 keV; $M-A$=–50821.4(7.7) keV											Nub269	**
$*O_2-^{65}$Fe$_{.492}$	D_M=16883.6(3.6) μu to ground state, D_M=16671.2(4.2) to ^{65}Fe^m at 393.6 keV											Nub269	**
$*^{65}$Ge H−^{85}Rb$_{.776}$	Combining their 3 results yields a precision for ^{65}Ge of 2.4 keV, while											GAu	**
*	their Table III gives 1.2 keV											GAu	**
$*^{65}$As^i(p)^{64}Ge	And E_p=2620(30) to 901.7 level											11Ro47	**
$*^{65}$As^i(p)^{64}Ge	And E_p=2642(15) to 901.7 level											17GoZT	**
^{66}Mn−u		−39860	860	−39453	11	0.5	U			MT1	1.0	11Es06	*
		−39453	28			−0.0	1	16	16 ^{66}Mn	TR1	1.0	25Ch42	
^{66}Mn−^{85}Rb$_{.776}$		28998	12	28998	11	0.0	1	84	84 ^{66}Mn	MA8	1.0	12Na15	
^{66}Fe−u		−52300	700	−53749	4	−1.4	U			TO3	1.5	90Tu01	
		−54020	350			0.5	U			TO5	1.5	94Se12	
		−52800	300			−2.1	U			TO6	1.5	98Ba.A	
		−53935	150			0.8	U			GT1	1.5	04Ma.A	
		−53742.7	10.7			−0.6	1	14	14 ^{66}Fe	TR1	1.0	22Po02	
^{66}Fe−^{28}Si F$_2$		−27482.9	4.4	−27482	4	0.2	1	86	86 ^{66}Fe	MS1	1.0	10Fe01	
^{66}Co−u		−60470	300	−60557	15	−0.2	U			TO5	1.5	94Se12	*
		−59870	290			−1.6	U			TO6	1.5	98Ba.A	*
^{66}Co−O C F$_2$		−52278	15	−52278	15	−0.0	o			MS1	1.0	08Bl05	
		−52278	15				2			MS1	1.0	10Fe01	
^{66}Ni−^{85}Rb$_{.776}$		−2409.5	1.5				2			MA8	1.0	07Gu09	
^{66}Cu−^{85}Rb$_{.776}$		−2680.6	2.2	−2679.8	0.7	0.4	1	10	10 ^{66}Cu	MA8	1.0	07Gu09	
^{66}Zn−u		−73977	29	−73966.0	0.8	0.4	U			TR1	1.0	25Ch42	
^{66}Ge ^{1}H−^{12}C$_5$ ^{1}H$_5$		−97438.0	2.6				2			MS1	1.0	07Sc24	
^{66}As−u		−55290	730	−55851	6	−0.5	U			GA6	1.5	02Li24	
^{66}As−^{85}Rb$_{.776}$		12607	32	12600	6	−0.2	U			MS1	1.0	07Sc24	
^{66}As O−^{85}Rb$_{.965}$		24186.3	6.1				2			MS1	1.0	19Sc11	
^{66}Se−u		−45070	65				2			LZ2	1.0	23Wa10	
^{65}Ge H−^{66}Zn		21182	24	21159.2	2.4	−1.0	U			RI1	1.0	18Ki21	
^{66}Ga−^{66}Zn		5528	11	5556.1	0.9	2.6	F			RI1	1.0	18Ki21	*
^{66}Ge−^{66}Zn		7824	14	7828.1	2.7	0.3	U			RI1	1.0	18Ki21	
^{66}Zn(p,α)^{63}Cu		1544.3	0.8	1544.6	0.7	0.3	1	74	65 ^{66}Zn	NDm		76Jo01	
^{63}Cu(α,n)^{66}Ga		−7670	30	−7502.4	1.1	5.6	B			Oak		64St01	
^{64}Ni(t,p)^{66}Ni		6559	25	6568.6	1.5	0.4	U			Ald		71Da16	
^{64}Zn(t,p)^{66}Zn		10582	15	10556.0	0.9	−1.7	U			Ald		72Hu06	
^{65}Cu(n,γ)^{66}Cu		7065.80	0.12	7065.93	0.09	1.1	–			BNn		83De29	Z
		7066.13	0.15			−1.3	–			Bdn		06Fi.A	
^{65}Cu(d,p)^{66}Cu		4837	8	4841.36	0.09	0.5	U			MIT		64Sp12	
^{65}Cu(n,γ)^{66}Cu	ave.	7065.93	0.09	7065.93	0.09	0.0	1	100	90 ^{66}Cu			average	
^{66}Zn(d,t)^{65}Zn		−4770	60	−4801.2	0.9	−0.5	U			ANL		60Ze02	
^{66}Co(β^-)^{66}Ni		9700	500	9598	14	−0.2	U					88Bo06	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{66}Ni(β^-)^{66}Cu	200	30	251.8	1.5	1.7	U					56Jo20	
^{66}Cu(β^-)^{66}Zn	2650	30	2640.8	0.9	−0.3	U					51Fr19	*
	2650	30			−0.3	U					56Jo20	
^{66}Ga(β^+)^{66}Zn	5175.0	3.0	5175.5	0.8	0.2	U					63Ca03	*
	5175.5	0.8				2					14Se12	
^{66}Zn(^{3}He,t)^{66}Ga^i	−9044	6				2					71Be29	*
^{66}Ge(β^+)^{66}Ga	2490	50	2116.4	2.7	−7.5	F					69Ba31	*
	2420	30			−10.1	F					69Sa08	*
	2100	30			0.5	U					70De39	*
^{66}As(β^+)^{66}Ge	9550	50	9582	6	0.6	U			ANB		79Da.A	
*^{66}Mn−u	$M-A$=–36900(790) keV for mixture gs+m at 464.5 keV (5$^-$)										Nub269	**
*^{66}Co−u	Original –60160(300) μu or $M-A$=–56040(280) keV										GAu	**
*^{66}Co−u	$M-A$=–55480(270) keV for mixture gs+m+n at 175.1 and 642(5) keV										Nub269	**
*	and assuming for first isomer a ratio R=0.5(0.2) to ground state,										GAu	**
*	from half-life=1.21 μs and TOF=1 μs										GAu	**
*^{66}Ga−^{66}Zn	Compromised measurement due to the influence of neighboring ToF peaks										Sar201	**
*^{66}Cu(β^-)^{66}Zn	E_{β^-}=2630(30) 1640(30) to ground state and 2$^+$ level at 1039.2279 keV										Ens104	**
*^{66}Ga(β^+)^{66}Zn	Original E_{β^+}=4153(3) corrected for recoil										14Se12	**
*^{66}Zn(^{3}He,t)^{66}Ga^i	CDE=9813(6) Q =–9049(6); recalibration +5 keV for ^{58}Ni(p,n)^{58}Cu^i from Ame1961										MMC124	**
*^{66}Ge(β^+)^{66}Ga	E_{β^+}=1440(50) to 43.9 level; F : probably distorted by annihilation pile up										AHW	**
*^{66}Ge(β^+)^{66}Ga	E_{β^+}=1370(30) to 43.9 level; F : probably distorted by annihilation pile up										AHW	**
*^{66}Ge(β^+)^{66}Ga	E_{β^+}=1028(30), 668(30), 558(50) to 43.812 1$^+$, 381.859 1$^+$, 536.618 1$^+$ level										Ens104	**
^{67}Mn−u	−36600	670	−36370	40	0.4	o			MT1	1.0	15Me.A	
	−36458	354			0.3	U			MT1	1.0	20Me06	
	−36365	42				2			TR1	1.0	25Ch42	
^{67}Fe−u	−50190	500	−49399	11	1.1	U			TO5	1.5	94Se12	*
	−48450	380			−1.7	U			TO6	1.5	98Ba.A	*
	−49641	440			0.4	U			GT1	1.5	04Ma.A	
	−49580	300			0.6	o			MT1	1.0	11Es06	*
	−48510	460			−1.9	o			MT1	1.0	15Me.A	
	−49220	370			−0.5	U			MT1	1.0	20Me06	*
	−49398.9	10.5				2			TR1	1.0	22Po02	*
^{67}Fe−^{84}Kr$_{.798}$	21554.1	4.1	21226	11	−80.0	B			JY1	1.0	20Ca08	*
^{67}Co−u	−59390	300	−59390	7	−0.0	U			TO5	1.5	94Se12	
	−58730	350			−1.3	U			TO6	1.5	98Ba.A	
^{28}Si F$_2$−^{67}Co$_{.985}$	32232.9	8.0	32232	7	−0.1	2			MS1	1.0	10Fe01	
	32231	13			0.1	2			MS1	1.0	10Fe01	*
^{67}Ni−u	−68370	430	−68431	3	−0.1	U			TO5	1.5	94Se12	*
	−68090	470			−0.5	U			TO6	1.5	98Ba.A	*
^{67}Ni−^{85}Rb$_{.788}$	1079.1	3.1				2			MA8	1.0	07Gu09	
^{67}Cu−^{85}Rb$_{.788}$	−2760.0	1.3	−2760.6	1.0	−0.5	1	54	54 ^{67}Cu	MA8	1.0	07Gu09	
^{67}Zn−u	−72870	43	−72872.2	0.8	−0.1	U			TR1	1.0	25Ch42	
^{67}As−u	−60500	260	−60748.9	0.5	−0.6	U			GA6	1.5	02Li24	
^{67}As−^{86}Kr$_{.779}$	8885.7	2.9	8885.4	0.5	−0.1	U			MS1	1.0	07Sc24	
	8808	30			2.6	U			MS1	1.0	07Sc24	
^{67}As−^{85}Rb$_{.788}$	8762.5	1.7	8760.8	0.5	−1.0	U			MS1	1.0	07Sc24	
	8760.87	0.54			−0.1	1	77	77 ^{67}As	MS1	1.0	19Sc11	
^{67}As O−^{85}Rb$_{.976}$	20258.7	1.0	20258.9	0.5	0.2	1	23	23 ^{67}As	MS1	1.0	19Sc11	
^{67}Se−u	−50006	72	−49975	21	0.4	2			LZ1	1.0	11Tu02	
	−49972.4	21.5			−0.1	2			LZ2	1.0	23Wa10	
^{67}Zn N−^{66}Zn ^{15}N	4060.21	0.32	4058.88	0.25	−1.7	U			H30	2.5	77Ba10	*
^{66}Zn H−^{67}Zn	6741	15	6731.26	0.25	−0.6	U			RI1	1.0	18Ki21	
^{67}Ga−^{67}Zn	1112	11	1074.7	1.2	−3.4	F			RI1	1.0	18Ki21	*
^{67}Ge−^{67}Zn	5587.9	4.9	5590	5	0.3	1	87	87 ^{67}Ge	RI1	1.0	18Ki21	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{67}As−^{67}Zn		12122	27	12123.3	0.9	0.0	U			RI1	1.0	18Ki21	
^{64}Zn(α,n)^{67}Ge		−9240	60	−8977	4	4.4	B			Oak		64St01	
		−8987.5	12.			0.9	1	13	13 ^{67}Ge	ANL		78Mu05	
		−8993	5			3.2	B					79Al04	
^{65}Cu(t,p)^{67}Cu		7716	25	7716.7	1.1	0.0	U			Ald		66Bj02	
^{65}Cu(^{3}He,p)^{67}Zn		8185	40	8258.8	0.9	1.8	U			MIT		74Is01	
^{67}Kr(2p)^{65}Se		1690	17				3					16Go26	
^{66}Zn(n,γ)^{67}Zn		7052.5	0.6	7052.48	0.23	−0.0	–					71Ot01	Z
		7052.5	0.5			−0.0	–					75De.A	Z
		7052.5	0.3			−0.1	–			Bdn		06Fi.A	
^{66}Zn(d,p)^{67}Zn		4827	10	4827.91	0.23	0.1	U			ANL		67Vo05	
		4820	5			1.6	U			MIT		74Is01	
^{67}Zn(d,t)^{66}Zn		−800	60	−795.25	0.23	0.1	U			ANL		60Ze02	
^{66}Zn(n,γ)^{67}Zn	ave.	7052.50	0.24	7052.48	0.23	−0.1	1	98	63 ^{67}Zn			average	
^{67}Ni(β^-)^{67}Cu		3830	90	3577	3	−2.8	U					75Re09	
^{67}Cu(β^-)^{67}Zn		577	8	560.7	0.8	−2.0	U					53Ea11	
		560.3	1.0			0.4	1	69	46 ^{67}Cu	ANB		15Ch57	
^{67}Zn(p,n)^{67}Ga		−1776	5	−1783.4	1.1	−1.5	U			Ric		57Ch30	Y
		−1783.3	1.4			−0.1	1	66	54 ^{67}Ga	Oak		64Jo11	Z
^{67}Ge(β^+)^{67}Ga		4330	100	4206	4	−1.2	U					59Ri35	*
		4370	150			−1.1	U					69Ba07	*
^{67}As(β^+)^{67}Ge		6010	100	6086	4	0.8	U			ANB		80Mu12	
*^{67}Fe−u	Original –50000(500) μu or $M-A$=–46570(470) keV											GAu	**
*^{67}Fe−u	$M-A$=–44930(330) keV for mixture gs+m at 403(9) keV											Nub269	**
*^{67}Fe−u	$M-A$=–45980(250) keV for mixture gs+m at 403(9) keV											Nub269	**
*^{67}Fe−u	$M-A$=–45560(320) keV for mixture gs+m+n at 403(9) and 450#(100#) keV											Nub269	**
*^{67}Fe−u	with laser on and off											22Po02	**
*^{67}Fe−^{84}Kr$_{.798}$	using updated frequency ratio r=0.797874191(49)											20Ca08	**
*^{28}Si F$_2$−^{67}Co$_{.985}$	$M-A$=–54829(12) for ^{67}Co^m at 491.55(0.11) keV											10Fe01	**
*^{67}Ni−u	Original –67840(300) μu or $M-A$=–63190(280) keV											GAu	**
*^{67}Ni−u	$M-A$=–62930(330) keV for mixture gs+m at 1006.6(0.2) keV											Nub269	**
*^{67}Zn N−^{66}Zn ^{15}N	Original 4060.21(0.25); plus syst 0.20 μu estimated by evaluator											GAu	**
*^{67}Ga−^{67}Zn	Compromised measurement due to the influence of neighboring ToF peaks											Sar201	**
*^{67}Ge(β^+)^{67}Ga	E_{β^+}=3140(100) 3180(100) respectively, to 1/2$^-$ level at 166.98 keV											Ens05c	**
^{68}Mn−u		−29748	1406	−31150	30	−1.0	B			MT1	1.0	20Me06	
		−31154	32				2			TR1	1.0	25Ch42	
^{68}Fe−u		−46300	500	−47344	6	−1.4	U			TO6	1.5	98Ba.A	
		−47330	460			−0.0	o			MT1	1.0	11Es06	
		−46830	460			−1.1	o			MT1	1.0	15Me.A	
		−47622	344			0.8	U			MT1	1.0	20Me06	
		−47344.3	6.0				2			TR1	1.0	22Po02	
^{68}Co−u		−55640	350	−55451.2	1.0	0.4	o			TO5	1.5	94Se12	
		−54750	300			−1.6	U			TO6	1.5	98Ba.A	
		−55690	110			0.9	U			GT2	2.5	08Kn.A	*
		−55760	250			1.2	U			MT1	1.0	11Es06	*
O ^{18}O−^{68}Co$_{.5}$		21794.5	2.4	21799.8	0.5	2.2	U			MS1	1.0	18Iz01	
^{68}Co−^{34}S$_2$		8825.2	8.2	8814.8	1.0	−1.3	U			MS1	1.0	18Iz01	
^{68}Ni−u		−68030	930	−68131	3	−0.1	U			TO5	1.5	94Se12	*
		−67530	930			−0.4	U			TO6	1.5	98Ba.A	*
^{68}Ni−^{85}Rb$_{.800}$		2437.0	3.2				2			MA8	1.0	07Gu09	
^{68}Cu−u		−70570	440	−70389.1	1.7	0.3	U			TO6	1.5	98Ba.A	*
^{68}Cu−^{85}Rb$_{.800}$		179.1	1.7				2			MA8	1.0	07Gu09	*
^{68}Zn−u		−75158	29	−75155.7	0.8	0.1	U			TR1	1.0	25Ch42	
^{68}Ga−^{85}Rb$_{.800}$		−1484	37	−1451.5	1.5	0.9	U			MA8	1.0	07Gu09	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{68}Ge−C$_5$ H$_8$		−134496.7	8.6	−134504.9	2.0	−1.0	U			CP1	1.0	04Cl03	
		−134506.3	2.8			0.5	2			CP1	1.0	04Cl03	
		−134503.5	2.9			−0.5	2			CP1	1.0	04Cl03	
^{68}As−u		−63221	107	−63225.9	2.0	−0.0	U			GT1	1.5	01Ha66	
^{68}As−C$_5$ H$_8$		−125839	13	−125826.1	2.0	1.0	U			CP1	1.0	04Cl03	
		−125827.7	9.9			0.2	U			CP1	1.0	04Cl03	
		−125827.1	2.9			0.3	–			CP1	1.0	04Cl03	
		−125824.4	3.1			−0.6	–			CP1	1.0	04Cl03	
	ave.	−125825.8	2.1			−0.1	1	88	88 ^{68}As			average	
C F$_3$−^{68}As$_{1.015}$		59385.8	5.7	59383.7	2.0	−0.4	1	12	12 ^{68}As	MS1	1.0	07Sc24	
^{68}Se−u		−56197	86	−58174.8	0.5	−23.0	F			CS1	1.0	01La31	*
		−57560	1070			−0.4	U			GA6	1.5	02Li24	
		−57900	300			−0.9	U			CS1	1.0	08Go23	
^{68}Se−C$_5$ H$_8$		−120801	31	−120775.0	0.5	0.8	U			CP1	1.0	04Cl03	
C F$_3$−^{68}Se$_{1.015}$		54256.87	0.54				2			MS1	1.0	09Sa12	
^{68}Zn ^{35}Cl−^{66}Zn ^{37}Cl		1757.9	1.0	1760.47	0.26	0.6	U			H18	4.0	64Ba03	
^{68}Co−^{68}Zn		19704.50	0.54				2			JY2	1.0	26Fl.A	
^{68}Co^m−^{68}Zn		19789.9	2.2				2			JY2	1.0	26Fl.A	
^{68}As−^{68}Ge		8698.8	9.9	8678.8	2.8	−2.0	U			CP1	1.0	04Cl03	
^{68}Se−^{68}Ge		13669	27	13729.9	2.1	2.3	U			CP1	1.0	04Cl03	
^{65}Cu(α,n)^{68}Ga		−5800	40	−5823.9	1.5	−0.6	U			Oak		64St01	
^{66}Ni(t,p)^{68}Ni−^{68}Zn()^{70}Zn		−2110	21	−2100	4	0.5	U			Hei		77Bh03	
^{66}Zn(t,p)^{68}Zn		8758	15	8769.00	0.24	0.7	U			Ald		72Hu06	
^{68}Zn(^{14}C,^{15}O)^{67}Ni		−6052	150	−6100	3	−0.3	U			Ors		84De33	
^{67}Zn(n,γ)^{68}Zn		10198.2	0.4	10198.32	0.04	0.3	U					71Ot01	Z
		10198.06	0.22			1.2	U			Bdn		06Fi.A	
		10198.32	0.04			0.0	1	100	99 ^{68}Zn	ILn		15BaZY	
^{68}Zn(d,t)^{67}Zn		−3930	60	−3941.09	0.04	−0.2	U			ANL		60Ze02	
^{68}Cu(β^-)^{68}Zn		4580	60	4440.0	1.8	−2.3	U					64Ba13	
		4590	50			−3.0	B					72Sw01	
^{68}Zn(t,^{3}He)^{68}Cu		−4410	20	−4421.4	1.8	−0.6	U			LAl		77Sh08	
^{68}Ga(β^+)^{68}Zn		2921.1	1.2				2					72Sl03	
		2915	10	2921.1	1.2	0.6	U					85Bo58	
^{68}Zn(p,n)^{68}Ga		−3693	6	−3703.4	1.2	−1.7	U			Ric		55Br16	Z
		−3703	5			−0.1	U			Ric		57Ch30	Z
		−3707	5			0.7	U			Oak		64Jo11	Z
^{68}As(β^+)^{68}Ge		8100	100	8084.3	2.6	−0.2	U			ANB		77Pa13	
		8073	54			0.2	U					02Cl.A	*
^{68}Se(β^+)^{68}As		4710	200	4705.1	1.9	−0.0	U					04Wo16	
*^{68}Co−u	$M-A$=–51838(96) keV for mixture gs+m at 79.5(2.1) keV											Nub269	**
*^{68}Co−u	$M-A$=–51860(210) keV for mixture gs+m at 79.5(2.1) keV											Nub269	**
*^{68}Ni−u	$M-A$=–61950(280) keV for mixture gs+n at 2849.1 keV											Nub269	**
*^{68}Ni−u	$M-A$=–61480(280) keV for mixture gs+n at 2849.1 keV											Nub269	**
*^{68}Cu−u	$M-A$=–65380(350) keV for mixture gs+m at 721.26 keV											Nub269	**
*^{68}Cu−^{85}Rb$_{.800}$	This result was first published in reference											04Bl16	**
*^{68}Cu−^{85}Rb$_{.800}$	Also 948.6(1.6) μu for ^{68}Cu^m−^{85}Rb$_{.800}$, yielding excit. of 716.7(2.2) keV											07Gu09	**
*^{68}Se−u	F : other results in same paper not trusted, see ^{80}Y and ^{80}Zr											GAu	**
*^{68}As(β^+)^{68}Ge	From mass difference 8667(64) μu											02Cl.A	**
^{69}Fe−u		−42240	640	−42409	12	−0.3	o			MT1	1.0	15Me.A	
		−43232	429			1.9	B			MT1	1.0	20Me06	
		−42409.3	11.8				2			TR1	1.0	22Po02	
^{69}Fe^m−u		−42170.6	9.8				2			TR1	1.0	22Po02	
^{69}Co−u		−54800	400	−54071	15	1.2	o			TO5	1.5	94Se12	
		−53050	300			−2.3	U			TO6	1.5	98Ba.A	
		−54070	230			−0.0	U			MT1	1.0	11Es06	
		−54117	223			0.2	U			LZ1	1.0	15Xu14	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{69}Co−^{84}Kr$_{.821}$	18570	92	18589	15	0.2	U			JY1	1.0	20Ca08	*
^{69}Co^{m}−^{39}K$_{1.769}$	10296	15	10297	14	0.1	1	93	93 ^{69}Co^{m}	MS1	1.0	18Iz01	*
^{69}Co^{m}−^{84}Kr$_{.821}$	18765	54	18754	14	−0.2	1	7	7 ^{69}Co^{m}	JY1	1.0	20Ca08	*
^{69}Ni−u	−64600	400	−64390	4	0.4	U			TO5	1.5	94Se12	*
	−64250	450			−0.2	U			TO6	1.5	98Ba.A	*
^{69}Ni−^{85}Rb$_{.812}$	7237.0	4.0				2			MA8	1.0	07Gu09	
^{69}Cu−^{85}Rb$_{.812}$	1056.0	1.5				2			MA8	1.0	07Gu09	
^{69}Zn−u	−73580	400	−73449.6	0.8	0.2	U			TO6	1.5	98Ba.A	*
C$_5$ H$_9$−^{69}Ga	144852.7	2.4	144851.8	1.2	−0.1	U			M15	2.5	63Ri07	
^{69}Ga−^{85}Rb$_{.812}$	−2799.8	1.6	−2799.8	1.2	−0.0	1	57	57 ^{69}Ga	MA8	1.0	07Gu09	
^{69}As−u	−67757.8	23.6	−67780	3	−0.9	U			GR1	1.0	21Ma22	
	−67780.1	3.1				2			LZ1	1.0	23Xi01	
C F$_3$−^{69}Se	55794.7	1.6				2			MS1	1.0	07Sc24	
^{69}Co−^{69}Co^{m}	−164.6	5.0				2			JY2	1.0	26Fl.A	
^{69}Co^{m}−^{69}Ga	20564.7	4.5	20520	15	−9.9	B			JY2	1.0	26Fl.A	
^{69}Ga(p,α)^{66}Zn	4440	10	4435.1	1.3	−0.5	U			ANL		67Ka11	
^{66}Zn(α,n)^{69}Ge	−7520	30	−7444.6	1.5	2.5	U			Oak		64St01	
^{67}Zn(t,p)^{69}Zn	8168	20	8198.67	0.05	1.5	U			Ald		72Hu06	
^{68}Zn(n,γ)^{69}Zn	6482.3	0.8	6482.15	0.03	−0.2	U					71Ot01	Z
	6481.8	0.5			0.7	U					75De.A	Z
	6482.07	0.16			0.5	U			Bdn		06Fi.A	
	6482.15	0.03				2			ILn		15BaZY	
^{68}Zn(d,p)^{69}Zn	4259	10	4257.58	0.03	−0.1	U			ANL		67Vo05	
	4243	10			1.5	U			MIT		75Is04	
^{68}Zn(^{3}He,d)^{69}Ga	1126	20	1116.3	1.3	−0.5	U					74Ri08	
^{69}Se(εp)^{68}Ge	3390	50	3255.1	2.4	−2.7	U			ChR		76Ha29	
	3370	70			−1.6	U			ChR		77Ma24	
^{69}Br(p)^{68}Se	789	37	640	40	−4.0	B			MSU		11Ro18	*
	641	42				3			MSU		14De41	*
^{69}Br^{i}(p)^{68}Se	4131	50	3804	14	−6.5	B					97Xu01	*
	3812	19			−0.4	3			MSU		11Ro47	*
	3793	22			0.5	3			MSU		14De41	*
^{69}Cu(β^{-})^{69}Zn	2480	70	2681.7	1.6	2.9	U					66Va12	
^{69}Zn(β^{-})^{69}Ga	897	5	910.0	1.3	2.6	U					53Du03	
^{69}Ge(β^{+})^{69}Ga	2225	15	2227.2	0.6	0.1	U					51Hu38	*
^{69}Ga(p,n)^{69}Ge	−3008.8	3.2	−3009.5	0.6	−0.2	U			Tkm		63Ok01	
	−3006.0	4.			−0.9	U			Oak		64Jo11	Z
	−3009.50	0.55				2			PTB		92Bo.B	Z
^{69}As(β^{+})^{69}Ge	3972	50	3964	3	−0.2	U					70Bo19	
	4067	50			−2.1	U			ChR		77Ma24	*
^{69}Se(β^{+})^{69}As	6817	75	6702	3	−1.5	U			ChR		77Ma24	*
*^{69}Co−^{84}Kr$_{.821}$	using the revised frequency ratio r=0.82164916(110)										20Ca08	**
*^{69}Co^{m}−^{39}K$_{1.769}$	Re-assigned the gs. to the short-lived iso. state suggested by 20Ca08										20Ca08	**
*^{69}Co^{m}−^{84}Kr$_{.821}$	using updated frequency ratio r=0.82165149(64)										20Ca08	**
*^{69}Ni−u	$M-A$=–59940(330) keV for mixture gs+m+n at 321(2) and 2700.1 keV										Nub269	**
*^{69}Ni−u	$M-A$=–59620(380) keV for mixture gs+m+n at 321(2) and 2700.1 keV										Nub269	**
*	and assuming for second isomer a ratio R=0.13(0.06) to ground state,										GAu	**
*	from half-life=439 ns and TOF=1 μs										GAu	**
*^{69}Zn−u	$M-A$=–68320(350) keV for mixture gs+m at 438.636 keV										Nub269	**
*^{69}Br(p)^{68}Se	Symmetrized from Q_p=785(+34–40) keV. E_p from excited (3/2^{-}) level										FKo248	**
*^{69}Br(p)^{68}Se	also Q_p=751(+132–82) from excited (3/2^{-}) level										14De41	**
*^{69}Br^{i}(p)^{68}Se	Q_p=2970(50) to (2^{+}) level at 853.75 (p-g coin) and Q_p=3810(20) to ground state										11Ro47	**
*^{69}Br^{i}(p)^{68}Se	Q_p=2939(22) to (2^{+}) level at 853.75 from p-g coin										14De41	**
*^{69}Ge(β^{+})^{69}Ga	$E_{\beta^{+}}$=1215, 610 to ground state 3/2^{-}, 574.220 5/2^{-} levels										Ens141	**
*^{69}As(β^{+})^{69}Ge	$E_{\beta^{+}}$=2812(50) to 3/2^{-} level at 232.70 keV										Ens141	**
*^{69}Se(β^{+})^{69}As	$E_{\beta^{+}}$=5006(75) to 789.46 (1/2^{-},3/2^{-}) level, and others										Ens141	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{70}Fe−u	−40483	526	−39778	13	1.3	B			MT1	1.0	20Me06	
	−39778.0	12.9				2			TR1	1.0	22Po02	
^{70}Co−u	−49000	600	−49929.2	2.1	−1.0	U			TO6	1.5	98Ba.A	
	−50410	310			1.6	U			MT1	1.0	11Es06	*
^{70}Co−^{84}Kr$_{.833}$	23776	13	23793.2	2.1	1.3	U			JY1	1.0	20Ca08	*
^{70}Ni−u	−63980	350	−63568.7	2.3	0.8	U			TO5	1.5	94Se12	*
	−63020	350			−1.0	U			TO6	1.5	98Ba.A	*
^{70}Cu−^{85}Rb$_{.824}$	5077.6	1.7	5077.3	1.2	−0.2	2			MA8	1.0	07Gu09	*
	5077.2	2.2			0.1	2			MA8	1.0	07Gu09	*
	5077.0	2.3			0.1	2			MA8	1.0	07Gu09	*
^{70}Ga−^{85}Rb$_{.824}$	−1293.0	2.3	−1292.9	1.2	0.0	1	28	28 ^{70}Ga	MA8	1.0	07Gu09	
C_5 H_{10}−^{70}Ge	154001.3	2.2	154001.7	0.9	0.1	U			M15	2.5	63Ri07	
C_4 H_6 O−^{70}Ge	117616.1	1.8	117616.2	0.9	0.0	U			M15	2.5	63Ri07	
^{70}As−^{85}Rb$_{.824}$	3619.9	1.5				2			MA8	1.0	20Ku19	
^{70}Se−u	−66890	490	−66483.4	1.5	0.6	o			GA6	1.5	98Ch20	
	−66635	75			1.3	U			GT1	1.5	01Ha66	
	−66520	140			0.2	U			GA6	1.5	02Li24	
	−66480.3	2.8			−1.1	1	27	27 ^{70}Se	GR1	1.0	21Ma22	
^{70}Se−^{13}C F_3	−65048.8	1.7	−65047.7	1.5	0.7	1	73	73 ^{70}Se	MS1	1.0	09Sa12	
^{70}Se−^{85}Rb$_{.824}$	6209	18	6201.9	1.5	−0.4	U			MA8	1.0	11He10	
^{70}Br−^{13}C F_3	−53771.7	16.1	−54318#	17#	−33.9	B			MS1	1.0	09Sa12	*
^{70}Kr−u	−44360	150				2			LZ2	1.0	23Wa10	
^{70}Ni−^{72}Ge$_{.972}$	12173.6	2.3				2			JY1	1.0	07Ra27	
^{70}Zn ^{35}Cl−^{68}Zn ^{37}Cl	3429.5	1.7	3424.7	1.8	−0.7	1	7	6 ^{70}Zn	H18	4.0	64Ba03	
^{70}Co−^{70}Zn	24751.9	1.3				2			JY2	1.0	26Fl.A	
^{70}Co^{m}−^{70}Zn	25040.6	4.8				2			JY2	1.0	26Fl.A	
^{70}Zn(^{3}He,^{8}B)^{65}Co	−18385	13	−18370.2	2.8	1.1	U			Pri		78Ko24	
^{70}Zn(α,^{7}Be)^{67}Ni	−19155	36	−19166	3	−0.3	U			Tex		78Co.A	
	−19164	22			−0.1	U			Pri		78Ko28	
^{70}Zn(^{14}C,^{17}O)^{67}Ni	−1661	100	−1994	3	−3.3	B			Ors		88Gi04	
^{70}Ge(p,α)^{67}Ga	1180.9	1.5	1181.0	1.2	0.1	1	60	46 ^{67}Ga	NDm		76Jo01	
^{70}Ge(^{3}He,^{6}He)^{67}Ge	−10572	30	−10549	4	0.8	U			MSU		78Pa11	
^{70}Zn(^{14}C,^{16}O)^{68}Ni	1727	30	1656	3	−2.4	U			Ors		88Gi04	
^{70}Zn(^{18}O,^{20}Ne)^{68}Ni	172	26	158	3	−0.5	U			Hei		84Ha31	
^{68}Zn(t,p)^{70}Zn	7196	15	7218.8	1.7	1.5	U			Ald		72Hu06	
^{70}Ge(p,t)^{68}Ge	−11251	13	−11244.0	2.0	0.5	U			ChR		72Hs01	
	−11242	7			−0.3	U			Ors		77Gu02	
^{70}Zn(^{14}C,^{15}O)^{69}Ni	−8936	150	−9422	4	−3.2	B			Ors		84De33	
^{70}Zn(d,^{3}He)^{69}Cu	−5605	10	−5624.3	2.1	−1.9	U			ANL		78Ze04	
	−5622	13			−0.2	U			Hei		84Ha31	
^{70}Zn(t,α)^{69}Cu	8682	20	8696.1	2.1	0.7	U			LAl		81Aj02	
^{69}Ga(n,γ)^{70}Ga	7654.0	1.0	7653.65	0.17	−0.3	U					71Ar12	Z
	7651.6	1.0			2.1	F					71Ve03	*
	7653.65	0.17			0.0	1	99	56 ^{70}Ga	Bdn		06Fi.A	
^{69}Ga(d,p)^{70}Ga	5430	10	5429.08	0.17	−0.1	U			Kop		71Ar12	
^{70}Ge(d,^{3}He)^{69}Ga	−3030	7	−3029.6	1.4	0.1	U			Ors		78Ro14	
^{70}Cu(β^-)^{70}Zn	6310	110	6588.6	1.9	2.5	U					75Re09	*
	5928	110			6.0	B					75Re09	*
^{70}Zn(t,^{3}He)^{70}Cu	−6559	20	−6570.0	1.9	−0.6	U			LAl		77Sh08	
	−6602	20			1.6	U			LAl		87Aj.A	
^{70}Zn(p,n)^{70}Ga	−1436.3	2.0	−1437.1	1.4	−0.4	–			Nvl		59Go68	Z
	−1439.1	3.0			0.7	–			Oak		64Jo11	Z
ave.	−1437.2	1.6			0.0	1	71	55 ^{70}Zn			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{70}Ga(β^-)^{70}Ge		1650	10	1651.8	1.4	0.2	U					57Bu41	
^{70}As(β^+)^{70}Ge		6220	50	6228.0	1.6	0.2	U					63Bo14	*
^{70}Se(β^+)^{70}As		2780	200	2405.1	1.9	−1.9	F					75La02	*
		2736	85			−3.9	B					01To06	
^{70}Br(β^+)^{70}Se		9970	170	9995#	16#	0.1	U			ANB		79Da.A	
		9898	80			1.2	U					04Ka38	*
*^{70}Co−u	$M-A$=–46820(280) keV for mixture gs+m at 269(5) keV											Nub269	**
*^{70}Co−^{84}Kr$_{.833}$	using the revised frequency ratio r=0.83361594(15)											20Ca08	**
*^{70}Ni−u	Original –63860(350) μu or $M-A$=–59490(330) keV											GAu	**
*^{70}Ni−u	$M-A$=–58590(330) keV for mixture gs+m at 2860.9(0.1) keV and											Nub269	**
*	assuming ratio R=0.04(2), from half-life=232 ns and TOF=1 μs											GAu	**
*^{70}Cu−^{85}Rb$_{.824}$	The three results for ^{70}Cu were first published in reference											04Va07	**
*^{70}Cu−^{85}Rb$_{.824}$	D_M=5185.7(2.2) μu for ^{70}Cu^m at 101.1(0.3) keV; $M-A$=–62875.4(2.0) keV											Nub269	**
*^{70}Cu−^{85}Rb$_{.824}$	D_M=5337.4(2.3) μu for ^{70}Cu^n at 242.6(0.5) keV; $M-A$=–62734.1(2.2) keV											Nub269	**
*^{70}Br−^{13}C F$_3$	D_M=–51310.8(16.1) μu for ^{70}Br^m at 2292.3(0.8) keV											Nub269	**
*^{70}Br−u	estimated from systematics of CDE											23Hu18	**
*^{69}Ga(n,γ)^{70}Ga	F : E(γ) systematically lower than for other authors; Z recalibrated											AHW	**
*^{70}Cu(β^-)^{70}Zn	E_{β^-}=4550(120), 3370(170) to 4$^+$ level at 1786.33, 5$^-$ at 3037.61 keV											Ens04c	**
*^{70}Cu(β^-)^{70}Zn	E_{β^-}=6170(110) from ^{70}Cu^n 1$^+$ at 242.6 keV											Nub269	**
*^{70}As(β^+)^{70}Ge	E_{β^+}=2144(50) to 3$^+$ level at 3046.427, 4$^+$ at 3058.707 keV											Ens04c	**
*^{70}Se(β^+)^{70}As	E_{β^+}=1500(200) to 1$^+$ level at 81.49, 1$^+$ at 234.70, 1$^+$ at 458.12 keV											Ens04c	**
*^{70}Se(β^+)^{70}As	F : author's half-life 20(2)m disagrees with Nubase 41.1(0.3)m											Nub269	**
*^{70}Br(β^+)^{70}Se	Q_{β^+}=12190(80) from 2292.3 ^{70}Br^m											04Ka38	**
^{71}Co−u		−47100	600	−47630	500	−0.6	2			TO6	1.5	98Ba.A	
		−47870	600			0.4	2			MT1	1.0	11Es06	
^{71}Ni−u		−60000	400	−59481.0	2.4	0.9	U			TO5	1.5	94Se12	
		−58700	350			−1.5	U			TO6	1.5	98Ba.A	
^{71}Cu−^{85}Rb$_{.835}$		6332.4	1.6				2			MA8	1.0	07Gu09	
^{71}Zn−u		−72080	380	−72277.7	1.6	−0.3	U			TO6	1.5	98Ba.A	*
^{71}Zn^m−^{85}Rb$_{.835}$		1544.3	2.6	1544.9	1.6	0.2	1	39	39 ^{71}Zn^m	MA8	1.0	08Ba54	
C$_5$ H$_{11}$−^{71}Ga		161370.2	3.2	161372.7	0.9	0.3	U			M15	2.5	63Ri07	
^{71}Ga−^{85}Rb$_{.835}$		−1641.6	3.0	−1641.8	0.9	−0.1	–			MA8	1.0	07Gu09	
		−1640.2	1.3			−1.2	–			MA8	1.0	07Ke09	
	ave.	−1640.4	1.2			−1.2	1	53	53 ^{71}Ga			average	
^{71}Se−u		−68160	340	−67791	3	0.7	o			GA6	1.5	98Ch20	
		−67687	75			−0.9	U			GT1	1.5	01Ha66	
		−67830	120			0.2	U			GA6	1.5	02Li24	
		−67791.1	24.7			0.0	U			GR1	1.0	21Ma22	
^{71}Se−^{85}Rb$_{.835}$		5865.0	3.0				2			MA8	1.0	11He10	
^{71}Br−u		−61260	610	−60658	6	0.7	U			GA6	1.5	02Li24	
		−60634.9	17.2			−1.3	U			GR1	1.0	21Ma22	
^{71}Br H$_2$−C$_4$ H$_9$ O		−110347.7	5.8				2			MS1	1.0	09Sa12	
^{71}Kr−u		−49727	151	−49443	26	1.9	U			LZ1	1.0	11Tu02	
		−49443.2	25.8				2			LZ2	1.0	23Wa10	
^{71}Ni−^{72}Ge$_{.986}$		17352.2	2.4				2			JY1	1.0	07Ra27	
^{71}Ga−^{71}Ge		−249.538	0.106	−249.57	0.10	−0.3	1	89	77 ^{71}Ge	JY1	1.0	16Al30	
^{68}Zn(α,n)^{71}Ge		−5630	40	−5746.9	1.0	−2.9	U			Oak		64St01	
^{70}Zn(^{18}O,^{17}F)^{71}Cu		−9529	35	−9588.4	2.1	−1.7	U			Ber		89Bo.A	
^{70}Zn(n,γ)^{71}Zn		5832.60	0.04	5832.60	0.04	0.0	–			ILn		15BaZY	
^{70}Zn(d,p)^{71}Zn		3609	10	3608.03	0.04	−0.1	–			ANL		67Vo05	
^{70}Zn(n,γ)^{71}Zn	ave.	5832.60	0.04	5832.60	0.04	0.0	1	100	61 ^{71}Zn			average	
^{70}Zn(^{3}He,d)^{71}Ga		2380	20	2369.6	1.7	−0.5	U					74Ri08	
^{71}Ga(γ,n)^{70}Ga		−9240	60	−9300.2	1.4	−1.0	U			Phi		60Ge01	
^{71}Ga(d,t)^{70}Ga		−3054	10	−3042.9	1.4	1.1	U			Kop		71Ar12	
^{70}Ge(n,γ)^{71}Ge		7415.3	1.5	7415.94	0.11	0.4	U					70Or.A	
		7415.1	2.			0.4	U					72Gr34	
		7415.95	0.15			−0.1	–			MMn		91Is01	Z
		7415.93	0.15			0.1	–			Bdn		06Fi.A	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{70}Ge(d,p)^{71}Ge		5182	10	5191.37	0.11	0.9	U			Kyu		73Ka03	
^{70}Ge(n,γ)^{71}Ge	ave.	7415.94	0.11	7415.94	0.11	−0.0	1	100	86 ^{70}Ge			average	
^{70}Ge(p,γ)^{71}As		4619	5	4620	4	0.2	R					75Li14	
^{71}Zn^{m}(IT)^{71}Zn		155.62	0.06	155.62	0.06	−0.0	1	100	61 ^{71}Zn^{m}			15BaZY	
^{71}Zn(β^-)^{71}Ga		2610	50	2812.8	1.7	4.1	B					61Th01	
		2786	50			0.5	U					61Th01	*
		2796	50			0.3	U					64So01	*
^{71}Ge(ε)^{71}Ga		233.0	0.5	232.47	0.09	−1.1	U			Hei		84Ha.A	
		229.3	1.0			3.2	F					91Zl01	*
		232.1	0.5			0.7	U					93Di03	*
		232.71	0.29			−0.8	1	10	9 ^{71}Ge			95Le19	
		233.5	1.2			−0.9	U			TT1		13Fr13	
^{71}Ga(p,n)^{71}Ge		−1018.4	2.0	−1014.82	0.09	1.8	U			Oak		64Jo11	Z
^{71}Ga(^{3}He,t)^{71}Ge−^{65}Cu()^{65}Zn		1122.0	0.9	1119.2	0.4	−3.1	B			Pri		84Ko10	
^{71}As(β^+)^{71}Ge		1997	20	2013	4	0.8	U					53St31	*
		2010	10			0.3	2					54Th36	*
		2012	10			0.1	2					55Gr08	*
^{71}Se(β^+)^{71}As		4428	125	4747	5	2.5	U					73Sc17	
		4762	35			−0.4	U					01To06	
^{71}Kr(ε)^{71}Br		10140	320	10446	25	1.0	U					97Oi01	
*^{71}Zn−u	$M-A$=–67060(350) keV for mixture gs+m at 155.62 keV											Nub269	**
*^{71}Zn(β^-)^{71}Ga	E_{β^-}=1450(50) 1460(50) respectively, from ^{71}Zn^{m} at 157.7(1.3) to 9/2$^+$ at 1493.74											Ens10c	**
*^{71}Ge(ε)^{71}Ga	F : sees 17 keV neutrino											AHW	**
*^{71}Ge(ε)^{71}Ga	Original error 0.1 increased for calibration uncertainty											GAu	**
*^{71}As(β^+)^{71}Ge	E_{β^+}=800(20) 813(10) 815(10) respectively, to 5/2$^-$ level at 174.943 keV											Ens10c	**
^{72}Ni−u		−58700	500	−58214.1	2.4	0.6	U			TO5	1.5	94Se12	
		−57400	400			−1.4	U			TO6	1.5	98Ba.A	
^{72}Cu−u		−64250	510	−64179.7	1.5	0.1	U			TO6	1.5	98Ba.A	*
^{72}Cu−^{85}Rb$_{.847}$		10534.4	1.5				2			MA8	1.0	07Gu09	
^{72}Zn−^{85}Rb$_{.847}$		1556.9	2.3				2			MA8	1.0	08Ba54	
^{72}Ga−^{85}Rb$_{.847}$		1079.5	1.5	1081.6	0.9	1.4	1	34	34 ^{72}Ga	MA8	1.0	07Gu09	
C_4 H_8 O−^{72}Ge		135438.4	2.1	135439.05	0.08	0.1	U			M15	2.5	63Ri07	
^{72}Ge−u		−77906	25	−77924.18	0.08	−0.7	U			MR1	1.0	15Wi.A	
		−77927	22			0.1	U			MR1	1.0	15Wi.A	
		−77898	20			−1.3	U			MR1	1.0	15Wi.A	
^{72}Se−^{85}Rb$_{.847}$		1854.6	2.1				2			MA8	1.0	11He10	
^{72}Br ^{27}Al−^{85}Rb$_{1.165}$		20892.1	7.2	20898.0	1.1	0.8	U			MA8	1.0	11He10	
^{72}Br−^{85}Rb$_{.847}$		11308.7	1.1				2			MS1	1.0	15Va05	*
^{72}Kr−^{85}Rb$_{.847}$		16806.5	8.6				2			MA8	1.0	06Ro11	
^{70}Ge H_2−^{72}Ge		17821.3	1.7	17822.8	0.9	0.4	U			M15	2.5	63Ri07	
^{72}Ge ^{35}Cl−^{70}Ge ^{37}Cl		779.8	5.9	777.4	0.9	−0.2	U			H40	2.5	85El01	
^{72}Ni−^{72}Ge		19710.1	2.4				2			JY1	1.0	07Ra27	
^{72}As−^{72}Ge		4663.037	0.079				2			JY2	1.0	21Ge04	
^{70}Zn(t,p)^{72}Zn		6231	20	6241.3	2.6	0.5	U			Ald		72Hu06	
^{71}Ga(n,γ)^{72}Ga		6521.1	1.0	6520.48	0.19	−0.6	U					70Li04	Z
		6519.8	1.0			0.7	F					71Ve03	*
		6520.44	0.19			0.2	1	99	66 ^{72}Ga	Bdn		06Fi.A	
^{72}Ge(d,^{3}He)^{71}Ga		−4241	7	−4242.3	0.8	−0.2	U			Ors		78Ro14	
^{72}Zn(β^-)^{72}Ga		422	20	442.7	2.3	1.0	U					63De11	*
		458	6			−2.5	B					63Th03	*
^{72}Ga(β^-)^{72}Ge		4000	20	3997.7	0.8	−0.1	U					55Jo09	*
		3984	10			1.4	U					60La04	*
^{72}As(β^+)^{72}Ge		4361	10	4343.59	0.07	−1.7	U					50Me55	
		4345	10			−0.1	U					68Vi05	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{72}Ge(p,n)^{72}As	−5140	5	−5125.94	0.07	2.8	U			Kyu		76Ki12	
^{72}Br(β^+)^{72}Se	8869	95	8806.4	2.2	−0.7	U					01To06	
^{72}Kr(β^+)^{72}Br	5040	80	5121	8	1.0	U					73Sc17	*
*^{72}Cu−u	$M-A$=–59710(470) keV for mixture gs+m at 270.3 keV										Nub269	**
*^{72}Br−^{85}Rb$_{.847}$	D_M=11308.4(1.1) μu, 11417.2(1.2) for ground state and ^{72}Br^m at 100.76(0.15) keV										Nub269	**
*	respectively $M-A$=–59062.0(1.0) and –58960.7(1.1) keV										15Va05	**
*^{71}Ga(n,γ)^{72}Ga	F : E(γ) systematically lower than for other authors; Z recalibrated										AHW	**
*^{72}Zn(β^-)^{72}Ga	E_{β^-}=260(20) 296(6) respectively, to 1$^+$ level at 161.53 keV										Ens102	**
*^{72}Ga(β^-)^{72}Ge	E_{β^-}=3166(20) 3150(10) respectively, to 2$^+$ level at 834.01 keV										Ens102	**
*^{72}Kr(β^+)^{72}Br	E_{β^+}=3794(180), 3626(105), 3682(80), 3364(155) to 162.67, 309.84 1$^+$,										73Sc17	**
*	415.05 1$^+$ and 576.74 1$^+$ levels										Ens102	**
^{73}Ni−u	−52500	500	−53792.9	1.7	−1.7	U			TO6	1.5	98Ba.A	
^{73}Cu−u	−62740	350	−63325.9	1.6	−1.1	U			TO6	1.5	98Ba.A	
^{73}Cu−^{85}Rb$_{.859}$	12447.9	4.2	12446.7	1.6	−0.3	1	15	15 ^{73}Cu	MA8	1.0	07Gu09	
^{73}Zn−u	−70100	380	−70417.4	2.0	−0.6	U			TO6	1.5	98Ba.A	*
^{73}Zn−^{85}Rb$_{.859}$	5355.2	2.0				2			MA8	1.0	08Ba54	
^{73}Ga−^{85}Rb$_{.859}$	947.3	1.8				2			MA8	1.0	07Gu09	
C$_4$ H$_9$ O−^{73}Ge	141878.4	2.1	141880.95	0.06	0.5	U			M15	2.5	63Ri07	
^{73}Se−^{85}Rb$_{.859}$	2511	11	2528	8	1.5	1	52	52 ^{73}Se	MA8	1.0	11He10	*
^{73}Br−u	−68428	97	−68323	6	0.7	U			GT1	1.5	01Ha66	
	−68315.3	19.4			−0.4	–			MR1	1.0	20Ku19	
	−68316.5	8.9			−0.7	–			LZ1	1.0	23Xi01	*
ave.	−68316	8			−0.8	1	48	48 ^{73}Br			average	
^{73}Br ^{27}Al−^{85}Rb$_{1.176}$	16945.3	7.8	16951	6	0.7	1	52	52 ^{73}Br	MA8	1.0	11He10	
^{73}Kr−^{85}Rb$_{.859}$	15061.8	7.1	15062	7	0.0	o			MA8	1.0	04Ro32	*
	15062.8	9.7			−0.1	2			MA8	1.0	06Ro11	
	15060.7	10.3			0.1	2			MA8	1.0	06Ro11	
^{73}Ni−^{72}Ge$_{1.014}$	25221.8	2.6	25222.2	1.7	0.2	1	41	40 ^{73}Ni	JY1	1.0	07Ra27	
^{73}Cu−^{72}Ge$_{1.014}$	15689.2	2.4	15689.2	1.6	0.0	1	47	47 ^{73}Cu	JY1	1.0	07Ra27	
^{72}Ge H−^{73}Ge	6443.9	1.3	6441.90	0.05	−0.6	U			M15	2.5	63Ri07	
^{73}Ni−^{73}Cu	9533.04	0.51	9533.0	0.5	−0.0	1	98	60 ^{73}Ni	JY2	1.0	24Fl.A	
	9538	24			−0.2	U			RKM	1.0	25Xi16	
^{73}Br−^{72}Br	−4610	330	−4917	6	−0.4	U			CR1	2.5	89Sh10	*
	−4709	166			−0.8	U			CR2	1.5	91Sh19	*
^{72}Ge(n,γ)^{73}Ge	6783.4	0.9	6782.94	0.05	−0.5	U					72Gr34	
	6780.9	2.			1.0	U					72Ha74	
	6782.94	0.05			0.0	1	100	100 ^{72}Ge	MMn		91Is01	Z
	6783.12	0.15			−1.2	U			Bdn		06Fi.A	
^{72}Ge(d,p)^{73}Ge	4571	4	4558.37	0.05	−3.2	B			Ald		69He05	
	4563	10			−0.5	U			Kop		72Ha74	
	4541	10			1.7	U			Kyu		73Ka03	
^{72}Ge(^{3}He,d)^{73}As	160	4	162	4	0.6	1	93	93 ^{73}As	Hei		76Sc13	
^{73}Kr(εp)^{72}Se	3700	150	4027	7	2.2	U			ChR		81Ha44	
^{73}Rb(p)^{72}Kr	640	40				3					20Ho17	
^{73}Rb^i(p)^{72}Kr	3803	40	3843	18	1.0	3					93Ba61	
	3853	20			−0.5	3					20Ho06	
^{73}Ga(β^-)^{73}Ge	1554	40	1598.2	1.7	1.1	U					58Yt22	*
	1564	80			0.4	U					70Wa21	*
^{73}Ge(p,n)^{73}As	−1121.6	15.	−1127	4	−0.4	U			Oak		64Jo11	*
^{73}Se(β^+)^{73}As	2740	10	2725	7	−1.5	1	55	48 ^{73}Se			56Ha10	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{73}Br(β^+)^{73}Se	4748	500	4585	9	−0.3	U					70Mu02	*
	4648	400			−0.2	U					74Ro11	*
	4688	140			−0.7	U					87He21	*
	4610	70			−0.4	U					01To06	
^{73}Kr(β^+)^{73}Br	6790	350	7090	8	0.9	U					73Sc17	*
	6860	220			1.0	U					97Oi01	
*^{73}Zn−u	$M-A$=–65200(350) keV for mixture gs+m at 195.5 keV										Nub269	**
*^{73}Se−^{85}Rb$_{.859}$	D_M=2524.6(7.3) μu for mixture gs+m at 25.71 keV; $M-A$=–68230.0(6.8) keV										Nub269	**
*^{73}Br−u	$M-A$=–63622.9(2.7) keV for mixture gs+m at 26.9(0.1) keV										HWJ235	**
*^{73}Kr−^{85}Rb$_{.859}$	Combined results of next two items										GAu	**
*^{73}Br−^{72}Br	D_M=–4660(330) μu corrected for ^{72}Br gs+m mixture at 100.76 keV										Nub269	**
*^{73}Br−^{72}Br	From ^{72}Br/^{73}Br=0.98635312(227)										AHW	**
*^{73}Ga(β^-)^{73}Ge	E_{β^-}=1190(40) 1200(80) respectively, to 3/2$^-$ level at 364.02 keV										Ens043	**
*^{73}Ge(p,n)^{73}As	T=1205(15) to 5/2$^-$ level at 67.039(0.008)keV										Ens043	**
*^{73}Se(β^+)^{73}As	E_{β^+}=1290(10) to 9/2$^+$ level at 427.906 keV										Ens043	**
*^{73}Br(β^+)^{73}Se	E_{β^+}=3700(500) 3600(400) 3640(140) respectively, to ^{73}Se^m at 25.71 keV										Nub269	**
*^{73}Kr(β^+)^{73}Br	E_{β^+}=5589(350) to 3/2$^-$ level at 178.08 keV										Ens043	**
^{74}Ni−u		−52830	1060	−52011.1	3.0	0.8	B			MT1	1.0	11Es06	
^{74}Ni−^{84}Kr$_{.881}$		25955.7	3.7	25959.4	3.0	1.0	1	65	65 ^{74}Ni	JY1	1.0	22Gi08	
^{74}Ni−^{39}K$_{1.897}$		16844.5	5.0	16837.7	3.0	−1.4	1	35	35 ^{74}Ni	RKM	1.0	25Xi16	
^{74}Cu−u		−59400	400	−60129	6	−1.2	U			TO6	1.5	98Ba.A	
^{74}Cu−^{85}Rb$_{.871}$		16706.0	6.6	16702	6	−0.5	1	74	74 ^{74}Cu	MA8	1.0	07Gu09	
^{74}Cu−^{39}K$_{1.897}$		8710	11	8720	6	0.9	1	26	26 ^{74}Cu	RKM	1.0	25Xi16	
^{74}Zn−^{85}Rb$_{.871}$		6238.4	2.7				2			MA8	1.0	08Ba54	
^{74}Zn−^{39}K$_{1.897}$		−1727	15	−1743.9	2.7	−1.1	U			RKM	1.0	25Xi16	
^{74}Ga−^{85}Rb$_{.871}$		3776.9	22.6	3777	3	−0.0	U			MA8	1.0	07Ke09	*
		3776.9	4.0			−0.0	2			MA8	1.0	07Gu09	
		3806.5	34.6			−0.9	U			MA8	1.0	07Ke09	*
		3776.8	5.4			0.0	2			TT1	1.0	15Ma30	
C ^{32}S$_2$−^{74}Ge H$_2$		7314.0	1.4	7314.523	0.014	0.1	U			M15	2.5	63Ri07	
^{74}Ge−^{84}Kr		9680.0337	0.0128	9680.034	0.013	0.0	1	100	100 ^{74}Ge	FS1	1.0	10Mo03	
C$_6$ H$_2$−^{74}Se		93173.8	3.8	93174.131	0.016	0.0	U			M15	2.5	63Ri07	
^{74}Se−^{84}Kr		10978.2066	0.0128	10978.207	0.015	0.0	o			FS1	1.0	10Mo03	*
^{74}Se−^{85}Rb$_{.871}$		−691.4	7.3	−692.927	0.016	−0.2	U			MA8	1.0	11He10	
^{74}Br ^{27}Al−^{85}Rb$_{1.188}$		16246.0	6.8	16242	6	−0.5	1	85	85 ^{74}Br	MA8	1.0	11He10	*
^{74}Kr−^{85}Rb$_{.871}$		9915.0	2.2	9915.2	2.2	0.1	o			MA8	1.0	04Ro32	*
		9916.8	2.6			−0.6	–			MA8	1.0	06Ro11	
		9909.7	4.4			1.2	–			MA8	1.0	06Ro11	
	ave.	9915.0	2.2			0.1	1	93	93 ^{74}Kr			average	
^{74}Rb−^{85}Rb$_{.871}$		21109	19	21097	3	−0.6	U			MA8	1.0	07Ke09	
		21097.9	4.3			−0.2	o			MA8	1.0	04Ke10	*
		21095.7	5.2			0.3	–			MA8	1.0	07Ke09	
		21102.7	7.5			−0.8	–			MA8	1.0	07Ke09	
		21096.4	6.5			0.1	–			TT1	1.0	15Ma30	
	ave.	21097	4			−0.1	1	83	83 ^{74}Rb			average	
^{74}Rb−u		−55765	125	−55734	3	0.2	U			P40	1.0	06Lu19	
^{74}Sr−u		−43983	33				2			TR1	1.0	25Ho01	
^{74}Ge ^{35}Cl−^{72}Ge ^{37}Cl		2047.5	1.1	2052.07	0.11	1.0	U			H18	4.0	64Ba03	
		2047.74	0.71			2.4	U			H40	2.5	85El01	
		2052.01	0.26			0.1	U			H44	1.5	91Hy01	
^{74}Se ^{35}Cl−^{72}Ge ^{37}Cl		3347.9	4.7	3350.24	0.11	0.2	U			H40	2.5	85El01	
^{73}Ge H−^{74}Ge		10105.1	1.7	10106.23	0.06	0.3	U			M15	2.5	63Ri07	
^{74}Se−^{74}Ge		1298.5	8.5	1298.173	0.008	−0.0	U			H40	2.5	85El01	
		1298.7	3.7			−0.1	U			H40	2.5	85El01	
		1298.096	0.053			1.5	U			JY1	1.0	10Ko15	
		1298.1729	0.0080			0.0	1	100	100 ^{74}Se	FS1	1.0	10Mo03	
^{74}Br−^{73}Br		−1244	410	−1767	8	−0.5	U			CR1	2.5	89Sh10	*
^{74}Se(p,t)^{72}Se		−11979	24	−12005.9	2.0	−1.1	U			Win		74De31	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{74}Ge(^{14}C,^{15}O)^{73}Zn		−8018	150	−7664.8	1.9	2.4	U				Ors		84De33	
^{74}Ge(d,^{3}He)^{73}Ga		−5515	7	−5518.6	1.7	−0.5	U				Ors		78Ro14	
		−5509	13			−0.7	U				Hei		84Ha31	
^{73}Ge(n,γ)^{74}Ge		10200.2	0.6	10196.24	0.06	−6.6	B						70Ha60	
		10198	2			−0.9	U						74Ch18	
		10195.90	0.15			2.2	–				ILn		85Ho.A	Z
		10196.32	0.14			−0.6	–						89Bu.A	
		10196.31	0.07			−1.0	–				MMn		91Is01	Z
		10196.06	0.20			0.9	–				Bdn		06Fi.A	
	ave.	10196.24	0.06			0.0	1	100	100	^{73}Ge			average	
^{74}Se(d,^{3}He)^{73}As		−3027	8	−3056	4	−3.6	B				Ors		83Ro08	*
^{74}Zn(β^-)^{74}Ga		2350	100	2293	4	−0.6	U						72Er05	*
^{74}Ga(β^-)^{74}Ge		5400	100	5372.8	3.0	−0.3	U						62Ei02	*
^{74}As(β^+)^{74}Ge		2558	4	2562.4	1.7	1.1	–						71Bo01	*
^{74}Ge(p,n)^{74}As		−3343.5	5.6	−3344.7	1.7	−0.2	–				Tkm		63Ok01	
		−3348.3	5.			0.7	–				Oak		64Jo11	Z
		−3346	5			0.3	–						70Fi03	Z
		−3347	3			0.8	–				Kyu		73Ki11	
^{74}As(β^+)^{74}Ge	ave.	2562.9	1.9	2562.4	1.7	−0.3	1	82	82	^{74}As			average	
^{74}As(β^-)^{74}Se		1351	4	1353.1	1.7	0.5	1	18	18	^{74}As			71Bo01	*
^{74}Br(β^+)^{74}Se		6857	100	6925	6	0.7	U						69La15	*
^{74}Se(p,n)^{74}Br		−7689	15	−7707	6	−1.2	1	15	15	^{74}Br			75Lu02	*
^{74}Kr(β^+)^{74}Br		3000	200	2956	6	−0.2	U						74Ro11	
		3327	125			−3.0	B						75Sc07	
^{74}Rb(β^+)^{74}Kr		10000	1500	10416	3	0.3	U						76Da.D	
		10413.8	7.0			0.3	1	24	17	^{74}Rb			03Pi08	*
*^{74}Ga−^{85}Rb$_{.871}$	D_M=3780.1(22.5) μu corrected –3.0(1.7) keV for gs+m mixture R<0.1 at 59.571 keV												Nub269	**
*^{74}Ga−^{85}Rb$_{.871}$	D_M=3809.7(34.6) μu corrected –3.0(1.7) keV for gs+m mixture R<0.1 at 59.571 keV												Nub269	**
*^{74}Se−^{84}Kr	Not independent measurement, use ^{74}Ge−^{74}Se below												10Mo03	**
*^{74}Br ^{27}Al−^{85}Rb$_{1.188}$	D_M=16253.4(5.3) μu for mixture gs+m at 13.58 keV; $M-A$=–82474.9(4.9) keV												Nub269	**
*^{74}Kr−^{85}Rb$_{.871}$	Combined results of next two items												GAu	**
*^{74}Rb−^{85}Rb$_{.871}$	Combined results of next two items												GAu	**
*^{74}Br−^{73}Br	D_M=–1230(410) μu for ^{74}Br^m at 13.58 keV												Nub269	**
*^{74}Se(p,t)^{72}Se	Original error 12; added systematic error 21 keV												GAu	**
*^{74}Se(d,^{3}He)^{73}As	Q =–3033(8) for Q(^{76}Se(d,^{3}He))=–4020.7(2.0), now 4014.5 keV												AHW	**
*^{74}Zn(β^-)^{74}Ga	E_{β^-}=2100(100) to 1$^+$ level at 251.787 keV												Ens067	**
*^{74}Ga(β^-)^{74}Ge	E_{β^-}=2450(100) to 3$^-$ level at 2949.48 keV												Ens067	**
*^{74}As(β^+)^{74}Ge	Original error increased: authors report E(2$^+$)=593.1(1.5) while												AHW	**
*	E(2$^+$)=595.850(0.006) keV; see also ^{84}Rb(β^+)												Ens067	**
*^{74}As(β^-)^{74}Se	Original value 1350.1(0.7), error increased, see ^{84}Rb(β^+)												AHW	**
*^{74}Br(β^+)^{74}Se	E_{β^+}=5200(100), 4500(100) to 634.76, 1363.21 levels												69La15	**
*	from ^{74}Br^m at 13.8(0.5) keV												93Do05	**
*^{74}Se(p,n)^{74}Br	T=7868(15) to (2$^-$) level at 72.65 keV												Ens067	**
*^{74}Rb(β^+)^{74}Kr	Deduced from measured half-life and branching ratio												GAu	**
*^{74}Rb(β^+)^{74}Kr	Original 10405(9) re-evaluated in reference												11To.A	**
^{75}Ni−^{84}Kr$_{.893}$		31736	16	31746	6	0.6	1	15	15	^{75}Ni	JY1	1.0	22Gi08	
^{75}Ni−^{39}K$_{1.923}$		22507.8	6.8	22506	6	−0.3	1	85	85	^{75}Ni	RKM	1.0	25Xi16	
^{75}Cu−u		−58100	700	−58476.2	0.8	−0.4	U				TO6	1.5	98Ba.A	
		−58304	311			−0.6	U				RR3	1.0	24Na17	*
^{75}Cu−^{85}Rb$_{.882}$		19325.40	0.81	19325.3	0.8	−0.2	1	91	91	^{75}Cu	MA8	1.0	17We16	
^{75}Cu−^{39}K$_{1.923}$		11316.5	3.6	11316.3	0.8	−0.1	U				RKM	1.0	25Xi16	
^{75}Zn−^{85}Rb$_{.882}$		10510.6	2.3	10507.9	1.2	−1.2	1	30	30	^{75}Zn	IS1	1.0	26Mu04	
^{75}Zn^m−^{85}Rb$_{.882}$		10641.7	2.1	10644.2	1.2	1.2	–				MA8	1.0	08Ba54	*
		10644.5	2.1			−0.1	–				IS1	1.0	26Mu04	
	ave.	10643.1	1.5			0.7	1	71	71	^{75}Zn^m			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$^{75}Zn^{m}-^{39}K_{1.923}$	2620	14	2635.2	1.2	1.1	U			RKM	1.0	25Xi16	*
$^{75}Ga-^{85}Rb_{.882}$	4301.7	2.6	4305.9	0.7	1.6	U			MA8	1.0	07Gu09	
	4305.94	0.72				2			MA8	1.0	19Hu15	
$C_3\ H_7\ O_2-^{75}As$	123009.8	2.6	123011.32	0.04	0.2	U			M15	2.5	63Ri07	
$^{75}As-u$	−78407.35	0.47	−78406.86	0.04	1.0	U			MS1	1.0	22Ga36	*
$^{75}As-^{85}Rb_{.882}$	−601.3	7.6	−605.40	0.04	−0.5	U			MA8	1.0	02Ke.A	
$^{75}Br\ ^{27}Al-^{85}Rb_{1.200}$	13201.3	4.6				2			MA8	1.0	11He10	
$^{75}Kr-u$	−69055.3	2.8	−69055.2	2.7	0.0	1	91	91 ^{75}Kr	LZ1	1.0	23Xi01	
$^{75}Kr-^{85}Rb_{.882}$	8747.2	8.7	8746.3	2.7	−0.1	1	9	9 ^{75}Kr	MA8	1.0	06Ro11	
$^{75}Rb-^{85}Rb_{.882}$	16371	8	16374.7	1.3	0.5	U			MA2	1.0	94Ot01	
	16374.7	1.7			−0.0	2			MA8	1.0	07Ke09	
	16368	21			0.3	U			MA8	1.0	07Ke09	
	16374.6	1.9			0.0	2			TT1	1.0	15Ma30	
$^{75}Sr-u$	−49600	160	−49676	11	−0.5	U			LZ2	1.0	23Wa10	
	−49676.1	10.7				2			TR1	1.0	25Ho01	
$^{75}Cu-^{72}Ge_{1.042}$	22719.6	2.5	22720.8	0.8	0.5	1	10	9 ^{75}Cu	JY1	1.0	07Ra27	
$^{75}As\ ^{35}Cl-^{73}Ge\ ^{37}Cl$	1079.6	5.0	1084.32	0.10	0.4	U			H40	2.5	85El01	
$^{75}As-^{76}Ge_{.987}$	−831.351	0.042	−831.35	0.04	0.1	1	86	86 ^{75}As	JY2	1.0	22Ra20	
$^{74}Ge(n,\gamma)^{75}Ge$	6505.9	1.1	6505.84	0.05	−0.1	U					72Gr34	
	6505.5	2.			0.2	U					72Ha74	
	6505.81	0.30			0.1	U					89Bu.A	*
	6505.26	0.08			7.2	B			MMn		91Is01	Z
	6505.45	0.14			2.8	C			Bdn		06Fi.A	
	6505.84	0.05				2					12Me04	
$^{74}Ge(d,p)^{75}Ge$	4265	15	4281.27	0.05	1.1	U			MIT		67Sp09	
	4282	10			−0.1	U			Kop		72Ha74	
	4268	10			1.3	U			Kyu		73Ka03	
$^{74}Ge(p,\gamma)^{75}As$	6901.6	5.	6902.05	0.04	0.1	U					74Wa08	
$^{74}Ge(^{3}He,d)^{75}As$	1414	4	1408.57	0.04	−1.4	U			Hei		76Sc13	
$^{75}As(\gamma,n)^{74}As$	−10259	31	−10246.8	1.7	0.4	U			Phi		60Ge01	
$^{74}Se(n,\gamma)^{75}Se$	8027.84	0.30	8027.60	0.07	−0.8	U			BNn		81En07	Z
	8027.60	0.08			−0.0	2			ILn		84To11	Z
	8027.59	0.16			0.0	2			Bdn		06Fi.A	
$^{75}Zn^{m}(IT)^{75}Zn$	126.94	0.09	126.94	0.09	−0.0	1	100	70 ^{75}Zn			Ens136	
$^{75}Zn(\beta^{-})^{75}Ga$	6060	80	5777.1	1.3	−3.5	B			Stu		86Ek01	
$^{75}Ga(\beta^{-})^{75}Ge$	3300	200	3396.3	0.7	0.5	U					60Mo01	
$^{75}Ge(\beta^{-})^{75}As$	1188	20	1178.55	0.07	−0.5	U					55Sc09	
$^{75}As(p,n)^{75}Se$	−1647.2	2.0	−1648.38	0.08	−0.6	U			Nvl		59Go68	Z
	−1647.3	1.1			−1.0	U			Oak		64Jo11	Z
	−1643	5			−1.1	U					70Fi03	
$^{75}Br(\beta^{+})^{75}Se$	3010	20	3062	4	2.6	U					52Fu04	*
	3030	50			0.6	U					61Ba43	*
	3050	20			0.6	U					69Ra24	*
$^{75}Kr(\beta^{+})^{75}Br$	4400	200	4783	5	1.9	U					74Ro12	*
$^{75}Sr(\varepsilon)^{75}Rb$	10600	220	10946	10	1.6	U					03Hu01	
$*^{75}Cu-u$	also ME=–54460(740) keV										24Na17	**
$*^{75}Zn^{m}-^{85}Rb_{.882}$	Assigned to the 126.94 keV isomeric state, originally identified as the gs.										26Mu04	**
$*^{75}Zn^{m}-^{39}K_{1.923}$	re-assigned to the isomeric state by the evaluator										HWJ264	*
$*^{75}As-u$	From frequency ratio $^{75}As^{+}/(C_6)^{+}$ =0.9610043784(60)										22Ga36	**
$*^{74}Ge(n,\gamma)^{75}Ge$	Original error 0.03 keV increased										GAu	**
$*^{75}Br(\beta^{+})^{75}Se$	$E_{\beta^{+}}$=1700(20) 1720(50) 1740(20) respectively, to $3/2^{-}$ level at 286.5714 keV										Ens139	**
$*^{75}Kr(\beta^{+})^{75}Br$	$E_{\beta^{+}}$=3200(200) to 132.46 $(5/2)^{+}$, 154.61 $(3/2)^{+}$ levels										Ens139	**
$^{76}Cu-^{85}Rb_{.894}$	24100	21	24096.9	2.2	−0.1	U			MA8	1.0	07Gu09	*
	24094	20			0.1	U			MA8	1.0	17We16	*
$^{76}Cu-^{39}K_{1.949}$	15970	20	15973.0	2.2	0.2	U			RI1	1.0	25Xi16	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{76}Zn−^{85}Rb$_{.894}$	11975.5	2.0	11974.9	1.6	−0.3	1	61	61 ^{76}Zn	MA8	1.0	08Ba54	
^{76}Zn−^{88}Rb$_{.864}$	9737.4	2.5	9738.3	1.6	0.4	1	39	39 ^{76}Zn	JY1	1.0	08Ha23	
^{76}Zn−^{39}K$_{1.949}$	3857.3	9.7	3851.0	1.6	−0.6	U			RKM	1.0	25Xi16	
^{76}Ga−^{85}Rb$_{.894}$	7687.6	2.1				2			MA8	1.0	07Gu09	
C ^{32}S$_2$−^{76}Ge	22741.6	1.5	22739.623	0.019	−0.5	U			M15	2.5	63Ri07	
^{76}Ge−u	−78597.242	0.096	−78597.276	0.019	−0.4	U			ST2	1.0	01Do08	
C$_6$ H$_4$−^{76}Se	112100	8	112086.426	0.017	−0.7	U			M15	2.5	63Ri07	
^{76}Se−u	−80786.205	0.081	−80786.298	0.017	−1.2	U			ST2	1.0	01Do08	
^{76}Kr−^{85}Rb$_{.894}$	4774.3	4.7	4771	4	−0.8	1	84	84 ^{76}Kr	MA8	1.0	06Ro11	
^{76}Rb−^{85}Rb$_{.894}$	13931	8	13933.0	1.0	0.3	U			MA2	1.0	94Ot01	
	13932.2	2.0			0.4	2			MA8	1.0	07Ke09	
	13923	15			0.7	U			MA8	1.0	07Ke09	
	13935.3	1.6			−1.4	2			MA8	1.0	07Ke09	
	13931.0	1.7			1.2	2			TT1	1.0	15Ma30	
	13925.2	4.5			1.7	U			TT1	1.0	15Ma30	
	13933.5	3.4			−0.1	o			TT1	1.0	15Ma30	*
^{76}Sr ^{19}F−^{85}Rb$_{1.118}$	38785	37	38776	9	−0.3	1	6	6 ^{76}Sr	MA8	1.0	05Si34	
^{76}Sr−u	−58813	107	−58247	9	5.3	F			CS1	1.0	01La31	*
	−58247.3	9.7			0.1	1	94	94 ^{76}Sr	TR1	1.0	25Ho01	
^{76}Cu−^{84}Kr$_{.905}$	25331.5	2.2				2			JY1	1.0	22Gi08	
^{76}Cu^m−^{84}Kr$_{.905}$	25401.1	1.5				2			JY2	1.0	24Ca11	
^{76}Ge−^{84}Kr	9904.9983	0.0175	9904.998	0.019	−0.0	o			FS1	1.0	10Mo03	*
^{76}Se−^{84}Kr	7715.9762	0.0169	7715.976	0.017	−0.0	1	100	100 ^{76}Se	FS1	1.0	10Mo03	
^{74}Ge H$_2$−^{76}Ge	15425.0	1.7	15425.099	0.023	0.0	U			M15	2.5	63Ri07	
^{76}Ge ^{35}Cl−^{74}Ge ^{37}Cl	3175.7	1.5	3175.09	0.07	−0.1	U			H18	4.0	64Ba03	
	3170.41	0.74			2.5	U			H40	2.5	85El01	
	3174.61	0.41			0.8	U			H44	1.5	91Hy01	
^{76}Se ^{35}Cl−^{74}Ge ^{37}Cl	986.30	0.65	986.07	0.07	−0.2	U			H44	1.5	91Hy01	
^{76}Ge−^{76}Se	2190.92	0.59	2189.022	0.008	−1.3	U			H40	2.5	85El01	
	2188.60	0.42			0.7	U			H44	1.5	91Hy01	
	2188.963	0.054			1.1	U			ST2	1.0	01Do08	
	2188.98	0.16			0.3	U			JY1	1.0	08Ra09	
	2189.0221	0.008			0.0	1	100	100 ^{76}Ge	FS1	1.0	10Mo03	
^{76}As−^{76}Se	3176.908	0.076	3176.90	0.06	−0.1	1	55	55 ^{76}As	JY2	1.0	22Ge07	
^{75}Rb−^{76}Rb$_{.493}$ ^{74}Rb$_{.507}$	−1140	170	−1081.1	2.0	0.1	U			P20	2.5	82Au01	
^{76}Ge(^{14}C,^{17}O)^{73}Zn	−3779	40	−3790.8	1.9	−0.3	U			Ors		84Be10	*
^{76}Ge(^{14}C,^{16}O)^{74}Zn	163	40	300.7	2.5	3.4	B			Ors		84Be10	
^{76}Ge(^{18}O,^{20}Ne)^{74}Zn	−1219	21	−1197.1	2.5	1.0	U			Hei		84Ha31	
^{76}Ge(^{14}C,^{15}O)^{75}Zn	−10354	150	−10365.1	1.3	−0.1	U			Ors		84De33	
^{76}Ge(d,^{3}He)^{75}Ga	−6545	7	−6547.8	0.7	−0.4	U			Ors		78Ro14	
	−6536	22			−0.5	U			Hei		84Ha31	
^{75}As(n,γ)^{76}As	7329	2	7328.49	0.05	−0.3	U					68Jo11	
	7328.421	0.075			0.9	–			ILn		90Ho10	Z
	7328.81	0.15			−2.1	–			Bdn		06Fi.A	
^{75}As(d,p)^{76}As	5105	5	5103.92	0.05	−0.2	U					76Mo32	
^{75}As(n,γ)^{76}As ave.	7328.50	0.07	7328.49	0.05	−0.1	1	59	45 ^{76}As			average	
^{75}Se(n,γ)^{76}Se	11154.15	0.30	11153.79	0.07	−1.2	U			ILn		83To20	Z
^{76}Zn(β^-)^{76}Ga	4160	80	3993.6	2.4	−2.1	U			Stu		86Ek01	
^{76}Ga(β^-)^{76}Ge	6770	150	6916.3	2.0	1.0	o			Stu		77Al17	
	7010	90			−1.0	U			Stu		86Ek01	
^{76}Ge(p,n)^{76}As	−1705	5	−1702.55	0.05	0.5	U					70Fi03	
^{76}As(β^-)^{76}Se	2970	2	2959.26	0.05	−5.4	B					69Na11	
^{76}Br(β^+)^{76}Se	5002	20	4963	9	−2.0	2					71Dz08	
^{76}Br(n,p)^{76}Se	5730	15	5745	9	1.0	2			ILL		78An14	
^{76}Se(p,n)^{76}Br	−5738.6	15.	−5745	9	−0.4	2					75Lu02	
^{76}Rb(β^+)^{76}Kr	8793	570	8535	4	−0.5	U					75We23	
	8063	44			10.7	F					82Mo10	*
	8094	162			2.7	F			BNL		83Li11	*
	8250	150			1.9	U			IRS		93Al03	
*^{76}Cu−^{85}Rb$_{.894}$	D_M=24135.0(7.2) μu and $M-A$=–50976.0(6.7) keV for a mixture										RCH261	**
*^{76}Cu−^{85}Rb$_{.894}$	of gs. and isomer at 64.8(2.5) keV										RCH261	**
*^{76}Cu−^{85}Rb$_{.894}$	D_M=24128.95(0.98) μu and $M-A$=–50981.63(0.91) for a mixture of										RCH261	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
$*^{76}$Cu−^{85}Rb$_{.894}$		gs. and isomer at 64.8(2.5) keV											RCH261	**
$*^{76}$Cu−^{39}K$_{1.949}$		D_M=16004.5(3.1) μu and $M-A$=–50982.1(2.9) keV for a mixture of gs.											RCH261	**
$*^{76}$Cu−^{39}K$_{1.949}$		and isomer at 64.8(2.5) keV											RCH261	**
$*^{76}$Rb−^{85}Rb$_{.894}$		no accuracy check was performed for this 12^+ charge state											15Ma30	**
$*^{76}$Sr−u		F : other results in same paper not trusted, see ^{80}Y											GAu	**
$*^{76}$Ge−^{84}Kr		Not independent measurement, use ^{76}Ge−^{76}Se below											10Mo03	**
$*^{76}$Ge(^{14}C,^{17}O)^{73}Zn		Q =–3974(40) $M-A$=–65410(40) to ^{73}Zn^m at 195.5 keV											GAu	**
$*^{76}$Rb(β^+)^{76}Kr		E_{β^+}=4558(30) to level (1^+,2^+) at 2570.95 keV and corrections by authors											Ens953	**
$*^{76}$Rb(β^+)^{76}Kr		F : 29.6% feeding above 2570.95 level from reference											84Mo22	**
^{77}Cu−u		−51850	540	−52456.3	1.3	−1.1	U				OR1	1.0	06Ha62	
^{77}Cu−^{85}Rb$_{.906}$		27462.1	1.3	27462.2	1.3	0.1	1	94	94	^{77}Cu	MA8	1.0	17We16	
^{77}Zn−u		−62790	780	−63112.8	2.1	−0.3	U				TO6	1.5	98Ba.A	*
		−63380	260			0.4	U				GT2	2.5	08Kn.A	*
^{77}Zn−^{88}Rb$_{.875}$		14485.7	4.5	14486.1	2.1	0.1	1	22	22	^{77}Zn	JY1	1.0	08Ha23	
C_6H_5−^{77}Se		119211.9	4.2	119210.99	0.05	−0.1	U				M15	2.5	63Ri07	
^{77}Rb−^{39}K$_{1.974}$		1912	27	2045.0	1.4	4.9	B				MA2	1.0	87Bo59	
^{77}Y−^{39}K$_{1.974}$		21285.8	4.1				2				MS2	1.0	26Er.A	
^{77}Zn−^{85}Rb$_{.906}$		16805.8	2.4	16805.7	2.1	−0.0	1	78	78	^{77}Zn	MA8	1.0	08Ba54	
^{77}Ga−^{85}Rb$_{.906}$		9072.8	2.6				2				MA8	1.0	07Gu09	
		9070.0	4.6	9072.8	2.6	0.6	U				MA8	1.0	19Hu15	
^{77}Kr−^{85}Rb$_{.906}$		4588.5	2.1				2				MA8	1.0	06Ro11	
^{77}Rb−^{85}Rb$_{.906}$		10327	8	10320.1	1.4	−0.9	U				MA2	1.0	94Ot01	
		10320.1	1.4				2				MA8	1.0	07Ke09	
^{77}Sr ^{19}F−^{85}Rb$_{1.129}$		35938.0	8.5				2				MA8	1.0	05Si34	
^{77}Cu−^{84}Kr$_{.917}$		28701.6	5.0	28700.3	1.3	−0.3	1	6	6	^{77}Cu	JY1	1.0	22Gi08	
^{77}Se−^{76}Se$_{1.013}$		1750.718	0.064	1750.69	0.05	−0.4	1	51	51	^{77}Se	JY2	1.0	22Ra20	
^{77}Cu−^{77}Zn		10664	25	10656.5	2.5	−0.3	U				RKM	1.0	25Xi16	
^{77}Zn^m−^{77}Zn		809	37	829.248	0.016	0.5	U				RKM	1.0	25Xi16	
^{77}As−^{77}Se		734.801	0.077	734.80	0.08	−0.0	1	100	100	^{77}As	JY2	1.0	24Ge01	
^{75}Rb−^{77}Rb$_{.325}$ ^{74}Rb$_{.676}$		−1340	380	−1053.6	2.4	0.3	U				P20	2.5	82Au01	
^{76}Rb−^{77}Rb$_{.494}$ ^{75}Rb$_{.507}$		525	30	557.1	1.3	0.4	U				P20	2.5	82Au01	
^{76}Ge(n,γ)^{77}Ge		6072.5	1.0	6071.29	0.05	−1.2	U						72Gr34	Z
		6071.7	1.2			−0.3	U						72Ha74	Z
		6072.3	0.4			−2.5	U				Bdn		06Fi.A	
		6071.29	0.05				2						12Me04	
^{76}Ge(d,p)^{77}Ge		3839	10	3846.72	0.05	0.8	U				Kop		72Ha74	
		3823	12			2.0	U				Kyu		73Ka03	
^{76}Ge(^{3}He,d)^{77}As		2497	3	2497.61	0.08	0.2	U				Hei		76Sc13	
^{76}Se(n,γ)^{77}Se		7418.87	0.20	7418.83	0.04	−0.2	–				BNn		81En07	
		7418.85	0.07			−0.2	–				ILn		85To10	Z
		7418.85	0.15			−0.1	–				Bdn		06Fi.A	
^{76}Se(d,p)^{77}Se		5192	10	5194.27	0.04	0.2	U				Ald		63Ma27	
^{76}Se(n,γ)^{77}Se	ave.	7418.85	0.06	7418.83	0.04	−0.3	1	49	49	^{77}Se			average	
^{77}Sr(εp)^{76}Kr		3850	200	3921	9	0.4	U				ChR		76Ha29	
^{77}Zn^m(IT)^{77}Zn		772.440	0.015				2						Ens209	
^{77}Zn(β^-)^{77}Ga		7270	120	7203	3	−0.6	U				Stu		86Ek01	
^{77}Ga(β^-)^{77}Ge		5340	60	5220.5	2.4	−2.0	U				Stu		77Al17	
		5690	300			−1.6	U				Stu		86Ek01	
^{77}Ge(β^-)^{77}As		2670	100	2702.14	0.10	0.3	U						52Sm13	*
^{77}As(β^-)^{77}Se		700	7	684.46	0.07	−2.2	U						51Ca04	
		679	4			1.4	U						51Je01	
^{77}Br(β^+)^{77}Se		1358	20	1364.7	2.8	0.3	U						51Ca28	
^{77}Se(p,n)^{77}Br		−2147	4	−2147.0	2.8	−0.0	2				Oak		58Jo01	
		−2147.0	4.			0.0	2				Tkm		63Ok01	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{77}Kr(β^+)^{77}Br	3012	30	3065	3	1.8	U					55Th01	*
	3027	40			1.0	U					73Ba22	*
	3300	100			−2.3	U					74Ro11	*
	2760	42			7.3	B					82Mo10	*
^{77}Rb(β^+)^{77}Kr	5180	390	5339.0	2.4	0.4	U					75We23	
	5272	26			2.6	U					82Mo10	
	5113	69			3.3	B			BNL		83Li11	
	5320	70			0.3	U			IRS		93Al03	
^{77}Sr(β^+)^{77}Rb	6986	227	7027	8	0.2	U			BNL		83Li11	
*^{77}Zn−u	$M-A$=–58100(700) keV for mixture gs+m at 772.440 keV										Nub269	**
*^{77}Zn−u	$M-A$=–58648(95) keV for mixture gs+m at 772.440 keV (1/2$^-$)										Nub269	**
*^{77}Ge(β^-)^{77}As	E_{β^-}=2196(100) to 9/2$^+$ ^{77}As^m at 475.48 keV										Nub269	**
*^{77}Kr(β^+)^{77}Br	Error not in 55Th01, estimated by evaluator										AHW	**
*^{77}Kr(β^+)^{77}Br	E_{β^+}=1860(30) 1875(40) respectively, to 5/2$^+$ level at 129.64 keV										Ens126	**
*^{77}Kr(β^+)^{77}Br	E_{β^+}=2000(100) 1528(36) respectively, to (3/2)$^+$ level at 276.22 keV										Ens126	**
^{78}Cu−u	−47770	540	−48081	6	−0.6	U			OR1	1.0	06Ha62	
	−48115.2	23.6			1.5	1	7	7 ^{78}Cu	MR1	1.0	17We16	
^{78}Cu−^{85}Rb$_{.918}$	32912	18	32896	6	−0.9	1	12	12 ^{78}Cu	MA8	1.0	17We16	
^{78}Cu−^{38}Ar ^{1}H	−18638	24	−18638	6	−0.0	1	7	7 ^{78}Cu	RKM	1.0	25Xi16	
^{78}Zn−^{88}Rb$_{.886}$	16863.8	2.9	16863.6	2.1	−0.1	1	51	51 ^{78}Zn	JY1	1.0	08Ha23	
^{78}Zn−^{38}Ar ^{1}H	−32247.3	5.7	−32267.9	2.1	−3.6	B			RKM	1.0	25Xi16	
^{78}Ga−^{88}Rb$_{.886}$	10184.3	3.3	10185.3	1.1	0.3	1	12	12 ^{78}Ga	JY1	1.0	08Ha23	
^{78}Ga−^{38}Ar ^{1}H	−38899	30	−38946.3	1.1	−1.6	U			RKM	1.0	25Xi16	
C_6 H_6−^{78}Se	129642.6	2.2	129640.91	0.19	−0.3	U			M15	2.5	63Ri07	
C_6 H_6−^{78}Kr	126548.3	3.6	126583.8	0.3	3.9	B			M15	2.5	63Ri07	
	126554	17			1.2	U			R11	1.5	78Di09	
	126560	7			2.3	U			R11	1.5	78Di09	
C_5 N H_4−^{78}Kr	113994	20	114007.8	0.3	0.5	U			R11	1.5	78Di09	
^{78}Sr−^{39}K$_2$	4773.8	9.4	4770	6	−0.4	1	42	42 ^{78}Sr	RKM	1.0	25Ki14	
^{78}Y−^{39}K$_{2.000}$	16403.3	3.5				2			MS2	1.0	26Er.A	
^{78}Y^m−^{39}K$_2$	16806	16	16807.2	1.0	0.1	U			RKM	1.0	25Ki14	
	16807.20	0.98				2			MS2	1.0	26Er.A	
^{78}Kr−^{86}Kr$_{.907}$	1441.2	1.0	1442.5	0.3	1.3	1	11	11 ^{78}Kr	MS1	1.0	06Ri15	
^{78}Zn−^{85}Rb$_{.918}$	19266.0	3.0	19266.3	2.1	0.1	1	48	48 ^{78}Zn	MA8	1.0	08Ba54	
^{78}Ga−^{85}Rb$_{.918}$	12585.2	2.6	12587.9	1.1	1.0	U			MA8	1.0	07Gu09	
	12588.0	1.2			−0.1	1	88	88 ^{78}Ga	MA8	1.0	19Hu15	
^{78}Kr−^{85}Rb$_{.918}$	1342.3	1.4	1343.4	0.3	0.8	U			MA8	1.0	06Ro11	
	1338.9	2.2			2.0	U			MA8	1.0	06Ro11	
^{78}Rb−^{85}Rb$_{.918}$	9118	8	9119	3	0.1	2			MA2	1.0	94Ot01	
	9121.3	7.5			−0.3	2			TT1	1.0	12Ga15	*
	9118.3	4.5			0.1	2			TT1	1.0	12Ga15	*
^{78}Sr−^{85}Rb$_{.918}$	13157	8	13160	6	0.4	1	58	58 ^{78}Sr	MA2	1.0	94Ot01	
^{78}Cu−^{84}Kr$_{.929}$	34139.1	8.1	34138	6	−0.2	1	58	58 ^{78}Cu	JY1	1.0	22Gi08	
^{78}Se ^{35}Cl$_2$−^{74}Ge ^{37}Cl$_2$	2030.4	2.2	2031.78	0.23	0.4	U			H44	1.5	91Hy01	
^{78}Se ^{35}Cl−^{76}Ge ^{37}Cl	−1147.60	0.92	−1143.32	0.20	1.9	U			H40	2.5	85El01	
	−1143.57	0.72			0.2	U			H44	1.5	91Hy01	
^{78}Se ^{35}Cl−^{76}Se ^{37}Cl	1042.03	1.35	1045.70	0.20	1.1	U			H40	2.5	85El01	
	1044.58	0.45			1.7	U			H44	1.5	91Hy01	
^{76}Se H_2−^{78}Kr	14440	25	14497.4	0.3	1.5	U			R11	1.5	78Di09	
^{77}Se H−^{78}Kr	7367	26	7372.8	0.3	0.1	U			R11	1.5	78Di09	
^{78}Cu−^{78}Zn	13628	15	13630	6	0.1	1	18	17 ^{78}Cu	RKM	1.0	25Xi16	
^{78}Ga−^{78}Zn	−6668	43	−6678.4	2.4	−0.2	U			RKM	1.0	25Xi16	
^{78}Kr−^{78}Se	3074	16	3057.10	0.28	−0.7	U			R11	1.5	78Di09	
	3098	20			−1.4	U			R11	1.5	78Di09	
	3057.18	0.29			−0.3	1	92	89 ^{78}Kr	MS1	1.0	13Bu17	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{78}Se H$-^{78}$Kr		4724	33	4767.94	0.28	0.9	U			R11	1.5	78Di09	
^{76}Rb$-^{78}$Rb$^{x}_{325}$ ^{75}Rb$_{.676}$		−130	40	−69	4	0.6	U			P20	2.5	82Au01	
^{77}Rb$-^{78}$Rb$^{x}_{494}$ ^{76}Rb$_{.507}$		−1192	19	−1138	6	1.1	U			P20	2.5	82Au01	
^{78}Kr(α,^{8}He)^{74}Kr		−41080	75	−41031.2	2.0	0.7	U			Tex		82Mo23	*
^{78}Se(p,α)^{75}As		870.9	2.3	873.66	0.18	1.2	U			NDm		82Zu04	
^{78}Kr(^{3}He,^{6}He)^{75}Kr		−12581	14	−12514.6	2.5	4.7	B					87Mo06	
^{76}Ge(t,p)^{78}Ge		6310	5	6310	4	0.0	2			LAl		78Ar12	
		6310	5			0.0	2			Phi		81St18	
^{78}Se(p,t)^{76}Se		−9433.7	4.3	−9434.80	0.17	−0.3	U			NDm		82Zu04	
^{78}Kr(α,^{6}He)^{76}Kr		−20351	10	−20332	4	1.9	o			Tex		82Mo23	*
^{78}Kr(p,t)^{76}Kr		−12840	15	−12825	4	1.0	U			Tky		81Ma30	
^{78}Se(d,^{3}He)^{77}As		−4904	4	−4906.40	0.18	−0.6	U			Ors		83Ro08	*
^{77}Se(n,γ)^{78}Se		10497.7	0.3	10497.76	0.17	0.2	–			BNn		81En07	Z
		10497.75	0.21			0.1	–			Bdn		06Fi.A	
^{78}Se(p,d)^{77}Se		−8271.9	4.0	−8273.20	0.17	−0.3	U			NDm		82Zu04	
^{77}Se(n,γ)^{78}Se	ave.	10497.73	0.17	10497.76	0.17	0.2	1	96	96 ^{78}Se			average	
^{78}Kr(d,t)^{77}Kr		−5804	7	−5822.9	2.0	−2.7	U					87Mo06	
^{78}Zn(β^-)^{78}Ga		6440	140	6220.9	2.2	−1.6	o			Stu		86Ek01	
		6364	90			−1.6	U			Stu		00Me.A	
^{78}Ga(β^-)^{78}Ge		8140	160	8158	4	0.1	o			Stu		77Al17	
		8200	80			−0.5	o			Stu		86Ek01	
		8054	43			2.4	U			Stu		00Me.A	
^{78}Ge(β^-)^{78}As		967	30	955	10	−0.4	R					65Fr04	*
		987	20			−1.6	R					65Kv01	*
^{78}As(β^-)^{78}Se		4270	100	4209	10	−0.6	U					70Mc01	
		4310	100			−1.0	U					71Mo20	*
^{78}Br(β^+)^{78}Se		3542	50	3574	4	0.6	U			Bar		61Ri02	
^{78}Se(p,n)^{78}Br		−4344	10	−4356	4	−1.2	2			Bar		61Ri02	
		−4370	10			1.4	2			LAl		61Sc11	
		−4355.5	7.4			−0.1	2			Tkm		63Ok01	Z
		−4356	5			−0.0	2					70Fi03	Z
^{78}Rb(β^+)^{78}Kr		7085	370	7243	3	0.4	U					75We23	*
		7240	50			0.1	U					81Ba40	
		7185	50			1.2	U			IRS		93Al03	*
^{78}Rb^{x}(IT)^{78}Rb		74	12				3					82Au01	*
*^{78}Rb$-^{85}$Rb$_{.918}$	Correction for e$^-$ binding=+97eV is negligible											12Ga15	**
*^{78}Rb$-^{85}$Rb$_{.918}$	D_M= 9237.6(4.5) μu $M-A$=–66824.9(4.2) keV corrected for e$^-$ binding=+97 eV											12Ga15	**
*	from ^{78}Rb^{n} at 111.19(0.22) keV											Nub269	**
*^{78}Kr(α,^{8}He)^{74}Kr	Original –41120(75) for 4 events included 1 background event											GAu	**
*^{78}Kr(α,^{6}He)^{76}Kr	Replaced by calibration free ^{80}Kr(α,^{6}He)^{78}Kr$-^{78}$Kr()^{76}Kr											GAu	**
*^{78}Se(d,^{3}He)^{77}As	Original value –4910(4) corrected, see ^{74}Se(d,^{3}He)											AHW	**
*^{78}Ge(β^-)^{78}As	E_{β^-}=690(30) 710(20) respectively, to 1$^+$ level at 277.3 keV											Ens09a	**
*^{78}As(β^-)^{78}Se	E_{β^-}=3000(100) to 2$^+$ level at 1308.644 keV											Ens09a	**
*^{78}Rb(β^+)^{78}Kr	E_{β^+}=3410(370) from ^{78}Rb^{n} 4$^{(-)}$ at 111.19(0.22) to (4)$^-$ level at 2764.10 keV											Ens09a	**
*^{78}Rb(β^+)^{78}Kr	Q_{β^+}=7180(80); and 7300(50) from ^{78}Rb^{n} at 111.19(0.22) keV											Ens09a	**
*^{78}Rb^{x}(IT)^{78}Rb	Corrected; using ^{78}Rb^{n}(IT)=111.2 keV											GAu	**
^{79}Cu−u		−46700	540	−45530	110	2.2	U			OR1	1.0	06Ha62	
		−45526.9	112.7				2			MR1	1.0	17We16	
^{79}Zn$-^{88}$Rb$_{.898}$		22278.1	2.9	22277.0	1.9	−0.4	1	43	43 ^{79}Zn	JY1	1.0	08Ha23	
^{79}Zn^{m}−u		−56349	15	−56351	8	−0.1	1	26	26 ^{79}Zn^{m}	MR1	1.0	23Ni10	
^{79}Ga$-^{88}$Rb$_{.898}$		12490.9	2.0	12490.2	1.3	−0.4	1	41	41 ^{79}Ga	JY1	1.0	08Ha23	
^{79}Ga−u		−67064	129	−67148.5	1.3	−0.3	U			GT2	2.5	08Su19	
C$_6$ H$_7-^{79}$Br		136444.3	2.4	136437.5	1.0	−1.1	U			M15	2.5	63Ri07	
		136444	15			−0.3	U			R11	1.5	78Di09	
		136449	12			−0.6	U			R11	1.5	78Di09	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$C_5\ ^{13}C\ H_6-{}^{79}Br$	131976	16	131967.3	1.0	−0.4	U			R11	1.5	78Di09	
	131974	17			−0.3	U			R11	1.5	78Di09	
$C_5\ N\ H_5-{}^{79}Br$	123870	7	123861.5	1.0	−0.8	U			R11	1.5	78Di09	
	123871	14			−0.5	U			R11	1.5	78Di09	
$C_5\ O\ H_3-{}^{79}Br$	100061	15	100052.0	1.0	−0.4	U			R11	1.5	78Di09	
	100057	20			−0.2	U			R11	1.5	78Di09	
$C_4\ N\ O\ H-{}^{79}Br$	87489	20	87476.0	1.0	−0.4	U			R11	1.5	78Di09	
$^{79}Kr-u$	−79981	52	−79917	4	1.2	U			GS2	1.0	05Li24	*
$^{79}Sr-u$	−70290.0	2.8	−70290.2	2.7	−0.1	1	91	91 ^{79}Sr	LZ1	1.0	23Xi01	
$^{79}Y-u$	−62054	86	−62253.3	0.5	−2.3	U			LZ1	1.0	18Xi04	
$^{79}Y-{}^{39}K_{2.026}$	11277.41	0.45				2			MS2	1.0	26Er.A	
$^{79}Zr-{}^{39}K_{2.026}$	23282.9	2.5				2			MS2	1.0	26Er.A	
$^{79}Zn-{}^{85}Rb_{.929}$	24582.4	4.2	24585.7	1.9	0.8	1	21	21 ^{79}Zn	MA8	1.0	08Ba54	
$^{79}Ga-{}^{85}Rb_{.929}$	14798.4	1.7	14798.9	1.3	0.3	1	57	57 ^{79}Ga	MA8	1.0	19Hu15	
$^{79}Ge-{}^{85}Rb_{.929}$	7365.76	0.77				2			JY2	1.0	24Fl.A	
$^{79}Rb-{}^{85}Rb_{.929}$	5934	8	5937.3	2.1	0.4	U			MA2	1.0	94Ot01	
	5937.2	2.3			0.0	1	83	83 ^{79}Rb	MA8	1.0	07Ke09	
$^{79}Sr-{}^{85}Rb_{.929}$	11655	9	11657.1	2.7	0.2	1	9	9 ^{79}Sr	MA2	1.0	94Ot01	
$^{79}Zn-{}^{84}Kr_{.940}$	25830.8	3.4	25830.5	1.9	−0.1	1	31	31 ^{79}Zn	JY1	1.0	22Gi08	
$^{79}Zn^{m}-{}^{84}Kr_{.940}$	26842	12	26841	8	−0.0	1	40	40 $^{79}Zn^{m}$	JY1	1.0	23Ni10	
$^{77}Se\ H_2-{}^{79}Br$	17239	8	17226.6	1.0	−1.0	U			R11	1.5	78Di09	
$^{78}Se\ H-{}^{79}Br$	6806	8	6796.6	1.0	−0.8	U			R11	1.5	78Di09	
$^{79}Zn-{}^{79}Ga$	9788.8	8.5	9786.8	2.3	−0.2	1	7	5 ^{79}Zn	TR1	1.0	25Ja07	
$^{79}Zn^{m}-{}^{79}Ga$	10796	13	10798	8	0.1	1	35	34 $^{79}Zn^{m}$	TR1	1.0	25Ja07	
$^{79}Br-{}^{79}Rb$	−5652.7	5.0	−5652.3	2.3	0.1	1	20	17 ^{79}Rb	RI1	1.0	18Ki21	
$^{79}Kr-{}^{79}Rb$	−3870	41	−3907	4	−0.9	U			RI1	1.0	18Ki21	*
$^{79}Sr-{}^{79}Rb$	5704	17	5720	3	0.9	U			RI1	1.0	18Ki21	
$^{79}Y-{}^{79}Sr$	8048	11	8036.9	2.7	−1.0	U			RKM	1.0	25Ki14	
$^{79}Br-{}^{78}Kr$	−2072	30	−2028.7	1.1	1.0	U			R11	1.5	78Di09	
$^{77}Rb-{}^{79}Rb_{.487}\ {}^{75}Rb_{.513}$	−1010	40	−996.3	1.7	0.1	U			P20	2.5	82Au01	
$^{77}Rb-{}^{79}Rb_{.325}\ {}^{76}Rb_{.675}$	−1060	40	−996.2	1.6	0.6	U			P20	2.5	82Au01	
	−990	70			−0.0	U			P20	2.5	82Au01	
$^{78}Rb^{x}-{}^{79}Rb_{.494}\ {}^{77}Rb_{.506}$	940	40	919	12	−0.2	U			P20	2.5	82Au01	
$^{78}Se(n,\gamma)^{79}Se$	6962.6	0.3	6962.83	0.13	0.8	2					79Br.A	Z
	6962.2	0.3			2.1	2			BNn		81En07	Z
	6963.11	0.17			−1.6	2			Bdn		06Fi.A	
$^{78}Se(d,p)^{79}Se$	4756	6	4738.27	0.13	−3.0	B			MIT		64Sp12	
$^{78}Kr(d,p)^{79}Kr$	5980	50	6111	3	2.6	U			Yal		56Bl10	
$^{78}Kr(^{3}He,d)^{79}Rb$	−1585	10	−1579.8	2.0	0.5	U			Phi		87St11	
$^{79}Zn(\beta^{-})^{79}Ga$	8550	240	9116.4	2.1	2.4	U			Stu		86Ek01	
$^{79}Ga(\beta^{-})^{79}Ge$	6770	80	6923.9	1.4	1.9	o			Stu		77Al17	
	7000	80			−1.0	o			Stu		86Ek01	
	6979	40			−1.4	U			Stu		00Me.A	
$^{79}Ge(\beta^{-})^{79}As$	4300	200	4164	5	−0.7	U					70Ka04	
	4110	100			0.5	U			Stu		81Al20	
$^{79}As(\beta^{-})^{79}Se$	2230	50	2281	5	1.0	U					61Ku09	*
$^{79}Se(\beta^{-})^{79}Br$	160	5	150.5	1.0	−1.9	U					49Pa.A	
$^{79}Kr(\beta^{+})^{79}Br$	1612	10	1626	3	1.4	2					52Be55	
	1620	5			1.2	2					54Th39	
	1635	5			−1.8	2					64Bo25	
$^{79}Rb(\beta^{+})^{79}Kr$	3530	50	3639	4	2.2	U					71Li02	*
	3720	90			−0.9	U					72Br31	*
	3650	70			−0.2	U			IRS		93Al03	
$^{79}Sr(\beta^{+})^{79}Rb$	5259	78	5328	3	0.9	U			BNL		81Li12	
	5059	67			4.0	B			Ors		82De36	
$^{79}Y(\beta^{+})^{79}Sr$	7120	450	7486.4	2.5	0.8	U					92Mu12	
$*^{79}Kr-u$	$M-A$=–74437(30) keV for mixture gs+m at 129.77 $7/2^{+}$ keV										Nub269	**
$*^{79}Zn-{}^{79}Ga$	typo in reference Actually frequency ratio 1.000124015(110)										25Ja07	**
$*^{79}Kr-{}^{79}Rb$	D_M=–3800(9) μu for ^{79}Kr gs+m mixture at 129.77 keV; $M-A$=–74342.7(9.3)keV										Nub269	**
$*^{79}As(\beta^{-})^{79}Se$	E_{β^-}=1700(50) to 527.93 $3/2^{-}$ level, and other E_{β^-}										Ens167	**
$*^{79}Rb(\beta^{+})^{79}Kr$	E_{β^+}=1825(50) 2010(90) respectively, to $3/2^{+}$ level at 688.17 keV										Ens167	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{80}Ga−u		−63441	129	−63579	3	−0.4	U			GT2	2.5	08Su19	
		−63567	52			−0.2	U			TR1	1.0	20Re04	
C_6 H_8−^{80}Se		146068.5	2.9	146078.4	1.0	1.4	U			M15	2.5	63Ri07	
C_6 H_8−^{80}Kr		146225.7	4.6	146222.0	0.7	−0.3	U			M15	2.5	63Ri07	
		146235	18			−0.5	U			R11	1.5	78Di09	
		146215	16			0.3	U			R11	1.5	78Di09	
C_5 O H_4−^{80}Kr		109834	20	109836.4	0.7	0.1	U			R11	1.5	78Di09	
^{80}Y−u		−65720	190	−65645	7	0.4	U			1.0	1.0	98Is06	
		−66664	86			11.8	F			CS1	1.0	01La31	*
		−65600	200			−0.2	U			CS1	1.0	08Go23	
^{80}Y O−^{96}Mo		24594.6	6.7				2			JY1	1.0	06Ka48	
^{80}Zr−u		−59600	1600	−58353.4	1.6	0.8	B			1.0	1.0	98Is06	
		−59740	161			8.6	F			CS1	1.0	01La31	*
^{80}Zr−^{39}$K_{2.051}$		16084.6	1.6				2			MS2	1.0	26Er.A	
^{80}Zn−^{88}$Rb_{.909}$		25165.2	7.3	25163.2	2.3	−0.3	1	10	10 ^{80}Zn	JY1	1.0	08Ha23	
^{80}Ga−^{88}$Rb_{.909}$		17034.9	3.1				2			JY1	1.0	08Ha23	
^{80}Ge−^{88}$Rb_{.909}$		5964.9	2.2				2			JY1	1.0	08Ha23	
^{80}Kr−^{86}$Kr_{.930}$		−488.9	1.1	−489.6	0.7	−0.6	1	45	45 ^{80}Kr	MS1	1.0	06Ri15	
^{80}Zn−^{85}$Rb_{.941}$		27559.1	3.0	27554.9	2.3	−1.4	1	60	60 ^{80}Zn	MA8	1.0	08Ba54	
^{80}Kr−^{85}$Rb_{.941}$		−614.5	1.7	−615.8	0.7	−0.8	1	19	19 ^{80}Kr	MA8	1.0	06Ro11	
		−627.1	9.6			1.2	U			MA8	1.0	10Na13	*
^{80}Rb−^{85}$Rb_{.941}$		5528	8	5524.3	1.8	−0.5	U			MA2	1.0	94Ot01	
		5522.3	2.0			1.0	1	85	85 ^{80}Rb	MA8	1.0	07Ke09	
^{80}Sr−^{85}$Rb_{.941}$		7531	8	7523	4	−1.0	2			MA2	1.0	94Ot01	
		7513	14			0.7	U			MA8	1.0	05Si34	
		7521.3	4.2			0.5	2			SH1	1.0	11Ha08	
^{80}Zr O−^{85}$Rb_{1.129}$		35322	84	36150.6	1.6	9.9	F			MS1	1.0	21Ha47	*
^{80}Se ^{35}Cl−^{78}Se ^{37}Cl		2164.8	1.4	2162.7	1.0	−0.4	U			H18	4.0	64Ba03	
		2160.8	9.2			0.1	U			H40	2.5	85El01	
^{80}Zn−^{80}Rb		22023.4	3.8	22030.6	2.5	1.9	1	45	29 ^{80}Zn	TR1	1.0	25Ja07	
^{80}As−^{80}Kr		6096.5	3.5				2			MS1	1.0	07Bo50	
^{80}Rb−^{80}Kr		6128	12	6140.1	2.0	1.0	U			RI1	1.0	18Ki21	
^{80}Sr−^{80}Kr		8110	16	8139	4	1.8	F			RI1	1.0	18Ki21	*
^{80}Kr−^{79}Br		−1955	28	−1959.4	1.2	−0.1	U			R11	1.5	78Di09	
^{80}Kr−^{78}Kr		−4046	30	−3988.1	0.8	1.3	U			R11	1.5	78Di09	
^{79}Rb−^{80}$Rb_{.658}$ ^{77}$Rb_{.342}$		−1218	27	−1140.6	2.3	1.1	U			P20	2.5	82Au01	
^{79}Rb−^{80}$Rb_{.494}$ ^{78}$Rb^{x}_{.506}$		−1313	24	−1317	7	−0.1	U			P20	2.5	82Au01	
^{80}Se(p,α)^{77}As		1020.0	2.8	1019.7	0.9	−0.1	1	10	10 ^{80}Se	NDm		82Zu04	
^{80}Kr(^{3}He,^{6}He)^{77}Kr		−10398	24	−10384.5	2.1	0.6	U					87Mo06	
^{80}Se(d,α)^{78}As		5755	12	5768	10	1.1	2			Phi		77Mo13	
^{80}Se(p,t)^{78}Se		−8395.1	3.0	−8394.3	0.9	0.3	–			NDm		82Zu04	
	ave.	−8394.1	2.1			−0.1	1	18	18 ^{80}Se			average	
^{80}Kr(α,^{6}He)^{78}Kr−^{78}Kr()^{76}Kr		1432	10	1450	4	1.8	1	17	16 ^{76}Kr			82Mo23	
^{78}Kr(^{3}He,n)^{80}Sr		2990	30	2993	3	0.1	U					79Al19	
^{80}Se(d,^{3}He)^{79}As		−5921	7	−5919	5	0.3	2			Ors		83Ro08	*
		−5921	13			0.2	2			Hei		83Wi14	
^{80}Se(t,α)^{79}As		8407	10	8401	5	−0.6	2			Phi		83Mo09	
^{80}Se(p,d)^{79}Se		−7687.6	3.0	−7688.7	0.9	−0.4	R			NDm		82Zu04	
^{79}Br(n,γ)^{80}Br		7892.11	0.20	7892.27	0.13	0.8	–			ILn		78Do06	Z
		7892.41	0.18			−0.8	–			Bdn		06Fi.A	
^{79}Br(d,p)^{80}Br		5640	20	5667.71	0.13	1.4	U			Mtr		72Ch33	
^{79}Br(n,γ)^{80}Br	ave.	7892.28	0.13	7892.27	0.13	−0.0	1	100	89 ^{79}Br			average	
^{80}Zn(β^-)^{80}Ga		7540	200	7571	4	0.2	U			Stu		86Ek01	
		7150	150			2.8	U			Trs		86Gi07	
^{80}Ga(β^-)^{80}Ge		10000	300	10312	4	1.0	o			Stu		81Al20	
		10380	120			−0.6	U			Stu		86Ek01	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{80}Ge(β^-)^{80}As	2640	70	2679	4	0.6	U			Stu		77Al17	
	2630	20			2.4	U			Trs		86Gi07	
^{80}As(β^-)^{80}Se	6000	200	5545	3	−2.3	U					59Me68	
	5470	90			0.8	U			Trs		86Gi07	
^{80}Se(t,^{3}He)^{80}As	−5560	25	−5527	3	1.3	U			LAl		79Aj02	
^{80}Br(β^+)^{80}Se	1884	10	1870.5	0.3	−1.4	U					54Li19	
	1872	7			−0.2	U					69Ka06	
^{80}Se(p,n)^{80}Br	−2655.2	2.8	−2652.8	0.3	0.9	U			Tkm		63Ok01	
	−2652.5	3.0			−0.1	U			Oak		64Jo11	Z
	−2653.2	5.			0.1	U					70Fi03	
	−2652.81	0.31			0.0	1	99	89 ^{80}Br	PTB		92Bo02	Z
^{80}Br(β^-)^{80}Kr	1970	30	2004.2	1.1	1.1	U					52Fu04	
	2040	20			−1.8	U					54Li19	
	1997	10			0.7	U					69Ka06	
^{80}Rb(β^+)^{80}Kr	5120	500	5719.5	1.8	1.2	U					61Ho13	
	5500	350			0.6	U					75We23	*
	5650	100			0.7	U			IRS		93Al03	
^{80}Kr(p,n)^{80}Rb	−6484.0	20.	−6501.9	1.8	−0.9	U					72Ja.A	
^{80}Y(β^+)^{80}Sr	6952	152	9163	7	14.5	F			BNL		81Li12	*
	6934	242			9.2	F			Ors		82De36	*
	6200	600			4.9	F					96Sh27	*
*^{80}Y−u	F : below lower limit M>–65890(90) μu –61376(83) keV determined in reference										03Ba18	**
*^{80}Zr−u	F : other results in same paper not trusted, see ^{80}Y and ^{68}Se										GAu	**
*^{80}Kr−^{85}Rb$_{.941}$	Only one measurement										GAu	**
*^{80}Zr O−^{85}Rb$_{1.129}$	conflict with new measurement by the same group 26Er.A										HWJ265	**
*^{80}Sr−^{80}Kr	Compromised measurement due to the influence of neighboring ToF peaks										Sar201	**
*^{80}Se(d,^{3}He)^{79}As	Originally –5927(7), see ^{74}Se(d,^{3}He)										AHW	**
*^{80}Rb(β^+)^{80}Kr	E_{β^+}=3860(350) to 2$^+$ level at 616.60 keV										Ens058	**
*^{80}Y(β^+)^{80}Sr	F : below lower limit Q_{β^-} > 8929(23) keV determined in reference										03Ba18	**
^{81}Ga−u	−61853	33	−61865	3	−0.4	U			TR1	1.0	20Re04	
^{81}Ge−u	−71710	240	−71167.1	2.2	0.9	U			GT2	2.5	08Kn.A	*
C_6 H_9−^{81}Br	154135.3	3.8	154137.1	1.0	0.2	U			M15	2.5	63Ri07	
	154143	17			−0.2	U			R11	1.5	78Di09	
	154134	10			0.2	U			R11	1.5	78Di09	
C_5 N H_7−^{81}Br	141561	10	141561.0	1.0	0.0	U			R11	1.5	78Di09	
	141553	18			0.3	U			R11	1.5	78Di09	
C_5 O H_5−^{81}Br	117742	12	117751.6	1.0	0.5	U			R11	1.5	78Di09	
C_4 O_2 H−^{81}Br	81356	20	81366.1	1.0	0.3	U			R11	1.5	78Di09	
C_4 ^{13}C O H_4−^{81}Br	113275	14	113281.4	1.0	0.3	U			R11	1.5	78Di09	
^{81}Rb−u	−80958	41	−81006	5	−1.2	U			GS2	1.0	05Li24	*
^{81}Y O−^{97}Mo	18352.0	5.8	18345.4	2.7	−1.1	1	21	21 ^{81}Y	JY1	1.0	06Ka48	
^{81}Y−u	−70554.1	3.0	−70552.3	2.7	0.6	1	79	79 ^{81}Y	LZ1	1.0	23Xi01	
^{81}Zr−u	−61755	99	−61789	11	−0.3	U			LZ1	1.0	18Xi04	
^{81}Zr−^{41}K$_{1.976}$	13644	11				2			MS1	1.0	21Ha47	
^{81}Ga−^{88}Rb$_{.920}$	19723.5	3.5	19725	3	0.3	1	90	90 ^{81}Ga	JY1	1.0	08Ha23	
^{81}Ge−^{88}Rb$_{.920}$	10422.6	2.2				2			JY1	1.0	08Ha23	
^{81}As−^{88}Rb$_{.920}$	3721.9	3.3	3722.0	2.8	0.0	1	74	74 ^{81}As	JY1	1.0	08Ha23	
^{81}Zn−^{85}Rb$_{.953}$	34467.0	5.4	34465	5	−0.4	1	76	76 ^{81}Zn	MA8	1.0	08Ba54	
^{81}Rb−^{85}Rb$_{.953}$	3063	8	3058	5	−0.6	–			MA2	1.0	94Ot01	
	3055.4	9.2			0.3	–			SH1	1.0	11Ha08	*
ave.	3060	6			−0.2	1	76	76 ^{81}Rb			average	
^{81}Sr−^{85}Rb$_{.953}$	7278	8	7276	3	−0.3	2			MA2	1.0	94Ot01	
	7272	12			0.3	U			MA8	1.0	05Si34	
	7275.3	3.7			0.1	2			SH1	1.0	11Ha08	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{81}Se−^{80}Kr$_{1.013}$		2704.2	2.4	2701.9	1.2	−0.9	1	24	16 ^{81}Se	MS1	1.0	07Bo50	*
^{80}Se H−^{81}Br		8023	8	8058.7	1.4	3.0	B			R11	1.5	78Di09	
^{80}Kr H−^{81}Br		7922	18	7915.1	1.3	−0.3	U			R11	1.5	78Di09	
^{81}Zn−^{81}Ga		12258.8	8.9	12265	5	0.7	1	34	24 ^{81}Zn	TR1	1.0	25Ja07	
^{81}Br−^{81}Sr		−6899.5	4.9	−6923	4	−4.8	B			RI1	1.0	18Ki21	
^{81}Rb−^{81}Sr		−4193	27	−4217	6	−0.9	U			RI1	1.0	18Ki21	*
^{81}Br−H ^{79}Br		−9865	13	−9874.5	1.5	−0.3	U			M15	2.5	63Ri07	
^{81}Br−^{80}Kr		−91	32	−90.1	1.3	0.0	U			R11	1.5	78Di09	
^{81}Br−^{79}Br		−2020	32	−2049.5	1.5	−0.6	U			R11	1.5	78Di09	
		−2014	35			−0.7	U			R11	1.5	78Di09	
^{79}Rb−^{81}Rb$_{.325}$ ^{78}Rb$^{x}_{.675}$		−1130	30	−1148	9	−0.2	U			P20	2.5	82Au01	Y
^{80}Rb−^{81}Rb$_{.494}$ ^{79}Rb$_{.506}$		927	29	928	3	0.0	U			P20	2.5	82Au01	Y
^{80}Se(n,γ)^{81}Se		6700.9	0.5	6700.8	0.3	−0.1	–			BNn		81En07	Z
		6700.9	0.5			−0.1	–			Bdn		06Fi.A	
^{80}Se(d,p)^{81}Se		4490	6	4476.3	0.3	−2.3	U			MIT		64Sp12	
		4477.5	3.0			−0.4	U			NDm		82Zu04	
^{80}Se(n,γ)^{81}Se	ave.	6700.9	0.4	6700.8	0.3	−0.2	1	97	75 ^{81}Se			average	
^{81}Br(γ,n)^{80}Br		−10130	35	−10159.4	1.4	−0.8	U			Phi		60Ge01	
^{80}Kr(d,p)^{81}Kr		5660	15	5649.8	1.2	−0.7	U			Tex		75Ch11	*
		5646	4			1.0	1	10	7 ^{81}Kr	Oak		86Bu18	
^{80}Kr(^{3}He,d)^{81}Rb		−637	10	−641	5	−0.4	1	24	24 ^{81}Rb	Phi		87St11	
^{81}Zr(εp)^{80}Sr		4700	200	5466	11	3.8	B					99Hu05	
^{81}Ga(β^-)^{81}Ge		8320	150	8665	4	2.3	U			Stu		81Al20	
^{81}Ge(β^-)^{81}As		6230	120	6242	3	0.1	U			Stu		81Al20	*
^{81}As(β^-)^{81}Se		3800	200	3855.6	2.8	0.3	U					60Mo01	
		3730	100			1.3	U			Stu		77Al17	
^{81}Se(β^-)^{81}Br		1600	50	1588.1	1.3	−0.2	U					60Ku06	
		1560	50			0.6	U					67Yt03	
^{81}Kr(ε)^{81}Br		280.7	0.5	280.9	0.5	0.3	1	89	84 ^{81}Kr			88Ax01	*
^{81}Br(p,n)^{81}Kr		−1062	4	−1063.2	0.5	−0.3	U					84Fi.A	
^{81}Br(^{3}He,t)^{81}Kr		−296	6	−299.4	0.5	−0.6	U					84Bu23	
^{81}Br(^{3}He,t)^{81}Kr−^{51}V()^{51}Cr		470.6	1.8	471.5	0.5	0.5	U			Pri		82Ko06	*
^{81}Rb(β^+)^{81}Kr		2260	30	2240	5	−0.7	U					75Va24	*
		2290	50			−1.0	U					77Li14	
^{81}Sr(β^+)^{81}Rb		3990	30	3928	6	−2.1	U					73Br32	*
^{81}Y(β^+)^{81}Sr		5408	86	5809	4	4.7	B			BNL		81Li12	
		5620	89			2.1	U			Ors		82De36	
^{81}Zr(β^+)^{81}Y		7160	290	8163	11	3.5	B			Ors		82De36	
*^{81}Ge−u	$M-A$=–66454(93) keV for mixture gs+m at 679.14 keV											Nub269	**
*^{81}Rb−u	$M-A$=–75369(29) keV for mixture gs+m at 86.31 keV											Nub269	**
*^{81}Rb−^{85}Rb$_{.953}$	D_M=3148.1(9.2) keV for ^{81}Rb^m at 86.31(0.07) keV; $M-A$=–75373.1(8.6) keV											Nub269	**
*^{81}Se−^{80}Kr$_{1.013}$	D_M=2814.8(2.4) μu for ^{81}Se^m at 103.97(0.01)keV; $M-A$=–76283.2(2.4) keV											Nub269	**
*^{81}Rb−^{81}Sr	D_M=4147.0(0.9) μu for mixture gs+m at 86.31 keV; $M-A$=–75391.5 keV											Nub269	**
*^{80}Kr(d,p)^{81}Kr	Original value 5610(15) reinterpreted as going to 49.57 level											76Me08	**
*^{81}Ge(β^-)^{81}As	Q_{β^-}=6230(120); and 6930(280) from ^{81}Ge^m at 679.14 keV											Nub269	**
*^{81}Kr(ε)^{81}Br	LM=0.42(0.05), $Q(\varepsilon)$=4.7(0.5) to 5/2$^-$ level at 275.985 keV											Ens08a	**
*^{81}Br(^{3}He,t)^{81}Kr−^{51}V()^{51}Cr	Q−Q to 456.89(0.03) level=13.7(1.8) keV											GAu	**
*^{81}Rb(β^+)^{81}Kr	E_{β^+}=1050(30) to 1/2$^-$ level at 190.64 keV											Ens08a	**
*^{81}Sr(β^+)^{81}Rb	E_{β^+}=2684(30) to 301.241 3/2$^-$ level, and other E_{β^+}											Ens08a	**
^{82}Ga−u		−56812	268	−56823.7	2.5	−0.0	U			GT1	1.5	04Ma.A	
		−56870	33			1.4	U			TR1	1.0	20Re04	*
^{82}Ge−u		−70400	129	−70225.0	1.0	0.5	U			GT2	2.5	08Su19	
C_6 H_{10}−^{82}Se		161545.0	4.6	161550.8	0.5	0.5	U			M15	2.5	63Ri07	
C_6 H_{10}−^{82}Kr		164769.8	3.4	164769.167	0.006	−0.1	U			M15	2.5	63Ri07	
		164787	14			−0.8	U			R11	1.5	78Di09	
		164784	16			−0.6	U			R11	1.5	78Di09	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
C_5 N $H_8-{}^{82}$Kr		152200	25	152193.107	0.006	−0.2	U			R11	1.5	78Di09	
C_5 O $H_6-{}^{82}$Kr		128396	20	128383.658	0.006	−0.4	U			R11	1.5	78Di09	
^{82}Rb−u		−81775	39	−81791	3	−0.4	U			GS2	1.0	05Li24	*
^{82}Sr−u		−81604	63	−81604	4	0.0	U			GS2	1.0	05Li24	
^{82}Y O−^{98}Mo		16441.2	5.9				2			JY1	1.0	06Ka48	
^{82}Zr−u		−68312	11	−68293.9	1.4	1.6	U			LZ1	1.0	18Xi04	
^{82}Nb−$^{41}K_2$		20187.7	2.4				2			MS2	1.0	26Er.A	
^{82}Ga−^{88}Rb$_{.932}$		25830.4	2.6	25830.2	2.5	−0.1	1	91	91 ^{82}Ga	JY1	1.0	08Ha23	
^{82}Ge−^{88}Rb$_{.932}$		12427.9	2.4	12428.9	1.0	0.4	1	18	17 ^{82}Ge	JY1	1.0	08Ha23	
^{82}As−^{88}Rb$_{.932}$		7392.6	4.0	7391.4	2.5	−0.3	1	41	40 ^{82}As	JY1	1.0	08Ha23	
$^{82}As^m-{}^{88}$Rb$_{.932}$		7532.5	4.0	7533.2	2.6	0.2	U			JY1	1.0	08Ha23	
^{82}Zr O−^{87}Rb$_{1.126}$		28879.9	2.7	28883.4	1.4	1.3	1	28	28 ^{82}Zr	MS1	1.0	21Ha47	
^{82}Kr−^{86}Kr$_{.953}$		−1329.4	1.1	−1330.771	0.005	−1.2	U			MS1	1.0	06Ri15	
		−1330.7684	0.0094			−0.3	1	26	25 ^{82}Kr	FS1	1.0	13Ho22	
^{82}Zn−^{85}Rb$_{.965}$		39697.0	3.3	39697	3	0.1	1	85	85 ^{82}Zn	MA8	1.0	13Wo06	
^{82}Kr−^{85}Rb$_{.965}$		−1394.9	2.6	−1395.944	0.006	−0.4	U			MA8	1.0	06Ro11	
$^{82}Rb^m-{}^{85}$Rb$_{.965}$		3407	9	3406.0	2.8	−0.1	U			MA2	1.0	94Ot01	
		3406.0	2.8				2			MA8	1.0	05Gu37	
^{82}Sr−^{85}Rb$_{.965}$		3517	8	3519	4	0.3	1	30	30 ^{82}Sr	MA2	1.0	94Ot01	
^{82}Zr−^{85}Rb$_{.965}$		16830.4	1.7	16829.0	1.4	−0.8	1	72	72 ^{82}Zr	JY1	1.0	19Vi05	
^{82}Se ^{35}Cl−^{80}Se ^{37}Cl		3128.92	0.63	3127.8	1.0	−0.7	1	37	32 ^{80}Se	H40	2.5	85El01	
^{80}Se $H_2-{}^{82}$Kr		18665	18	18690.8	1.0	1.0	U			R11	1.5	78Di09	
		18671	19			0.7	U			R11	1.5	78Di09	
^{82}Kr−^{84}Kr$_{.976}$		−140.6277	0.0051	−140.627	0.005	0.1	1	78	75 ^{82}Kr	FS1	1.0	13Ho22	
^{81}Br H−^{82}Se		7419	8	7413.7	1.2	−0.4	U			R11	1.5	78Di09	
^{81}Br H−^{82}Kr		10662	20	10632.1	1.0	−1.0	U			R11	1.5	78Di09	
^{82}Zn−^{82}Ga		11399.5	7.5	11398	4	−0.2	1	24	15 ^{82}Zn	TR1	1.0	25Ja07	
^{82}Ge−^{82}Kr		16294.1	1.1	16293.9	1.0	−0.2	1	83	83 ^{82}Ge	RKM	1.0	24Xi06	
^{82}As−^{82}Kr		11255.5	3.3	11256.3	2.5	0.3	1	60	60 ^{82}As	RKM	1.0	24Xi06	*
^{82}Se−^{82}Kr		3222	16	3218.4	0.5	−0.2	U			R11	1.5	78Di09	
		3218	22			0.0	U			R11	1.5	78Di09	
		3216.1	1.6			1.0	U			H45	1.5	93Nx01	
		3218.36	0.52			0.1	1	92	92 ^{82}Se	MS1	1.0	13Li01	
		3223.1	3.4			−1.4	U			RKM	1.0	24Xi06	
^{82}Kr−^{78}Se H_3		−27269	35	−27303.22	0.19	−0.7	U			R11	1.5	78Di09	
^{80}Se $H_3-{}^{82}$Kr		26466	32	26515.8	1.0	1.0	U			R11	1.5	78Di09	
^{82}Kr−^{81}Br		−2805	32	−2807.1	1.0	−0.0	U			R11	1.5	78Di09	
^{82}Se H−^{82}Kr		11082	40	11043.4	0.5	−0.6	U			R11	1.5	78Di09	
^{79}Rb−^{82}Rb$_{.241}$ ^{78}Rb$^x_{.760}$		−1536	29	−1627	10	−1.3	U			P20	2.5	82Au01	Y
^{81}Rb−^{82}Rb$_{.741}$ ^{78}Rb$^x_{.260}$		−1680	40	−1617	6	0.6	U			P20	2.5	82Au01	Y
^{80}Rb−^{82}Rb$_{.325}$ ^{79}Rb$_{.675}$		440	40	379.4	2.4	−0.6	U			P20	2.5	82Au01	Y
^{82}Kr(^{3}He,^{6}He)^{79}Kr		−8822	31	−8810	3	0.4	U					87Mo06	
^{82}Se(^{14}C,^{16}O)^{80}Ge		−449	60	−301.7	2.1	2.5	U			Ors		83Be.C	
^{82}Se(^{18}O,^{20}Ne)^{80}Ge		−2020	40	−1799.4	2.1	5.5	B			Hei		83Wi14	*
^{82}Se(p,t)^{80}Se		−7496.1	3.0	−7495.3	0.9	0.3	1	9	8 ^{80}Se	NDm		82Zu04	
^{82}Se(d,^{3}He)^{81}As		−6864	10	−6856.1	2.7	0.8	–			Ors		83Ro08	*
		−6861	18			0.3	U			Hei		83Wi14	
^{82}Se(t,α)^{81}As		7467	6	7464.3	2.7	−0.4	–			Phi		82Mo04	
^{82}Se(d,^{3}He)^{81}As	ave.	−6856	5	−6856.1	2.7	0.0	1	27	26 ^{81}As			average	
^{82}Se(p,d)^{81}Se		−7051.8	2.8	−7051.7	0.9	0.0	1	11	10 ^{81}Se	NDm		82Zu04	
^{81}Br(n,γ)^{82}Br		7592.80	0.20	7592.94	0.12	0.7	–			ILn		78Do06	Z
		7593.02	0.15			−0.5	–			Bdn		06Fi.A	
^{81}Br(d,p)^{82}Br		5400	20	5368.38	0.12	−1.6	U			Mtr		72Ch33	
^{81}Br(n,γ)^{82}Br	ave.	7592.94	0.12	7592.94	0.12	0.0	1	100	94 ^{81}Br			average	
^{82}Ge(β^-)^{82}As		4700	140	4692.4	2.5	−0.1	U			Stu		81Al20	
$^{82}As^m$(IT)^{82}As		132.1	0.2				2					14Mi16	
		131.2	0.6	132.10	0.20	1.5	U					15Et01	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{82}As(β^-)^{82}Se	7270	200	7487.3	2.4	1.1	U					70Va31	*
	6360	200			5.6	B					70Ka04	*
	7740	30			−8.4	C			Stu		00Me.A	*
	7531	21			−2.1	U			Stu		04Ga44	*
^{82}Se(t,^{3}He)^{82}As	−7500	25	−7468.7	2.4	1.3	U			LAl		79Aj02	
^{82}Br(β^-)^{82}Kr	3092.9	1.0	3093.1	1.0	0.2	1	94	94 ^{82}Br			56Wa24	*
^{82}Rb(β^+)^{82}Kr	4400	15	4404	3	0.3	U					69Be74	*
	4420	60			−0.3	U			IRS		93Al03	*
^{82}Kr(p,n)^{82}Rb	−5161	20	−5186	3	−1.3	U					72Ja.A	
^{82}Rb^m(IT)^{82}Rb	69.0	1.5				3					Ens196	
^{82}Y(β^+)^{82}Sr	7868	185	7949	7	0.4	U			BNL		81Li12	
	7793	123			1.3	U			Ors		82De36	
^{82}Zr(β^+)^{82}Y	4000	500	4449	6	0.9	F			Ors		82De36	*
*^{82}Ga−u	corr. for gs+m of ^{82}Rb as calibrant, -52939(23) keV assuming mainly iso.										20Re04	**
*^{82}Rb−u	$M-A$=–76138(30) keV for mixture gs+m at 69.0(1.5) keV										Nub269	**
*^{82}As−^{82}Kr	could be the gs.										HWJ256	**
*^{82}Se(^{18}O,^{20}Ne)^{80}Ge	Recalibrated to ^{64}Ni()^{62}Fe=–1938(15) keV										AHW	**
*^{82}Se(d,^{3}He)^{81}As	Originally –6870(10), see ^{74}Se(d,^{3}He)										AHW	**
*^{82}As(β^-)^{82}Se	E_{β^-}=7200(200) to ground state (80%) and 654.75 2^+ level (10%)										Ens035	**
*^{82}As(β^-)^{82}Se	E_{β^-}=3600(200) from ^{82}As^m at 132.1 to 5^- level at 2893.70 and higher levels										Ens035	**
*^{82}As(β^-)^{82}Se	and E_{β^-}=7625(22) from ^{82}As^m at 132.1 keV										00Me.A	**
*^{82}As(β^-)^{82}Se	Average of 3 branches; and 7677(17) average of 2 branches from ^{82}As^m at 132.1										04Ga44	**
*^{82}Br(β^-)^{82}Kr	E_{β^-}=444(1) to 4^- level at 2648.360 keV										Ens035	**
*^{82}Rb(β^+)^{82}Kr	E_{β^+}=3350(60); and 800(15) from ^{82}Rb^m at 69.0(1.5) to 4^- level at 2648.360										Ens035	**
*^{82}Rb(β^+)^{82}Kr	Q_{β^+}=4360(100); and 4510(60) of ^{82}Rb^m at 69.0(1.5) keV										Nub269	**
*^{82}Zr(β^+)^{82}Y	F : for 2.5(0.1) m activity, but Ensdf adopts 32(5) s										Ens035	**
^{83}Ga−u	−52881	27	−52880.2	2.6	0.0	U			TR1	1.0	20Re04	
	−52883.6	7.6			0.4	1	12	12 ^{83}Ga	RKM	1.0	24Xi06	
^{83}Ge−u	−65626	268	−65460.9	2.6	0.4	o			GT1	1.5	04Ma.A	
	−65270	320			−0.6	U			OR1	1.0	06Ha62	
	−65276	129			−0.6	U			GT2	2.5	08Su19	
	−65460.0	9.9			−0.1	U			RKM	1.0	24Xi06	
^{83}As−u	−74677	129	−74793	3	−0.4	U			GT2	2.5	08Su19	
	−74793.8	11.8			0.1	U			RKM	1.0	24Xi06	
C_6 H_{11}−^{83}Kr	171946.8	3.4	171948.836	0.010	0.2	U			M15	2.5	63Ri07	
	171948	16			0.0	U			R11	1.5	78Di09	
C_5 N H_9−^{83}Kr	159344	25	159372.776	0.010	0.8	U			R11	1.5	78Di09	
	159360	19			0.4	U			R11	1.5	78Di09	
C_5 O H_7−^{83}Kr	135543	25	135563.327	0.010	0.5	U			R11	1.5	78Di09	
^{83}Y O−^{98}Mo$_{1.010}$	12941	20				2			JY1	1.0	06Ka48	*
^{83}Zr−^{41}K$_{2.024}$	6501.57	0.70	6501.4	0.7	−0.2	1	99	99 ^{83}Zr	MS1	1.0	21Ha47	
^{83}Zr O−^{98}Mo$_{1.010}$	19697.9	6.9	19692.8	0.7	−0.7	U			JY1	1.0	06Ka48	
^{83}Nb−u	−61850	174	−61889	3	−0.2	U			LZ1	1.0	18Xi04	
^{83}Nb−^{41}K$_{2.024}$	15372.6	3.8	15377	3	1.1	1	80	80 ^{83}Nb	MS2	1.0	26Er.A	
^{83}Ga−^{88}Rb$_{.943}$	30749.7	2.8	30749.2	2.6	−0.2	1	88	88 ^{83}Ga	JY1	1.0	08Ha23	
^{83}Ge−^{88}Rb$_{.943}$	18168.5	2.6				2			JY1	1.0	08Ha23	
^{83}As−^{88}Rb$_{.943}$	8836.3	3.0				2			JY1	1.0	08Ha23	
^{80}Se H_3−^{83}Kr	25825	25	25870.5	1.0	1.2	U			R11	1.5	78Di09	
^{83}Kr−^{86}Kr$_{.965}$	386.6	1.1	387.261	0.009	0.6	U			MS1	1.0	06Ri15	
^{83}Rb−^{85}Rb$_{.976}$	1207	8	1207.4	2.5	0.1	U			MA2	1.0	94Ot01	
	1207.4	2.5			0.0	1	100	100 ^{83}Rb	MA8	1.0	07Ke09	
^{82}Se H−^{83}Kr	10380	18	10398.1	0.5	0.7	U			R11	1.5	78Di09	
	10368	16			1.3	U			R11	1.5	78Di09	
^{82}Kr H−^{83}Kr	7160	18	7179.669	0.010	0.7	U			R11	1.5	78Di09	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{83}Kr−^{84}Kr$_{.988}$	1566.7601	0.0089	1566.760	0.009	0.0	1	100	100 ^{83}Kr	FS1	1.0	13Ho22	
^{83}Zn−^{83}Ga	13899	18				2			TR1	1.0	25Ja07	
^{83}Sr−^{83}Rb	2447	9	2440	7	−0.8	1	59	59 ^{83}Sr	MA2	1.0	94Ot01	
^{83}Y^{m}−^{83}Zr	−6681	13	−6685	20	−0.3	U			RKM	1.0	25Ki14	
^{83}Nb−^{83}Zr	8891.6	7.6	8875	3	−2.2	1	21	20 ^{83}Nb	RKM	1.0	25Ki14	
^{83}Kr−^{80}Se H$_2$	−18022	36	−18045.4	1.0	−0.4	U			R11	1.5	78Di09	
^{85}Rb−^{83}Kr H	−10211	45	−10161.812	0.010	0.7	U			R11	1.5	78Di09	
^{83}Kr−^{82}Se	−2572	35	−2573.0	0.5	−0.0	U			R11	1.5	78Di09	
^{83}Kr−^{82}Kr	648	12	645.363	0.010	−0.1	U			M15	2.5	63Ri07	
^{81}Rb−^{83}Rb$_{.488}$ ^{79}Rb$_{.513}$	−529	26	−548	5	−0.3	U			P20	2.5	82Au01	Y
^{81}Rb−^{83}Rb$_{.325}$ ^{80}Rb$_{.675}$	−1054	27	−1041	5	0.2	U			P20	2.5	82Au01	Y
^{82}Rb−^{83}Rb$_{.659}$ ^{80}Rb$_{.342}$	627	24	603	3	−0.4	U			P20	2.5	82Au01	Y
^{82}Rb−^{83}Rb$_{.494}$ ^{81}Rb$_{.506}$	1098	23	1054	4	−0.8	U			P21	2.5	82Au01	Y
^{82}Ge(d,p)^{83}Ge	1470	70	1409.1	2.6	−0.9	U			NDm		05Th03	
^{82}Se(d,p)^{83}Se	3593.4	3.0				2			NDm		78Mo12	
^{82}Se(^{3}He,d)^{83}Br	3207.4	5.6	3215	4	1.4	1	46	46 ^{83}Br	NDm		83Zu01	
^{82}Kr(^{3}He,d)^{83}Rb	288	10	274.3	2.3	−1.4	U			Phi		87St11	
^{83}Zr(εp)^{82}Sr	2750	100	2808	4	0.6	U					83Ha06	
^{83}As(β^-)^{83}Se	5460	220	5671	4	1.0	U			Stu		77Al17	
^{83}Se(β^-)^{83}Br	3610	40	3673	5	1.6	U					67Ma35	
	3681	20			−0.4	U					68Sc10	*
^{83}Br(β^-)^{83}Kr	982	10	977	4	−0.5	–					51Du03	*
	967	15			0.7	U					63Pa09	*
	966	6			1.8	–					69Ph03	*
ave.	970	5			1.3	1	54	54 ^{83}Br			average	
^{83}Rb(ε)^{83}Kr	750	20	920.0	2.3	8.5	B					70Go45	*
^{83}Sr(β^+)^{83}Rb	2264	10	2273	6	0.9	1	41	41 ^{83}Sr			68Et01	*
^{83}Y(β^+)^{83}Sr	4509	85	4592	20	1.0	U			BNL		81Li12	*
	4455	50			2.7	B			Ors		82De36	*
^{83}Y^{m}(IT)^{83}Y	62.04	0.10				3					87Ra06	*
^{83}Zr(β^+)^{83}Y	5868	85	6289	19	5.0	B			Ors		82De36	*
^{83}Nb(β^+)^{83}Zr	7500	300	8267	3	2.6	B					88Ku14	
*^{83}Y O−^{98}Mo$_{1.010}$	D_M=12973.8(5.9) μu for mixture gs+m at 62.04(0.10) keV; $M-A$=–72172.9(5.8) keV										Nub269	**
*^{83}Se(β^-)^{83}Br	Q_{β^-}=3910(20) from ^{83}Se^{m} at 228.92 keV										Nub269	**
*^{83}Br(β^-)^{83}Kr	E_{β^-}=940(10) 925(15) 924(6) respectively, to ^{83}Kr^{n} at 41.5575 keV										Nub269	**
*^{83}Rb(ε)^{83}Kr	LK=0.132(0.002) to 5/2$^-$ level at 561.9585, recalculated Q										Ens153	**
*^{83}Sr(β^+)^{83}Rb	E_{β^+}=1227(8) 24% to ground state, 20% to ^{83}Rb^{m} at 42.0780, and other E_{β^+}										Nub269	**
*^{83}Y(β^+)^{83}Sr	E_{β^+}=2868(85) from ^{83}Y^{m} at 62.04 keV to (3/2$^-$) level at 681.11 keV										Ens153	**
*^{83}Y(β^+)^{83}Sr	E_{β^+}=3353(50) to 9/2$^+$ level at 35.47 keV										Ens153	**
*	and E_{β^+}=2941(84) from ^{83}Y^{m} at 62.04 to (3/2$^-$) level at 681.11 keV										Ens153	**
*^{83}Y^{m}(IT)^{83}Y	From least-squares fit										Ens14c	**
*^{83}Zr(β^+)^{83}Y	Q_{β^+}=5806(85) to ^{83}Y^{m} at 62.04 keV										Nub269	**
*^{83}Zr(β^+)^{83}Y	Recalculated value 5802(50) of reference not accepted										87Ra06	**
^{84}Ga−u	−47337	32	−47331	8	0.2	2			TR1	1.0	20Re04	
	−47331.0	8.1			−0.0	2			RKM	1.0	24Xi06	
^{84}Ge−u	−62270	430	−62424.6	2.3	−0.4	U			OR1	1.0	06Ha62	
	−62424.4	3.2			−0.1	1	53	53 ^{84}Ge	RKM	1.0	24Xi06	
^{84}As−u	−70530	320	−70696.7	2.7	−0.5	U			OR1	1.0	06Ha62	*
	−70710	128			0.0	U			GT2	2.5	08Su19	
	−70696.6	4.4			−0.0	1	37	37 ^{84}As	RKM	1.0	24Xi06	
^{84}Se−u	−81538.4	5.0	−81534.0	1.9	0.9	1	15	15 ^{84}Se	RKM	1.0	24Xi06	
C$_6$ H$_{12}$−^{84}Kr	182399.4	2.5	182402.657	0.004	0.5	U			M15	2.5	63Ri07	
	182392	6			1.2	U			R11	1.5	78Di09	
C$_5$ N H$_{10}$−^{84}Kr	169819	18	169826.597	0.004	0.3	U			R11	1.5	78Di09	
	169819	13			0.4	U			R11	1.5	78Di09	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
C_5 O $H_8-{}^{84}$Kr	146010	20	146017.148	0.004	0.2	U			R11	1.5	78Di09	
C_4 ^{13}C O $H_7-{}^{84}$Kr	141543	18	141546.952	0.004	0.1	U			R11	1.5	78Di09	
^{84}Kr$-N_6$	−106946.3154	0.0152	−106946.299	0.004	1.1	o			FS1	1.0	05Sh38	*
	−106946.2971	0.0086			−0.2	1	22	21 ^{84}Kr	FS1	1.0	09Re03	
^{84}Kr H$-{}^{85}$Rb	7515	18	7533.023	0.004	0.7	U			R11	1.5	78Di09	
C_6 $H_{12}-{}^{84}$Sr	180470.8	2.6	180480.89	0.04	1.6	U			M15	2.5	63Ri07	
^{84}Y O$-{}^{97}$Mo$_{1.031}$	12483.5	5.1	12482	5	−0.2	1	82	82 ^{84}Y	JY1	1.0	08We10	
^{84}Zr O$-{}^{98}$Mo$_{1.020}$	14728.6	5.9				2			JY1	1.0	06Ka48	
^{84}Nb$-$u	−65721	13	−65694.3	0.4	2.1	U			LZ1	1.0	18Xi04	
^{84}Ge$-{}^{88}$Rb$_{.955}$	22268.7	3.4	22269.0	2.3	0.1	1	47	47 ^{84}Ge	JY1	1.0	08Ha23	
^{84}As$-{}^{88}$Rb$_{.955}$	13996.9	3.4	13996.9	2.7	0.0	1	63	63 ^{84}As	JY1	1.0	08Ha23	
^{84}Se$-{}^{88}$Rb$_{.955}$	3160.4	2.1	3159.6	1.9	−0.4	1	85	85 ^{84}Se	JY1	1.0	08Ha23	
^{82}Se $H_2-{}^{84}$Kr	20834	16	20851.9	0.5	0.7	U			R11	1.5	78Di09	
^{84}Kr$-{}^{86}$Kr$_{.977}$	−1168.3	1.0	−1168.8535	0.0022	−0.6	U			MS1	1.0	06Ri15	
^{84}Br$-{}^{85}$Rb$_{.988}$	3665.3	1.7				2			JY2	1.0	24Ja10	
^{83}Kr H$-{}^{84}$Kr	10465	16	10453.821	0.009	−0.5	U			R11	1.5	78Di09	
^{84}Kr$-{}^{85}$Rb$_{.988}$	−1349.4	1.5	−1350.534	0.004	−0.8	U			MA8	1.0	06De36	
^{84}Rb$-{}^{85}$Rb$_{.988}$	1536	8	1527.1	2.3	−1.1	U			MA2	1.0	94Ot01	
^{84}Sr$-{}^{85}$Rb$_{.988}$	570.9	1.5	571.23	0.04	0.2	U			MA8	1.0	07Ke09	
	572.1	4.3			−0.2	U			SH1	1.0	11Ha08	
^{84}Nb$-{}^{85}$Rb$_{.988}$	21457.45	0.43				2			JY1	1.0	19Vi05	
^{84}Br$^m-{}^{84}$Br	207.8	1.6				3			JY2	1.0	24Ja10	
^{84}Sr$-{}^{84}$Kr	1921.768	0.040	1921.77	0.04	−0.0	1	100	100 ^{84}Sr	JY2	1.0	24Ge05	
^{84}Zr$-{}^{84}$Nb	−10998	31	−10980	6	0.6	U			RKM	1.0	25Ki14	
^{84}Mo$-{}^{84}$Nb	7576	24				3			RKM	1.0	25Ki14	
^{84}Kr$-{}^{80}$Se H_3	−28505	48	−28499.3	1.0	0.1	U			R11	1.5	78Di09	
^{84}Kr$-{}^{83}$Kr	−2628	12	−2628.790	0.009	−0.0	U			M15	2.5	63Ri07	
	−2646	30			0.4	U			R11	1.5	78Di09	
^{84}Kr$-{}^{40}$Ar$_2$	−13268.5136	0.0171	−13268.509	0.006	0.3	1	11	6 ^{84}Kr	FS1	1.0	05Sh38	*
C_2 $O_4-{}^{84}$Kr	68160.7359	0.0205	68160.750	0.004	0.7	o			FS1	1.0	05Sh38	*
	68160.7516	0.0131			−0.1	1	10	9 ^{84}Kr	FS1	1.0	09Re03	
^{82}Se(t,p)^{84}Se	6016	15	6015.4	1.9	−0.0	U			LAl		74Kn02	
^{84}Sr(p,t)^{82}Sr	−12310	10	−12297	4	1.3	1	17	17 ^{82}Sr	Oak		73Ba56	
	−12295	24			−0.1	U			Win		74De31	*
^{83}Kr(n,γ)^{84}Kr	10519.5	1.8	10520.020	0.008	0.3	U					72Ma42	Z
	10520.6	0.3			−1.9	U			Bdn		06Fi.A	
^{84}Sr(d,t)^{83}Sr	−5720	30	−5666	7	1.8	U					70Be24	*
^{84}As(β^-)^{84}Se	7195	200	10095	3	14.5	F			Trs		94Gi07	*
	9120	880			1.1	U					96WaZX	
^{84}Se(β^-)^{84}Br	1818	50	1818.7	2.4	0.0	U					68Re12	*
	1808	100			0.1	U					70Ei02	*
^{84}Br(β^-)^{84}Kr	4650	30	4672.2	1.6	0.7	U					70Ha21	*
^{84}Br^m(β^-)^{84}Kr	4970	100	4865.8	2.2	−1.0	U					70Ha21	*
^{84}Rb(β^+)^{84}Kr	2679	3	2680.5	2.2	0.5	–					64La03	
	2682	5			−0.3	–					71Bo01	*
^{84}Rb(n,p)^{84}Kr	3450	10	3462.8	2.2	1.3	U			ILL		76An05	
^{84}Rb(β^+)^{84}Kr ave.	2679.8	2.6	2680.5	2.2	0.3	1	71	71 ^{84}Rb			average	
^{84}Rb(β^-)^{84}Sr	892	4	890.4	2.2	−0.4	1	29	29 ^{84}Rb			71Bo01	*
^{84}Y(β^+)^{84}Sr	6950	30	6755	4	−6.5	F					70Va.A	*
	6750	10			0.5	1	18	18 ^{84}Y			70Re.A	*
	6423	135			2.5	U			BNL		81Li12	*
	6408	124			2.8	U			Ors		82De36	*
^{84}Nb(β^+)^{84}Zr	7200	300	10228	6	10.1	B					96Sh27	
*^{84}As$-$u	Erroneously reported as –75700(300) keV in the publication										08Ha.A	**
*^{84}Kr$-N_6$	Corrected in reference of same group										09Re03	**
*^{84}Kr$-{}^{40}$Ar$_2$	Corrected in reference of same group										09Re03	**
*C_2 $O_4-{}^{84}$Kr	Corrected in reference of same group										09Re03	**
*^{84}Sr(p,t)^{82}Sr	Original error 12; added systematic error 21 keV										GAu	**
*^{84}Sr(d,t)^{83}Sr	Q =–5755(30) to 9/2$^+$ level at 35.47 keV										Ens153	**
*^{84}As(β^-)^{84}Se	F : observed (β^-n) decay implies $Q_{\beta^-} > 8681(15)$ keV										93Ru01	**
*^{84}Se(β^-)^{84}Br	E_{β^-}=1410(50) 1400(100) respectively, to 1$^+$ level at 408.2 keV										Ens09a	**
*^{84}Br(β^-)^{84}Kr	E_{β^-}=4626(15),3810(50),2700(50) to ground state, 881.615 2$^+$, 1897.784 0$^+$										Ens09a	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value	Adjusted value	v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{84}Br^m(\beta^-)^{84}Kr$	E_{β^-}=2200(100) to 5^- level at 2770.94 keV								Ens09a	**
$*^{84}Rb(\beta^+)^{84}Kr$	Original error increased: authors report E(2^+)=877.2(1.5) while								AHW	**
*	E(2^+)=881.615(0.003) keV; see also $^{74}As(\beta^+)$								Ens09a	**
$*^{84}Rb(\beta^-)^{84}Sr$	Originally 891.8(2.0), error increased, see $^{84}Rb(\beta^+)$								AHW	**
$*^{84}Y(\beta^+)^{84}Sr$	F : possibly additional with e^+e^-								AHW	**
$*^{84}Y(\beta^+)^{84}Sr$	E_{β^+}=1641(10) and 2242(17) to levels at 4062 and 3511 keV								70Re.A	**
$*^{84}Y(\beta^+)^{84}Sr$	Q_{β^+}=6409(170), and 6499(135) from $^{84}Y^m$ at 67 keV								00Do10	**
$*^{84}Y(\beta^+)^{84}Sr$	Q_{β^+}=6475(124) from $^{84}Y^m$ at 67 keV								00Do10	**
$^{85}Ga-u$	−42667	40	−42645	12	0.6	1	10	10 ^{85}Ga	TR1	1.0	20Re04	
$^{85}Ge-u$	−57220	540	−57028	3	0.4	U			OR1	1.0	06Ha62	
$^{85}As-u$	−68095	225	−67845.0	1.1	0.7	o			GT1	1.5	04Ma.A	
	−67887	129			0.1	U			GT2	2.5	08Su19	
$C_6\ H_{13}-^{85}Rb$	189927.6	3.9	189935.680	0.005	0.8	U			M15	2.5	63Ri07	
	189930	15			0.3	U			R11	1.5	78Di09	
$C_4\ N\ O\ H_7-^{85}Rb$	140985	18	140974.112	0.005	−0.4	U			R11	1.5	78Di09	
$^{85}Rb-^{39}K_{2.179}$	−9124.6	2.7	−9126.689	0.012	−0.8	U			MA8	1.0	09Na.A	
$^{85}Rb-^{120}Sn_{.708}$	−18970.8	2.2	−18969.6	0.7	0.6	U			JY1	1.0	11Ha48	
$^{85}Y-u$	−83559	31	−83567	20	−0.3	2			GS2	1.0	05Li24	*
$^{85}Zr\ O-^{98}Mo_{1.031}$	13886.7	6.9				2			JY1	1.0	06Ka48	
$^{85}Nb\ O-^{98}Mo_{1.031}$	21246	26	21289	4	1.7	U			JY1	1.0	06Ka48	*
$^{85}Ge-^{88}Rb_{.966}$	28638.8	4.0	28641	3	0.7	1	65	65 ^{85}Ge	JY1	1.0	08Ha23	
$^{85}As-^{88}Rb_{.966}$	17832.8	3.3	17824.1	1.1	−2.6	U			JY1	1.0	08Ha23	
$^{85}Se-^{88}Rb_{.966}$	7929.9	2.8	7928.3	1.6	−0.6	1	34	34 ^{85}Se	JY1	1.0	08Ha23	
$^{85}Br-^{88}Rb_{.966}$	1314.9	3.3	1313	3	−0.5	1	93	93 ^{85}Br	JY1	1.0	07Ra23	
$^{85}Ge-^{84}Kr\ H$	23654.5	5.5	23650	3	−0.9	1	35	35 ^{85}Ge	RKM	1.0	24Xi06	
$^{85}As-^{84}Kr\ H$	12832.2	1.1				2			RKM	1.0	24Xi06	
$^{85}Se-^{84}Kr\ H$	2935.6	2.0	2936.4	1.6	0.4	1	66	66 ^{85}Se	RKM	1.0	24Xi06	
$^{85}Br-^{84}Kr\ H$	−3702	12	−3679	3	1.9	1	7	7 ^{85}Br	RKM	1.0	24Xi06	
$^{85}Ga-^{85}Rb$	45568	13	45566	12	−0.2	1	90	90 ^{85}Ga	TR1	1.0	25Ja07	
$^{85}Nb-^{85}Rb$	17056.1	4.4				2			SH1	1.0	11Ha08	*
$^{85}Mo-^{85}Rb$	26471	17				2			SH1	1.0	11Ha08	
$C_6\ H_{14}-^{85}Rb$	197760.706	0.014	197760.712	0.005	0.4	U			MI2	1.0	99Br47	
$^{85}Rb-C_6\ H_{12}$	−182110.662	0.024	−182110.648	0.005	0.6	U			MI2	1.0	99Br47	
$^{85}Rb-^{84}Kr$	300	32	292.009	0.004	−0.2	U			R11	1.5	78Di09	
	292.0121	0.0064			−0.5	1	39	34 ^{85}Rb	FS1	1.0	10Mo30	
$^{83}Rb-^{85}Rb_{.488}\ ^{81}Rb_{.512}$	−351	22	−339	3	0.2	U			P21	2.5	82Au01	Y
$^{84}Kr(d,p)^{85}Kr$	4895	8	4887.7	2.0	−0.9	U			MIT		63Ho.A	
$^{85}Rb(\gamma,n)^{84}Rb$	−10650	80	−10479.8	2.2	2.1	U			Phi		60Ge01	
$^{85}Rb(p,d)^{84}Rb$	−8275	6	−8255.2	2.2	3.3	B			Bld		78Sh11	
$^{84}Sr(d,p)^{85}Sr$	6303	8	6300.8	2.8	−0.3	1	12	12 ^{85}Sr			71Mo02	
$^{85}Mo(\varepsilon p)^{84}Zr$	5100	200	6623	17	7.6	B					99Hu05	
$^{85}Se(\beta^-)^{85}Br$	6185	90	6162	3	−0.3	o			Bwg		87Gr.A	
	6182	23			−0.9	U			Bwg		92Gr.A	
$^{85}Br(\beta^-)^{85}Kr$	2870	19	2903	4	1.7	U			Stu		79Al05	
$^{85}Kr(\beta^-)^{85}Rb$	687	2				2					70Wo08	
$^{85}Sr(\varepsilon)^{85}Rb$	1007	30	1064.1	2.8	1.9	U					69Mc05	
$^{85}Rb(p,n)^{85}Sr$	−1890	30	−1846.4	2.8	1.5	U			BNL		58El44	
$^{85}Rb(^3He,t)^{85}Sr$	−1083	3	−1082.7	2.8	0.1	1	88	88 ^{85}Sr	Pri		82Ko06	
$^{85}Y(\beta^+)^{85}Sr$	3255	25	3261	19	0.2	R					63Do07	*
$^{85}Zr(\beta^+)^{85}Y$	4693	99	4667	20	−0.3	U			Ors		82De36	
$^{85}Nb(\beta^+)^{85}Zr$	6000	200	6896	8	4.5	F					88Ku14	*
$*^{85}Y-u$	$M-A$=–77824(28) keV for mixture gs+m at 19.68 keV								Nub269	**
$*^{85}Nb\ O-^{98}Mo_{1.031}$	D_M=21292.2(6.9) μu for mixture gs+m at 150#80 keV; $M-A$=–66274.8(6.6) keV								Nub269	**
$*^{85}Nb-^{85}Rb$	Misprint in publication 1.000200869 (not 1.00200869)								GAu	**
$*^{85}Y(\beta^+)^{85}Sr$	E_{β^+}=1540(20) to $3/2^-$ level at 743.25 keV								Ens148	**
*	and E_{β^+}=2240(10) from $^{85}Y^m$ at 19.68 (conflicting → outer error used)								Nub269	**
$*^{85}Nb(\beta^+)^{85}Zr$	F : see discussion of this result in reference								06Ka48	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{86}Ge−u		−54750	540	−53244.4	1.2	2.8	B			OR1	1.0	06Ha62	
		−53033	188			−0.4	U			GT3	2.5	16Kn03	
^{86}As−u		−63586	247	−63300.9	1.5	0.8	o			GT1	1.5	04Ma.A	
		−63189	129			−0.3	U			GT2	2.5	08Su19	
^{86}Se−u		−75702	128	−75687.9	0.8	0.0	U			GT2	2.5	08Su19	
C_6 H_{14}−^{86}Kr		198936.7	2.7	198939.823	0.004	0.5	U			M15	2.5	63Ri07	
		198933	15			0.3	U			R11	1.5	78Di09	
C_5 N H_{12}−^{86}Kr		186366	20	186363.763	0.004	−0.1	U			R11	1.5	78Di09	
^{86}Kr−u		−89389.271	0.110	−89389.376	0.004	−1.0	U			ST2	1.0	02Bf02	
^{86}Kr−^{120}Sn$_{.717}$		−19269.6	2.2	−19268.5	0.7	0.5	U			JY1	1.0	11Ha48	
C_6 H_{14}−^{86}Sr		200264.9	3.6	200289.723	0.006	2.8	U			M15	2.5	63Ri07	
^{86}Y−u		−85019	75	−85114	15	−1.3	U			GS2	1.0	05Li24	*
^{86}Zr O−^{98}Mo$_{1.041}$		9692.8	6.9	9686	4	−0.9	1	31	31 ^{86}Zr	JY1	1.0	06Ka48	
^{86}Nb O−^{98}Mo$_{1.041}$		19171.0	5.9				2			JY1	1.0	06Ka48	
^{86}As−^{88}Rb$_{.977}$		23346.2	3.7	23343.8	1.6	−0.6	1	18	17 ^{86}As	JY1	1.0	08Ha23	
^{86}Se−^{88}Rb$_{.977}$		10956.4	2.7	10956.8	0.8	0.1	1	9	9 ^{86}Se	JY1	1.0	08Ha23	
^{86}Br−^{88}Rb$_{.977}$		5450.1	3.3	5444.7	1.4	−1.6	1	19	19 ^{86}Br	JY1	1.0	07Ra23	
^{86}Kr−^{84}Kr$_{1.024}$		1236.9544	0.0042	1236.9566	0.0023	0.5	1	30	20 ^{84}Kr	FS1	1.0	12Ra34	
^{86}Sr−^{84}Kr$_{1.024}$		−112.9463	0.0054	−112.943	0.004	0.5	1	59	54 ^{86}Sr	FS1	1.0	12Ra34	
^{86}Kr−^{85}Rb$_{1.012}$		−120.3	3.6	−120.591	0.004	−0.1	U			MA8	1.0	06Ro11	
		−119.1	1.6			−0.9	U			MA8	1.0	06De36	
		−54	88			−0.8	U			MA9	1.0	10Na13	*
^{86}Rb−^{85}Rb$_{1.012}$		441	9	436.23	0.21	−0.5	U			MA2	1.0	94Ot01	
^{86}Sr ^{19}F−^{85}Rb$_{1.235}$		16608	13	16603.564	0.006	−0.3	U			MA8	1.0	05Si34	
^{86}Zr−^{85}Rb$_{1.012}$		5562.7	4.6	5566	4	0.6	1	69	69 ^{86}Zr	SH1	1.0	11Ha08	
^{86}Mo−^{85}Rb$_{1.012}$		20443.6	4.0	20443	3	−0.2	2			SH1	1.0	11Ha08	
		20441.7	5.1			0.2	2			JY1	1.0	19Vi05	
^{86}Ga−^{86}Rb		52580	20				3			TR1	1.0	25Ja07	
^{86}Ge−^{86}Kr		36145.0	1.2				2			RKM	1.0	24Xi06	
^{86}As−^{86}Kr		26088.0	1.7	26088.5	1.5	0.3	1	83	83 ^{86}As	RKM	1.0	24Xi06	
^{86}Se−^{86}Kr		13701.49	0.83	13701.5	0.8	−0.0	1	91	91 ^{86}Se	RKM	1.0	24Xi06	
^{86}Br−^{86}Kr		8188.1	1.6	8189.4	1.4	0.8	1	81	81 ^{86}Br	RKM	1.0	24Xi06	
^{86}Sr−^{86}Kr		−1349.8965	0.0060	−1349.900	0.004	−0.6	1	49	46 ^{86}Sr	FS1	1.0	12Ra34	
^{86}Kr−^{85}Rb		−1206	42	−1179.111	0.004	0.4	U			R11	1.5	78Di09	
		−1179.1083	0.0071			−0.4	–			FS1	1.0	10Mo30	*
		−1179.1109	0.0059			−0.1	–			FS1	1.0	10Mo30	*
	ave.	−1179.110	0.005			−0.3	1	69	66 ^{85}Rb			average	
^{86}Kr−N_6		−107833.3986	0.0074	−107833.401	0.004	−0.4	1	28	27 ^{86}Kr	FS1	1.0	09Re03	
C_2 O_4−^{86}Kr		69047.8337	0.0155	69047.853	0.004	1.2	o			FS1	1.0	05Sh38	*
		69047.8440	0.0113			0.8	1	12	11 ^{86}Kr	FS1	1.0	09Re03	
^{86}Kr−^{84}Kr		−908	32	−887.1023	0.0023	0.4	U			R11	1.5	78Di09	
		−887.1041	0.0125			0.1	o			FS1	1.0	05Sh38	*
		−887.1080	0.0069			0.8	–			FS1	1.0	09Re03	*
		−887.0954	0.0060			−1.2	–			FS1	1.0	10Mo30	*
	ave.	−887.101	0.005			−0.3	1	25	15 ^{84}Kr			average	
^{86}Sr(p,t)^{84}Sr		−11535	10	−11534.71	0.04	0.0	U			Oak		73Ba56	
^{85}Rb(n,γ)^{86}Rb		8651.1	1.0	8650.98	0.20	−0.1	U					69Da15	Z
		8651.3	1.5			−0.2	U					70Or.A	
		8650.98	0.20				2			Bdn		06Fi.A	
^{85}Rb(d,p)^{86}Rb		6433	10	6426.41	0.20	−0.7	U			Tal		69Da15	
^{86}Se(β^-)^{86}Br		5095	100	5134.5	1.5	0.4	o			Bwg		87Gr.A	
		5099	11			3.2	C			Bwg		92Gr.A	
^{86}Br(β^-)^{86}Kr		7620	60	7628.4	1.3	0.1	U			Stu		79Al05	
		7626	11			0.2	U			Bwg		92Gr.A	
^{86}Rb(β^-)^{86}Sr		1774	5	1776.10	0.20	0.4	U					64Da16	
		1770	3			2.0	U					66An10	
		1779.2	2.5			−1.2	U					75Be21	
		1775	3			0.4	U					75Ra09	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{86}Y(β^+)^{86}Sr	5220	20	5240	14	1.0	2					62Ya01	*
	5260	20			−1.0	2					65Va02	*
^{86}Nb(β^+)^{86}Zr	7978	80	8835	7	10.7	F					82De43	*
^{86}Mo(β^+)^{86}Nb	5019	430	5023	6	0.0	U					94Sh07	*
*^{86}Y−u	$M-A$=–79086(29) keV for mixture gs+m at 218.21 keV										Nub269	**
*^{86}Kr−^{85}Rb$_{1.012}$	Typo in original paper, ratio should read 1.011 763 90(1 03)										GAu	**
*^{86}Kr−^{85}Rb	Different charge states : 3^+ and 2^+										10Mo30	**
*C$_2$ O$_4$−^{86}Kr	Corrected in reference of same group										09Re03	**
*^{86}Kr−^{84}Kr	Corrected in reference of same group										09Re03	**
*^{86}Kr−^{84}Kr	Different charge states in these two results: 3^+ and 2^+										10Mo30	**
*^{86}Y(β^+)^{86}Sr	E_{β^+}=2019(20) 1960(20) respectively, to 2229.81 4^+ level, and other E_{β^+}										Ens14c	**
*^{86}Nb(β^+)^{86}Zr	F : see discussion of this result in reference										06Ka48	**
*^{86}Mo(β^+)^{86}Nb	E_{β^+}=3900(400) to (1^+) level at 97.1 keV										94Sh07	**
^{87}Se−u	−71357	128	−71310.1	1.9	0.1	U			GT2	2.5	08Su19	
^{87}Kr−u	−86622	30	−86645.24	0.26	−0.8	U			GS2	1.0	05Li24	
C$_4$ H$_7$ O$_2$−^{87}Rb	135417.8	2.7	135423.933	0.006	0.9	U			M15	2.5	63Ri07	
C$_5$ ^{13}C H$_{14}$−^{87}Rb	203767	15	203724.754	0.006	−1.9	U			R11	1.5	78Di09	
C$_4$ O N H$_9$−^{87}Rb	159277	15	159233.382	0.006	−1.9	U			R11	1.5	78Di09	
C$_3$ ^{13}C O N H$_8$−^{87}Rb	154809	25	154763.185	0.006	−1.2	U			R11	1.5	78Di09	
C$_4$ H$_7$ O$_2$−^{87}Sr	135722.2	3.5	135726.968	0.005	0.5	U			M15	2.5	63Ri07	
^{87}Y−u	−89153	30	−89123.9	1.2	1.0	U			GS2	1.0	03Li.A	*
^{87}Zr−u	−85222	30	−85183	4	1.3	U			GS2	1.0	05Li24	
^{87}Zr O−^{97}Mo$_{1.062}$	9543.3	5.2	9542	4	−0.2	1	73	73 ^{87}Zr	JY1	1.0	08We10	
^{87}Nb$_{1.069}$−C$_7$ H$_9$	−155224	30	−155205	8	0.6	U			CP1	1.0	11Fa10	
^{87}Nb O−^{98}Mo$_{1.051}$	15027.9	7.3				2			JY1	1.0	06Ka48	*
^{87}Mo$_{1.069}$−C$_7$ H$_9$	−147186.1	4.8	−147182.6	2.4	0.7	–			CP1	1.0	11Fa10	
^{87}Mo−u	−71801.9	3.5	−71802.9	2.2	−0.3	–			LZ1	1.0	23Xi01	
^{87}Mo$_{1.069}$−C$_7$ H$_9$	ave. −147183.2	2.8	−147182.6	2.4	0.2	1	72	72 ^{87}Mo			average	
^{87}Mo^m−u	−71469	32				2			LZ1	1.0	23Xi01	
^{87}Sr−^{84}Kr$_{1.036}$	565.8516	0.0059	565.851	0.004	−0.2	1	47	41 ^{87}Sr	FS1	1.0	12Ra34	
^{87}Kr−^{85}Rb$_{1.024}$	3683.0	2.9	3682.07	0.26	−0.3	U			MA8	1.0	06De36	
	3684.1	4.7			−0.4	U			MA8	1.0	10Na13	
^{87}Rb−^{85}Rb$_{1.024}$	−490	9	−492.156	0.007	−0.2	U			MA2	1.0	94Ot01	
	−493.0	2.7			0.3	U			MA8	1.0	06De36	
	−492.33	0.80			0.2	U			MA8	1.0	07Ke09	
	−492.4	1.4			0.2	U			MA8	1.0	09Na.A	
	−492.04	0.87			−0.1	U			MA8	1.0	11He10	
	−492.166	0.056			0.2	U			JY2	1.0	18Ne09	*
^{87}Sr−^{85}Rb$_{1.024}$	−780	9	−795.191	0.005	−1.7	U			MA2	1.0	94Ot01	
^{87}Mo−^{85}Rb$_{1.024}$	18525.6	4.2	18524.4	2.2	−0.3	1	28	28 ^{87}Mo	SH1	1.0	11Ha08	
^{87}Tc−^{85}Rb$_{1.024}$	28394.5	4.5				2			SH1	1.0	11Ha08	*
^{87}As−^{86}Kr H	21856.1	2.8	21856.1	2.1	−0.0	1	57	57 ^{87}As	RKM	1.0	24Xi06	
^{87}As−^{88}Rb$_{.989}$	28000.6	3.2	28000.6	2.1	0.0	1	43	43 ^{87}As	JY1	1.0	08Ha23	
^{87}Se−^{86}Kr H	10256.3	3.1	10254.2	1.9	−0.7	1	38	38 ^{87}Se	RKM	1.0	24Xi06	
^{87}Se−^{88}Rb$_{.989}$	16397.5	2.4	16398.8	1.9	0.5	1	63	62 ^{87}Se	JY1	1.0	08Ha23	
^{87}Br−^{88}Rb$_{.989}$	8382.9	3.4				2			JY1	1.0	07Ra23	
^{86}Kr H−^{87}Rb	9309	16	9255.127	0.006	−2.2	U			R11	1.5	78Di09	
^{87}Sr−^{86}Kr$_{1.012}$	−660.4616	0.0050	−660.461	0.004	0.2	1	62	59 ^{87}Sr	FS1	1.0	12Ra34	
C$_6$ H$_{16}$−^{87}Rb	216019.966	0.023	216019.982	0.007	0.7	U			MI2	1.0	99Br47	
^{87}Rb−C$_6$ H$_{14}$	−200369.931	0.015	−200369.918	0.007	0.8	1	19	19 ^{87}Rb	MI2	1.0	99Br47	
^{87}Rb−^{86}Kr	−1477	30	−1430.095	0.006	1.0	U			R11	1.5	78Di09	
	−1430.0932	0.0059			−0.4	1	87	81 ^{87}Rb	FS1	1.0	10Mo30	
^{87}Sr−^{86}Sr	−382	12	−383.230	0.006	−0.0	U			M15	2.5	63Ri07	
^{87}Rb−^{85}Rb	−2620	35	−2609.207	0.007	0.2	U			R11	1.5	78Di09	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{85}Rb−^{87}Rb$_{.489}$ ^{83}Rb$_{.512}$	−310	30	−314.8	1.2	−0.1	U			P21	2.5	82Au01	
^{84}Rb−^{87}Rb$_{.241}$ ^{83}Rb$_{.759}$	850	72	643.8	2.8	−1.1	U			P21	2.5	82Au01	*
^{87}Sr(p,t)^{85}Sr	−11440	10	−11437.7	2.8	0.2	U			Oak		73Ba56	
^{87}Br(β^-n)^{86}Kr	1335	25	1303	3	−1.3	U					84Kr.B	
^{86}Kr(n,γ)^{87}Kr	5515.04	0.6	5515.17	0.25	0.2	2					77Je03	Z
	5515.20	0.27			−0.1	2			Bdn		06Fi.A	
^{86}Kr(d,p)^{87}Kr	3286	8	3290.61	0.25	0.6	U			MIT		63Ho.A	
^{87}Rb(γ,n)^{86}Rb	−9990	70	−9922.12	0.20	1.0	U			Phi		60Ge01	
^{87}Rb(d,t)^{86}Rb	−3659	15	−3664.89	0.20	−0.4	U			Tal		69Da15	
^{86}Sr(n,γ)^{87}Sr	8428.12	0.17	8428.295	0.005	1.0	U			ILn		86Wi16	Z
	8428.17	0.17			0.7	U			Bdn		06Fi.A	
^{86}Sr(d,p)^{87}Sr	6203	8	6203.728	0.005	0.1	U					71Mo02	
^{86}Sr(p,γ)^{87}Y	5785.4	3.3	5784.3	1.1	−0.3	R					71Um03	
^{86}Sr(^{3}He,d)^{87}Y	346	15	290.8	1.1	−3.7	B			ANL		71Ma11	
^{87}Mo(εp)^{86}Zr	3700	300	3796	4	0.3	U					83Ha06	
^{87}Se(β^-)^{87}Br	7250	150	7467	4	1.4	o			Bwg		87Gr.A	
	7275	35			5.5	B			Bwg		92Gr.A	
^{87}Br(β^-)^{87}Kr	6830	120	6818	3	−0.1	U			Stu		79Al05	
	6750	150			0.5	o			Bwg		87Gr.B	
	6855	25			−1.5	U			Bwg		92Gr.A	
^{87}Kr(β^-)^{87}Rb	3888	7	3888.27	0.25	0.0	U					73Wo01	
^{87}Rb(β^-)^{87}Sr	272	3	282.275	0.006	3.4	B					59Fl40	
	274	3			2.8	B					61Be41	
^{87}Rb(^{3}He,t)^{87}Sr−^{81}Br()^{81}Kr	564.0	1.5	563.1	0.5	−0.6	1	10	9 ^{81}Kr	Pri		82Ko06	
^{87}Y(β^+)^{87}Sr	2190	50	1861.7	1.1	−6.6	B					67Mi13	*
	1791	40			1.8	U					69Zo04	*
^{87}Sr(p,n)^{87}Y	−2644.2	1.2	−2644.0	1.1	0.1	2					71Um03	Z
^{87}Zr(β^+)^{87}Y	3663	40	3671	4	0.2	U					65Ba48	*
^{87}Nb(β^+)^{87}Zr	5165	60	5473	8	5.1	B					82De43	*
^{87}Mo(β^+)^{87}Nb	6382	308	6991	7	2.0	U					82De43	*
	6589	300			1.3	U					91Mi15	*
*^{87}Y−u	$M-A$=–82665(28) keV for ^{87}Y^m at 380.82 keV										Nub269	**
*^{87}Nb O−^{98}Mo$_{1.051}$	D_M=15030.0(6.9) μu for mixture gs+m at 3.9(0.1) keV; $M-A$=–73870.2(6.7) keV										Nub269	**
*^{87}Rb−^{85}Rb$_{1.024}$	from ^{85}Rb$^+$/^{87}Rb$^+$ =1.023523278677(654), alt.in reference 1.023523278629(662)										18Ne09	**
*^{87}Tc−^{85}Rb$_{1.024}$	Most probably the high-spin isomer										11Ha08	**
*^{84}Rb−^{87}Rb$_{.241}$ ^{83}Rb$_{.759}$	D_M=1080(40) keV corrected –230(60) for mixture gs+m at 463.59 keV										Nub269	**
*^{87}Y(β^+)^{87}Sr	E_{β^+}=780(50) to ^{87}Sr^m at 388.53 keV										Nub269	**
*^{87}Y(β^+)^{87}Sr	E_{β^+}=1150(40) from ^{87}Y^m at 380.82 keV										Nub269	**
*^{87}Zr(β^+)^{87}Y	E_{β^+}=2260(40) to ^{87}Y^m at 380.82 keV										Nub269	**
*^{87}Nb(β^+)^{87}Zr	Q_{β^+}=5169(60) from ^{87}Nb^m at 3.9(0.1) keV										Nub269	**
*^{87}Mo(β^+)^{87}Nb	Q_{β^+}=6378(308) to ^{87}Nb^m at 3.9(0.1) keV										Nub269	**
*^{87}Mo(β^+)^{87}Nb	E_{β^+}=5300(300) to (7/2)$^+$ level at 266.9 keV										Ens15a	**
^{88}Se−u	−68555	129	−68581.3	2.5	−0.1	U			GT2	2.5	08Su19	
^{88}Br−u	−75832	100	−75917.8	2.5	−0.3	o			GT2	2.5	08Kn.A	
	−75823	129			−0.3	U			GT2	2.5	08Su19	
C_4 H_8 O_2−^{88}Sr	146789.1	4.7	146817.241	0.006	2.4	U			M15	2.5	63Ri07	
^{88}Y−u	−90500	31	−90498.7	1.6	0.0	U			GS2	1.0	05Li24	
^{88}Zr O−^{98}Mo$_{1.061}$	5502.3	6.9	5502	6	−0.0	1	71	71 ^{88}Zr	JY1	1.0	06Ka48	
^{88}Nb O−^{98}Mo$_{1.061}$	13458	76	13510	60	0.7	1	67	67 ^{88}Nb	JY1	1.0	06Ka48	*
^{88}Sr−^{84}Kr$_{1.048}$	−1637.3606	0.0069	−1637.366	0.005	−0.8	1	46	42 ^{88}Sr	FS1	1.0	12Ra34	
^{88}Kr−^{85}Rb$_{1.035}$	5745.5	2.8				2			MA8	1.0	06De36	
^{88}Rb−^{85}Rb$_{1.035}$	2615	9	2613.19	0.17	−0.2	U			MA4	1.0	02Ra23	
^{88}Sr−^{85}Rb$_{1.035}$	−3108	20	−3090.126	0.006	0.9	U			MA8	1.0	07Ke09	
	−3088	11			−0.2	U			MA8	1.0	05Si34	
^{88}Mo−^{85}Rb$_{1.035}$	13265.4	4.1				2			JY1	1.0	08We10	
^{88}Tc−^{85}Rb$_{1.035}$	25045	23	25092	4	2.0	U			JY1	1.0	08We10	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{88}Sr−^{86}Kr$_{1.023}$	−2942.4188	0.0059	−2942.415	0.005	0.7	1	61	58 ^{88}Sr	FS1	1.0	12Ra34	
^{88}Tc−^{87}Rb$_{1.011}$	25612.7	4.4				2			JY1	1.0	19Vi05	
^{88}As−^{88}Br	21508.9	2.4				2			RKM	1.0	24Xi06	
^{88}Se−^{88}Br	7336.66	0.94	7336.6	0.9	−0.1	1	97	51 ^{88}Se	RKM	1.0	24Xi06	
^{88}Se−^{88}Rb	20101.9	3.6	20103.2	2.5	0.3	1	49	49 ^{88}Se	JY1	1.0	08Ha23	
^{88}Br−^{88}Rb	12767.7	3.4	12766.6	2.5	−0.3	1	54	54 ^{88}Br	JY1	1.0	07Ra23	
^{88}Tc^{m}−^{88}Tc	75.6	3.5				3			JY1	1.0	19Vi05	
^{88}Sr−^{87}Sr	−3260	12	−3265.241	0.006	−0.2	U			M15	2.5	63Ri07	
^{88}Ru−^{12}C ^{32}S	−30311	20				2			RKM	1.0	25Ki14	
^{86}Kr(t,p)^{88}Kr	4091	15	4086.5	2.6	−0.3	U			LAl		76Fl02	
^{88}Sr(p,t)^{86}Sr	−11060	10	−11059.369	0.006	0.1	U			Oak		73Ba56	
^{87}Rb(n,γ)^{88}Rb	6082.52	0.16	6082.54	0.16	0.1	1	96	96 ^{88}Rb	Bdn		06Fi.A	
^{87}Rb(d,p)^{88}Rb	3858	15	3857.97	0.16	−0.0	U			Oak		71Ra17	
	3837	20			1.0	U					71To05	
^{87}Sr(n,γ)^{88}Sr	11112.63	0.22	11112.870	0.006	1.1	U			ILn		87Wi15	Z
	11112.64	0.22			1.0	U			Bdn		06Fi.A	
^{87}Sr(d,p)^{88}Sr	8865	5	8888.304	0.005	4.7	B			MIT		68Co20	
^{88}Se(β^-)^{88}Br	6854	31	6834.0	0.9	−0.6	U			Bwg		92Gr.A	
^{88}Br(β^-)^{88}Kr	8970	130	8974	4	0.0	U			Stu		79Al05	
	8880	200			0.5	o			Bwg		87Gr.B	
	8960	36			0.4	U			Bwg		92Gr.A	
^{88}Kr(β^-)^{88}Rb	2930	30	2917.7	2.6	−0.4	U			Trs		78Wo15	
^{88}Rb(β^-)^{88}Sr	5310	60	5312.61	0.16	0.0	U			Trs		78Wo15	
	5318	9			−0.6	U			Gsn		80De02	*
	5313	5			−0.1	U			Trs		82Br23	
^{88}Y(β^+)^{88}Sr	3622.6	1.5				2					79An36	*
^{88}Sr(p,n)^{88}Y	−4402	6	−4404.9	1.5	−0.5	U			Bar		61Sh23	
	−4406	10			0.1	U			Tal		62Ne08	
^{88}Nb(β^+)^{88}Zr	7550	100	7460	60	−0.9	1	34	33 ^{88}Nb			84Ox01	
^{88}Nb^{m}(β^+)^{88}Zr	7590	100				2					84Ox01	
^{88}Tc(β^+)^{88}Mo	8600	1300	11016	6	1.9	U					96Od01	
	7800	600			5.4	B					96Sh27	
*^{88}Nb O−^{98}Mo$_{1.061}$	D_M=13527.5(6.9) μu for mixture gs+m at 130(120) keV; $M-A$=–76150.9(6.7) keV										Nub269	**
*^{88}Tc−^{85}Rb$_{1.035}$	Most probably the low-spin isomer										08We10	**
*^{88}Tc−^{85}Rb$_{1.035}$	D_M=25082.4(4.1) μu for mixture gs+m at 70(3) keV; $M-A$=–61679.1(3.8) keV										Nub269	**
*^{88}Rb(β^-)^{88}Sr	Original error 4 corrected in reference										94Ha.A	**
*^{88}Y(β^+)^{88}Sr	E_{β^+}=764.6(1.5) to 2^+ level at 1836.090 keV										Ens141	**
^{89}Se−u	−63285	225	−63329	3	−0.1	o			GT1	1.5	04Ma.A	
	−63291	129			−0.1	U			GT2	2.5	08Su19	
C_7 H_5−^{89}Y	133247.0	3.4	133287.0	0.4	4.7	B			M15	2.5	63Ri07	
^{89}Zr−u	−91126	38	−91119.9	3.0	0.2	U			TR1	1.0	25Cz.1	
^{89}Nb−u	−86588	34	−86529	8	1.7	U			GS2	1.0	05Li24	*
^{89}Kr−^{85}Rb$_{1.047}$	10191.6	2.3				2			MA8	1.0	06De36	
^{89}Rb−^{85}Rb$_{1.047}$	4628	9	4634	3	0.6	1	12	12 ^{89}Rb	MA4	1.0	02Ra23	
^{89}Y−^{85}Rb$_{1.047}$	−1806.32	0.52	−1805.7	0.4	1.2	1	49	49 ^{89}Y	MS1	1.0	19Sa39	
^{89}Mo−^{85}Rb$_{1.047}$	11824.3	4.2	11822	3	−0.5	1	61	61 ^{89}Mo	JY1	1.0	08We10	
^{89}Tc−^{85}Rb$_{1.047}$	20007	17	20007	3	−0.0	U			SH1	1.0	08We10	
	20004.8	4.1			0.5	1	62	62 ^{89}Tc	JY1	1.0	08We10	
^{89}Ru−^{85}Rb$_{1.047}$	29694	26				2			JY1	1.0	19Vi05	
^{89}Y−^{87}Rb$_{1.023}$	−1252.96	0.51	−1253.5	0.4	−1.1	1	51	51 ^{89}Y	MS1	1.0	19Sa39	
^{89}Se−^{88}Rb$_{1.011}$	26329.0	4.0	26331	3	0.4	1	62	62 ^{89}Se	JY1	1.0	08Ha23	
^{89}Br−^{88}Rb$_{1.011}$	16364.5	3.5	16363.5	2.9	−0.3	1	70	70 ^{89}Br	JY1	1.0	07Ra23	
^{89}As−^{89}Rb	37601.3	3.5				2			RKM	1.0	24Xi06	
^{89}Se−^{89}Rb	24394.4	3.4	24393.2	2.9	−0.4	1	73	38 ^{89}Se	RKM	1.0	24Xi06	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{89}Br−^{89}Rb		14424.8	3.6	14425.9	3.0	0.3	1	68	38 ^{89}Rb	RKM	1.0	24Xi06	
^{89}Kr−^{89}Rb		5557	11	5558	4	0.1	U			RKM	1.0	24Xi06	
^{89}Nb−^{89}Tc		−14179.4	7.3				2			RI1	1.0	26Ho.A	
^{89}Mo−^{89}Tc		−8185.6	3.2	−8184.4	2.8	0.4	1	77	39 ^{89}Mo	RI1	1.0	26Ho.A	
^{88}Rb−^{89}Rb$_{.494}$ ^{87}Rb$_{.506}$		545	23	563.6	1.4	0.3	U			P21	2.5	82Au01	
^{89}Y(d,α)^{87}Sr		7889	15	7879.7	0.3	−0.6	U			Mtr		72Br13	
^{88}Sr(n,γ)^{89}Sr		6358.70	0.13	6358.72	0.09	0.1	–			ILn		89Wi05	Z
		6358.73	0.13			−0.1	–			Bdn		06Fi.A	
^{88}Sr(d,p)^{89}Sr		4133	5	4134.15	0.09	0.2	U			MIT		67Sp09	
^{88}Sr(n,γ)^{89}Sr	ave.	6358.72	0.09	6358.72	0.09	0.0	1	100	100 ^{89}Sr			average	
^{88}Sr(p,γ)^{89}Y		7078	4	7078.5	0.3	0.1	U					75Be.B	Z
^{89}Y(γ,n)^{88}Y		−11540	40	−11483.5	1.5	1.4	U			Phi		63Ge02	
^{89}Br(β^-)^{89}Kr		8140	140	8261	3	0.9	U			Stu		81Ho17	
		8120	120			1.2	o			Bwg		87Gr.B	
		8155	30			3.5	C			Bwg		92Gr.A	
^{89}Kr(β^-)^{89}Rb		5150	30	5177	4	0.9	U					67Ki01	
		5191	60			−0.2	U			Trs		78Wo15	*
		5140	120			0.3	U			Stu		81Ho17	*
^{89}Rb(β^-)^{89}Sr		4486	12	4496.1	2.9	0.8	–					66Ki06	
		4491	15			0.3	o			Gsn		80Bl.A	
		4510	9			−1.5	–			Gsn		80De02	*
	ave.	4501	7			−0.7	1	16	16 ^{89}Rb			average	
^{89}Sr(β^-)^{89}Y		1463	5	1502.2	0.4	7.8	B					49La06	
		1488	4			3.5	B					70Wo05	
^{89}Zr(β^+)^{89}Y		2841	10	2833.5	2.8	−0.7	U					51Hy24	*
		2832	10			0.2	U					53Sh48	*
		2828	7			0.8	–					60Ha26	*
^{89}Y(p,n)^{89}Zr		−3612.8	4.	−3615.9	2.8	−0.8	–			Tkm		63Ok01	Z
		−3619.4	6.			0.6	–			Oak		64Jo11	Z
^{89}Zr(β^+)^{89}Y	ave.	2832	3	2833.5	2.8	0.6	1	85	85 ^{89}Zr			average	
^{89}Nb(β^+)^{89}Zr		3870	100	4277	8	4.1	B					55Ma13	
		4340	50			−1.3	U					74Vo08	
^{89}Mo(β^+)^{89}Nb		5970	300	5584	7	−1.3	U					64Bu12	
^{89}Tc(β^+)^{89}Mo		7510	210	7623.7	2.6	0.5	U					91He04	*
*^{89}Nb−u	$M-A$=–80656(28) keV for mixture gs+m at 0#30 keV											Nub269	**
*^{89}Kr(β^-)^{89}Rb	E_{β^-}=4970(60) to 220.948 level											GAu	**
*^{89}Rb(β^-)^{89}Rb	Splitting Table 3a into 3 groups yields 4610(120) 4867(152) 5135(123) keV											GAu	**
*^{89}Rb(β^-)^{89}Sr	Original error 8 corrected in reference											94Ha.A	**
*^{89}Zr(β^+)^{89}Y	E_{β^+}=910(10) 901(10) 897(7) respectively, to ^{89}Y^m at 908.97 keV											Nub269	**
*^{89}Tc(β^+)^{89}Mo	E_{β^+}=6370(210) to 118.8 level; no Fermi−Kurie plot											91He04	**
^{90}Se−u		−59904	236	−59992	5	−0.2	U			GT1	1.5	04Ma.A	
^{90}Rb−u		−85420	210	−85194	4	1.1	U			TG2	1.0	20Gr.1	
^{90}Y−u		−92859	43	−92858.2	0.4	0.0	U			RT1	1.0	25Cz.1	
C_4 H_{10} O_2−^{90}Zr		163377	6	163380.63	0.10	0.2	U			M15	2.5	63Ri07	
^{90}Zr−u		−95301.36	0.23	−95301.07	0.10	1.3	1	20	20 ^{90}Zr	MS1	1.0	15Gu09	
^{90}Nb−u		−88872	50	−88742	3	2.6	U			GS2	1.0	05Li24	*
^{90}Mo−C_7 H_6		−133018.9	4.7	−133019	3	−0.0	1	44	44 ^{90}Mo	CP1	1.0	11Fa10	
^{90}Mo$_{1.033}$−C_7 H_9		−159318	23	−159334	3	−0.7	U			CP1	1.0	11Fa10	
^{90}Tc−C_7 H_6		−122880.3	6.7	−122876.0	1.1	0.6	U			CP1	1.0	11Fa10	
^{90}Tc$_{1.033}$−C_7 H_9		−148835	22	−148856.7	1.1	−1.0	U			CP1	1.0	11Fa10	
		−148854.2	8.5			−0.3	U			CP1	1.0	11Fa10	
^{90}Ru$_{1.033}$−C_7 H_9		−142382	11	−142382.2	2.1	−0.0	U			CP1	1.0	11Fa10	
^{90}Kr−^{85}Rb$_{1.059}$		12942.6	2.0	12941.7	1.8	−0.4	1	84	84 ^{90}Kr	MA8	1.0	06De36	
^{90}Rb−^{85}Rb$_{1.059}$		8211	9	8221	4	1.1	1	21	21 ^{90}Rb	MA4	1.0	02Ra23	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{90}Tc−^{85}Rb$_{1.059}$		17489.2	8.0	17488.9	1.1	−0.0	U				SH1	1.0	08We10	
		17489.8	4.2			−0.2	U				JY1	1.0	08We10	
^{90}Ru−^{85}Rb$_{1.059}$		23775	11	23756.5	2.1	−1.7	–				SH1	1.0	08We10	
		23756.6	4.7			−0.0	–				JY1	1.0	08We10	
	ave.	23759	4			−0.7	1	23	23	^{90}Ru			average	
^{90}Tc−^{86}Kr$_{1.047}$		17664.6	1.1	17664.9	1.1	0.2	1	92	92	^{90}Tc	JY1	1.0	12Ka12	
^{90}Tc^{m}−^{86}Kr$_{1.047}$		17819.2	1.4				2				JY1	1.0	12Ka12	
^{90}Zr−^{87}Rb$_{1.034}$		−1393.79	0.16	−1393.73	0.10	0.4	1	41	41	^{90}Zr	MS1	1.0	15Gu09	
^{90}Br−^{88}Rb$_{1.023}$		22017.0	3.6	22017.4	2.9	0.1	1	63	63	^{90}Br	JY1	1.0	07Ra23	
^{90}Se−^{90}Kr		20481.3	4.2				2				RKM	1.0	24Xi06	
^{90}Br−^{90}Kr		11766.8	4.3	11766.2	3.0	−0.1	1	48	37	^{90}Br	RKM	1.0	24Xi06	
^{90}Rb−^{90}Kr		−4716.6	4.7	−4721	4	−0.9	1	70	65	^{90}Rb	RKM	1.0	24Xi06	*
^{90}Nb−^{90}Tc		−12833	15	−12816	3	1.2	1	5	5	^{90}Nb	RI1	1.0	26Ho.A	
^{90}Mo−^{90}Tc		−10142.4	5.8	−10143	3	−0.1	1	30	28	^{90}Mo	RI1	1.0	26Ho.A	
^{90}Ru−^{90}Tc		6266.9	2.1	6267.6	1.9	0.3	1	82	77	^{90}Ru	RI1	1.0	26Ho.A	
^{89}Rb−^{90}Rb$^{x}_{791}$ ^{85}Rb$_{.209}$		−1826	24	−1824	10	0.0	U				P21	2.5	82Au01	
^{90}Zr(α,^{8}He)^{86}Zr		−40136	30	−39988	4	4.9	B				INS		90Ka01	
^{90}Zr(^{3}He,^{6}He)^{87}Zr		−12083	8	−12086	4	−0.4	1	27	27	^{87}Zr	MSU		78Pa11	
^{90}Zr(p,t)^{88}Zr		−12805	10	−12804	5	0.1	1	29	29	^{88}Zr	Oak		71Ba43	
^{89}Y(n,γ)^{90}Y		6857.1	1.0	6857.03	0.10	−0.1	U				ORn		81Ra07	
		6857.26	0.30			−0.8	–						83De17	
		6856.98	0.17			0.3	–				ILn		93Mi04	Z
		6857.01	0.14			0.1	–				Bdn		06Fi.A	
^{89}Y(d,p)^{90}Y		4635	5	4632.46	0.10	−0.5	U						64Wa14	
^{89}Y(n,γ)^{90}Y	ave.	6857.03	0.10	6857.03	0.10	−0.0	1	100	100	^{90}Y			average	
^{89}Y(p,γ)^{90}Zr		8351	4	8350.2	0.4	−0.2	U						75Be.B	
^{90}Zr(γ,n)^{89}Zr		−11940	50	−11966.0	2.8	−0.5	U				Phi		63Ge02	
^{90}Zr(p,d)^{89}Zr		−9728	10	−9741.5	2.8	−1.3	U				Oak		71Ba43	
^{90}Zr(d,t)^{89}Zr		−5719.2	7.1	−5708.8	2.8	1.5	1	15	15	^{89}Zr	SPa		79Bo37	
^{90}Zr(^{3}He,α)^{89}Zr		8580	50	8611.6	2.8	0.6	U				Phi		67Fo04	
^{90}Br(β^-)^{90}Kr		9800	400	10960.1	2.8	2.9	U				Stu		81Ho17	
		10280	110			6.2	C				Bwg		87Gr.B	
		10350	75			8.1	C				Bwg		92Gr.A	
^{90}Kr(β^-)^{90}Rb		4410	30	4397	4	−0.4	U						70Ma11	
		4390	40			0.2	U				Trs		78Wo15	
		4380	25			0.7	U				Bwg		87Gr.A	
^{90}Rb^{x}(IT)^{90}Rb		71	12				2						82Au01	
^{90}Rb(β^-)^{90}Sr		6550	60	6593	4	0.7	U				Trs		78Wo15	
		6560	150			0.2	U				Bwg		78St02	
		6585	15			0.6	o				Gsn		80Bl.A	
		6578	15			1.0	o				Gsn		80De02	
		6587	10			0.6	1	16	14	^{90}Rb	Gsn		92Pr03	
^{90}Sr(β^-)^{90}Y		546	2	546.1	1.4	0.1	–						64Da16	
		546	2			0.1	–						83Ha35	
	ave.	546.0	1.4			0.1	1	98	98	^{90}Sr			average	
^{90}Y(β^-)^{90}Zr		2271	2	2275.5	0.4	2.2	B						61Ni02	
		2284	5			−1.7	U						64Da16	
		2273	5			0.5	U						64La13	
		2280	5			−0.9	U						66Ri01	
		2278	8			−0.3	U				Gsn		80Bl.A	
		2279.5	2.9			−1.4	U						83Ha35	
^{90}Nb(β^+)^{90}Zr		6111	4	6110	3	−0.2	1	60	60	^{90}Nb			68Pe01	*
^{90}Mo(β^+)^{90}Nb		2489	4	2490	3	0.1	1	62	35	^{90}Nb			66Pe10	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{90}Tc(β^+)^{90}Mo	8920	410	9448.2	3.0	1.3	U					74Ia01	*
	9270	300			0.6	U					81Ox01	*
	8726	300			2.4	U					81Ox01	*
^{90}Rh(ε)^{90}Ru	13410	1330	13540#	200#	0.1	D					19Pa16	*
*^{90}Nb−u	$M-A$=–82721(29) keV for mixture gs+n at 124.67 keV										Nub269	**
*^{90}Rb−^{85}Rb$_{1.059}$	D_M=8326(9) μu for ^{90}Rb^m at 106.90 keV; $M-A$=–79260(9) keV										Nub269	**
*^{90}Rb−^{90}Kr	could be the gs.										HWJ25a	**
*^{90}Nb(β^+)^{90}Zr	E_{β^+}=1500(4) to 6$^+$ level at 3589.419 keV										Ens981	**
*^{90}Mo(β^+)^{90}Nb	E_{β^+}=1085(4) to 1$^+$ level at 382.01 keV										Ens981	**
*^{90}Tc(β^+)^{90}Mo	E_{β^+}=7900(400) from ^{90}Tc^m at 144.1 to ground state (85%) and 947.97 (15%) level										74Ia01	**
*^{90}Tc(β^+)^{90}Mo	E_{β^+}=5300(300) to 2946.82 level										Ens981	**
*^{90}Tc(β^+)^{90}Mo	E_{β^+}=6900(300) from ^{90}Tc^m at 144.0(1.7) to 2$^+$ level a 947.97 keV										Ens981	**
*^{90}Rh(ε)^{90}Ru	Symmetrized from 13190(+1500–1160)										HWJ199	**
*^{90}Rh(ε)^{90}Ru	Trends in Mass Surface TMS suggest ^{90}Rh 130 keV less bound										HWJ268	**
^{91}Se−u		−54300	186	−53964.1	2.5	0.7	U			GT3	2.5	16Kn03	
^{91}Se−^{79}Br ^{12}C		27698.2	2.3				2			RKM	1.0	24Xi06	
^{91}Br−^{79}Br ^{12}C		16061.3	2.1	16061.2	1.9	−0.0	1	78	72 ^{91}Br	RKM	1.0	24Xi06	
^{91}Kr−^{79}Br ^{12}C		5471.7	5.9	5469.1	2.4	−0.4	1	16	14 ^{91}Kr	RKM	1.0	24Xi06	
^{91}Rb−^{79}Br ^{12}C		−1817.3	2.0				2			RKM	1.0	24Xi06	
^{91}Rb−u		−83532	21	−83479.6	2.2	1.0	U			Pb1	2.5	89Al33	
		−83330	170			−0.9	U			TG2	1.0	20Gr.1	
^{91}Y−u		−92702	35	−92702.2	2.0	−0.0	U			RT1	1.0	25Cz.1	
C_7 H_7−^{91}Zr		149143.1	4.4	149135.13	0.09	−0.7	U			M15	2.5	63Ri07	
^{91}Zr−u		−94359.85	0.25	−94359.91	0.09	−0.2	1	13	13 ^{91}Zr	MS1	1.0	15Gu09	
^{91}Nb−u		−93064	46	−93010	3	1.2	U			GS2	1.0	05Li24	*
^{91}Mo−C_7 H_7		−143031.3	8.3	−143038	6	−0.9	1	46	46 ^{91}Mo	CP1	1.0	11Fa10	
^{91}Tc−C_7 H_7		−136340.9	6.7	−136351.1	2.1	−1.5	–			CP1	1.0	11Fa10	
		−136353.6	4.6			0.6	–			CP1	1.0	11Fa10	
	ave.	−136350	4			−0.4	1	31	31 ^{91}Tc			average	
^{91}Ru$_{1.011}$−C_7 H_8		−136730	620	−136663.2	2.0	0.1	U			CP1	1.0	11Fa10	
^{91}Ru−C_7 H_7		−128035.6	3.9	−128032.3	2.0	0.8	–			CP1	1.0	11Fa10	
^{91}Ru$_{1.022}$−C_7 H_9		−145260	23	−145294.0	2.0	−1.5	U			CP1	1.0	11Fa10	
^{91}Ru−u		−73259.7	6.3	−73257.1	2.0	0.4	–			LZ1	1.0	23Xi01	
^{91}Ru−C_7 H_7	ave.	−128035	3	−128032.3	2.0	0.9	1	35	35 ^{91}Ru			average	
^{91}Ru^m−u		−72796	32				2			LZ1	1.0	23Xi01	
^{91}Kr−^{85}Rb$_{1.071}$		18279.5	2.4	18279.9	2.2	0.2	1	86	86 ^{91}Kr	MA8	1.0	06De36	
^{91}Rb−^{85}Rb$_{1.071}$		11003	10	10993.6	2.2	−0.9	U			MA4	1.0	02Ra23	
^{91}Sr−^{85}Rb$_{1.071}$		4702	9	4676.2	1.1	−2.9	B			MA4	1.0	02Ra23	
		4676.1	1.1			0.1	1	100	100 ^{91}Sr	JY2	1.0	24Ja03	
^{91}Tc−^{85}Rb$_{1.071}$		12898.3	5.4	12897.4	2.1	−0.2	1	15	15 ^{91}Tc	SH1	1.0	08We10	
^{91}Ru−^{85}Rb$_{1.071}$		21223	11	21216.1	2.0	−0.6	–			SH1	1.0	08We10	
		21215.5	4.2			0.1	–			JY1	1.0	08We10	
	ave.	21216	4			−0.1	1	25	25 ^{91}Ru			average	
^{91}Zr−^{87}Rb$_{1.046}$		637.32	0.19	637.26	0.09	−0.3	1	22	22 ^{91}Zr	MS1	1.0	15Gu09	
^{91}Br−^{88}Rb$_{1.034}$		26098.3	3.8	26098.6	2.0	0.1	1	28	28 ^{91}Br	JY1	1.0	07Ra23	
^{91}Tc−^{94}Mo$_{.968}$		10303.0	4.4	10303.3	2.1	0.1	1	23	23 ^{91}Tc	JY1	1.0	08We10	
^{91}Ru−^{94}Mo$_{.968}$		18620.9	4.7	18622.0	2.0	0.2	1	17	17 ^{91}Ru	JY1	1.0	08We10	
^{91}Mo−^{91}Tc		−6706	10	−6687	6	1.9	1	33	30 ^{91}Mo	RI1	1.0	26Ho.A	
^{91}Ru−^{91}Tc		8322.1	3.3	8318.7	2.3	−1.0	1	51	28 ^{91}Tc	RI1	1.0	26Ho.A	
^{91}Rh−^{91}Tc		18820.2	7.5				2			RI1	1.0	26Ho.A	
^{91}Zr−^{90}Zr		942	12	941.17	0.10	−0.0	U			M15	2.5	63Ri07	
^{90}Rb^x−^{91}Rb$_{.824}$ ^{85}Rb$_{.176}$		−686	24	−750	13	−1.1	U			P21	2.5	82Au01	
^{91}Zr(p,t)^{89}Zr		−10677	10	−10678.9	2.8	−0.2	U			Oak		71Ba43	
^{90}Zr(n,γ)^{91}Zr		7194.4	0.5	7194.63	0.09	0.5	U					81Lo.A	Z
		7192.7	0.8			2.4	U			Bdn		06Fi.A	
		7194.75	0.12			−1.0	1	63	39 ^{90}Zr	JY2		25Ru03	*
^{90}Zr(d,p)^{91}Zr		4959	20	4970.06	0.09	0.6	U			Pit		64Co11	
		4969	8			0.1	U			MIT		72Gr12	
		4970.3	2.2			−0.1	U			SPa		79Bo37	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{91}Zr(p,d)^{90}Zr		−4977	10	−4970.06	0.09	0.7	U			Oak		71Ba43	
^{91}Zr(d,t)^{90}Zr		−932	20	−937.40	0.09	−0.3	U			Pit		64Co11	
		−940.3	3.7			0.8	U			SPa		79Bo37	
^{90}Zr(p,γ)^{91}Nb		5167	5	5154.9	2.9	−2.4	B					71Ra08	
		5167	4			−3.0	C					75Be.B	Z
^{90}Zr(^{3}He,d)^{91}Nb		−227	20	−338.6	2.9	−5.6	B			Hei		70Kn05	
^{90}Zr(α,t)^{91}Nb		−14643	27	−14659.0	2.9	−0.6	U			Brk		71Zi03	
^{91}Ru^m(εp)^{90}Mo		4300	500	5075	30	1.5	U					83Ha06	
^{91}Br(β^-)^{91}Kr		9790	100	9866.5	2.8	0.8	U			Bwg		89Gr03	
		9805	50			1.2	U			Bwg		92Gr.A	
^{91}Kr(β^-)^{91}Rb		6420	80	6787.2	2.9	4.6	B			Trs		78Wo15	
		6450	80			4.2	B			Bwg		89Gr03	
^{91}Rb(β^-)^{91}Sr		5830	45	5884.6	2.3	1.2	U			Trs		78Wo15	*
		5927	24			−1.8	o			Gsn		80De02	*
		5920	28			−1.3	U			McG		83Ia02	*
		5930	22			−2.1	U			Gsn		92Pr03	*
^{91}Sr(β^-)^{91}Y		2669	10	2706.2	2.1	3.7	B					53Am08	*
		2684	10			2.2	U					73Ha11	*
		2704	8			0.3	U			Gsn		80De02	*
		2709	15			−0.2	U			McG		83Ia02	
^{91}Y(β^-)^{91}Zr		1545	5	1544.1	1.9	−0.2	2					64La13	
		1544	2			0.1	2					75Ra08	
^{91}Zr(p,n)^{91}Nb		−2045	6	−2039.7	2.9	0.9	–			Oak		70Ki01	
		−2038.8	3.4			−0.3	–			Kyu		71Ma47	
	ave.	−2040.3	3.0			0.2	1	98	98 ^{91}Nb			average	
^{91}Mo(β^+)^{91}Nb		4460	30	4422	6	−1.3	–					56Sm96	
		4435	23			−0.6	–					93Os06	
	ave.	4444	18			−1.2	1	10	8 ^{91}Mo			average	
^{91}Tc(β^+)^{91}Mo		6220	200	6229	5	0.0	U					74Ia01	
^{91}Pd(ε)^{91}Rh		11800	2200	12240#	300#	0.2	D					18Pa20	*
*^{91}Nb−u	$M-A$=–86636(30) keV for mixture gs+m at 104.60 keV											Nub269	**
*^{90}Zr(n,γ)^{91}Zr	constructed from frequency ratio $^{90}ZrO^+/(^{91}ZrO)^+$ =0.9906366606(12)											HWJ25a	**
*^{91}Rb(β^-)^{91}Sr	E_{β^-}=5760(40) to ^{91}Sr ground state <8% and 93.628 keV $(3/2)^+$ level 25%											Ens13a	**
*^{91}Rb(β^-)^{91}Sr	Original error 8 corrected to 13 keV in reference											94Ha.A	**
*^{91}Rb(β^-)^{91}Sr	E_{β^-}=5857 to mixture ^{91}Sr ground state <8% and 93.628 $(3/2)^+$ level 25%											Ens13a	**
*^{91}Rb(β^-)^{91}Sr	E_{β^-}=5850(20) and E_{β^-}=5860(10) respectively, to ^{91}Sr ground state <8% and 93.628 level 25%											Ens13a	**
*^{91}Sr(β^-)^{91}Y	E_{β^-}=2665(10), 2030(20), 1359(10), 1093(10) to											53Am08	**
*	ground state, $3/2^-$ level at 653.02 keV, $(5/2)^+$ at 1305.39, $(5/2)^-$ at 1545.90											Ens13a	**
*^{91}Sr(β^-)^{91}Y	Original error 4 increased: in disagreement with other results											AHW	**
*^{91}Sr(β^-)^{91}Y	Original error 3 corrected in reference											94Ha.A	**
*^{91}Pd(ε)^{91}Rh	Trends in Mass Surface TMS suggest ^{91}Pd 440 keV less bound											GAu	**
^{92}Br−^{78}Se ^{12}C H_2		6663.9	6.6	6668	5	0.6	1	54	54 ^{92}Br	RKM	1.0	24Xi06	
^{92}Br−u		−60711	103	−60373	5	1.3	U			GT2	2.5	08Kn.A	
^{92}Kr−^{78}Se ^{12}C H_2		−6789.2	9.0	−6786.5	2.8	0.3	1	9	9 ^{92}Kr	RKM	1.0	24Xi06	
^{92}Rb−u		−80323	32	−80270	5	0.7	U			Pb1	2.5	89Al33	
C_7 H_8−^{92}Zr		157569.4	3.8	157565.02	0.09	−0.5	U			M15	2.5	63Ri07	
^{92}Zr−u		−94964.66	0.18	−94964.76	0.09	−0.6	1	25	25 ^{92}Zr	MS1	1.0	15Gu09	
^{92}Nb−u		−92851	56	−92811.5	1.9	0.7	U			GS2	1.0	05Li24	*
C_7 H_8−^{92}Mo		155790.0	3.2	155793.10	0.17	0.4	U			M15	2.5	63Ri07	
$^{92}Mo_{1.011}$−C_7 H_9		−164641.3	7.0	−164643.26	0.17	−0.3	U			CP1	1.0	11Fa10	
^{92}Mo−u		−93193.14	0.47	−93192.85	0.17	0.6	1	13	13 ^{92}Mo	MS1	1.0	15Gu09	
^{92}Tc−C_7 H_8		−147328	13	−147332.3	2.8	−0.3	U			CP1	1.0	11Fa10	
$^{92}Tc_{.989}$−C_7 H_7		−138569	10	−138575.2	2.8	−0.6	–			CP1	1.0	11Fa10	
$^{92}Tc_{1.011}$−C_7 H_9		−156087.6	6.1	−156089.4	2.9	−0.3	–			CP1	1.0	11Fa10	
$^{92}Tc_{.989}$−C_7 H_7	ave.	−138572	5	−138575.2	2.8	−0.6	1	29	29 ^{92}Tc			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{92}Ru−C$_7$ H$_8$		−142352	18	−142366.3	2.3	−0.8	o			CP1	1.0	08Fa11	
		−142352	18			−0.8	U			CP1	1.0	11Fa10	
		−142377	10			1.1	o			CP1	1.0	08Fa11	
		−142378	10			1.2	U			CP1	1.0	11Fa10	
^{92}Ru$_{1.011}$−C$_7$ H$_9$		−151074.5	5.6	−151068.8	2.3	1.0	o			CP1	1.0	08Fa11	
		−151074.5	5.6			1.0	1	17	17 ^{92}Ru	CP1	1.0	11Fa10	
^{92}Rh$_{1.011}$−C$_7$ H$_9$		−138825	23	−138795.8	2.8	1.3	U			CP1	1.0	11Fa10	
		−138818	17			1.3	U			CP1	1.0	11Fa10	
^{92}Rh−u		−67622	11	−67626.6	2.7	−0.4	1	6	6 ^{92}Rh	GR1	1.0	25Kr04	
^{92}Kr−^{85}Rb$_{1.082}$		21616.6	2.9	21616.3	2.8	−0.1	1	91	91 ^{92}Kr	MA8	1.0	06De36	
^{92}Rb−^{85}Rb$_{1.082}$		15176	9	15173	5	−0.3	1	33	33 ^{92}Rb	MA4	1.0	02Ra23	
^{92}Sr−^{85}Rb$_{1.082}$		6482	9	6482	4	−0.0	–			MA4	1.0	02Ra23	
		6484.0	4.3			−0.5	–			MA8	1.0	05Gu37	
	ave.	6484	4			−0.5	1	84	84 ^{92}Sr			average	
^{92}Mo−^{85}Rb$_{1.082}$		2251.43	0.85	2250.66	0.17	−0.9	U			JY1	1.0	12Ka13	
^{92}Tc−^{85}Rb$_{1.082}$		10728	12	10711.4	2.8	−1.4	U			SH1	1.0	08We10	
		10712.5	4.3			−0.2	1	43	43 ^{92}Tc	JY1	1.0	08We10	
^{92}Ru−^{85}Rb$_{1.082}$		15684.3	5.7	15677.4	2.3	−1.2	–			SH1	1.0	08We10	
		15677.9	4.3			−0.1	–			JY1	1.0	08We10	
	ave.	15680	3			−0.8	1	45	45 ^{92}Ru			average	
^{92}Rh−^{85}Rb$_{1.082}$		27841	37	27816.9	2.7	−0.7	U			SH1	1.0	08We10	
		27811.2	4.7			1.2	1	34	34 ^{92}Rh	JY1	1.0	08We10	
^{92}Zr−^{87}Rb$_{1.057}$		1031.51	0.21	1031.42	0.09	−0.4	1	19	19 ^{92}Zr	MS1	1.0	15Gu09	
^{92}Mo−^{87}Rb$_{1.057}$		2803.38	0.18	2803.34	0.17	−0.2	1	87	87 ^{92}Mo	MS1	1.0	15Gu09	
^{92}Br−^{88}Rb$_{1.045}$		32306.8	7.2	32302	5	−0.6	1	46	46 ^{92}Br	JY1	1.0	07Ra23	
^{92}Zr ^{35}Cl−^{90}Zr ^{37}Cl		3285	2	3286.44	0.13	0.2	U			H13	4.0	63Ba20	
^{92}Mo−^{92}Ru		−13430	11	−13426.8	2.3	0.3	U			RI1	1.0	26Ho.A	
^{92}Tc−^{92}Ru		−4969.7	4.7	−4966	3	0.8	1	44	28 ^{92}Tc	RI1	1.0	26Ho.A	
^{92}Rh−^{92}Ru		12140.7	2.4	12139.4	2.2	−0.5	1	82	60 ^{92}Rh	RI1	1.0	26Ho.A	
^{92}Pd−^{92}Ru		21102	27				2			RI1	1.0	26Ho.A	
^{92}Zr−^{91}Zr		−603	12	−604.86	0.03	−0.1	U			M15	2.5	63Ri07	
^{88}Rb−^{92}Rb$_{.410}$ ^{85}Rb$_{.592}$		−3258	22	−3309.6	2.0	−0.9	U			P21	2.5	82Au01	
^{89}Rb−^{92}Rb$_{.553}$ ^{85}Rb$_{.449}$		−3457	24	−3471	4	−0.2	U			P21	2.5	82Au01	
^{91}Rb−^{92}Rb$_{.848}$ ^{85}Rb$_{.153}$		−1703	25	−1783	5	−1.3	U			P21	2.5	82Au01	
^{90}Rb^x−^{92}Rb$_{.699}$ ^{85}Rb$_{.303}$		−2059	24	−2124	13	−1.1	U			P21	2.5	82Au01	
^{90}Rb^x−^{92}Rb$_{.326}$ ^{89}Rb$_{.674}$		209	24	164	13	−0.8	U			P21	2.5	82Au01	
^{92}Mo(α,^{8}He)^{88}Mo		−43278	20	−43307	4	−1.4	U			INS		90Ka01	
^{92}Mo(p,α)^{89}Nb		−1306	50	−1343	7	−0.7	U			ANL		75Se.A	
^{92}Mo(^{3}He,^{6}He)^{89}Mo		−14465	15	−14453	3	0.8	U			MSU		80Pa02	
^{92}Zr(p,t)^{90}Zr		−7372	14	−7347.57	0.10	1.7	U			Bld		66St15	
		−7350	10			0.2	U			Oak		71Ba43	
^{92}Mo(p,t)^{90}Mo		−14330	30	−14296.7	2.9	1.1	U			VUn		76Ka08	
^{92}Rb(β^-n)^{91}Sr		785	15	803	5	1.2	1	11	10 ^{92}Rb			84Kr.B	
^{91}Zr(n,γ)^{92}Zr		8634.91	0.20	8634.74	0.03	−0.8	U			ILn		79Br25	Z
		8634.64	0.15			0.7	U					81Su.A	Z
		8635.00	0.24			−1.1	U			Bdn		06Fi.A	
		8634.75	0.04			−0.2	–			ILn		18Rz01	
		8634.74	0.05			0.0	–			JY2		25Ru03	*
^{91}Zr(d,p)^{92}Zr		6470	30	6410.17	0.03	−2.0	U					62Ma06	
		6395	20			0.8	U			Pit		64Co11	
		6410.9	4.3			−0.2	U			SPa		79Bo37	
^{92}Zr(p,d)^{91}Zr		−6410	11	−6410.17	0.03	−0.0	U			Bld		66St15	
		−6410	10			−0.0	U			Oak		71Ba43	
^{92}Zr(d,t)^{91}Zr		−2363	25	−2377.51	0.03	−0.6	U			Pit		64Co11	
^{91}Zr(n,γ)^{92}Zr	ave.	8634.75	0.03	8634.74	0.03	−0.2	1	97	56 ^{92}Zr			average	
^{92}Mo(p,d)^{91}Mo		−10446	15	−10439	5	0.5	–			Tex		73Ko03	
		−10432	25			−0.3	–			Grn		73Mo03	
	ave.	−10442	13			0.3	1	16	16 ^{91}Mo			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{92}Br(β^-)^{92}Kr	12155	100	12533	5	3.8	B			Bwg		89Gr03	
	12220	55			5.7	C			Bwg		92Gr.A	
^{92}Kr(β^-)^{92}Rb	6160	80	6002	5	−2.0	U			Trs		78Wo15	
	6045	80			−0.5	o			Bwg		89Gr03	
	5987	10			1.5	o			Bwg		92Gr.A	
	5993	27			0.3	U			Bwg		92Gr06	
^{92}Rb(β^-)^{92}Sr	8080	160	8096	4	0.1	U			Trs		78Wo15	
	8091	15			0.3	o			Gsn		80Bl.A	
	8111	15			−1.0	o			Gsn		80De02	
	8080	30			0.5	U			McG		83Ia02	
	8095	25			0.0	o			Bwg		87Gr.A	
	8096	16			−0.0	–			Bwg		92Gr.A	
	8107	15			−0.8	–			Gsn		92Pr03	
	8095	6			0.1	–					25Al22	
ave.	8097	5			−0.2	1	70	57 ^{92}Rb			average	
^{92}Sr(β^-)^{92}Y	1929	50	1949	9	0.4	U					57He39	*
	1930	30			0.6	–			Trs		78Wo15	*
	1920	20			1.5	–			McG		83Ia02	
ave.	1923	17			1.6	1	32	29 ^{92}Y			average	
^{92}Y(β^-)^{92}Zr	3640	20	3643	9	0.1	–					62Bu16	
	3630	15			0.8	–			McG		83Ia02	
ave.	3634	12			0.8	1	58	58 ^{92}Y			average	
^{92}Nb(β^+)^{92}Zr	2005	6	2005.7	1.8	0.1	U					59We30	*
	2008	6			−0.4	U					62Bu16	*
^{92}Zr(p,n)^{92}Nb	−2790.7	2.3	−2788.1	1.8	1.1	–			Kyu		74Ku01	
	−2792	5			0.8	–					75Ke12	
ave.	−2790.9	2.1			1.4	1	73	73 ^{92}Nb			average	
^{92}Tc(β^+)^{92}Mo	7880	100	7881.2	2.6	0.0	U					64Va05	
^{92}Mo(p,n)^{92}Tc	−8672	50	−8663.5	2.6	0.2	U			Tal		66Mo06	*
^{92}Mo(^{3}He,t)^{92}Tc	−7882	30	−7899.8	2.6	−0.6	U			ChR		73Ha02	
^{92}Pd(ε)^{92}Rh	8220	345	8349	25	0.4	U					19Pa16	*
*^{92}Nb−u	$M-A$=–86422(34) keV for mixture gs+m at 135.5 keV										Nub269	**
*^{91}Zr(n,γ)^{92}Zr	constructed from frequency ratio (^{92}ZrO+)/(^{91}ZrO+)=1.00934887819(52)										25Ru03	**
*^{92}Sr(β^-)^{92}Y	E_{β^-}=545(50) 546(30) respectively, to 1^+ level at 1383.91 keV										Ens12a	**
*^{92}Nb(β^+)^{92}Zr	p^+ =56(6)×10^{-5}, 60(6)×10^{-5} respectively, to 2^+ level at 934.51 keV										Ens12a	**
*	recalculated Q_{β^+}=2140(6) 2143(6) respectively, from ^{92}Nb^m at 135.5 keV										AHW	**
*^{92}Mo(p,n)^{92}Tc	T=9040(50) to (4^+) level at 270.09 keV										Ens12a	**
*^{92}Pd(ε)^{92}Rh	Q_ε=8170(330) keV decays to the daughter excited state at 50#(100#) keV										HWJ202	**
^{93}Br−u	−56866	322	−56780	460	0.2	o			GT1	1.5	04Ma.A	
	−56780	185				2			GT3	2.5	16Kn03	
^{93}Rb−u	−78036	21	−77960	9	1.4	U			Pb1	2.5	89Al33	
	−77868	100			−0.4	U			GT2	2.5	08Kn.A	
	−77940	130			−0.2	U			TG2	1.0	20Gr.1	
^{93}Sr−u	−87010	420	−85976	8	2.5	U			TG2	1.0	20Gr.1	
C_7 H_9−^{93}Nb	164046.9	3.5	164052.2	1.6	0.6	U			M15	2.5	63Ri07	
^{93}Mo−u	−93194	30	−93191.23	0.19	0.1	U			GS2	1.0	05Li24	*
^{93}Tc−u	−89729	31	−89754.9	1.1	−0.8	U			GS2	1.0	05Li24	
^{93}Tc−C_7 H_9	−160189.5	7.7	−160180.1	1.1	1.2	U			CP1	1.0	11Fa10	
	−160170	22			−0.5	U			CP1	1.0	11Fa10	
	−160189.4	8.5			1.1	U			CP1	1.0	11Fa10	
^{93}Tc$_{.989}$−C_7 H_8	−151270	190	−151367.8	1.1	−0.5	U			CP1	1.0	11Fa10	
^{93}Ru−C_7 H_9	−153318.2	6.4	−153319.7	2.1	−0.2	–			CP1	1.0	11Fa10	
	−153307	23			−0.6	U			CP1	1.0	11Fa10	
	−153324.0	4.8			0.9	–			CP1	1.0	11Fa10	
	−153321.9	3.5			0.6	–			CP1	1.0	11Fa10	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{93}Ru−u		−82853	47	−82894.4	2.1	−0.9	U			GR1	1.0	19An10	
^{93}Ru−C_7 H_9	ave.	−153321.9	2.6	−153319.7	2.1	0.9	1	68	68 ^{93}Ru			average	
^{93}Rh−C_7 H_9		−144485	25	−144514.4	2.7	−1.2	o			CP1	1.0	08Fa11	
		−144485	26			−1.1	U			CP1	1.0	11Fa10	
		−144527.7	5.3			2.5	U			CP1	1.0	08Fa11	
		−144527.7	5.2			2.6	U			CP1	1.0	11Fa10	
		−144512.9	3.8			−0.4	1	49	49 ^{93}Rh	CP1	1.0	11Fa10	
^{93}Rh−u		−74081	21	−74089.1	2.7	−0.4	U			LZ1	1.0	23Xi01	
^{93}Rh^{m}−u		−73794	47				2			LZ1	1.0	23Xi01	
^{93}Pd−u		−63475	38	−63431	7	1.2	U			GR1	1.0	25Kr04	
^{93}Kr−^{85}Rb$_{1.094}$		27649.2	2.7				2			MA8	1.0	06De36	
^{93}Rb−^{85}Rb$_{1.094}$		18549	10	18542	9	−0.7	1	72	72 ^{93}Rb	MA4	1.0	02Ra23	
^{93}Sr−^{85}Rb$_{1.094}$		10526	10	10526	8	0.0	1	66	66 ^{93}Sr	MA4	1.0	02Ra23	
^{93}Ru−^{85}Rb$_{1.094}$		13609.4	4.3	13607.6	2.1	−0.4	1	25	25 ^{93}Ru	JY1	1.0	08We10	
^{93}Rh−^{85}Rb$_{1.094}$		22428	12	22412.9	2.7	−1.3	–			SH1	1.0	08We10	
		22413.5	4.5			−0.1	–			JY1	1.0	08We10	
	ave.	22415	4			−0.6	1	40	40 ^{93}Rh			average	
^{93}Tc−^{93}Rh		−15686	16	−15665.7	2.9	1.3	U			RI1	1.0	26Ho.A	
^{93}Ru−^{93}Rh		−8792.3	7.4	−8805	3	−1.8	1	19	12 ^{93}Rh	RI1	1.0	26Ho.A	
^{93}Pd−^{93}Rh		10658.6	7.0				2			RI1	1.0	26Ho.A	
^{91}Rb−^{93}Rb$_{.489}$ ^{89}Rb$_{.511}$		−471	9	−495	5	−1.0	U			P31	2.5	86Au02	
^{91}Rb−^{93}Rb$_{.326}$ ^{90}Rb$^{x}_{.674}$		−656	23	−648	9	0.1	U			P21	2.5	82Au01	
^{92}Rb−^{93}Rb$_{.495}$ ^{91}Rb$_{.505}$		465	23	444	6	−0.4	U			P21	2.5	82Au01	
^{93}Rb(β^-n)^{92}Sr		2220	30	2176	9	−1.5	U					84Kr.B	
^{92}Sr(n,γ)^{93}Sr		5230	6	5290	8	10.0	B					80Kr07	
^{92}Zr(n,γ)^{93}Zr		6733.7	1.1	6734.3	0.4	0.6	–					72Gr23	Z
		6734.0	0.7			0.5	–					79Ke.D	Z
		6735.3	0.7			−1.4	–			Bdn		06Fi.A	
^{92}Zr(d,p)^{93}Zr		4493	20	4509.8	0.4	0.8	U			Pit		64Co11	
^{92}Zr(n,γ)^{93}Zr	ave.	6734.5	0.5	6734.3	0.4	−0.4	1	98	98 ^{93}Zr			average	
^{93}Nb(γ,n)^{92}Nb		−8780	60	−8830.9	2.0	−0.8	U			Phi		60Ge01	
		−8825	3			−2.0	1	44	27 ^{92}Nb	McM		79Ba06	
^{93}Nb(d,t)^{92}Nb		−2581	20	−2573.6	2.0	0.4	U			Pit		64Co11	
		−2571	10			−0.3	U			Tal		64Sh04	
^{92}Mo(n,γ)^{93}Mo		8067.4	1.5	8069.81	0.09	1.6	U					73Wa17	
		8066	2			1.9	U					77Ri04	
		8069.81	0.09				2			MMn		91Is02	Z
		8070.0	0.3			−0.6	U			Bdn		06Fi.A	
^{92}Mo(d,p)^{93}Mo		5853	20	5845.24	0.09	−0.4	U			Pit		64Co11	
^{92}Mo(p,γ)^{93}Tc		4081	5	4086.5	1.0	1.1	U					75Be.B	
		4086.5	1.0				2					83Ay01	
^{92}Mo(^{3}He,d)^{93}Tc		−1411	4	−1407.0	1.0	1.0	U			Hei		83Wi.A	
^{93}Kr(β^-)^{93}Rb		8700	500	8483	8	−0.4	U			Trs		78Wo15	
		8600	100			−1.2	U			Bwg		87Gr.A	
^{93}Rb(β^-)^{93}Sr		7560	120	7466	9	−0.8	U			Trs		78Wo15	
		7488	15			−1.4	o			Gsn		80Bl.A	
		7485	15			−1.2	o			Gsn		80De02	
		7440	30			0.9	–			McG		83Ia02	
		7455	35			0.3	–			Bwg		87Gr.A	
		7456	15			0.7	–			Gsn		92Pr03	
	ave.	7453	13			1.1	1	51	27 ^{93}Rb			average	
^{93}Sr(β^-)^{93}Y		4130	100	4141	12	0.1	U			Bwg		78St02	
		4110	20			1.6	1	34	24 ^{93}Y	McG		83Ia02	
^{93}Y(β^-)^{93}Zr		2890	20	2895	10	0.2	–					59Kn38	
		2880	15			1.0	–			McG		83Ia02	
	ave.	2884	12			0.9	1	76	76 ^{93}Y			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{93}Zr(β^-)^{93}Nb	93.8	2.	90.8	1.5	−1.5	1	55	53 ^{93}Nb			53Gl.A	*
^{93}Mo(ε)^{93}Nb	158	15	405.8	1.5	16.5	B					64Ho08	*
^{93}Nb(p,n)^{93}Mo	−1188	10	−1188.2	1.5	−0.0	U					68Fi01	
	−1190	5			0.4	U					75Ch05	
^{93}Tc(β^+)^{93}Mo	3185.1	5.	3201.0	1.0	3.2	B					51Bo48	*
	3192.1	3.			3.0	B					74An24	*
^{93}Ru(β^+)^{93}Tc	6337	85	6390.5	2.2	0.6	U					83Ay01	
^{93}Pd(ε)^{93}Rh	10030	370	9928	7	−0.3	U					19Pa16	
*^{93}Mo−u	$M-A$=–84385(28) keV for ^{93}Mo^m at 2424.95 keV										Ens115	**
*^{93}Zr(β^-)^{93}Nb	E_{β^-}=63(2) to 1/2$^-$ level at 30.77 keV										Ens115	**
*^{93}Mo(ε)^{93}Nb	L/K=0.36(0.04) to 1/2$^-$ level at 30.82 keV, recalculated Q										Ens115	**
*^{93}Tc(β^+)^{93}Mo	E_{β^+}=800(5) 807(3) respectively, to 7/2$^+$ level 1363.048 keV										Ens115	**
^{94}Br−u	−50242	429	−51150#	220#	−0.9	D			GT3	2.5	16Kn03	*
^{94}Kr−u	−66238	247	−65860	13	1.0	U			GT1	1.5	04Ma.A	
^{94}Kr−^{85}Rb$_{1.106}$	31701	13				2			MA8	1.0	06De36	
	31665	24	31701	13	1.5	U			MA8	1.0	10Na13	
	31649	97			0.5	U			MA9	1.0	10Na13	*
^{94}Rb−u	−73602	54	−73605.2	2.2	−0.0	F			Pb1	2.5	89Al33	*
^{94}Rb−^{85}Rb$_{1.106}$	23958	10	23955.4	2.2	−0.3	U			MA4	1.0	02Ra23	
	23955.6	2.6			−0.1	1	70	70 ^{94}Rb	TT1	1.0	12Si10	*
^{94}Sr−^{85}Rb$_{1.106}$	12924	10	12916.2	1.8	−0.8	U			MA4	1.0	02Ra23	
	12916.0	1.8			0.1	1	98	98 ^{94}Sr	TT1	1.0	12Si10	*
C$_7$ H$_{10}$−^{94}Zr	171929.4	3.9	171937.80	0.18	0.9	U			M15	2.5	63Ri07	
^{94}Zr−u	−93687.34	0.20	−93687.48	0.18	−0.7	1	77	77 ^{94}Zr	MS1	1.0	15Gu09	
^{94}Mo−^{85}Rb$_{1.106}$	2645.6	1.0	2644.11	0.14	−1.5	U			JY1	1.0	12Ka13	
C$_7$ H$_{10}$−^{94}Mo	173159.6	3.2	173166.76	0.14	0.9	U			M15	2.5	63Ri07	
^{94}Mo−u	−94916.31	0.42	−94916.44	0.14	−0.3	1	11	11 ^{94}Mo	MS1	1.0	15Gu09	
^{94}Tc−u	−90362	39	−90347	4	0.4	U			GS2	1.0	05Li24	*
^{94}Ru−^{85}Rb$_{1.106}$	8891	25	8903	3	0.5	U			SH1	1.0	08We10	
	8907.1	4.5			−0.8	1	56	56 ^{94}Ru	JY1	1.0	08We10	
^{94}Ru−C$_7$ H$_{10}$	−166912.2	5.1	−166907	3	0.9	1	44	44 ^{94}Ru	CP1	1.0	11Fa10	
^{94}Ru−u	−88618	28	−88657	3	−1.4	U			GR1	1.0	19An10	
	−88628	37			−0.8	U			GR1	1.0	23Mo07	
^{94}Rh−^{85}Rb$_{1.106}$	19291.2	4.6	19289	3	−0.6	1	57	57 ^{94}Rh	JY1	1.0	08We10	
^{94}Rh−C$_7$ H$_{10}$	−156520.2	5.9	−156522	3	−0.4	1	35	35 ^{94}Rh	CP1	1.0	11Fa10	
^{94}Rh$_{.989}$−C$_7$ H$_9$	−147834	30	−147836	3	−0.1	U			CP1	1.0	11Fa10	
^{94}Rh−u	−78376	103	−78272	3	1.0	U			GR1	1.0	19An10	*
	−78245.3	21.5			−1.2	U			GR1	1.0	23Mo07	
^{94}Rh^m−u	−77930.7	22.5				2			GR1	1.0	23Mo07	
^{94}Kr−^{86}Kr$_{1.093}$	31710	110	31843	13	1.2	U			MA9	1.0	10Na13	
^{94}Zr−^{87}Rb$_{1.080}$	4397.13	0.37	4397.55	0.18	1.1	1	23	23 ^{94}Zr	MS1	1.0	15Gu09	
^{94}Mo−^{87}Rb$_{1.080}$	3168.68	0.35	3168.59	0.14	−0.3	1	15	15 ^{94}Mo	MS1	1.0	15Gu09	
^{94}Rb−^{88}Rb$_{1.068}$	21109.1	4.0	21109.8	2.2	0.2	1	30	30 ^{94}Rb	JY1	1.0	07Ra23	
^{94}Zr ^{35}Cl−^{92}Zr ^{37}Cl	4235.0	2.	4227.41	0.21	−0.9	U			H13	4.0	63Ba20	
^{94}Mo ^{35}Cl−^{92}Mo ^{37}Cl	1234.0	2.	1226.53	0.23	−0.9	U			H11	4.0	63Bi12	
^{94}Mo−^{95}Mo$_{0.989}$	−1789.675	0.069	−1789.66	0.06	0.2	1	68	53 ^{94}Mo	JY2	1.0	22Ra20	
^{94}Rh−^{94}Pd	−7335	11	−7312	5	2.1	1	22	14 ^{94}Pd	RI1	1.0	26Ho.A	
^{94}Pd−^{94}Mo	23952.7	4.6	23957	4	0.9	1	86	86 ^{94}Pd	JY1	1.0	08We10	
^{92}Rb−^{94}Rb$_{.587}$ ^{89}Rb$_{.413}$	−764	24	−778	5	−0.2	U			P21	2.5	82Au01	Y
^{92}Rb−^{94}Rb$_{.489}$ ^{90}Rb$^x_{.511}$	−717	23	−729	8	−0.2	U			P21	2.5	82Au01	Y
^{93}Rb−^{94}Rb$_{.742}$ ^{90}Rb$^x_{.258}$	−1296	25	−1290	9	0.1	U			P21	2.5	82Au01	Y
^{93}Rb−^{94}Rb$_{.495}$ ^{92}Rb$_{.505}$	−840	40	−921	8	−0.8	F			P31	2.5	86Au02	*
^{94}Zr(d,α)^{92}Y	8278	25	8258	9	−0.8	1	13	13 ^{92}Y	Grn		74Gi09	
^{94}Zr(p,t)^{92}Zr	−6466	12	−6471.06	0.18	−0.4	U			Bld		66St15	
	−6470	10			−0.1	U			Oak		71Ba43	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{94}Ag^{n}(2p)^{92}Rh	3449	100	2411	18	−10.4	F					06Mu03	*
^{94}Rb(β^-n)^{93}Sr	3580	80	3452	8	−1.6	U					84Kr.B	
^{94}Zr(p,d)^{93}Zr	−5983	15	−5994.0	0.5	−0.7	U			Bld		66St15	
	−6000	10			0.6	U			Oak		71Ba43	
^{94}Zr(d,t)^{93}Zr	−1969	20	−1961.3	0.5	0.4	U			Pit		64Co11	
	−1960.2	2.4			−0.5	U			SPa		79Bo37	
^{93}Nb(n,γ)^{94}Nb	7229.13	0.12	7227.54	0.08	−13.2	C					84Bo.C	
	7227.51	0.09			0.3	–			MMn		88Ke09	Z
	7227.63	0.15			−0.6	–			Bdn		06Fi.A	
ave.	7227.54	0.08			−0.0	1	100	69 ^{94}Nb			average	
^{94}Mo(d,t)^{93}Mo	−3441	20	−3421.11	0.22	1.0	U			Pit		64Co11	
^{94}Ag^{n}(p)^{93}Pd	5784	30	5791	17	0.2	3					05Mu15	*
	5794	20			−0.2	3					09Ce04	*
^{94}Rb(β^-)^{94}Sr	10304	20	10282.9	2.6	−1.1	o			Gsn		80Bl.A	
	10322	100			−0.4	o			Gsn		80De02	*
	10353	140			−0.5	U			Trs		82Br23	*
	10335	45			−1.2	U			Bwg		82Pa24	*
	10312	20			−1.5	U			Gsn		92Pr03	*
^{94}Sr(β^-)^{94}Y	3512	10	3506	6	−0.6	1	41	40 ^{94}Y	Gsn		80De02	*
^{94}Y(β^-)^{94}Zr	4920	9	4918	6	−0.2	1	50	50 ^{94}Y	Gsn		80De02	*
^{94}Nb(β^-)^{94}Mo	2043.3	6.	2045.0	1.5	0.3	–					66Sn02	*
	2046.3	3.			−0.4	–					68Ho10	*
ave.	2045.7	2.7			−0.3	1	31	31 ^{94}Nb			average	
^{94}Tc(β^+)^{94}Mo	4261	5	4256	4	−1.0	2					64Ha29	*
^{94}Mo(p,n)^{94}Tc	−5028.8	7.	−5038	4	−1.4	2					73Mc04	*
^{94}Rh(β^+)^{94}Ru	9930	400	9674	5	−0.6	U					80Ox01	*
	9750	320			−0.2	U					06Ba55	
^{94}Pd(β^+)^{94}Rh	6700	320	6811	5	0.3	U					06Ba55	
^{94}Ag(ε)^{94}Pd	13400	650	13700#	400#	0.5	D					19Pa16	*
^{94}Ag^{n}(β^+)^{94}Pd	17700	500	20093	18	4.8	B					04Mu30	*
*^{94}Br−u	Trends in Mass Surface TMS suggest ^{94}Br 850 keV more bound										GAu	**
*^{94}Kr−^{85}Rb$_{1.106}$	Typo in original paper, ratio should read 1.00625564(114)										GAu	**
*^{94}Rb−u	F : possibly isomeric mixture										92Al.B	**
*^{94}Rb−^{85}Rb$_{1.106}$	D_M=23956.4(2.6) μu $M-A$=–68561.8(2.4)keV corrected for e$^-$ binding=–775 eV										12Si10	**
*^{94}Sr−^{85}Rb$_{1.106}$	D_M=12916.7(1.8) μu $M-A$=–78845.2(1.7)keV corrected for e$^-$ binding=–625 eV										12Si10	**
*^{94}Tc−u	$M-A$=–84133(29) keV for mixture gs+m at 74.9(2.4) keV										Nub269	**
*^{94}Rh−u	$M-A$=–72848(24) keV for mixture gs+m at 318(21) keV										Nub269	**
*^{93}Rb−^{94}Rb$_{.495}$ ^{92}Rb$_{.505}$	F : rejection based on line-shape analysis										86Au02	**
*^{94}Ag^{n}(2p)^{92}Rh	Q_{2p}=1900(100) to (11$^+$) level at 1548.6 keV										Ens12a	**
*^{94}Ag^{n}(2p)^{92}Rh	F : no evidence from He-jet experiment										09Ce04	**
*^{94}Ag^{n}(2p)^{92}Rh	F : possibly misidentified ^{92}Rh γ rays										09Je05	**
*^{94}Ag^{n}(p)^{93}Pd	E_p=790(30), 1010(30) to (3/2$^+$) at 4995.6, (3/2$^-$,35/2$^-$) at 4752.7										Ens115	**
*^{94}Ag^{n}(p)^{93}Pd	E_p=790(20) to level (3/2$^+$) at 4995.6 keV										Ens115	**
*^{94}Rb(β^-)^{94}Sr	Original value 10304(30) corrected in reference										94Ha.A	**
*^{94}Rb(β^-)^{94}Sr	Original error 100 keV increased by 100 in reference for lower level feeding										94Ha.A	**
*^{94}Rb(β^-)^{94}Sr	As corrected in reference										87Gr.A	**
*^{94}Rb(β^-)^{94}Sr	E_{β^-}=9475(20) to 2$^+$ level at 836.91 keV										Ens116	**
*^{94}Sr(β^-)^{94}Y	Original error 6 corrected in reference										94Ha.A	**
*^{94}Y(β^-)^{94}Zr	Original error 5 corrected in reference										94Ha.A	**
*^{94}Nb(β^-)^{94}Mo	E_{β^-}=470(6) 473(3) respectively, to level 4$^+$ at 1573.76 keV										Ens069	**
*^{94}Tc(β^+)^{94}Mo	E_{β^+}=816(5) to 6$^+$ level at 2423.45 keV										Ens069	**
*^{94}Mo(p,n)^{94}Tc	T=5158(7) to ^{94}Tc^m at 74.9(2.4) keV										Nub269	**
*^{94}Rh(β^+)^{94}Ru	E_{β^+}=6400(400) to (3,4,5) level at 2503.2 keV										Ens069	**
*^{94}Ag(ε)^{94}Pd	Symmetrized from 13350(+690–610)										HWJ199	**
*^{94}Ag(ε)^{94}Pd	Trends in Mass Surface TMS suggest ^{94}Ag 300 keV less bound										HWJ199	**
*^{94}Ag^{n}(β^+)^{94}Pd	Q_{β^+} larger than 17.7 MeV, uncertainty not given										04Mu30	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{95}Kr−u		−60183	150	−60289	20	−0.5	U				GT1	1.5	04Ma.A	
^{95}Kr−^{85}Rb$_{1.118}$		38330	20				2				MA8	1.0	06De36	
^{95}Rb−u		−70618	86	−70739	22	−0.6	U				Pb1	2.5	89Al33	
^{95}Sr−^{85}Rb$_{1.118}$		17987	10	17972	6	−1.5	1	34	34	^{95}Sr	MA4	1.0	02Ra23	
^{95}Y−^{85}Rb$_{1.118}$		11418.9	1.1	11419.1	1.1	0.2	1	99	99	^{95}Y	JY2	1.0	24Ja03	
^{95}Mo−^{85}Rb$_{1.118}$		4457.6	1.0	4456.51	0.13	−1.1	U				JY1	1.0	12Ka13	
C_7 H_{11}−^{95}Mo		180236.5	3.5	180237.92	0.13	0.2	U				M15	2.5	63Ri07	
^{95}Mo−u		−94161.96	0.38	−94162.57	0.13	−1.6	1	12	12	^{95}Mo	MS1	1.0	15Gu09	
^{95}Tc−u		−92417	32	−92341.92	0.19	2.3	U				GS2	1.0	05Li24	*
^{95}Rh−^{85}Rb$_{1.118}$		14515.1	4.5	14517	4	0.4	1	86	86	^{95}Rh	JY1	1.0	08We10	
^{95}Rh$_{.989}$−C_7 H_{10}		−161416	11	−161427	4	−1.0	1	14	14	^{95}Rh	CP1	1.0	11Fa10	
^{95}Pd−u		−75113	24	−75112	3	0.1	U				LZ1	1.0	23Xi01	
^{95}Pd^{m}−u		−74250	35				2				LZ1	1.0	23Xi01	
^{95}Mo−^{87}Rb$_{1.092}$		5012.33	0.44	5012.30	0.13	−0.1	U				MS1	1.0	15Gu09	
^{95}Sr−^{97}Zr$_{.979}$		6529	10	6519	6	−1.0	1	34	34	^{95}Sr	JY1	1.0	06Ha03	
^{95}Y−^{97}Zr$_{.979}$		−32.4	6.7	−33.5	1.1	−0.2	U				JY1	1.0	07Ha32	
^{95}Pd−^{94}Mo$_{1.011}$		20848.9	4.7	20849	3	0.0	2				JY1	1.0	08We10	
		20849.1	4.5			−0.0	2				JY1	1.0	08We10	*
^{95}Tc−^{95}Mo		1820.65	0.14	1820.65	0.14	0.0	1	100	100	^{95}Tc	JY2	1.0	24Ge09	
^{95}Ag−^{95}Mo		30025.6	1.5				2				JY2	1.0	24Ge07	
^{95}Mo−^{94}Mo		757	12	753.88	0.06	−0.1	U				M15	2.5	63Ri07	
^{93}Rb−^{95}Rb$_{.653}$ ^{89}Rb$_{.348}$		−1323	25	−1155	15	2.7	U				P21	2.5	82Au01	
^{93}Rb−^{95}Rb$_{.587}$ ^{90}Rb$^{x}_{.413}$		−1376	24	−1195	15	3.0	B				P21	2.5	82Au01	
^{94}Rb−^{95}Rb$_{.792}$ ^{90}Rb$^{x}_{.209}$		−16	28	195	16	3.0	B				P21	2.5	82Au01	Y
^{92}Rb−^{95}Rb$_{.242}$ ^{91}Rb$_{.758}$		80	23	117	7	0.6	U				P21	2.5	82Au01	
^{93}Rb−^{95}Rb$_{.489}$ ^{91}Rb$_{.511}$		−654	12	−662	13	−0.3	F				P31	2.5	86Au02	*
^{94}Rb−^{95}Rb$_{.660}$ ^{92}Rb$_{.341}$		433	15	424	14	−0.2	1	13	13	^{95}Rb	P31	2.5	86Au02	
		462	28			−0.5	U				P31	2.5	86Au02	
^{95}Mo(n,α)^{92}Zr		6405	30	6393.64	0.15	−0.4	U				ILL		75Em04	
^{95}Rb(β^-n)^{94}Sr		5120	100	4881	20	−2.4	U						84Kr.B	
^{94}Zr(n,γ)^{95}Zr		6461.6	1.0	6462.0	0.9	0.4	–						79Ke.D	Z
		6357.8	0.3			347.3	F				Bdn		06Fi.A	*
		6460.3	0.5			3.4	C				Bdn		08Fi.A	*
^{94}Zr(d,p)^{95}Zr		4223	20	4237.4	0.9	0.7	U				Pit		64Co11	
		4237.4	2.0			0.0	–				SPa		79Bo37	
^{94}Zr(n,γ)^{95}Zr	ave.	6461.7	0.9	6462.0	0.9	0.4	1	92	92	^{95}Zr			average	
^{94}Mo(n,γ)^{95}Mo		7367	2	7369.09	0.05	1.0	U						77Ri04	
		7369.10	0.10			−0.1	1	28	21	^{94}Mo	MMn		91Is02	Z
		7368.4	0.5			1.4	U				Bdn		06Fi.A	
^{94}Mo(d,p)^{95}Mo		5137	20	5144.52	0.05	0.4	U				Pit		64Co11	
^{95}Pd(εp)^{94}Ru		5116	300	5329	4	0.7	U						82Ku15	*
^{95}Rb(β^-)^{95}Sr		9224	30	9230	20	0.2	o				Gsn		80Bl.A	
		9276	100			−0.5	o				Gsn		80De02	*
		9300	30			−2.3	C				Gsn		84Bl.A	
		9280	45			−1.1	–				Bwg		87Gr.A	
		9272	35			−1.2	–				Gsn		92Pr03	
	ave.	9275	28			−1.6	1	53	52	^{95}Rb			average	
^{95}Sr(β^-)^{95}Y		6110	150	6104	5	−0.0	U						70Ma.A	
		6060	100			0.4	U				Bwg		78St02	
		6082	10			2.2	1	30	29	^{95}Sr	Gsn		84Bl.A	
		6052	25			2.1	U						90Ma03	
^{95}Y(β^-)^{95}Zr		4445	9	4433.8	1.3	−1.2	U				Gsn		80De02	*
^{95}Zr(β^-)^{95}Nb		1125	8	1126.3	1.0	0.2	U						54Za05	
		1119	5			1.5	U						55Dr43	
		1122.7	3.			1.2	1	11	8	^{95}Zr			74An22	*
^{95}Nb(β^-)^{95}Mo		925.5	0.5	925.6	0.5	0.2	1	98	97	^{95}Nb			63La06	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{95}Tc(β^+)^{95}Mo	1683	10	1695.92	0.13	1.3	U					65Cr04	*
	1693	6			0.5	U					74An05	*
^{95}Mo(p,n)^{95}Tc	−2440	30	−2478.27	0.13	−1.3	U					57Le27	
	−2490	6			2.0	B			Oak		70Ki01	
^{95}Ru(β^+)^{95}Tc	2558	30	2559	9	0.0	1	10	10 ^{95}Ru			68Pi03	*
^{95}Rh(β^+)^{95}Ru	5110	150	5117	10	0.0	U					75We03	*
^{95}Ag(ε)^{95}Pd	10560	480	10223	3	−0.7	U					19Pa16	
^{95}Cd(ε)^{95}Ag	10200	1700	12850#	400#	1.6	D					18Pa20	*
*^{95}Tc−u	$M-A$=–86066(28) keV for mixture gs+m at 38.91 keV										Nub269	**
*^{95}Pd−^{94}Mo$_{1.011}$	D_M=22862.1(4.5) μu for ^{95}Pd^m at 1875.13 keV; $M-A$=–68090.2(4.4) keV										Nub269	**
*^{93}Rb−^{95}Rb$_{.489}$ ^{91}Rb$_{.511}$	F : Rejected by authors										86Au02	**
*^{94}Zr(n,γ)^{95}Zr	F : value from 06Fi.A retracted										08Fi.A	**
*^{94}Zr(n,γ)^{95}Zr	Weak evidence										08Fi.A	**
*^{95}Pd(εp)^{94}Ru	E_p=4300(300) from ^{95}Pd^m at 1875.13 to ^{94}Ru^m at 2644.1 keV										Nub269	**
*	same E_p; both from figures										82No06	**
*^{95}Rb(β^-)^{95}Sr	E_{β^-}=8595(100) to (3/2$^+$,5/2$^+$) level at 680.70, corrected in reference										92Pr03	**
*^{95}Y(β^-)^{95}Zr	Original error 5 corrected in reference										94Ha.A	**
*	Q_{β^-}=4417(10) given by same group, not used										84Bl.A	**
*^{95}Zr(β^-)^{95}Nb	E_{β^-}=887(3) to 1/2$^-$ level at 235.69 keV										Ens10a	**
*^{95}Nb(β^-)^{95}Mo	E_{β^-}=159.7(0.5) to 7/2$^+$ level at 765.803 keV										Ens10a	**
*^{95}Tc(β^+)^{95}Mo	E_{β^+}=700(10) 710(6) respectively, from ^{95}Tc^m at 38.91 keV										Nub269	**
*^{95}Ru(β^+)^{95}Tc	E_{β^+}=1200(30) to 7/2$^+$ level at 336.413 keV										Ens10a	**
*^{95}Rh(β^+)^{95}Ru	E_{β^+}=3150(150) to 7/2$^+$ level at 941.79 keV										Ens10a	**
*^{95}Cd(ε)^{95}Ag	Trends in Mass Surface TMS suggest ^{95}Cd 2650 keV less bound										GAu	**
^{96}Kr−u	−57002	61	−57006	10	−0.1	U			TR1	1.0	20Sm.1	
	−57011.6	10.7			0.5	1	81	81 ^{96}Kr	MR1	1.0	25Lu16	
^{96}Kr−^{85}Rb$_{1.129}$	42606	22	42583	10	−1.0	1	19	19 ^{96}Kr	MA8	1.0	10Na13	
^{96}Rb−u	−65508	43	−65867	4	−3.3	F			Pb1	2.5	89Al33	*
^{96}Zr−^{87}Rb$_{1.103}$	8451.49	0.34	8451.49	0.12	0.0	1	13	13 ^{96}Zr	MS1	1.0	15Gu09	
C$_7$ H$_{12}$−^{96}Zr	185628	6	185622.77	0.12	−0.3	U			M15	2.5	63Ri07	
^{96}Zr−u	−91691	43	−91722.39	0.12	−0.7	U			JY0	1.0	04Ri12	
	−91722.60	0.17			1.3	1	51	51 ^{96}Zr	MS1	1.0	15Gu09	
^{96}Mo−^{85}Rb$_{1.129}$	4265.7	1.1	4264.15	0.13	−1.4	U			JY1	1.0	12Ka13	
^{96}Mo−^{87}Rb$_{1.103}$	4848.96	0.43	4848.65	0.13	−0.7	U			MS1	1.0	15Gu09	
C$_7$ H$_{12}$−^{96}Mo	189226.9	3.0	189225.62	0.13	−0.2	U			M15	2.5	63Ri07	
^{96}Mo−u	−95324.94	0.47	−95325.23	0.13	−0.6	U			MS1	1.0	15Gu09	
^{96}Tc−u	−92192	32	−92133	6	1.8	U			GS2	1.0	05Li24	*
C$_7$ H$_{12}$−^{96}Ru	186304.6	3.8	186311.40	0.14	0.7	U			M16	2.5	63Da10	
^{96}Ru−u	−92405.3	30.1	−92411.02	0.14	−0.2	U			GR1	1.0	23Mo07	
^{96}Rh−u	−85513.2	42.9	−85550	3	−0.9	U			GR1	1.0	23Mo07	
^{96}Pd−u	−81853	41	−81795.9	2.7	1.4	U			GR1	1.0	19An10	
^{96}Cd−u	−59776	32				2			MR1	1.0	26La.A	
^{96}Rb−^{88}Rb$_{1.091}$	30887.7	3.6	30888	4	0.1	1	100	100 ^{96}Rb	JY1	1.0	07Ra23	
^{96}Zr ^{35}Cl−^{94}Zr ^{37}Cl	4929	3	4915.22	0.22	−1.1	U			H13	4.0	63Ba20	
^{96}Mo ^{35}Cl−^{94}Mo ^{37}Cl	2539	2	2541.34	0.10	0.3	U			H11	4.0	63Bi12	
^{96}Pd−^{94}Mo$_{1.021}$	15123.4	4.5	15113.8	2.7	−2.1	1	35	35 ^{96}Pd	JY1	1.0	08We10	
^{96}Sr−^{97}Zr$_{.990}$	9868	10	9865	9	−0.3	1	82	82 ^{96}Sr	JY1	1.0	06Ha03	
^{96}Y−^{97}Zr$_{.990}$	4053.7	6.8	4055	5	0.1	1	47	47 ^{96}Y	JY1	1.0	07Ha32	
^{96}Y^m−^{97}Zr$_{.990}$	5708.1	6.7	5709	5	0.1	1	49	49 ^{96}Y^m	JY1	1.0	07Ha32	
^{96}Mo−^{97}Mo$_{.990}$	−2280.5	5.8	−2281.93	0.17	−0.2	U			JY1	1.0	06Ka48	
^{96}Zr−^{96}Nb	176.02	0.13	176.03	0.11	0.1	1	68	63 ^{96}Nb	JY1	1.0	16Al03	
^{96}Zr−^{96}Mo	3602.919	0.092	3602.85	0.08	−0.8	1	75	45 ^{96}Mo	JY1	1.0	16Al03	
^{96}Nb−^{96}Mo	3426.80	0.17	3426.82	0.11	0.1	1	46	37 ^{96}Nb	JY1	1.0	16Al03	
^{96}Ru−^{96}Mo	2914.14	0.13	2914.21	0.05	0.6	–			SH1	1.0	11El04	
	2914.225	0.053			−0.2	–			JY2	1.0	22Ne04	
ave.	2914.21	0.05			0.0	1	100	100 ^{96}Ru			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{96}Ru−^{96}Pd	−10613.7	5.1	−10615.1	2.7	−0.3	1	27	27 ^{96}Pd	RI1	1.0	26Ho.A	
^{96}Rh−^{96}Pd	−3753.8	1.5				2			RI1	1.0	26Ho.A	
^{96}Ag−^{96}Pd	12391.8	4.2	12384.4	2.7	−1.8	1	41	38 ^{96}Pd	RI1	1.0	26Ho.A	
^{96}Cd−^{96}Pd	22162	17	22020	30	−8.4	B			RI1	1.0	26Ho.A	
^{96}Ag−^{96}Mo	25913.41	0.91	25913.8	0.9	0.4	1	97	97 ^{96}Ag	JY2	1.0	24Ge07	
^{96}Ag^{m}−^{96}Mo	26036.8	1.6				2			JY2	1.0	24Ge07	
^{96}Mo−^{95}Mo	−1161	12	−1162.67	0.05	−0.1	U			M15	2.5	63Ri07	
^{93}Rb−^{96}Rb$_{.554}$ ^{89}Rb$_{.448}$	−2210	27	−2022	8	2.8	U			P21	2.5	82Au01	
^{95}Rb−^{96}Rb$_{.848}$ ^{89}Rb$_{.152}$	−1590	30	−1444	20	1.9	U			P21	2.5	82Au01	
^{94}Rb−^{96}Rb$_{.699}$ ^{89}Rb$_{.302}$	−1250	30	−999	3	3.3	B			P21	2.5	82Au01	Y
^{94}Rb−^{96}Rb$_{.588}$ ^{91}Rb$_{.413}$	−380	25	−371.2	3.0	0.1	U			P21	2.5	82Au01	
^{95}Rb−^{96}Rb$_{.742}$ ^{92}Rb$_{.258}$	−1116	27	−1077	20	0.6	U			P21	2.5	82Au01	
	−1143	16			1.6	1	26	25 ^{95}Rb	P31	2.5	86Au02	
^{96}Zr(d,α)^{94}Y	7609	20	7623	6	0.7	1	10	10 ^{94}Y	Grn		74Gi09	
^{96}Zr(p,t)^{94}Zr	−5825	10	−5830.37	0.20	−0.5	U			Oak		71Ba43	
^{96}Ru(p,t)^{94}Ru	−11165	10	−11158	3	0.7	U			Oak		71Ba01	
^{96}Zr(t,α)^{95}Y	8294	20	8312.3	1.0	0.9	U			LAl		83Fl06	
^{96}Zr(p,d)^{95}Zr	−5440	20	−5625.6	0.9	−9.3	B			Bld		67St24	
	−5630	10			0.4	U			Oak		71Ba43	
^{96}Zr(d,t)^{95}Zr	−1603	20	−1592.9	0.9	0.5	U			Pit		64Co11	
	−1595.8	2.8			1.0	U			SPa		79Bo37	
^{96}Mo(t,α)^{95}Nb	10524	20	10516.3	0.5	−0.4	U			LAl		83Fl06	
^{95}Mo(n,γ)^{96}Mo	9154.2	0.5	9154.34	0.05	0.3	U					70He27	
	9154.32	0.05			0.3	1	96	65 ^{95}Mo	MMn		91Is02	Z
	9153.90	0.20			2.2	U			Bdn		06Fi.A	
^{96}Mo(d,t)^{95}Mo	−2923	20	−2897.11	0.05	1.3	U			Pit		64Co11	
^{96}Ru(p,d)^{95}Ru	−8470	10	−8470	9	0.0	1	90	90 ^{95}Ru	Oak		71Ba01	
^{96}Ag(εp)^{95}Rh	6540	90	6395	4	−1.6	U					03Ba39	*
^{96}Rb(β^-)^{96}Sr	10800	220	11564	9	3.5	B					79Pe17	
	11303	250			1.0	o			Gsn		80De02	
	11547	100			0.2	U			Trs		82Br23	
	11553	45			0.2	U			Gsn		84Bl.A	
	11590	80			−0.3	U			Bwg		87Gr.A	
	11709	40			−3.6	B			Gsn		92Pr03	*
^{96}Sr(β^-)^{96}Y	5332	30	5412	9	2.7	F					79Pe17	*
	5413	22			−0.0	–			Gsn		80De02	*
	5345	50			1.3	U			Bwg		87Gr.A	
	5354	40			1.5	–					90Ma03	
ave.	5399	19			0.7	1	22	18 ^{96}Sr			average	
^{96}Y^{m}(IT)^{96}Y	1540.6	0.4	1540.6	0.4	−0.0	1	100	51 ^{96}Y^{m}			20Is08	*
^{96}Y(β^-)^{96}Zr	7120	50	7108	4	−0.2	U			Gsn		80De02	*
	7030	70			1.1	U			Bwg		87Gr.A	
	7067	30			1.4	U					90Ma03	*
^{96}Y^{m}(β^-)^{96}Zr	8030	150	8649	4	4.1	C			Bwg		87Gr.A	
	8600	200			0.2	U					88St.A	
	8237	21			19.6	C			Bwg		92Gr.A	
^{96}Nb(β^-)^{96}Mo	3186.8	3.2	3192.06	0.11	1.6	U					68An03	*
^{96}Mo(p,n)^{96}Tc	−3760	10	−3756	5	0.4	2					74Do09	
	−3754	6			−0.3	2					78Ke10	
^{96}Rh(β^+)^{96}Ru	6472	200	6391.3	2.8	−0.4	U					75Gu01	*
^{96}Ru(p,n)^{96}Rh	−7175	10	−7173.6	2.8	0.1	U					70As08	Z
^{96}Pd(β^+)^{96}Rh	3450	150	3496.6	1.4	0.3	U					85Ry02	*
^{96}Cd(ε)^{96}Ag^{m}	8480	460	8861	30	0.8	U					19Pa16	
*^{96}Rb−u	F : possibly isomeric mixture										92Al.B	**
*^{96}Tc−u	$M-A$=–85860(28) keV for mixture gs+m at 34.23 keV										Nub269	**
*^{96}Cd−u	Laser on and off										26La.A	*G
*^{96}Ag(εp)^{95}Rh	Original 6430(60) corrected by –110 keV for mixture of two β-decaying ^{96}Ag										03Ba39	**
*	states, to two isomeric states in ^{95}Rh										03Ba39	**
*^{96}Rb(β^-)^{96}Sr	E_{β^-}=10894(40) to 2^+ level at 814.93 keV										Ens08a	**
*^{96}Sr(β^-)^{96}Y	E_{β^-}=4400(30) to 1^+ level at 931.70 keV, and other E_{β^-}										Ens08a	**
*^{96}Sr(β^-)^{96}Y	F : all other results of reference are strongly conflicting										79Pe17	**
*^{96}Sr(β^-)^{96}Y	Original error 20 corrected in reference										94Ha.A	**
*	Q_{β^-}=5362(10) given by same group, not used										84Bl.A	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
$*^{96}Y^m(IT)^{96}Y$		lsq fit to Eg in 20Is08											Nub255	**
$*^{96}Y(\beta^-)^{96}Zr$		Q_{β^-}=7079(15) given by same group, not used											84Bl.A	**
$*^{96}Y(\beta^-)^{96}Zr$		E_{β^-}=5326(36) to 2^+ level at 1750.497 keV, and other E_{β^-}											Ens08a	**
$*^{96}Nb(\beta^-)^{96}Mo$		E_{β^-}=748.4(3.1) to 5^+ level at 2438.477 keV											Ens08a	**
$*^{96}Rh(\beta^+)^{96}Ru$		E_{β^+}=3300(200) to 6^+ level at 2149.74 keV											Ens08a	**
$*^{96}Pd(\beta^+)^{96}Rh$		p^+ =0.257(0.03) to 1^+ level at 1274.78 keV											Ens08a	**
$^{97}Kr-^{86}Kr_{1.128}$		49920	140	49850	30	−0.5	1	6	6	^{97}Kr	MA9	1.0	10Na13	
$^{97}Kr-^{39}K_{2.487}$		39260	38	39280	30	0.4	1	76	76	^{97}Kr	MA8	1.0	25Lu16	
$^{97}Kr-u$		−50941	76	−50990	30	−0.6	1	19	19	^{97}Kr	MR1	1.0	25Lu16	
$^{97}Rb-u$		−62512	64	−62822.9	2.1	−1.9	U				Pb1	2.5	89Al33	
$^{97}Rb-^{88}Rb_{1.102}$		34908.0	5.7	34907.4	2.1	−0.1	1	13	13	^{97}Rb	JY1	1.0	07Ra23	
$^{97}Rb-^{85}Rb_{1.141}$		37824.9	2.2	37825.0	2.1	0.1	1	87	87	^{97}Rb	TT1	1.0	12Si10	*
$^{97}Sr-^{85}Rb_{1.141}$		27022.9	3.9	27024	4	0.2	1	87	87	^{97}Sr	TT1	1.0	12Si10	*
$^{97}Sr-u$		−73599	99	−73624	4	−0.1	U				GT2	2.5	08Kn.A	
$^{97}Mo-^{85}Rb_{1.141}$		6666.9	1.2	6664.78	0.17	−1.8	U				JY1	1.0	12Ka13	
$^{97}Mo-^{87}Rb_{1.115}$		7280.86	0.39	7280.57	0.17	−0.7	1	20	20	^{97}Mo	MS1	1.0	15Gu09	
$C_5\ H_5\ O_2-^{97}Mo$		122937.6	2.3	122937.54	0.17	−0.0	U				M15	2.5	63Ri07	
$^{97}Mo-u$		−93982.95	0.36	−93983.14	0.17	−0.5	1	24	24	^{97}Mo	MS1	1.0	15Gu09	
$^{97}Ru-u$		−92471	30	−92454.2	3.0	0.6	U				GS2	1.0	05Li24	
$^{97}Rh-u$		−88621.1	20.4	−88634	4	−0.6	U				GR1	1.0	23Mo07	*
$^{97}Pd-^{85}Rb_{1.141}$		17119.9	5.2	17118.8	1.7	−0.2	1	10	10	^{97}Pd	JY1	1.0	09El08	
$^{97}Pd-u$		−83511	40	−83529.2	1.7	−0.5	U				GR1	1.0	19An10	
		−83547	63			0.3	U				GR1	1.0	23Mo07	
$^{97}Ag-u$		−76118.6	12.9	−76152.1	1.5	−2.6	U				GR1	1.0	20Ho03	
		−76157.2	20.4			0.3	U				GR1	1.0	23Mo07	
$^{97}Ag^m-u$		−75455.1	38.6				2				GR1	1.0	20Ho03	*
		−75505	81	−75460	40	0.6	U				GR1	1.0	23Mo07	*
$^{97}Cd-u$		−64971	26	−64963	12	0.3	1	23	23	^{97}Cd	MR1	1.0	26La.A	
$^{97}Mo\ ^{35}Cl-^{95}Mo\ ^{37}Cl$		3138	2	3129.56	0.19	−1.1	U				H11	4.0	63Bi12	
$^{97}Sr-^{97}Zr$		15416	10	15412	4	−0.4	1	13	13	^{97}Sr	JY1	1.0	06Ha03	
$^{97}Y-^{97}Zr$		7322.9	7.2				2				JY1	1.0	07Ha32	*
$^{97}Tc-^{97}Mo$		348.70	0.22				2				JY2	1.0	25Ge08	
$^{97}Ru-^{97}Pd$		−8913	28	−8925	3	−0.4	U				RI1	1.0	26Ho.A	
$^{97}Rh-^{97}Pd$		−5104.7	3.3				2				RI1	1.0	26Ho.A	
$^{97}Ag-^{97}Pd$		7377.11	0.72	7377.1	0.7	−0.0	1	98	89	^{97}Pd	RI1	1.0	26Ho.A	
$^{97}Cd-^{97}Pd$		18569	14	18567	12	−0.2	1	78	77	^{97}Cd	RI1	1.0	26Ho.A	
$^{97}Ag-^{97}Mo$		17831.0	1.6	17831.1	1.5	0.0	1	91	91	^{97}Ag	JY1	1.0	24Ge07	
$^{97}Cd^n-^{97}Cd$		2411	25				2				IS1	1.0	26La.A	
$^{97}Mo-^{96}Mo$		1346	12	1342.09	0.17	−0.1	U				M15	2.5	63Ri07	
$^{94}Rb-^{97}Rb_{.485}\ ^{91}Rb_{.516}$		−21	25	−56.5	2.5	−0.6	U				P21	2.5	82Au01	Y
$^{96}Rb-^{97}Rb_{.792}\ ^{92}Rb_{.209}$		650	30	620	4	−0.4	U				P21	2.5	82Au01	
$^{95}Rb-^{97}Rb_{.490}\ ^{93}Rb_{.511}$		−165	25	−110	21	0.9	1	11	10	^{95}Rb	P21	2.5	82Au01	
$^{96}Rb-^{97}Rb_{.742}\ ^{93}Rb_{.258}$		848	19	803	4	−1.0	U				P31	2.5	86Au02	
$^{96}Zr(n,\gamma)^{97}Zr$		5574	5	5569.15	0.04	−1.0	U						77Ba33	
		5575.1	0.4			−14.9	C				Bdn		06Fi.A	
		5569.15	0.04			0.0	1	100	100	^{97}Zr	ILn		18Rz01	
$^{96}Zr(d,p)^{97}Zr$		3338	20	3344.58	0.04	0.3	U				Pit		64Co11	
$^{96}Mo(n,\gamma)^{97}Mo$		6821.1	1.0	6821.16	0.16	0.1	U						73De39	
		6820	2			0.6	U						77Ri04	
		6821.15	0.25			0.1	–				MMn		91Is02	Z
		6821.5	0.4			−0.8	–				Bdn		06Fi.A	
$^{96}Mo(d,p)^{97}Mo$		4582	20	4596.60	0.16	0.7	U				Pit		64Co11	
$^{96}Mo(n,\gamma)^{97}Mo$	ave.	6821.25	0.21	6821.16	0.16	−0.4	1	58	43	^{97}Mo			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{96}Mo(^{3}He,d)^{97}Tc	229	8	220.53	0.26	−1.1	U			ANL		74Co27	
	220	8			0.1	U			Pit		74Co27	
^{96}Ru(d,p)^{97}Ru	5886	3	5886.9	2.8	0.3	2			Can		77Ho02	
	5892	7			−0.7	2			ANL		77Me04	
^{97}Rb(β^-)^{97}Sr	10020	50	10062	4	0.8	U			Gsn		80De02	
	10450	30			−12.9	C			Gsn		84Bl.A	
	10440	60			−6.3	C			Bwg		87Gr.A	
	10462	40			−10.0	B			Gsn		92Pr03	
^{97}Sr(β^-)^{97}Y	7452	40	7535	8	2.1	U			Gsn		84Bl.A	
	7420	80			1.4	o			Bwg		87Gr.A	
	7480	18			3.0	C			Bwg		92Gr.A	
^{97}Y(β^-)^{97}Zr	6702	25	6821	7	4.8	C			Gsn		84Bl.A	
	6640	70			2.6	o			Bwg		87Gr.A	*
	6689	13			10.2	C			Bwg		92Gr.A	*
^{97}Zr(β^-)^{97}Nb	2657.3	6.	2666	4	1.5	1	50	50 ^{97}Nb			74Ra.A	*
^{97}Nb(β^-)^{97}Mo	1933.1	6.	1942	4	1.5	1	50	50 ^{97}Nb			74Ra.A	*
^{97}Mo(p,n)^{97}Tc	−1128	9	−1107.16	0.20	2.3	B			Oak		70Ki01	
	−1102	6			−0.9	U			ANL		74Co27	
^{97}Ru(β^+)^{97}Tc	1150	100	1099.4	2.8	−0.5	U					70Ho01	*
^{97}Rh(β^+)^{97}Ru	3533	50	3559	4	0.5	U					62Ba28	*
	3513	50			0.9	U					62Ch21	*
^{97}Pd(β^+)^{97}Rh	4790	300	4755	3	−0.1	U					80Go11	*
^{97}Ag(β^+)^{97}Pd	6980	110	6871.7	0.7	−1.0	U					99Hu10	
^{97}Cd(ε)^{97}Ag	10170	420	10423	11	0.6	U					19Pa16	
^{97}In(ε)^{97}Cd	10000	3000	12820#	400#	0.9	U					18Pa20	
*^{97}Rb−^{85}Rb$_{1.141}$	D_M=37825.7(2.2) μu, $M-A$=–58518.5(2.1)keV corrected for e$^-$ binding=–703eV										12Si10	**
*^{97}Sr−^{85}Rb$_{1.141}$	D_M=27023.5(3.9) μu, $M-A$=–68580.7(3.7)keV corrected for e$^-$ binding=–553eV										12Si10	**
*^{97}Rh−u	isomer-to-gs. ratio is measured.										23Mo07	**
*^{97}Ag^m−u	reconstructed from E_x =618(38) keV										20Ho03	**
*^{97}Ag^m−u	reconstructed from E_x =608(73) keV										23Mo07	**
*^{97}Y−^{97}Zr	D_M=8039.5(7.2) μu for ^{97}Y^m at 667.52(0.23) keV; $M-A$=–75453.9(6.7) keV										Nub269	**
*^{97}Y(β^-)^{97}Zr	E_{β^-}=6645(70); and 7280(150) from ^{97}Y^m at 667.52 keV										Nub269	**
*^{97}Y(β^-)^{97}Zr	E_{β^-}=6688(13); and 7361(26) from ^{97}Y^m at 667.52 keV										Nub269	**
*^{97}Zr(β^-)^{97}Nb	E_{β^-}=1914(2) to 1/2$^-$ level at 743.35 keV; error increased										Ens104	**
*^{97}Nb(β^-)^{97}Mo	E_{β^-}=1275(2) to 7/2$^+$ level at 658.13 keV; error increased										Ens104	**
*^{97}Ru(β^+)^{97}Tc	p$^+$ <0.0001 (see fig 1), decay to 7/2$^+$ level at 970.03 keV										Ens104	**
*^{97}Rh(β^+)^{97}Ru	E_{β^+}=2090(50) 2070(50) respectively, to 7/2$^+$ level at 421.54 keV										Ens104	**
*^{97}Pd(β^+)^{97}Rh	E_{β^+}=3500(300) to 7/2$^+$ level at 265.36 keV										Ens104	**
^{98}Kr−u		−47503	56			2			MR1	1.0	25Lu16	
^{98}Rb−^{85}Rb$_{1.153}$		43331	32	43339	17	0.2	1	29	29 ^{98}Rb	TT1	1.0	12Si10	*
^{98}Rb−u		−58359	29	−58368	17	−0.3	–			MA8	1.0	13Ma81	
		−58370	29			0.1	–			TT1	1.0	16Kl04	*
	ave.	−58365	21			−0.2	1	71	71 ^{98}Rb			average	
^{98}Rb^m−u		−58289	22			2			TT1	1.0	12Si10	*
^{98}Sr−^{85}Rb$_{1.153}$		30396.7	4.3	30399	3	0.6	–			TT1	1.0	12Si10	*
		30405.3	7.2			−0.9	–			TT1	1.0	16Kl04	*
	ave.	30399	4			0.0	1	88	88 ^{98}Sr			average	
^{98}Y^n−u		−77110.6	2.0	−77110.3	1.9	0.1	1	95	95 ^{98}Y^n	JY2	1.0	25St02	*
^{98}Zr−u		−87247	43	−87260	9	−0.3	U			JY0	1.0	04Ri12	
^{98}Mo−^{85}Rb$_{1.153}$		7104.1	5.7	7110.00	0.18	1.0	U			MA8	1.0	11He10	
		7111.6	1.3			−1.2	U			JY1	1.0	12Ka13	
^{98}Mo−^{87}Rb$_{1.126}$		7665.96	0.64	7666.29	0.18	0.5	U			MS1	1.0	15Gu09	
C_5 H_6 O_2−^{98}Mo		131375.4	2.8	131375.87	0.18	0.1	U			M15	2.5	63Ri07	
^{98}Mo−u		−94596.21	0.53	−94596.44	0.18	−0.4	1	12	12 ^{98}Mo	MS1	1.0	15Gu09	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$C_7\ H_{14}-{}^{98}Ru$		204263.5	2.9	204268.91	0.20	0.7	U			M16	2.5	63Da10	
${}^{98}Rh-u$		−89300	37	−89297	11	0.1	U			GS2	1.0	05Li24	*
${}^{98}Pd-{}^{85}Rb_{1.153}$		14404.5	5.1	14405	5	0.0	1	100	100 ${}^{98}Pd$	JY1	1.0	09El08	
${}^{98}Pd-u$		−87307.1	22.5	−87302	5	0.2	U			GR1	1.0	23Mo07	
${}^{98}Ag-{}^{85}Rb_{1.153}$		23283	40	23256	28	−0.7	1	48	48 ${}^{98}Ag$	MA8	1.0	11He10	
${}^{98}Cd-u$		−72706.0	64.4	−72637	12	1.1	U			GR1	1.0	23Mo07	
		−72638	12			0.1	1	96	96 ${}^{98}Cd$	MR1	1.0	26La.A	
${}^{98}Mo\ {}^{35}Cl-{}^{96}Mo\ {}^{37}Cl$		3690	2	3678.92	0.20	−1.4	U			H11	4.0	63Bi12	
${}^{98}Sr-{}^{97}Zr_{1.010}$		18620	10	18619	3	−0.1	1	12	12 ${}^{98}Sr$	JY1	1.0	06Ha03	
${}^{98}Y-{}^{97}Zr_{1.010}$		12321.4	8.5	12316.3	2.1	−0.6	1	6	6 ${}^{98}Y$	JY1	1.0	07Ha32	
${}^{98}Zr-{}^{97}Zr_{1.010}$		2668	10	2667	9	−0.1	1	82	82 ${}^{98}Zr$	JY1	1.0	06Ha03	
${}^{98}Mo-{}^{97}Mo_{1.010}$		327.9	5.8	326.53	0.07	−0.2	U			JY1	1.0	06Ka48	
${}^{98}Y^{n}-{}^{98}Y$		499.99	0.78	499.9	0.8	−0.1	1	99	94 ${}^{98}Y$	JY1	1.0	17Ur03	
${}^{98}Mo-{}^{98}Ru$		122.028	0.073				2			JY2	1.0	22Ne04	
${}^{98}Mo-{}^{97}Mo$		−614	12	−613.30	0.07	0.0	U			M15	2.5	63Ri07	
${}^{94}Rb-{}^{98}Rb_{.411}\ {}^{91}Rb_{.590}$		−290	40	−338	7	−0.5	U			P21	2.5	82Au01	Y
${}^{97}Rb-{}^{98}Rb_{.792}\ {}^{93}Rb_{.209}$		−250	60	−281	13	−0.2	U			P21	2.5	82Au01	
${}^{96}Rb-{}^{98}Rb_{.490}\ {}^{94}Rb_{.511}$		330	30	322	9	−0.1	U			P21	2.5	82Au01	Y
${}^{97}Rb-{}^{98}Rb_{.660}\ {}^{95}Rb_{.340}$		−300	50	−232	13	0.5	U			P21	2.5	82Au01	
		−232	27			0.0	U			P31	2.5	86Au02	
${}^{96}Zr(t,p){}^{98}Zr$		3508	20	3504	8	−0.2	1	18	18 ${}^{98}Zr$	LAl		69Bl01	
${}^{96}Zr({}^{3}He,p){}^{98}Nb$		5728	5				2			Phi		75Me13	
${}^{96}Ru({}^{16}O,{}^{14}C){}^{98}Pd$		−12529	20	−12516	5	0.6	U			BNL		82Th01	
${}^{98}Mo(t,\alpha){}^{97}Nb$		10019	20	10012	4	−0.4	U			LAl		83Fl06	
${}^{97}Mo(n,\gamma){}^{98}Mo$		8642.4	0.5	8642.60	0.06	0.4	U					71He10	
		8642.60	0.07			0.0	–			MMn		91Is02	Z
		8642.57	0.18			0.2	–			Bdn		06Fi.A	
${}^{98}Mo(d,t){}^{97}Mo$		−2379	20	−2385.37	0.06	−0.3	U			Pit		64Co11	
${}^{97}Mo(n,\gamma){}^{98}Mo$	ave.	8642.60	0.07	8642.60	0.06	0.1	1	98	85 ${}^{98}Mo$			average	
${}^{97}Mo({}^{3}He,d){}^{98}Tc$		680	8	684	3	0.5	–			ANL		74Co27	
		686	10			−0.2	–			McM		76Ma16	
	ave.	682	6			0.3	1	29	29 ${}^{98}Tc$			average	
${}^{98}Rb(\beta^-){}^{98}Sr$		11200	110	12053	16	7.8	B					79Pe17	
		12343	150			−1.9	U			Trs		82Br23	
		12519	60			−7.8	C			Gsn		84Bl.A	
		12270	30			−7.2	C			McG		84Ia.A	
		12440	75			−5.2	C			Bwg		87Gr.A	
		12380	65			−5.0	B			Gsn		92Pr03	
${}^{98}Rb^{m}(\beta^-){}^{98}Sr$		12710	120	12127	21	−4.9	C			Bwg		87Gr.A	
${}^{98}Sr(\beta^-){}^{98}Y$		5730	40	5871	4	3.5	C					79Pe17	
		5821	10			5.0	C			Gsn		84Bl.A	
		5815	40			1.4	U			Bwg		87Gr.A	
${}^{98}Y(\beta^-){}^{98}Zr$		8974	100	8988	9	0.1	U					79Pe17	*
		8780	30			6.9	C			Gsn		84Bl.A	
		8840	55			2.7	C			Bwg		87Gr.A	*
		8963	41			0.6	U					88Ma.A	
		8830	17			9.3	C			Bwg		92Gr.A	
		9062	27			−2.7	C			Bwg		92Gr.A	*
${}^{98}Zr(\beta^-){}^{98}Nb$		2300	200	2243	10	−0.3	U					76He10	
${}^{98}Nb(\beta^-){}^{98}Mo$		4300	200	4591	5	1.5	U					66Gu05	
		4300	200			1.5	U					67Hu07	
		4800	200			−1.0	U					76He10	
		4580	100			0.1	U			Bwg		78St02	
${}^{98}Mo(p,n){}^{98}Tc$		−2458	10	−2465	3	−0.7	1	11	11 ${}^{98}Tc$	ANL		74Co27	
${}^{98}Tc(\beta^-){}^{98}Ru$		1795	22	1797	3	0.1	U					73Ok.A	*
${}^{98}Rh(\beta^+){}^{98}Ru$		5120	100	5050	10	−0.7	U					72Ba37	*
		5151	50			−2.0	U					94Ba06	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{98}Ru(p,n)^{98}Rh	−5832	10				3					70As08	Z
^{98}Ag(β^+)^{98}Pd	8420	150	8245	26	−1.2	U					79Ve.A	*
	8200	70			0.6	1	14	14 ^{98}Ag			00Hu17	
^{98}Cd(β^+)^{98}Ag	5330	140	5416	26	0.6	U					92Pl01	
^{98}Cd(ε)^{98}Ag	5430	40			−0.4	1	43	39 ^{98}Ag			01St.A	
^{98}In(ε)^{98}Cd	12930	400	13730#	300#	2.0	D					19Pa16	*
*^{98}Rb−^{85}Rb$_{1.153}$	Original D_M=43394.0(3.7) μu $M-A$=–54317.7(3.4) keV corrected for gs+m mixture										16Kl04	**
*^{98}Rb−^{85}Rb$_{1.153}$	Original $M-A$=–54309.4(4.0) corrected for isom mixture E_x =80(30) R=0.65(0.15)										16Kl04	**
*^{98}Rb−u	Original $M-A$=–54319.5(5.5) corrected for isom mixture E_x =80(30) R=0.65(0.15)										16Kl04	**
*^{98}Rb^m−u	Data re-analysis										16Kl04	**
*^{98}Sr−^{85}Rb$_{1.153}$	D_M=30397.3(4.3) μu $M-A$=–66424.0(4.0)keV corrected for e$^-$ binding=–529eV										12Si10	**
*^{98}Sr−^{85}Rb$_{1.153}$	Former measured using 15$^+$ ion; latter with 10$^+$ ion										16Kl04	**
*^{98}Y^n−u	from $M-A$=–76565.1(1.9) for ^{98}Y$_{16}$O										HWJ25a	**
*^{98}Rh−u	$M-A$=–83154(30) keV for mixture gs+m at 56.3(1.0) keV										Nub269	**
*^{98}Y(β^-)^{98}Zr	E_{β^-}=4810(100) to 1$^-$ level at 4164.60 keV										Ens035	**
*^{98}Y(β^-)^{98}Zr	also Q_{β^-}=9780(200) from ^{98}Y^m at 170.76 keV										87Gr.A	**
*^{98}Y(β^-)^{98}Zr	Q_{β^-}=9233(27) from ^{98}Y^m at 170.76 keV										HWJ189	**
*^{98}Tc(β^-)^{98}Ru	E_{β^-}=397(22) to 4$^+$ level at 1397.82 keV;										FKo261	**
*^{98}Tc(β^-)^{98}Ru	unc=22 in 73Ok.A, 95% CL (2sigma)										FKo261	**
*^{98}Rh(β^+)^{98}Ru	E_{β^+}=3450(100) to 2$^+$ level at 652.44 keV, and others										Ens035	**
*^{98}Rh(β^+)^{98}Ru	E_{β^+}=3476(50) to 2$^+$ level at 652.44 keV										Ens035	**
*^{98}Ag(β^+)^{98}Pd	Q_{β^+}=6880(150) to 4$^+$ level at 1541.40 keV										Ens035	**
*^{98}In(ε)^{98}Cd	Trends in Mass Surface TMS suggest ^{98}In 600 keV less bound										HWJ199	**
^{99}Rb−^{85}Rb$_{1.165}$	47885.1	4.8	47884	4	−0.2	2			MA8	1.0	13Ma81	
	47880	10			0.4	2			TT1	1.0	16Kl04	
^{99}Rb−u	−54859	33	−54881	4	−0.7	U			TR1	1.0	21Mu05	
^{99}Sr−^{85}Rb$_{1.165}$	35663	21	35649	5	−0.7	o			TT1	1.0	12Si10	*
	35644.5	7.0			0.6	1	53	53 ^{99}Sr	TT1	1.0	16Kl04	
^{99}Sr−u	−67106	33	−67116	5	−0.3	U			TR1	1.0	21Mu05	
^{99}Zr−u	−83323	19	−83325	11	−0.1	1	35	35 ^{99}Zr	JY0	1.0	04Ri12	
C$_7$ H$_{15}$−^{99}Ru	211442.8	3.0	211445.0	0.4	0.3	U			M16	2.5	63Da10	
^{99}Ag−^{85}Rb$_{1.165}$	20401.0	8.5	20414	4	1.5	–			SH1	1.0	07Ma92	
	20427	11			−1.2	–			MA8	1.0	11He10	
ave.	20411	7			0.5	1	31	31 ^{99}Ag			average	
^{99}Ag−u	−82350.5	35.4	−82351	4	−0.0	U			GR1	1.0	23Mo07	
^{99}Cd−^{85}Rb$_{1.165}$	27690.8	1.7	27690.3	1.6	−0.3	1	93	93 ^{99}Cd	MA8	1.0	09Br09	
^{99}In−u	−65946	83	−65939	7	0.1	o			MR1	1.0	21Mo23	
	−65948.9	15.5			0.7	1	23	23 ^{99}In	MR1	1.0	23Ni07	
^{99}Pd−^{96}Mo$_{1.031}$	10052.8	5.5	10055	5	0.4	1	81	81 ^{99}Pd	JY1	1.0	09El08	
^{99}Sr−^{97}Zr$_{1.021}$	23794.1	7.4	23790	5	−0.6	1	47	47 ^{99}Sr	JY1	1.0	06Ha03	
^{99}Y−^{97}Zr$_{1.021}$	15066.8	7.1				2			JY1	1.0	07Ha32	
^{99}Zr−^{97}Zr$_{1.021}$	7580	14	7581	11	0.1	1	65	65 ^{99}Zr	JY1	1.0	06Ha03	
^{99}Rh−^{99}Cd	−16728	25	−16773	16	−1.8	1	41	41 ^{99}Rh	RI1	1.0	26Ho.A	
^{99}Pd−^{99}Cd	−13151	12	−13150	5	0.0	1	18	17 ^{99}Pd	RI1	1.0	26Ho.A	
^{99}Ag−^{99}Cd	−7274.9	4.2	−7276	4	−0.3	1	73	69 ^{99}Ag	RI1	1.0	26Ho.A	
^{99}In−^{99}Cd	9139.0	8.2	9136	7	−0.4	1	78	77 ^{99}In	RI1	1.0	26Ho.A	
^{99}Ru−^{98}Ru	652	11	648.9	0.4	−0.1	U			M16	2.5	63Da10	
^{97}Rb−^{99}Rb$_{.653}$ ^{93}Rb$_{.348}$	100	100	135	4	0.1	U			P21	2.5	82Au01	
^{98}Rb−^{99}Rb$_{.742}$ ^{95}Rb$_{.258}$	690	180	563	17	−0.3	U			P21	2.5	82Au01	
^{97}Rb−^{99}Rb$_{.490}$ ^{95}Rb$_{.511}$	350	60	202	11	−1.0	U			P31	2.5	86Au02	
^{99}Ru(n,α)^{96}Mo	6822	5	6816.1	0.3	−1.2	U					01Wa50	
^{96}Ru(^{16}O,^{13}C)^{99}Pd	−11723	20	−11761	5	−1.9	U			BNL		82Th01	
^{98}Mo(n,γ)^{99}Mo	5927.7	1.	5925.48	0.15	−2.2	U					73De39	
	5927	1			−1.5	U					74Er.A	
	5924.6	0.6			1.5	U					76Ch02	
	5923	2			1.2	U					77Ri04	
	5925.42	0.15			0.4	1	98	96 ^{99}Mo	MMn		91Is02	Z
	5927.7	0.5			−4.4	C			Bdn		06Fi.A	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{98}Mo(d,p)^{99}Mo		3687	20	3700.91	0.15	0.7	U			Pit		64Co11	
^{98}Mo(^{3}He,d)^{99}Tc		1010	20	1008.8	0.4	−0.1	U			McM		77Ch06	
^{99}Tc(p,d)^{98}Tc		−6740	5	−6743	3	−0.6	–					76Sl06	
		−6755	9			1.3	–			Bld		77Em02	
	ave.	−6744	4			0.1	1	60	59 ^{98}Tc			average	
^{99}Rb(β^-)^{99}Sr		11340	120	11397	6	0.5	U			McG		84Ia.A	
		10960	130			3.4	C			Bwg		87Gr.A	
^{99}Sr(β^-)^{99}Y		8030	80	8125	8	1.2	U			McG		84Ia.A	
		8360	75			−3.1	C			Bwg		87Gr.A	
^{99}Y(β^-)^{99}Zr		7605	60	6973	12	−10.5	C			Bwg		87Gr.A	
		7568	14			−42.5	C			Bwg		92Gr.A	
^{99}Zr(β^-)^{99}Nb		4550	35	4719	16	4.8	C			Bwg		87Gr.A	
		4559	15			10.6	C			Bwg		92Gr.A	
^{99}Nb(β^-)^{99}Mo		3740	200	3635	12	−0.5	U					70Ei02	*
^{99}Mo(β^-)^{99}Tc		1356.7	1.0	1359.1	0.4	2.4	1	16	12 ^{99}Tc			71Na01	*
^{99}Tc(β^-)^{99}Ru		292	3	295.88	0.16	1.3	U					51Ta05	
		290	4			1.5	U					52Fe16	
		293.5	2.0			1.2	U					80Al02	*
		295.82	0.16			0.4	1	98	88 ^{99}Tc			24Pa41	
^{99}Rh(β^+)^{99}Ru		2038	10	2070	15	3.2	C					52Sc11	*
		2053	10			1.7	U					59To25	*
		2170	30			−3.3	U					74An23	*
		2111	58			−0.7	1	7	7 ^{99}Rh			averag	*
^{99}Pd(β^+)^{99}Rh		3410	20	3374	15	−1.8	1	55	53 ^{99}Rh			69Ph01	*
^{99}Ag(β^+)^{99}Pd		5430	150	5472	6	0.3	U					81Hu03	
^{99}In^m(IT)^{99}In		671	37				2			MR1		23Ni07	*
^{99}Sn(ε)^{99}In		14700	3600	13400#	500#	−0.4	D					18Pa20	*
*^{99}Sr−^{85}Rb$_{1.165}$	Original D_M=35661.6(4.4) μu $M-A$=–62506.3(4.1)keV re-evaluated											16Kl04	**
*^{99}Nb(β^-)^{99}Mo	E_{β^-}=3500(200) to 7/2$^+$ level at 235.508 keV											Ens112	**
*^{99}Mo(β^-)^{99}Tc	E_{β^-}=1214(1) to 1/2$^-$ level at 142.6832 keV											Ens112	**
*^{99}Tc(β^-)^{99}Ru	E_{β^+}=434.8(2.6), 346.7(2.0) from ^{99}Tc^m at 142.6832 to ground state, 89.57 level											Ens112	**
*^{99}Rh(β^+)^{99}Ru	E_{β^+}=740(10) from ^{99}Rh^m at 64.5 to 340.90 7/2$^+$ level											Ens112	**
*^{99}Rh(β^+)^{99}Ru	E_{β^+}=1030(10), 710(10), 590(10), 420(20) keV											59To25	**
*	to ground state, 321.99 3/2$^+$, 442.59 3/2$^+$, 618.13 1/2$^+$ levels											Ens112	**
*^{99}Rh(β^+)^{99}Ru	Q_{β^+} larger than 2059.34 keV because that state populated via β decay											Ens112	**
*^{99}Rh(β^+)^{99}Ru	Ref. proposed that E_{β^+}=1030(10) keV decays to 89.57 3/2$^+$ level											74An23	**
*^{99}Rh(β^+)^{99}Ru	E_{β^+}=1100(50), 680(30) and 540(30) to 89.57, 442.59 and 618.13 levels											Ens112	**
*^{99}Rh(β^+)^{99}Ru	unweighted average 59To25=2053(10) 74An23=2170(30) keV											WgM20b	**
*^{99}Pd(β^+)^{99}Rh	E_{β^+}=2180(20), 1930(20), 1510(20) keV											69Ph01	**
*	to 200.7 7/2$^+$, 464.4 (5/2,7/2)$^+$, 874.5 5/2$^+$ levels above (1/2$^-$) ground state											Ens112	**
*^{99}In^m(IT)^{99}In	from 23Ni07=671(33,stat)(16,syst)											23Ni07	**
*^{99}Sn(ε)^{99}In	Trends in Mass Surface TMS suggest ^{99}Sn 1300 keV more bound											HWJ202	**
^{100}Rb−u		−49685	19	−49664	13	1.1	–			MA8	1.0	17De18	*
		−49643	32			−0.7	–			TR1	1.0	21Mu05	
	ave.	−49674	16			0.6	1	62	62 ^{100}Rb			average	
^{100}Rb−^{85}Rb$_{1.176}$		54087	21	54071	13	−0.8	1	38	38 ^{100}Rb	MA8	1.0	13Ma81	
		54150	140			−0.6	U			TT1	1.0	16Kl04	
^{100}Sr−^{85}Rb$_{1.176}$		39520	12	39519	7	−0.1	–			TT1	1.0	16Kl04	
		39515	29			0.1	–			MA8	1.0	17De18	
	ave.	39519	11			−0.1	1	45	45 ^{100}Sr			average	
^{100}Sr−u		−64223	31	−64217	7	0.2	U			TR1	1.0	21Mu05	
^{100}Y−u		−72270	110	−72274	9	−0.0	o			GT2	2.5	08Kn.A	*
		−72290	140			0.0	U			GT2	2.5	08Su19	*
^{100}Zr−u		−82016	18	−81990	9	1.5	1	24	24 ^{100}Zr	JY0	1.0	04Ri12	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{100}Mo−^{85}Rb$_{1.176}$		11216	27	11203.4	0.3	−0.5	U			MA8	1.0	11He10	
		11205.5	1.4			−1.5	U			JY1	1.0	12Ka13	
^{100}Mo−^{87}Rb$_{1.149}$		11820.25	0.57	11819.7	0.3	−1.0	1	30	30 ^{100}Mo	MS1	1.0	15Gu09	
C_7 H_{16}−^{100}Mo		217730.3	4.2	217732.4	0.3	0.2	U			M15	2.5	63Ri07	
^{100}Mo−u		−92532.51	0.40	−92531.9	0.3	1.6	1	61	61 ^{100}Mo	MS1	1.0	15Gu09	
C_7 H_{16}−^{100}Ru		220983.8	3.7	220989.9	0.4	0.7	U			M16	2.5	63Da10	
^{100}Rh−u		−91855	46	−91886	19	−0.7	1	18	18 ^{100}Rh	GS2	1.0	05Li24	*
^{100}Pd−u		−91468.1	64.4	−91483	14	−0.2	U			GR1	1.0	23Mo07	
^{100}Ag−^{85}Rb$_{1.176}$		19849.9	7.1	19850	5	0.0	–			JY1	1.0	09El08	*
		19851.8	8.2			−0.2	–			MA8	1.0	11He10	*
	ave.	19851	5			−0.1	1	87	87 ^{100}Ag			average	
^{100}Ag−u		−83902	44	−83885	5	0.4	o			GR1	1.0	19An10	*
		−83847	34			−1.1	U			GR1	1.0	23Mo07	*
^{100}Cd−u		−79636	214	−79651.3	1.8	−0.1	U			CS1	1.0	96Ch32	
		−79650.5	12.9			−0.1	U			GR1	1.0	23Mo07	
^{100}Cd−^{85}Rb$_{1.176}$		24084.1	1.8	24084.0	1.8	−0.1	1	96	96 ^{100}Cd	MA8	1.0	09Br09	
^{100}In−u		−69405	322	−68897.8	2.3	1.6	U			CS1	1.0	96Ch32	
^{100}In−^{85}Rb$_{1.176}$		34837.2	2.4	34837.5	2.3	0.1	1	96	96 ^{100}In	MA8	1.0	21Mo23	
^{100}In−u		−68906	22	−68897.8	2.3	0.4	U			MR1	1.0	21Mo23	
		−68911.9	12.9			1.1	U			MR1	1.0	23Ni07	
^{100}Sn−u		−62020	1020	−61350	260	0.7	U			CS1	1.0	96Ch32	
^{100}Sr−^{97}Zr$_{1.031}$		27579	10	27580	7	0.1	1	55	55 ^{100}Sr	JY1	1.0	06Ha03	
^{100}Y−^{97}Zr$_{1.031}$		19524	12	19522	9	−0.1	1	50	50 ^{100}Y	JY1	1.0	07Ha32	
^{100}Y^m−^{97}Zr$_{1.031}$		19679	12	19681	9	0.1	1	50	50 ^{100}Y^m	JY1	1.0	07Ha32	
^{100}Zr−^{97}Zr$_{1.031}$		9815	10	9807	9	−0.8	1	76	76 ^{100}Zr	JY1	1.0	06Ha03	
^{100}Nb^m−^{97}Zr$_{1.031}$		6472.6	2.1				2			JY1	1.0	07Ha32	
^{100}Mo ^{35}Cl−^{98}Mo ^{37}Cl		5019	2	5014.7	0.4	−0.5	U			H11	4.0	63Bi12	
^{100}Y^m−^{100}Y		158.6	1.4	158.6	1.4	−0.0	1	99	50 ^{100}Y	JY2	1.0	22We.A	*
^{100}Nb−^{100}Nb^m		−335.7	8.3				3			JY1	1.0	07Ha32	
^{100}Mo−^{100}Ru		3257.55	0.18	3257.48	0.18	−0.4	1	97	88 ^{100}Ru	JY1	1.0	08Ra09	
^{100}Pd−^{100}Cd		−11835	20	−11831	14	0.2	1	47	47 ^{100}Pd	RI1	1.0	26Ro.A	*
^{100}Ag−^{100}Cd		−4237	14	−4234	5	0.2	1	14	13 ^{100}Ag	RI1	1.0	26Ro.A	*
^{100}In−^{100}Cd		10760	11	10753.5	2.9	−0.6	1	7	4 ^{100}In	RI1	1.0	26Ro.A	*
^{100}Ru−^{99}Ru		−1718	11	−1719.826	0.028	−0.1	U			M16	2.5	63Da10	
^{96}Ru(^{16}O,^{12}C)^{100}Pd		−5599	26	−5602	13	−0.1	1	24	24 ^{100}Pd	BNL		82Th01	
^{100}Mo(d,^{3}He)^{99}Nb		−5639	15	−5653	12	−0.9	2			Tex		74Bi08	
^{100}Mo(t,α)^{99}Nb		8642	20	8667	12	1.3	2			LAl		83Fl06	
^{100}Mo(d,t)^{99}Mo		−2038	20	−2036.8	0.4	0.1	U			Pit		64Co11	
^{99}Tc(n,γ)^{100}Tc		6764.4	1.				2					79Pi08	
		6765.20	0.04	6764.4	1.0	−20.0	C					04Fu.A	
^{99}Ru(n,γ)^{100}Ru		9673.2	0.7	9673.325	0.026	0.2	U					74Ri03	
		9672.65	0.06			11.3	B			ILn		88Co18	Z
		9673.39	0.05			−1.3	–			MMn		91Is02	Z
		9673.30	0.03			0.8	–			ILn		00Ge01	
		9673.41	0.19			−0.4	U			Bdn		06Fi.A	
	ave.	9673.324	0.026			0.1	1	100	90 ^{99}Ru			average	
^{100}Sr(β^-)^{100}Y		7460	140	7505	11	0.3	U			McG		84Ia.A	*
		7075	100			4.3	C			Bwg		87Gr.A	
^{100}Y(β^-)^{100}Zr		7920	100	9050	11	11.3	C			McG		84Ia.A	*
		9310	70			−3.7	C			Bwg		87Gr.A	
^{100}Zr(β^-)^{100}Nb		3335	25	3419	11	3.3	C			Bwg		87Gr.A	
^{100}Nb(β^-)^{100}Mo		6245	25	6402	8	6.3	C			Bwg		87Gr.A	
^{100}Nb^m(β^-)^{100}Mo		6745	75	6714.3	2.0	−0.4	U			Bwg		87Gr.A	
^{100}Mo(t,^{3}He)^{100}Nb^m		−6690	30	−6695.8	2.0	−0.2	U			LAl		79Aj03	
^{100}Rh(β^+)^{100}Ru		3630	20	3636	18	0.3	1	82	82 ^{100}Rh			53Ma64	
^{100}Ag(β^+)^{100}Pd		7075	90	7077	14	0.0	U					79Ve.A	*
		7022	200			0.3	U					80Ha20	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{100}Cd(β^+)^{100}Ag	3890	70	3944	5	0.8	U					89Ry02	
^{100}In(β^+)^{100}Cd	10900	930	10016.8	2.7	−0.9	U			Lvp		95Sz01	*
	10080	230			−0.3	U					02Pl03	
^{100}Sn(β^+)^{100}In	7390	660	7030	240	−0.5	o			GSI		97Su06	*
	7840	660			−1.2	o			GSI		02Fa13	*
	7030	240				2			GSI		12Hi07	*
	7690	160			−4.1	B					19Lu08	
*^{100}Rb−u	MR−TOF data in the paper recalculated by evaluators										HWJ187	**
*^{100}Y−u	$M-A$=–67245(93) keV for mixture gs+m at 147.7(1.3) keV										Nub269	**
*^{100}Y−u	$M-A$=–67264(119) keV for mixture gs+m at 147.7(1.3) keV										Nub269	**
*^{100}Rh−u	$M-A$=–85508(29) keV for mixture gs+m at 107.6 keV										Nub269	**
*^{100}Ag−^{85}Rb$_{1.176}$	D_M=19858.2(5.2) μu for mixture gs+m at 15.52 keV; $M-A$=–78131.0(4.9) keV										Nub269	**
*^{100}Ag−^{85}Rb$_{1.176}$	D_M=19860.2(6.6) μu for mixture gs+m at 15.52 keV; $M-A$=–78129.1(6.2) keV										Nub269	**
*^{100}Ag−u	$M-A$=–78146(41) keV for mixture gs+m at 15.52 keV										Nub269	**
*^{100}Ag−u	$M-A$=–78095(31) keV for mixture gs+m at 15.52 keV										Nub269	**
*^{100}Y^m−^{100}Y	quadrature 1.45e−8 (stat) and 4.23e−8 (syst) unc.										22We.A	**
*^{100}Pd−^{100}Cd	from frequency ratio ^{100}Pd^{++}/(^{100}Cd)$^{++}$ =0.999881558(201)										26Ro.A	**
*^{100}Pd−^{100}Cd	Original error 19; added systematic error 3 keV										SNa265	**
*^{100}Ag−^{100}Cd	from frequency ratio ^{100}Ag^{++}/(^{100}Cd)$^{++}$ =0.999957688(126)										26Ro.A	**
*^{100}Ag−^{100}Cd	D_M=–4228(13) μu for mixture gs+m at 15.52 keV; $M-A$=–78133(12)										SNa265	**
*^{100}In−^{100}Cd	from frequency ratio ^{100}In^{++}/(^{100}Cd)$^{++}$ =1.0001076893(76)										26Ro.A	**
*^{100}In−^{100}Cd	Original error 1.8; added systematic error 10 keV										SNa265	**
*^{100}Sr(β^-)^{100}Y	E_{β^-}=7450(140) assumed by evaluator to 1$^+$ level at 10.70 keV										Ens082	**
*^{100}Y(β^-)^{100}Zr	Not unambiguously ground state transition										GAu	**
*^{100}Ag(β^+)^{100}Pd	From 5$^+$ ground state to high spin level at 2920.4 keV										79Ve.A	**
*^{100}Ag(β^+)^{100}Pd	E_{β^+}=5350(200) from ^{100}Ag^m 2$^+$ at 15.52 to 2$^+$ level at 665.50 keV										Ens082	**
*^{100}In(β^+)^{100}Cd	From lower and upper limits 9300–12500										GAu	**
*^{100}Sn(β^+)^{100}In	Q_{β^+}=7200(+800–500); also E_{β^+}=3400(+700–300) keV to 2760(430) level										97Su06	**
*^{100}Sn(β^+)^{100}In	E_{β^+}=3800(+700–300) to 2760(430) level										96Ki23	**
*^{100}Sn(β^+)^{100}In	E_{β^+}=3290(200) to 1$^+$ level at 2721+x, with x<80 keV										12Hi07	**
^{101}Rb−u		−45698	22	−45667	18	1.4	2			IS1	1.0	17De18	
		−45604	31			−2.0	2			TR1	1.0	21Mu05	
^{101}Sr−^{85}Rb$_{1.188}$		45410	22	45401	9	−0.4	–			MA8	1.0	17De18	
		45398	10			0.3	–			TT1	1.0	16Kl04	
	ave.	45400	9			0.1	1	92	92 ^{101}Sr			average	
^{101}Sr−u		−59379	31	−59393	9	−0.4	1	8	8 ^{101}Sr	TR1	1.0	21Mu05	
^{101}Zr−u		−78573	20	−78542	9	1.6	1	20	20 ^{101}Zr	JY0	1.0	04Ri12	
C_8 H_5−^{101}Ru		133549.5	2.2	133551.9	0.4	0.4	U			M16	2.5	63Da10	
^{101}Rh−u		−93821	58	−93841	6	−0.3	U			GS2	1.0	05Li24	*
^{101}Pd−u		−91816	30	−91715	5	3.4	C			GS2	1.0	05Li24	
^{101}Ag−^{85}Rb$_{1.188}$		17470.5	7.2	17478	5	1.0	2			SH1	1.0	07Ma92	
		17485.6	7.5			−1.0	2			MA8	1.0	11He10	
^{101}Cd−^{85}Rb$_{1.188}$		23367	11	23380.0	1.6	1.2	U			SH1	1.0	07Ma92	
		23380.0	1.6				2			MA8	1.0	09Br09	
^{101}In−u		−73591.4	15.4	−73588	4	0.2	U			LZ1	1.0	19Xu13	
		−73575.4	21.5			−0.6	U			GR1	1.0	20Ho03	
		−73594.2	10.4			0.6	1	13	13 ^{101}In	MR1	1.0	23Ni07	
		−73601	27			0.5	U			GR1	1.0	23Mo07	
^{101}In−^{85}Rb$_{1.188}$		31207.3	5.0	31206	4	−0.3	1	55	55 ^{101}In	MA8	1.0	21Mo23	
^{101}In−^{82}Sr ^{19}F		9613.6	4.4	9613	4	−0.2	1	72	50 ^{82}Sr	IS1	1.0	21Mo23	
^{101}In^m−u		−72884.0	52.0	−72871	7	0.2	U			LZ1	1.0	19Xu13	
		−72922.6	57.3			0.9	U			GR1	1.0	20Ho03	*
^{101}In^m−^{82}Sr ^{19}F		10330.7	6.7	10329	6	−0.2	1	87	84 ^{101}In^m	IS1	1.0	21Mo23	
^{101}Sn−^{50}Ti$_{2.020}$		47238.6	1.0	47238.7	1.0	0.1	1	99	99 ^{101}Sn	MS2	1.0	26Ir.1	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{101}Pd−^{96}Mo$_{1.052}$		8567.4	5.1	8567	5	−0.1	1	93	93 ^{101}Pd	JY1	1.0	09El08	
^{101}Cd−^{96}Mo$_{1.052}$		18872.7	5.5	18868.4	1.6	−0.8	U			JY1	1.0	09El08	
^{101}Y−^{97}Zr$_{1.041}$		22847.5	7.6				2			JY1	1.0	07Ha32	
^{101}Zr−^{97}Zr$_{1.041}$		14153	10	14145	9	−0.8	1	80	80 ^{101}Zr	JY1	1.0	06Ha03	
^{101}Nb−^{102}Ru$_{.990}$		10009.6	4.0				2			JY1	1.0	07Ha32	
^{101}Sn−^{101}In		9305	12	9293	4	−1.0	1	10	9 ^{101}In	RI1	1.0	26Ro.A	*
^{101}Ru−^{100}Ru		1368	11	1362.63	0.25	−0.2	U			M16	2.5	63Da10	
^{100}Mo(n,γ)^{101}Mo		5398.4	0.5	5398.24	0.07	−0.3	U			ILn		75Ka.A	
		5399.6	0.7			−1.9	U			ORn		79We07	Z
		5398.23	0.08			0.1	2			ILn		90Se17	Z
		5398.27	0.13			−0.2	2			Bdn		06Fi.A	
^{100}Mo(d,p)^{101}Mo		3161	6	3173.67	0.07	2.1	U					72Si25	
^{100}Ru(n,γ)^{101}Ru		6802.0	0.7	6802.04	0.23	0.1	–					82Ba69	
		6802.04	0.25			−0.0	–			Bdn		06Fi.A	
^{100}Ru(d,p)^{101}Ru		4581	4	4577.47	0.23	−0.9	U					77Ho02	
^{100}Ru(n,γ)^{101}Ru	ave.	6802.04	0.24	6802.04	0.23	0.0	1	99	98 ^{101}Ru			average	
^{101}Sn(εp)^{100}Cd		6600	300	7015.9	1.9	1.4	U					10St.A	
^{101}Rb(β^-)^{101}Sr		11810	110	12786	19	8.9	B			Bwg		92Ba28	
^{101}Sr(β^-)^{101}Y		9505	80	9731	11	2.8	B			Bwg		92Ba28	
^{101}Y(β^-)^{101}Zr		8545	90	8106	11	−4.9	B			Bwg		92Ba28	
^{101}Zr(β^-)^{101}Nb		5520	45	5730	9	4.7	B			Bwg		87Gr18	
		5485	25			9.8	C			Bwg		92Gr.A	
^{101}Nb(β^-)^{101}Mo		4575	30	4629	4	1.8	U			Bwg		87Gr.A	
		4590	30			1.3	U			Bwg		87Gr18	
		4569	18			3.3	C			Bwg		92Gr.A	
^{101}Mo(β^-)^{101}Tc		2836	40	2825	24	−0.3	R					57Ok.A	*
^{101}Tc(β^-)^{101}Ru		1620	30	1614	24	−0.2	2					71Ar23	*
^{101}Pd(β^+)^{101}Rh		1980	4	1980	4	0.1	1	95	88 ^{101}Rh			71Ib01	*
^{101}Ag(β^+)^{101}Pd		4100	200	4098	7	−0.0	U					72We.A	
		4350	200			−1.3	U					78Ha11	
		4180	150			−0.5	U					79Ve.A	*
^{101}Cd(β^+)^{101}Ag		5530	130	5498	5	−0.2	U					70Be.A	*
		5350	200			0.7	U					72We.A	
^{101}In^m(IT)^{101}In		658	16	667	7	0.6	1	17	16 ^{101}In^m	MR1		23Ni07	
*^{101}Rh−u	$M-A$=–87315(29) keV for mixture gs+m at 157.32 keV											Nub269	**
*^{101}In^m−u	reconstructed from E_x =608(57) keV											20Ho03	**
*^{101}Sn−^{101}In	from frequency ratio ^{101}Sn^{++}/(^{101}In)$^{++}$ =1.000092196(107)											26Ro.A	**
*^{101}Sn−^{101}In	Original error 11; added systematic error 3 keV											SNa265	**
*^{101}Mo(β^-)^{101}Tc	E_{β^-}=2230(40) to (1/2$^+$,3/2$^+$) level at 606.47 keV											Ens06a	**
*^{101}Tc(β^-)^{101}Ru	E_{β^-}=1320(30) to 306.858 7/2$^+$ and 1070(30) to 545.115 7/2$^+$ levels											Ens06a	**
*^{101}Pd(β^+)^{101}Rh	E_{β^+}=776(4) to 7/2$^+$ level at 181.78 keV											Ens06a	**
*^{101}Ag(β^+)^{101}Pd	E_{β^+}=2895(150) to 7/2$^+$ level at 261.0 keV, and others											Ens06a	**
*^{101}Cd(β^+)^{101}Ag	Measured E_{β^+} may go to excited state											70Be.A	**
^{102}Rb−u		−39992	89	−39981	29	0.1	2			MA8	1.0	17De18	
		−39980	31			−0.0	2			TR1	1.0	21Mu05	
^{102}Sr−^{85}Rb$_{1.200}$		49857	72	49843	28	−0.2	1	16	16 ^{102}Sr	MA8	1.0	17De18	
^{102}Sr−u		−56012	31	−56009	28	0.1	1	84	84 ^{102}Sr	TR1	1.0	21Mu05	
^{102}Y−^{120}Sn$_{.850}$		17456.3	4.3				2			JY1	1.0	11Ha48	*
^{102}Zr−u		−76780	43	−76846	9	−1.5	U			JY0	1.0	04Ri12	
C_8 H_6−^{102}Ru		142604.8	3.2	142609.7	0.4	0.6	U			M16	2.5	63Da10	
C_8 H_6−^{102}Pd		141324	19	141317.7	0.4	−0.1	U			M16	2.5	63Da10	
		141346	18			−1.0	U			R12	1.5	83De51	
C_7 N H_4−^{102}Pd		128775	19	128741.6	0.4	−1.2	U			R12	1.5	83De51	
^{102}Pd−u		−94370.9	15.6	−94367.5	0.4	0.2	U			GS4	1.0	14Ya.A	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{102}Ag−u		−88315	30	−88295	9	0.7	U				GS2	1.0	05Li24	*
^{102}Cd−^{85}Rb$_{1.200}$		20320.9	7.3	20334.1	1.8	1.8	U				SH1	1.0	07Ma92	
		20334.2	1.9			−0.0	1	88	88	^{102}Cd	MA8	1.0	09Br09	
^{102}In−^{85}Rb$_{1.200}$		29959	13	29958	5	−0.1	1	14	14	^{102}In	SH1	1.0	07Ma92	
^{102}Cd−^{96}Mo$_{1.063}$		15811.9	5.2	15812.5	1.8	0.1	1	12	12	^{102}Cd	JY1	1.0	09El08	
^{102}In−^{96}Mo$_{1.063}$		25436.5	5.3	25437	5	0.0	1	86	86	^{102}In	JY1	1.0	09El08	
^{102}Zr−^{97}Zr$_{1.052}$		16822.0	9.8	16820	9	−0.2	1	92	92	^{102}Zr	JY1	1.0	06Ha03	
^{102}Nb−^{97}Zr$_{1.052}$		11756.4	2.7	11756.5	2.7	0.0	1	99	99	^{102}Nb	JY1	1.0	07Ha32	
^{102}Mo−^{97}Zr$_{1.052}$		3961.0	9.8	3960	9	−0.1	1	83	83	^{102}Mo	JY1	1.0	06Ha03	
^{102}Y^{m}−^{102}Y		13.2	1.7				3				JY2	1.0	22We.A	*
^{102}Nb^{m}−^{102}Nb		100.2	7.9	101	8	0.1	1	95	94	^{102}Nb^{m}	JY1	1.0	07Ha32	
^{102}Pd−^{102}Ru		1291.76	0.39	1291.99	0.04	0.6	U				SH1	1.0	11Go23	
		1291.980	0.048			0.2	–				JY1	1.0	19Ne08	
		1292.04	0.10			−0.5	–				JY2	1.0	25Ru01	
	ave.	1291.99	0.04			−0.0	1	100	100	^{102}Pd			average	
^{102}Sn−^{102}In		6307	17				2				RI1	1.0	26Ro.A	*
^{102}Ru−^{101}Ru		−1233	11	−1232.77	0.05	0.0	U				M16	2.5	63Da10	
^{100}Mo(t,p)^{102}Mo		5034	20	5029	8	−0.3	1	17	17	^{102}Mo	LAl		72Ca10	
^{100}Mo(^{3}He,p)^{102}Tc		6054	20	6023	9	−1.6	1	20	20	^{102}Tc	Pri		82De03	
^{102}Pd(p,t)^{100}Pd		−10356	24	−10348	13	0.3	1	29	29	^{100}Pd	Win		74De31	*
^{101}Ru(n,γ)^{102}Ru		9220.4	0.9	9219.64	0.05	−0.8	U						74Ri03	
		9219.64	0.05			0.0	1	100	97	^{102}Ru	MMn		91Is02	Z
		9219.63	0.19			0.1	U				Bdn		06Fi.A	
^{102}In(εp)^{101}Ag		3420	310	3351	7	−0.2	o				Lvp		91Re.A	*
^{102}Sr(β^-)^{102}Y		8815	70	9000	27	2.6	B				Bwg		92Ba28	
^{102}Y(β^-)^{102}Zr		9850	70	10409	10	8.0	B				Bwg		92Ba28	
^{102}Zr(β^-)^{102}Nb^{m}		4605	30	4622	11	0.6	1	14	8	^{102}Zr	Bwg		87Gr18	
^{102}Nb(β^-)^{102}Mo		7300	50	7263	9	−0.7	o				Bwg		87Gr.A	
		7335	40			−1.8	U				Bwg		87Gr18	
^{102}Nb^{m}(β^-)^{102}Mo		7215	40	7357	11	3.5	C				Bwg		87Gr.A	
		7210	35			4.2	B				Bwg		87Gr18	
^{102}Tc(β^-)^{102}Ru		4420	100	4533	9	1.1	U						69Bl16	
^{102}Rh(β^+)^{102}Ru		2317	10	2323	6	0.6	2						61Hi06	
		2325	10			−0.2	2						63Bo17	
^{102}Ru(p,n)^{102}Rh		−3115	15	−3105	6	0.6	2						83Do11	
^{102}Rh(β^-)^{102}Pd		1150	6	1120	6	−5.1	B						61Hi06	
^{102}Ag(β^+)^{102}Pd		5800	200	5656	8	−0.7	U						67Ch05	
		5966	100			−3.1	B						67Ch05	*
		4910	140			5.3	C						70Be.A	*
		5350	200			1.5	U						72We.A	
		5880	110			−2.0	U						79Ve.A	
^{102}Cd(β^+)^{102}Ag		2554	57	2587	8	0.6	U						72We.A	
		2587	8				2				GSI		91Ke08	*
^{102}In(β^+)^{102}Cd		9250	380	8965	5	−0.8	U				Lvp		95Sz01	*
		8970	150			−0.0	U				GSI		98Ka.A	
		8910	170			0.3	U				GSI		03Gi06	*
^{102}Sn(β^+)^{102}In		5780	70	5875	16	1.4	o				GSI		01St.A	
		5760	100			1.1	U				GSI		02Fa13	

*^{102}Y−^{120}Sn$_{.850}$	Associated with low-spin isomer	11Ha48 **
*^{102}Ag−u	$M-A$=–82260(28) keV for mixture gs+m at 9.40 keV	Nub269 **
*^{102}Y^{m}−^{102}Y	include in quadrature 1.64e−8 (stat) and 4.23e−8 (syst) unc.	22We.A **
*^{102}Sn−^{102}In	from frequency ratio ^{102}Sn^{++}/(^{102}In)$^{++}$ =1.000061876(164)	26Ro.A **
*^{102}Sn−^{102}In	Original error 16; added systematic error 3 keV	SNa265 **
*^{102}Pd(p,t)^{100}Pd	Original error 12; added systematic error 21 keV	GAu **
*^{102}In(εp)^{101}Ag	Estimated using proton spectrum from 1450 to 3200 keV	GAu **
*^{102}Ag(β^+)^{102}Pd	E_{β^+}=3340(100), 3070(130) from ^{102}Ag^{m} at 9.40 to 1534.48, 2017.8 levels	Ens098 **
*^{102}Ag(β^+)^{102}Pd	Q_{β^+}=4920(100) from ^{102}Ag^{m} at 9.40 keV	Nub269 **
*^{102}Cd(β^+)^{102}Ag	E_{β^+}=1075(8) to 1$^+$ level at 490.44 keV	Ens098 **
*^{102}In(β^+)^{102}Cd	From 9900 keV upper and 8600 keV lower limits	GAu **
*^{102}In(β^+)^{102}Cd	Good agreement with authors' earlier measurement, average=8950(120) keV	03Gi06 **

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{103}Rb−u		−35480	34				2			TR1	1.0	21Mu05	
^{103}Sr−u		−50693	31				2			TR1	1.0	21Mu05	
^{103}Y−u		−63060	183	−62756	12	1.1	o			GT1	1.5	04Ma.A	
		−62803	106			0.2	U			GT2	2.5	08Kn.A	
^{103}Y−^{120}Sn$_{.858}$		21154	12				2			JY1	1.0	11Ha48	
^{103}Zr−u		−72765	64	−72796	10	−0.5	U			JY0	1.0	04Ri12	
C_8 H_7−^{103}Rh		149263.5	3.3	149281.0	2.5	2.1	U			M16	2.5	63Da10	
		149261	19			0.7	U			R12	1.5	83De51	
C_7 N H_5−^{103}Rh		136681	18	136704.9	2.5	0.9	U			R12	1.5	83De51	
^{103}Pd−u		−93894.8	14.5	−93888.7	0.9	0.4	U			GS4	1.0	14Ya.A	
^{103}Ag−u		−91091	52	−91039	4	1.0	U			GS2	1.0	05Li24	*
^{103}Ag−^{85}Rb$_{1.212}$		15875	14	15871	4	−0.3	U			SH1	1.0	07Ma92	
		15871.4	4.4				2			MA8	1.0	11He10	
^{103}Cd−^{85}Rb$_{1.212}$		20328	11	20327.6	1.9	−0.0	U			SH1	1.0	07Ma92	
		20328.2	2.1			−0.3	1	83	83 ^{103}Cd	MA8	1.0	09Br09	
^{103}In−^{85}Rb$_{1.212}$		26785	11	26790	10	0.4	1	77	77 ^{103}In	SH1	1.0	07Ma92	
^{103}In−u		−80119.7	26.8	−80121	10	−0.1	1	13	13 ^{103}In	GR1	1.0	20Ho03	
^{103}Sn−u		−72075	73	−72062.4	2.9	0.2	U			LZ1	1.0	23Xi01	
		−72062.6	3.0			0.1	1	91	91 ^{103}Sn	MS1	1.0	25Ir01	*
		−72039.1	19.3			−1.2	U			IS1	1.0	25Ni02	
^{103}Cd−^{96}Mo$_{1.073}$		15699.2	5.2	15700.7	1.9	0.3	1	14	13 ^{103}Cd	JY1	1.0	09El08	
^{103}Zr−^{97}Zr$_{1.062}$		21760.5	9.9				2			JY1	1.0	06Ha03	
^{103}Mo−^{97}Zr$_{1.062}$		7648.4	9.9				2			JY1	1.0	06Ha03	
^{103}Nb−^{102}Ru$_{1.010}$		16069.7	4.2				2			JY1	1.0	07Ha32	
^{103}Sn−^{103}Cd		14526	10	14521	3	−0.5	1	11	8 ^{103}Sn	RI1	1.0	26Ro.A	*
^{103}Cd−^{102}Cd		−1534	154	−1065.1	2.6	2.0	U			CR2	1.5	92Sh.A	*
^{103}Rh(p,t)^{101}Rh		−8275	17	−8280	6	−0.3	1	13	12 ^{101}Rh	Pri		64Th05	
^{102}Ru(n,γ)^{103}Ru		6232.2	0.3	6232.05	0.15	−0.5	–					82Ba69	Z
		6232.00	0.17			0.3	–			Bdn		06Fi.A	
^{102}Ru(d,p)^{103}Ru		4005	15	4007.49	0.15	0.2	U			ANL		71Fo01	
^{102}Ru(n,γ)^{103}Ru	ave.	6232.05	0.15	6232.05	0.15	0.0	1	100	98 ^{103}Ru			average	
^{103}Rh(γ,n)^{102}Rh		−9307	32	−9320	7	−0.4	U			Phi		60Ge01	
^{103}Rh(p,d)^{102}Rh		−7144	16	−7095	7	3.1	B			Pri		64Th05	
^{102}Pd(n,γ)^{103}Pd		7624.6	1.5	7625.3	0.8	0.5	2					70Bo29	
		7625.6	0.9			−0.3	2			Bdn		06Fi.A	
^{103}Zr(β^-)^{103}Nb		6945	85	7219	10	3.2	B			Bwg		87Gr18	
^{103}Nb(β^-)^{103}Mo		5535	35	5926	10	11.2	C			Bwg		87Gr.A	
		5530	30			13.2	B			Bwg		87Gr18	
^{103}Mo(β^-)^{103}Tc		3750	60	3650	13	−1.7	U			Bwg		87Gr18	
^{103}Tc(β^-)^{103}Ru		2615	45	2662	10	1.1	U			Bwg		87Gr.A	
^{103}Ru(β^-)^{103}Rh		764	4	764.5	2.3	0.1	–					58Ro09	*
		760	6			0.8	–					65Mu09	*
		762	5			0.5	–					70Pe04	*
		769	4			−1.1	–					82Oh04	
	ave.	764.6	2.3			−0.0	1	98	98 ^{103}Rh			average	
^{103}Pd(ε)^{103}Rh		564	20	574.7	2.4	0.5	U					54Ri09	*
		543.0	0.8			39.7	B					86Be53	
^{103}Ag(β^+)^{103}Pd		2580	100	2654	4	0.7	U					62Pa05	*
		2700	100			−0.5	U					66Ja12	
		2320	50			6.7	B					69Ba02	
		2622	34			0.9	U			Dlf		88Bo28	*
^{103}Cd(β^+)^{103}Ag		4200	130	4151	4	−0.4	U					70Be.A	
		4250	200			−0.5	U					72We.A	
		4131	23			0.9	U			Dlf		88Bo28	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{103}In(β^+)^{103}Cd	5380	200	6019	9	3.2	B			Brk		83Wo04	
	6050	28			−1.1	1	11	10 ^{103}In	Dlf		88Bo28	*
	6040	60			−0.3	U					98Ka42	
^{103}Sn(β^+)^{103}In	7500	600	7507	9	0.0	o			GSI		04Mu32	
	7660	70			−2.2	B			GSI		05Ka34	*
*^{103}Ag−u	$M-A$=–84784(29) keV for mixture gs+m at 134.45 keV										Nub269	**
*^{103}Sn−u	recalculated from frequency ratio $^{103}Sn^{++}/(^{12}C_3H_2N)^+$ =1.010 779 605(29)										HWJ257	**
*^{103}Sn−^{103}Cd	from frequency ratio $^{103}Sn^{++}/(^{103}Cd)^{++}$ =1.0001411493(904)										26Ro.A	**
*^{103}Sn−^{103}Cd	Original error 8.9; added systematic error 3 keV										SNa265	**
*^{103}Cd−^{102}Cd	From ^{102}Cd/^{103}Cd=0.99029800(150)										AHW	**
*^{103}Ru(β^-)^{103}Rh	E_{β^-}=227(4) to 5/2$^+$ level at 536.840 keV, and other E_{β^-}										Ens09a	**
*^{103}Ru(β^-)^{103}Rh	E_{β^-}=112(6) to 5/2$^+$ level at 650.064 keV, and other E_{β^-}										Ens09a	**
*^{103}Ru(β^-)^{103}Rh	E_{β^-}=225(5) to 5/2$^+$ level at 536.840 keV, and other E_{β^-}										Ens09a	**
*^{103}Pd(ε)^{103}Rh	IBE=500(20) to ^{103}Rh^m 7/2$^+$ at 39.753 keV										Nub269	**
*^{103}Ag(β^+)^{103}Pd	E_{β^+}=1290(100) to 5/2$^+$ level at 266.861 keV, and other E_{β^+}										Ens09a	**
*^{103}Ag(β^+)^{103}Pd	E_{β^+}=1601(27), plus systematic error 20 keV estimated by evaluator										GAu	**
*^{103}Cd(β^+)^{103}Ag	E_{β^+}=3109(11), plus systematic error 20 keV estimated by evaluator										GAu	**
*^{103}In(β^+)^{103}Cd	E_{β^+}=4841(20), plus systematic error 20 keV estimated by evaluator										Ens09a	**
*^{103}Sn(β^+)^{103}In	Original 7640(70) recalibrated										05Ka34	**
*^{103}Sn(β^+)^{103}In	Trends in Mass Surface TMS suggest ^{103}Sn 120 keV more bound										HWJ20a	**
^{104}Sr−u	−46604	35				2			TR1	1.0	21Mu05	
^{104}Y−u	−59274	1343	−57967	17	1.0	B			MT1	1.0	24Wa19	*
^{104}Y−^{133}Cs$_{.782}$	15970	17				2			JY2	1.0	24Hu16	
^{104}Zr−u	−70661	54	−70551	10	2.0	U			JY0	1.0	04Ri12	
^{104}Nb−u	−77355	100	−77093.2	1.2	1.0	U			GT2	2.5	08Kn.A	*
^{104}Nb−^{52}Cr$_{2.000}$	41898.3	1.9	41897.4	1.2	−0.5	1	42	41 ^{104}Nb	CP2	1.0	18Or.A	
^{104}Nb−^{133}Cs$_{.782}$	−3157.2	1.9	−3156.7	1.2	0.3	1	41	41 ^{104}Nb	JY2	1.0	24Hu16	*
^{104}Nb^m−^{52}Cr$_{2.000}$	41908.8	2.0	41908.7	1.9	−0.0	1	94	94 ^{104}Nb^m	CP2	1.0	18Or.A	
^{104}Nb^m−^{133}Cs$_{0.782}$	−3146.6	7.7	−3145.3	1.9	0.2	1	6	6 ^{104}Nb^m	JY2	1.0	24Hu16	*
^{104}Tc−^{133}Cs$_{.782}$	−14715.0	2.4				2			JY2	1.0	25Ru08	
C_8 H_8−^{104}Ru	157171.5	3.4	157175.7	1.3	0.5	U			M16	2.5	63Da10	
C_8 H_8−^{104}Pd	158612	10	158568.9	1.3	−1.7	U			M16	2.5	63Da10	
	158599	12			−1.7	U			R12	1.5	83De51	
C_7 N H_6−^{104}Pd	146013	8	145992.8	1.3	−1.7	U			R12	1.5	83De51	
C_6 ^{13}C N H_5−^{104}Pd	141552	20	141522.6	1.3	−1.0	U			R12	1.5	83De51	
^{104}Pd−u	−95938	30	−95968.6	1.3	−1.0	U			GS2	1.0	05Li24	
^{104}Ag−u	−91410	30	−91375	4	1.2	U			GS2	1.0	05Li24	*
^{104}Cd−u	−90147	30	−90143.8	1.8	0.1	U			GS2	1.0	05Li24	
^{104}Cd−^{85}Rb$_{1.224}$	17813.7	5.5	17825.6	1.8	2.2	U			SH1	1.0	07Ma92	
	17825.5	1.9			0.0	1	89	89 ^{104}Cd	MA8	1.0	09Br09	
^{104}In−^{85}Rb$_{1.224}$	26183.9	6.2				2			SH1	1.0	07Ma92	
	26140.3	29.6	26184	6	1.5	U			JY1	1.0	09El08	*
^{104}Sn−^{87}Rb$_{1.195}$	31636.9	6.4	31635	5	−0.3	1	53	53 ^{104}Sn	JY1	1.0	09El07	
^{104}Sn−u	−76896.9	10.7	−76894	5	0.3	–			IS1	1.0	25Ni02	
	−76867	54			−0.5	U			RT1	1.0	25Cz.1	
	−76889.6	9.7			−0.5	–			RI1	1.0	26Ro.A	*
ave.	−76893	7			−0.2	1	42	42 ^{104}Sn			average	
^{104}Sb−u	−63723	25	−63702	13	0.8	1	28	28 ^{104}Sb	RI1	1.0	26Ro.A	*
^{104}Cd−^{96}Mo$_{1.083}$	13094.2	5.5	13093.4	1.8	−0.1	1	11	11 ^{104}Cd	JY1	1.0	09El08	
^{104}Zr−^{97}Zr$_{1.072}$	24896	10				2			JY1	1.0	06Ha03	
^{104}Nb−^{97}Zr$_{1.072}$	18352.8	2.9	18353.6	1.2	0.3	1	18	18 ^{104}Nb	JY1	1.0	07Ha32	*
^{104}Mo−^{97}Zr$_{1.072}$	9194.0	9.7				2			JY1	1.0	06Ha03	
^{104}Ru−^{104}Pd	1393.15	0.04	1393.15	0.04	0.0	1	100	80 ^{104}Ru	JY2	1.0	25Ru01	
^{104}In−^{103}In	−1241	231	−1664	11	−1.2	U			CR2	1.5	91Sh19	*
^{104}Pd−^{102}Pd	−1617	30	−1601.1	1.4	0.4	U			R12	1.5	83De51	
^{104}Te(α)^{100}Sn	5086	177	5200	80	0.6	3			Ara		18Au04	*
	5067.2	166.4			0.8	o			ORa		19Xi06	
	5231.6	93.6			−0.3	3			ORa		26Co.1	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{104}Ru(d,α)^{102}Tc	7180	10	7188	9	0.8	1	80	80 ^{102}Tc	Pri		82De03	
^{102}Ru(t,p)^{104}Ru	6648	30	6651.1	1.3	0.1	U			LAl		72Ca10	
^{104}Ru(d,^{3}He)^{103}Tc	−5289	10	−5287	9	0.2	2			VUn		83De20	
^{104}Ru(t,α)^{103}Tc	9048	30	9033	9	−0.5	2			LAl		81Fl02	
^{104}Ru(d,t)^{103}Ru−^{148}Gd()^{147}Gd	85	3	83.0	1.6	−0.7	1	28	14 ^{104}Ru	Jul		86Ru04	*
^{103}Rh(n,γ)^{104}Rh	6998.96	0.10	6998.96	0.08	−0.0	2			MMn		81Ke03	Z
	6998.95	0.14			0.0	2			Bdn		06Fi.A	
^{103}Rh(d,p)^{104}Rh	4786	10	4774.39	0.08	−1.2	U			MIT		64Sp12	
^{104}Pd(d,t)^{103}Pd	−3762	25	−3751.5	1.5	0.4	U			Pit		64Co11	
^{104}Sb(p)^{103}Sn	510	100	499	12	−0.1	U					94Pa12	*
^{104}Nb(β^-)^{104}Mo	8105	90	8532	9	4.7	B			Bwg		87Gr18	
^{104}Nb^m(β^-)^{104}Mo	8320	80	8543	9	2.8	B			Bwg		87Gr18	*
^{104}Mo(β^-)^{104}Tc	4800	200	2234	9	−12.8	B					64Te02	
	2155	40			2.0	B			Bwg		87Gr18	
	2160	40			1.9	B			Jyv		94Jo.A	
^{104}Tc(β^-)^{104}Ru	5620	70	5518.1	2.5	−1.5	U					78Su03	
	5590	60			−1.2	U			Bwg		87Gr18	
^{104}Rh(β^-)^{104}Pd	2440	30	2435.0	2.6	−0.2	U					55Bu.A	
^{104}Ag(β^+)^{104}Pd	4276	15	4279	4	0.2	U					60Nu02	*
	4350	200			−0.4	U					72We.A	
	4306	31			−0.9	U			Dlf		88Bo28	*
^{104}Pd(p,n)^{104}Ag	−5061	4				2					79De44	
^{104}Cd(β^+)^{104}Ag	1587	60	1147	5	−7.3	B					70Mu17	*
	1403	26			−9.8	B			GSI		79Pl06	*
^{104}In(β^+)^{104}Cd	7100	200	7786	6	3.4	B					78Hu06	*
	7260	250			2.1	U			Brk		83Wo04	*
	7800	250			−0.1	U			Dlf		88Bo28	
	7890	120			−0.9	U			Dlf		90Re08	
	7880	100			−0.9	U			GSI		98Ka.A	
^{104}Sn(β^+)^{104}In	4550	300	4556	7	0.0	U					88Ba10	*
	4515	60			0.7	U			GSI		91Ke11	
*^{104}Y−u	Trends in Mass Surface TMS suggest ^{104}Y 1200 keV less bound										HWJ24b	**
*^{104}Nb−u	D_M=–77350(100) μu for mixture gs+m at 11(2) keV; $M-A$=–72051(93) keV										Nub269	**
*^{104}Nb−^{133}Cs$_{.782}$	only one state observed										24Hu16	**
*^{104}Nb^m−^{133}Cs$_{0.782}$	only one state observed										24Hu16	**
*^{104}Ag−u	$M-A$=–85144(28) keV for mixture gs+m at 6.90 keV										Nub269	**
*^{104}In−^{85}Rb$_{1.224}$	D_M=26190.5(5.5) μu for mixture gs+m at 93.48 keV; $M-A$=–76176.5(5.1) keV										Nub269	**
*^{104}Sn−u	from frequency ratio ^{104}Sn^{++}/(^{40}Ar^{12}C)$^+$ =0.9999840670(870)										26Ro.A	**
*^{104}Sn−u	Original error 8.4; added systematic error 3 keV										SNa265	**
*^{104}Sb−u	from frequency ratio ^{104}Sb^{++}/(^{40}Ar^{12}C)$^+$ =1.000110755(242)										26Ro.A	**
*^{104}Sb−u	Original error 23; added systematic error 3 keV										SNa265	**
*^{104}Nb−^{97}Zr$_{1.072}$	assigned to the gs. by the evaluator										HWJ266	**
*^{104}In−^{103}In	From ^{103}In/^{104}In=0.99038900(222)										AHW	**
*^{104}Te(α)^{100}Sn	average of two events with E_α=4.73(0.24) and 5.06(0.25) MeV										FKo256	**
*^{104}Ru(d,t)^{103}Ru−^{148}Gd()^{147}Gd	Q =82(3) to 5/2$^+$ level at 2.81 keV										Ens09a	**
*^{104}Sb(p)^{103}Sn	Below 550 keV; value and error estimated by evaluator										94Pa12	**
*^{104}Nb^m(β^-)^{104}Mo	Better use the difference of the two Q_{β^-} 's										GAu	**
*^{104}Ag(β^+)^{104}Pd	E_{β^+}=2705(15) from ^{104}Ag^m at 6.90 to 2$^+$ level at 555.81 keV										Ens079	**
*^{104}Ag(β^+)^{104}Pd	E_{β^+}=2012(71) to 4$^+$ level at 1323.59 keV										Ens079	**
*	and E_{β^+}=2729(24) from ^{104}Ag^m at 6.90 to 2$^+$ level at 555.81 keV										Ens079	**
*	plus systematic error 20 keV estimated by evaluator										GAu	**
*^{104}Cd(β^+)^{104}Ag	p$^+$ =0.011(0.003) 0.0019(0.0005) respectively, to 1$^+$ level at 90.6 keV; recalculated E_{β^+}										Ens079	**
*^{104}In(β^+)^{104}Cd	E_{β^+}=4600(200) 4750(250) respectively, to 4$^+$ level at 1492.1 keV										Ens079	**
*^{104}Sn(β^+)^{104}In	p$^+$ =0.71(0.07) to 1139.25 level										Ens079	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{105}Sr−u		−40672	47				2			TR1	1.0	21Mu05	
^{105}Y−u		−55041	574	−54290#	210#	0.5	D			GT3	2.5	16Kn03	*
^{105}Mo−^{133}Cs$_{.789}$		−8421.7	1.9				2			JY2	1.0	24Ja10	
^{105}Tc−^{85}Rb$_{1.235}$		20698.5	9.3				2			JY1	1.0	25Ru08	
^{105}Rh−u		−94378	53	−94311.9	2.6	1.2	U			GS2	1.0	05Li24	*
C_8 H_9−^{105}Pd		165357	14	165345.0	1.2	−0.3	U			M16	2.5	63Da10	
		165360	9			−1.1	U			R12	1.5	83De51	
C_7 N H_7−^{105}Pd		152773	18	152769.0	1.2	−0.1	U			R12	1.5	83De51	
C_6 ^{13}C N H_6−^{105}Pd		148309	26	148298.8	1.2	−0.3	U			R12	1.5	83De51	
C_7 O H_5−^{105}Pd		128970	18	128959.5	1.2	−0.4	U			R12	1.5	83De51	
^{105}Ag−u		−93534	31	−93475	5	1.9	U			GS2	1.0	05Li24	*
^{105}Cd−^{96}Mo$_{1.094}$		13748.5	5.4	13749.7	1.5	0.2	U			JY1	1.0	09El08	
^{105}Cd−^{85}Rb$_{1.235}$		18403.4	1.5	18403.5	1.5	0.1	1	99	99 ^{105}Cd	MA8	1.0	09Br09	
^{105}In−^{85}Rb$_{1.235}$		23442	11				2			SH1	1.0	07Ma92	
^{105}In−u		−85536.8	33.3	−85498	11	1.2	U			GR1	1.0	20Ho03	
^{105}Sn−^{85}Rb$_{1.235}$		30204.1	7.1	30207	4	0.4	1	33	33 ^{105}Sn	SH1	1.0	07Ma92	
^{105}Sn−^{87}Rb$_{1.207}$		30890.0	5.6	30886	4	−0.6	1	52	52 ^{105}Sn	JY1	1.0	09El07	
^{105}Sn−u		−78742.3	14.0	−78733	4	0.7	–			IS1	1.0	25Ni02	
		−78743	37			0.3	–			RT1	1.0	25Cz.1	
	ave.	−78742	13			0.7	1	10	10 ^{105}Sn			average	
^{105}Zr−^{97}Zr$_{1.082}$		30359	13				2			JY1	1.0	06Ha03	
^{105}Mo−^{97}Zr$_{1.082}$		13319.4	9.8	13317.1	1.9	−0.2	U			JY1	1.0	06Ha03	
^{105}Nb−^{102}Ru$_{1.029}$		23376.4	4.3				2			JY1	1.0	07Ha32	
^{105}Pd−^{104}Pd		1049	35	1048.9	0.7	−0.0	U			R12	1.5	83De51	
^{105}In−^{104}In		−3618	144	−3712	13	−0.4	U			CR2	1.5	91Sh19	*
^{105}Te(α)^{101}Sn		5079	50	5069	3	−0.2	U					06Se08	*
		5061.1	5.			1.6	o					06Li41	*
		5069.2	3.				2					10Da17	*
^{104}Ru(n,γ)^{105}Ru		5909.9	0.5	5910.09	0.11	0.4	–					74Hr01	
		5910.1	0.2			−0.0	–					78Gu14	
		5910.11	0.14			−0.1	–			Bdn		06Fi.A	
^{104}Ru(d,p)^{105}Ru		3684	15	3685.53	0.11	0.1	U			ANL		71Fo01	
		3684.5	1.0			1.0	U			Mun		76Ma49	
^{104}Ru(n,γ)^{105}Ru	ave.	5910.10	0.11	5910.09	0.11	−0.0	1	100	94 ^{105}Ru			average	
^{104}Pd(n,γ)^{105}Pd		7094.1	0.7	7094.3	0.7	0.3	1	93	80 ^{104}Pd			70Bo29	
^{104}Pd(d,p)^{105}Pd		4867	20	4869.7	0.7	0.1	U			Pit		64Co11	
^{105}Pd(d,t)^{104}Pd		−851	30	−837.1	0.7	0.5	U			Pit		64Co11	
^{105}Sb(p)^{104}Sn		482.6	15.	322	22	−10.7	F			Bkp		94Ti03	*
^{105}Nb(β^-)^{105}Mo		6485	70	7417	4	13.3	B			Bwg		87Gr18	
^{105}Mo(β^-)^{105}Tc		4950	45	4863	9	−1.9	U			Bwg		87Gr18	
^{105}Tc(β^-)^{105}Ru		3640	55	3739	9	1.8	U			Bwg		87Gr18	
^{105}Ru(β^-)^{105}Rh		1916	4	1915.7	2.3	−0.1	1	32	30 ^{105}Rh			67Sc01	
^{105}Rh(β^-)^{105}Pd		570	5	566.2	2.2	−0.8	–					51Du03	
		560	5			1.2	–					56La24	
		568	4			−0.5	–					64Ka23	
	ave.	566.3	2.6			−0.0	1	70	70 ^{105}Rh			average	
^{105}Ag(ε)^{105}Pd		1347	25	1346	5	−0.0	U					67Pi03	
		1310	25			1.4	U					67Sc26	*
^{105}Cd(β^+)^{105}Ag		2738	5	2737	4	−0.2	–					53Jo20	*
		2600	200			0.7	U					72We.A	
		2742	11			−0.4	–					86Bo28	*
	ave.	2739	5			−0.3	1	92	91 ^{105}Ag			average	
^{105}In(β^+)^{105}Cd		5140	200	4693	10	−2.2	U			Brk		83Wo04	
		4849	13			−12.0	B					86Bo28	*
^{105}Sn(β^+)^{105}In		6230	80	6302	11	0.9	U			GSI		06Ka74	
*^{105}Y−u	Trends in Mass Surface TMS suggest ^{105}Y 700 keV less bound											GAu	**
*^{105}Rh−u	$M-A$=–87847(32) keV for mixture gs+m at 129.742 keV											Nub269	**
*^{105}Ag−u	$M-A$=–87113(28) keV for mixture gs+m at 25.468 keV											Nub269	**
*^{105}In−^{104}In	From ^{104}In/^{105}In=0.99050293(139)											AHW	**
*^{105}Te(α)^{101}Sn	E_α=4720(50), 4703(5) to 7/2$^+$ level at 171.7 keV (same group as next)											Ens07a	**
*^{105}Te(α)^{101}Sn	E_α=4711(3) to 7/2$^+$ level at 171.7 keV; also E_α=4880(20) to ground state											Ens07a	**
*^{105}Sb(p)^{104}Sn	F : expected 150 protons, no proton peak observed											05Li47	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{105}$Ag(ε)^{105}Pd		L/K=0.152(0.002) → Q =222(12+theor.error) to 3/2$^-$ level at 1087.96 keV										Ens05b	**
$*^{105}$Cd(β^+)^{105}Ag		E_{β^+}=1691(5) to ^{105}Ag^m at 25.468 keV										Nub269	**
$*^{105}$Cd(β^+)^{105}Ag		E_{β^+}=1695(11) to ^{105}Ag^m at 25.468 keV										Nub269	**
$*^{105}$In(β^+)^{105}Cd		E_{β^+}=3696(13) to 7/2$^+$ level at 131.11 keV										Ens05b	**
^{106}Zr−u		−62674	322	−62891	5	−0.4	o			GT1	1.5	04Ma.A	
		−62856	186			−0.1	U			GT3	2.5	16Kn03	
		−62384	750			−0.7	B			MT1	1.0	24Wa19	
^{106}Zr−^{133}Cs$_{.797}$		12464.1	5.3				2			JY2	1.0	24Hu16	
^{106}Nb−u		−70843	258	−71070.9	1.3	−0.6	U			GT1	1.5	04Ma.A	
^{106}Nb−^{52}Cr$_{2.038}$		50180.0	1.6	50180.5	1.4	0.3	o			CP2	1.0	18Or.A	
^{106}Nb−u		−71070.8	1.4	−71070.9	1.3	−0.1	1	91	91 ^{106}Nb	CP2	1.0	21Mi07	*
^{106}Mo−^{97}Zr$_{1.093}$		15589.8	9.8				2			JY1	1.0	06Ha03	
^{106}Tc−^{133}Cs$_{.797}$		−10245.5	2.0				2			JY2	1.0	25Ru08	
C$_8$ H$_{10}$−^{106}Pd		174764.0	4.3	174769.3	1.1	0.5	U			M16	2.5	63Da10	
		174751	32			0.4	U			R12	1.5	83De51	
		174766	8			0.3	U			R12	1.5	83De51	
C$_7$ ^{13}C H$_9$−^{106}Pd		170285	32	170299.1	1.1	0.3	U			R12	1.5	83De51	
		170298	30			0.0	U			R12	1.5	83De51	
C$_7$ N H$_8$−^{106}Pd		162186	18	162193.2	1.1	0.3	U			R12	1.5	83De51	
C$_7$ O H$_6$−^{106}Pd		138378	20	138383.8	1.1	0.2	U			R12	1.5	83De51	
^{106}Pd−u		−96495	30	−96519.0	1.1	−0.8	U			GS2	1.0	05Li24	
		−96521.0	1.9			0.7	–			TG1	1.5	12Sm01	
		−96524.9	4.7			0.8	–			TG1	1.5	12Sm01	
	ave.	−96521.5	2.6			1.0	1	18	18 ^{106}Pd			average	
^{106}Ag−u		−93318	44	−93336	3	−0.4	U			GS2	1.0	05Li24	*
C$_8$ H$_{10}$−^{106}Cd		171789.3	2.7	171789.8	1.1	0.1	U			M16	2.5	63Da10	
		171841	17			−2.0	U			R12	1.5	83De51	
C$_7$ N H$_8$−^{106}Cd		159210	15	159213.7	1.1	0.2	U			R12	1.5	83De51	
^{106}Cd−u		−93540.6	1.7	−93539.5	1.1	0.4	–			TG1	1.5	12Sm01	
		−93545.7	3.47			1.2	–			TG1	1.5	12Sm01	
	ave.	−93541.6	2.3			0.9	1	24	24 ^{106}Cd			average	
^{106}Cd−^{85}Rb$_{1.247}$		16459.8	1.8	16458.7	1.1	−0.6	1	39	39 ^{106}Cd	MA8	1.0	09Br09	
^{106}In−u		−86516	32	−86536	13	−0.6	U			GS2	1.0	05Li24	*
^{106}Sn−^{85}Rb$_{1.247}$		26959.4	8.7	26957	5	−0.3	1	34	34 ^{106}Sn	SH1	1.0	07Ma92	
^{106}Sn−^{87}Rb$_{1.218}$		27578.0	7.6	27577	5	−0.1	1	45	45 ^{106}Sn	JY1	1.0	09El07	
^{106}Sn−u		−83034	15	−83041	5	−0.5	–			IS1	1.0	25Ni02	
		−83014	40			−0.7	–			RT1	1.0	25Cz.1	
	ave.	−83032	14			−0.7	1	13	13 ^{106}Sn			average	
^{106}Sb−^{87}Rb$_{1.218}$		39256.1	8.0				2			JY1	1.0	09El07	
^{106}Nb−^{102}Ru$_{1.039}$		28318.2	4.4	28319.3	1.4	0.3	1	10	9 ^{106}Nb	JY1	1.0	07Ha32	
^{106}Tc−^{102}Ru$_{1.039}$		13780.7	4.7	13789.9	2.0	2.0	o			JY1	1.0	07Ha20	*
^{106}Ru−^{105}Ru$_{1.010}$		511.8	9.1	506	6	−0.6	1	39	38 ^{106}Ru	JY1	1.0	07Ha20	
^{106}Cd−^{106}Pd		2979.50	0.11	2979.50	0.11	−0.0	1	100	63 ^{106}Pd	SH1	1.0	11Go23	
		2979.08	0.60			0.5	U			TG1	1.5	12Sm01	
^{106}Pd−^{105}Pd		−1608	25	−1599.2	0.3	0.2	U			R12	1.5	83De51	
^{106}Pd−^{104}Pd		−508	32	−550.3	0.8	−0.9	U			R12	1.5	83De51	
^{106}Te(α)^{102}Sn		4323.5	30.	4290	9	−1.1	U					81Sc17	
		4290.2	9.				3					94Pa11	
		4323.5	30.			−1.1	U					02Ma19	
		4261.0	62.4			0.5	U					16Ca33	
^{106}Cd(^{3}He,^{6}He)^{103}Cd		−9173	17	−9140.5	2.1	1.9	U			MSU		78Pa11	*
^{104}Ru(t,p)^{106}Ru		5892	20	5887	5	−0.2	U			LAl		72Ca10	
^{104}Pd(^{3}He,p)^{106}Ag		5153	6	5190.0	2.9	6.2	F			Bld		75An07	*
^{106}Cd(p,t)^{104}Cd		−10802	15	−10823.9	2.0	−1.5	U			MSU		82Cr01	
		−10829	12			0.4	U			Pri		83De03	
		−10819	12			−0.4	U			Ors		84Ro.A	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{105}Pd(n,γ)^{106}Pd		9562.8	1.1	9560.98	0.28	−1.7	U					70Bo29	
		9560.5	0.4			1.2	–			BNn		87Fo20	*
		9561.4	0.4			−1.0	–			Bdn		06Fi.A	
^{105}Pd(d,p)^{106}Pd		7349	30	7336.42	0.28	−0.4	U			Pit		64Co11	
^{106}Pd(d,t)^{105}Pd		−3311	25	−3303.75	0.28	0.3	U			Pit		64Co11	
^{105}Pd(n,γ)^{106}Pd	ave.	9560.95	0.28	9560.98	0.28	0.1	1	99	87 ^{105}Pd			average	
^{105}Pd(^{3}He,d)^{106}Ag		322	8	320.2	2.8	−0.2	1	12	12 ^{106}Ag	Bld		75An07	
^{106}Cd(d,t)^{105}Cd		−4661	50	−4611.7	1.7	1.0	U					73De16	
^{106}Cd(^{3}He,α)^{105}Cd		9728	25	9708.7	1.7	−0.8	U			Man		75Ch21	
^{106}Mo(β^-)^{106}Tc		3510	45	3608	9	2.2	B			Bwg		87Gr18	
		3520	17			5.2	C			Bwg		92Gr.A	
		3520	17			5.2	C			Jyv		94Jo.A	
^{106}Tc(β^-)^{106}Ru		6545	45	6587	6	0.9	o			Bwg		87Gr18	
		6547	11			3.6	B			Bwg		92Gr.A	
^{106}Ru(β^-)^{106}Rh		39.2	0.3	39.40	0.21	0.7	–					50Ag01	
		39.6	0.3			−0.7	–					58Gr07	
	ave.	39.40	0.21			0.0	1	100	62 ^{106}Ru			average	
^{106}Rh(β^-)^{106}Pd		3530	10	3544	5	1.4	–					52Al06	
		3550	10			−0.6	–					58Gr07	
		3550	20			−0.3	–					60Se05	
	ave.	3541	7			0.5	1	62	62 ^{106}Rh			average	
^{106}Rh^{m}(β^-)^{106}Pd		3677	10				2					66De11	*
^{106}Ag(β^+)^{106}Pd		2980	20	2964.9	2.8	−0.8	U					53Be42	
		3011	72			−0.6	U					86Bo28	
^{106}Ag(ε)^{106}Pd		2961	4			1.0	–					78Ge01	*
^{106}Pd(p,n)^{106}Ag		−3754	13	−3747.3	2.8	0.5	U			Oak		64Jo11	
		−3756	5			1.7	–					79De44	
^{106}Ag(ε)^{106}Pd	ave.	2966	3	2964.9	2.8	−0.3	1	81	81 ^{106}Ag			average	
^{106}In(β^+)^{106}Cd		6516	30	6524	12	0.3	2					66Ca09	*
		6507	29			0.6	2					86Bo28	*
^{106}Cd(p,n)^{106}In		−7312.9	15.	−7306	12	0.4	2			ANL		84Fi05	*
^{106}Sn(β^+)^{106}In		3195	60	3255	13	1.0	U			GSI		79Pl06	*
		3200	100			0.6	U					88Ba10	
*^{106}Nb−u	Possible mixture of ground state and m at 100(50)# keV											21Mi07	**
*^{106}Ag−u	$M-A$=–86880(32) keV for mixture gs+m at 89.66 keV											Nub269	**
*^{106}In−u	$M-A$=–80575(29) keV for mixture gs+m at 28.6 keV											Nub269	**
*^{106}Tc−^{102}Ru$_{1.039}$	no impurities were found by reference											25Ru08	**
*^{106}Cd(^{3}He,^{6}He)^{103}Cd	First excited state at 187.89 keV yields Q =–9146 keV											GAu	**
*^{104}Pd(^{3}He,p)^{106}Ag	F : withdrawn by authors											AHW	**
*^{105}Pd(n,γ)^{106}Pd	Calculated from 13 γ energies in 2 keV n-capture											AHW	**
*	to levels in ^{106}Pd; corrected for recoil											Ens086	**
*^{106}Rh^{m}(β^-)^{106}Pd	E_{β^-}=920(10) to 5$^+$ level at 2757.06 keV											Ens086	**
*^{106}Ag(ε)^{106}Pd	L/K=0.203(0.003) gives Q_{β^+}=99(4), recalculated Q											AHW	**
*	from ^{106}Ag^{m} 6$^+$ at 89.66 keV to 5$^+$ level at 2951.84 keV											Ens086	**
*^{106}In(β^+)^{106}Cd	E_{β^+}=4890(30) from ^{106}In^{m} 2$^+$ at 28.6 to 2$^+$ level at 632.64 keV											Ens086	**
*^{106}In(β^+)^{106}Cd	E_{β^+}=2965(30) to 6$^+$ level at 2491.66 keV and 4908(29) from											Ens086	**
*	^{106}In^{m} (2)$^+$ at 28.6 to 2$^+$ level at 632.64 keV											Ens086	**
*^{106}Cd(p,n)^{106}In	T=7535(15) to 2$^+$ level at 151.1 keV											Ens086	**
*^{106}Sn(β^+)^{106}In	p$^+$ =0.253(0.021) to 1$^+$ level at 893.0 keV											Ens086	**
^{107}Zr−u		−58379	482	−57760#	220#	0.5	D			GT3	2.5	16Kn03	*
^{107}Nb−u		−68326	279	−68410	9	−0.2	U			GT1	1.5	04Ma.A	
^{107}Mo−^{97}Zr$_{1.103}$		20326.7	9.9				2			JY1	1.0	06Ha03	
^{107}Pd−u		−95013	95	−94871.1	1.2	1.5	U			GS2	1.0	05Li24	*
C$_8$ H$_{11}$−^{107}Ag		180986.4	3.1	180986.7	1.2	0.0	U			M16	2.5	63Da10	
		180994	17			−0.3	U			R12	1.5	83De51	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
C_7 N $H_9-{}^{107}$Ag		168415	8	168410.7	1.2	−0.4	U			R12	1.5	83De51	
C_7 O $H_7-{}^{107}$Ag		144595	18	144601.2	1.2	0.2	U			R12	1.5	83De51	
C_6 ^{13}C O $H_6-{}^{107}$Ag		140131	16	140131.0	1.2	0.0	U			R12	1.5	83De51	
C_6 N O $H_5-{}^{107}$Ag		132025	16	132025.2	1.2	0.0	U			R12	1.5	83De51	
^{107}Cd−u		−93410	30	−93388.4	1.7	0.7	U			GS2	1.0	05Li24	
		−93359	97			−0.3	U			GR1	1.0	19An10	
^{107}Cd−^{85}Rb$_{1.259}$		17668.7	1.9	17668.3	1.7	−0.2	1	84	84 ^{107}Cd	MA8	1.0	09Br09	
^{107}In−u		−89710	30	−89713	10	−0.1	–			GS2	1.0	05Li24	
		−89730.0	29.0			0.6	–			GR1	1.0	20Ho03	
	ave.	−89720	21			0.4	1	25	25 ^{107}In			average	
^{107}Sn−^{87}Rb$_{1.230}$		27421.6	5.7				2			JY1	1.0	09El07	
^{107}Sn−u		−84285	37	−84286	6	−0.0	U			RT1	1.0	25Cz.1	
^{107}Sb−^{87}Rb$_{1.230}$		35866.2	5.8	35859	4	−1.3	1	59	59 ^{107}Sb	JY1	1.0	09El07	
^{107}Sb−^{133}Cs$_{.805}$		251.8	9.7	262	4	1.0	1	21	21 ^{107}Sb	SH1	1.0	07Ma92	
^{107}Nb−^{102}Ru$_{1.049}$		31936.7	8.6				2			JY1	1.0	07Ha32	
^{107}Tc−^{105}Ru$_{1.019}$		9465.8	8.9				2			JY1	1.0	07Ha20	
^{107}Ru−^{105}Ru$_{1.019}$		3977.2	8.9				2			JY1	1.0	07Ha20	
^{107}Sn−^{106}Sn		−1148	86	−1245	8	−0.8	U			CR2	1.5	92Sh.A	*
^{107}Te(α)^{103}Sn		3982.2	15.	4007	4	1.6	–					79Sc22	
		4011.3	5.			−0.8	–					91He21	
		4007	10			0.0	–			Ara		19Au02	
		4000.8	17.7			0.3	–					20Ca01	
	ave.	4008	4			−0.2	1	98	97 ^{107}Te			average	
^{107}Ag(p,t)^{105}Ag		−9015	15	−8999	5	1.1	1	10	9 ^{105}Ag	Min		75Ku14	*
^{106}Pd(n,γ)^{107}Pd		6536.4	0.5	6536.4	0.5	−0.1	1	99	92 ^{107}Pd	Bdn		06Fi.A	
^{107}Ag(γ,n)^{106}Ag		−9353	34	−9538.8	2.9	−5.5	B			Phi		60Ge01	
^{107}Ag(p,d)^{106}Ag		−7305	11	−7314.2	2.9	−0.8	1	7	7 ^{106}Ag	Bld		75An07	
^{107}Mo(β^-)^{107}Tc		6160	60	6206	12	0.8	U			Bwg		89Gr23	
^{107}Tc(β^-)^{107}Ru		4820	85	5113	12	3.4	B			Bwg		89Gr23	
^{107}Ru(β^-)^{107}Rh		3140	300	3000	15	−0.5	U					62Pi02	*
		2900	135			0.7	U			Bwg		89Gr23	
^{107}Rh(β^-)^{107}Pd		1510	40	1508	12	−0.0	U					62Pi02	*
^{107}Pd(β^-)^{107}Ag		33	3	37.49	0.09	1.5	U					49Pa.B	
		37.5	0.1			−0.1	–			Pri		93Lo.1	*
		37.46	0.19			0.2	–			Pri		25Ko08	*
	ave.	37.49	0.09			−0.0	1	100	92 ^{107}Ag			average	
^{107}Cd(β^+)^{107}Ag		1417	4	1418.6	1.9	0.4	1	22	15 ^{107}Cd			62La10	*
^{107}In(β^+)^{107}Cd		3426	11	3424	10	−0.2	1	76	75 ^{107}In			86Bo28	*
*^{107}Zr−u	Trends in Mass Surface TMS suggest ^{107}Zr 360 keV less bound											GAu	**
*^{107}Pd−u	$M-A$=–88397(62) keV for mixture gs+n at 214.6 keV											Nub269	**
*^{107}Sn−^{106}Sn	From ^{107}Sn/^{106}Sn=1.00943053(81)											AHW	**
*^{107}Ag(p,t)^{105}Ag	Recalibrated with (p,t) results on ^{104}Pd, ^{105}Pd, ^{106}Pd and ^{108}Pd											AHW	**
*^{107}Ru(β^-)^{107}Rh	E_{β^-}=2100(300) to (5/2$^+$,7/2$^+$) level at 1041.950 keV											Ens085	**
*^{107}Rh(β^-)^{107}Pd	E_{β^-}=840(40) to 5/2$^+$ level at 670.06 keV											Ens085	**
*^{107}Pd(β^-)^{107}Ag	from 93Lo.1=37.5(0.01,stat)(0.10,syst), using micro-calorimeter											93Lo.1	**
*^{107}Pd(β^-)^{107}Ag	using micro-calorimeter; other E_{β^-}=37.4(1.7) using liquid scintillation											25Ko08	**
*^{107}Cd(β^+)^{107}Ag	E_{β^+}=302(4) to ^{107}Ag^m at 93.125 keV											Nub269	**
*^{107}In(β^+)^{107}Cd	E_{β^+}=2199(11) to 5/2$^+$ level at 204.98 keV											Ens085	**
^{108}Nb−u		−64413	440	−63925	9	0.7	o			GT1	1.5	04Ma.A	
		−63945	112			0.1	U			GT2	2.5	08Kn.A	
^{108}Nb−^{120}Sn$_{.900}$		24093.3	8.8				2			JY1	1.0	11Ha48	
^{108}Mo−^{97}Zr$_{1.113}$		23144.8	9.9				2			JY1	1.0	06Ha03	
^{108}Mo−u		−76039	215	−75952	10	0.3	U			GT1	1.5	04Ma.A	
^{108}Rh−^{120}Sn$_{.900}$		−3267	15	−3204.2	1.9	4.2	B			JY1	1.0	07Ha20	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{108}Rh^m−^{120}Sn$_{.900}$		−3144	13	−3133.7	2.0	0.8	U			JY1	1.0	07Ha20	
C_8 H_{12}−^{108}Pd		190014	6	190008.4	1.1	−0.4	U			M16	2.5	63Da10	
		190005	19			0.1	U			R12	1.5	83De51	
C_7 ^{13}C H_{11}−^{108}Pd		185532	30	185538.2	1.1	0.1	U			R12	1.5	83De51	
C_7 N H_{10}−^{108}Pd		177422	17	177432.3	1.1	0.4	U			R12	1.5	83De51	
C_6 ^{13}C N H_9−^{108}Pd		172943	18	172962.1	1.1	0.7	U			R12	1.5	83De51	
C_7 O H_8−^{108}Pd		153611	17	153622.9	1.1	0.5	U			R12	1.5	83De51	
C_6 N O H_6−^{108}Pd		141031	16	141046.8	1.1	0.7	U			R12	1.5	83De51	
^{108}Pd−u		−96109.6	1.3	−96108.0	1.1	0.8	–			TG1	1.5	12Sm01	
		−96108.1	4.7			0.0	–			TG1	1.5	12Sm01	
	ave.	−96109.5	1.9			0.8	1	36	36 ^{108}Pd			average	
^{108}Ag−u		−93973	50	−94052.6	1.2	−1.6	U			GS2	1.0	05Li24	*
^{108}Ag−^{133}Cs$_{.812}$		−17218	84	−17279.6	1.2	−0.7	U			MA8	1.0	08Br.A	*
C_8 H_{12}−^{108}Cd		189715.6	2.9	189716.6	1.2	0.1	U			M16	2.5	63Da10	
		189695	16			0.9	U			R12	1.5	83De51	
C_7 N H_{10}−^{108}Cd		177140	30	177140.6	1.2	0.0	U			R12	1.5	83De51	
C_6 ^{13}C N H_9−^{108}Cd		172653	15	172670.4	1.2	0.8	U			R12	1.5	83De51	
C_6 N O H_6−^{108}Cd		140746	15	140755.1	1.2	0.4	U			R12	1.5	83De51	
^{108}Cd−^{85}Rb$_{1.271}$		16298.5	2.3	16299.0	1.2	0.2	1	26	26 ^{108}Cd	MA8	1.0	09Br09	
^{108}Cd−u		−95818.3	1.7	−95816.2	1.2	0.8	–			TG1	1.5	12Sm01	
		−95817.1	4.87			0.1	–			TG1	1.5	12Sm01	
	ave.	−95818.2	2.4			0.8	1	24	24 ^{108}Cd			average	
^{108}In−u		−90277	31	−90306	9	−0.9	U			GS2	1.0	05Li24	*
^{108}Sn−u		−88102	32	−88106	6	−0.1	U			GS2	1.0	05Li24	
^{108}Sn−^{87}Rb$_{1.241}$		24599.8	5.9	24601	6	0.2	1	96	96 ^{108}Sn	JY1	1.0	09El07	
^{108}Sb−^{87}Rb$_{1.241}$		34933.7	5.9				2			JY1	1.0	09El07	
^{108}Te−^{87}Rb$_{1.241}$		42085.3	6.0	42087	6	0.4	1	94	94 ^{108}Te	JY1	1.0	09El07	
^{108}Tc−^{105}Ru$_{1.029}$		13423.4	9.0				2			JY1	1.0	07Ha20	
^{108}Ru−^{105}Ru$_{1.029}$		5115.7	8.9	5118.3	1.8	0.3	U			JY1	1.0	07Ha20	
^{108}Ru−^{108}Pd		6295.627	0.043				2			CP2	1.0	24Po14	
^{108}Rh−^{108}Pd		4886.0	1.2				2			CP2	1.0	25Li19	
^{108}Rh^m−^{108}Pd		4956.5	1.4				2			CP2	1.0	25Li19	
^{108}Pd−^{108}Cd		−292.05	0.59	−291.8	0.8	0.3	1	86	48 ^{108}Cd	TG1	1.5	12Sm01	
^{108}Sn−^{107}Sn		−3650	76	−3819	8	−1.5	U			CR2	1.5	92Sh.A	*
^{108}Pd−^{106}Pd		425	40	411.0	1.6	−0.2	U			R12	1.5	83De51	
^{108}Cd−^{106}Cd		−2232	32	−2276.8	1.6	−0.9	U			R12	1.5	83De51	
^{108}Te(α)^{104}Sn		3406.4	30.	3420	7	0.4	U					65Ma12	
		3448.0	20.			−1.4	1	11	6 ^{108}Te			81Sc17	
		3444.9	4.			−6.3	B					91He21	
		3406.4	25.			0.5	U					91Pa05	
^{108}I(α)^{104}Sb		4034.7	25.	4099	5	1.2	F					91Pa05	*
		4099.1	5.			0.1	–					94Pa12	
		4097	10			0.2	–			Ara		19Au02	
	ave.	4099	5			0.2	1	97	72 ^{104}Sb			average	
^{108}Xe(α)^{104}Te		4569.6	207.7	4780	90	1.0	4			Ara		18Au04	
		4818.8	93.5			−0.4	4			ORa		26Co.1	
^{108}Pd(d,^{3}He)^{107}Rh		−4456	12				2			Grn		86Ka43	
^{108}Pd(d,t)^{107}Pd		−2977	30	−2966.2	1.6	0.4	U			Pit		64Co11	
^{107}Ag(n,γ)^{108}Ag		7269.6	0.6	7271.41	0.17	3.0	B			ILn		85Ma54	Z
		7271.41	0.17				2			Bdn		06Fi.A	
^{107}Ag(d,p)^{108}Ag		5051	7	5046.84	0.17	−0.6	U			MIT		67Sp09	
^{108}I(p)^{107}Te		597	13	591	11	−0.5	1	78	74 ^{108}I	Ara		19Au02	
^{108}Mo(β^-)^{108}Tc		5135	60	5174	13	0.7	U			Bwg		92Gr.A	
		5120	40			1.4	o					94Jo.A	
		5100	60			1.2	U					95Jo02	
^{108}Tc(β^-)^{108}Ru		7720	50	7736	9	0.3	U			Bwg		89Gr23	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{108}Ru(β^-)^{108}Rh		1315	100	1313.1	1.1	−0.0	U					62Pi02	*
		1420	185			−0.6	U			Bwg		89Gr23	
		1380	80			−0.8	o			Jyv		92Jo05	
		1350	60			−0.6	U			Jyv		94Jo.A	
^{108}Rh(β^-)^{108}Pd		4500	600	4551.3	1.1	0.1	U					62Pi02	
		4505	105			0.4	U			Bwg		89Gr23	
^{108}Rh^m(β^-)^{108}Pd		4434	50	4617.0	1.3	3.7	B					69Pi08	*
		4510	100			1.1	U					84Bh02	*
^{108}Ag(β^+)^{108}Pd		1902	25	1914.6	1.6	0.5	U					62Fr07	
^{108}Pd(p,n)^{108}Ag		−2675	100	−2696.9	1.6	−0.2	U			Oak		64Jo11	
^{108}Ag(β^-)^{108}Cd		1650	40	1642.8	1.6	−0.2	U					60Wa10	
^{108}In(β^+)^{108}Cd		5124	50	5133	9	0.2	U					62Ka23	*
		5125	14			0.5	–					86Bo28	*
^{108}Cd(p,n)^{108}In		−5927	12	−5915	9	1.0	–			ANL		84Fi05	*
^{108}In(β^+)^{108}Cd	ave.	5136	9	5133	9	−0.4	1	89	89 ^{108}In			average	
^{108}Sn(β^+)^{108}In		2078	25	2050	10	−1.1	1	15	11 ^{108}In	GSI		79Pl06	*
*^{108}Ag−u	$M-A$=–87480(34) keV for mixture gs+m at 109.466 keV											Nub269	**
*^{108}Ag−^{133}Cs$_{.812}$	D_M=–17159(77) μu for mixture gs+m at 109.466 keV; $M-A$=–87497(72) keV											Nub269	**
*^{108}In−u	$M-A$=–84078(28) keV for mixture gs+m at 29.75 keV											Nub269	**
*^{108}Sn−^{107}Sn	From ^{107}Sn/^{108}Sn=0.99076701(70)											AHW	**
*^{108}I(α)^{104}Sb	F : Same authors say: Consistent with new value, if recalibrated											94Pa12	**
*^{108}Ru(β^-)^{108}Rh	E_{β^-}=1150(100) to 1^+ level at 164.98 keV											Ens08a	**
*^{108}Rh^m(β^-)^{108}Pd	E_{β^-}=1570(50) to (4^+,5^+,6^+) level at 2863.70 keV											Ens08a	**
*^{108}Rh^m(β^-)^{108}Pd	E_{β^-}=1970(100) to 4^+ level at 2540.2 keV											Ens08a	**
*^{108}In(β^+)^{108}Cd	E_{β^+}=1290(80) to 6^+ level at 2807.81 keV, and E_{β^+}=3500(50) from ^{108}In^m											Ens08a	**
*	at 29.75 to 2^+ level at 632.986 keV											Nub269	**
*^{108}In(β^+)^{108}Cd	E_{β^+}=1887(28) to 4^+ level at 2239.35 keV, and E_{β^+}=3494(14) from ^{108}In^m											Ens08a	**
*	at 29.75 to 2^+ level at 632.986 keV											Nub269	**
*^{108}Cd(p,n)^{108}In	Q(not T, PrvCom Fi)=–6191(8), –6244(9), errors statistical only,											AHW	**
*	to 3^+ levels at respectively, 198.36 and 266.06 keV											Ens08a	**
*^{108}Sn(β^+)^{108}In	p^+ =35(6)×10^{-4} to 1^+ level at 698.85 keV											Ens08a	**
^{109}Nb−u		−60784	376	−60960	210	−0.3	o			GT1	1.5	04Ma.A	
		−60859	185			−0.2	1	21	21 ^{109}Nb	GT3	2.5	16Kn03	
^{109}Nb−^{133}Cs$_{.820}$		16546	239	16570	210	0.1	1	79	79 ^{109}Nb	JY1	1.0	24Hu16	
^{109}Mo−^{97}Zr$_{1.124}$		28515	12				2			JY1	1.0	06Ha03	
^{109}Mo−u		−71552	247	−71562	12	−0.0	U			GT1	1.5	04Ma.A	
^{109}Rh−^{120}Sn$_{.908}$		−2463.6	7.2	−2450	4	1.9	1	37	36 ^{109}Rh	JY1	1.0	07Ha20	
C_8 H_{13}−^{109}Ag		196972.1	3.8	196969.1	0.4	−0.3	U			M16	2.5	63Da10	
		196972	6			−0.3	U			R12	1.5	83De51	
C_6 ^{13}C O H_8−^{109}Ag		156110	16	156113.4	0.4	0.1	U			R12	1.5	83De51	
C_6 N O H_7−^{109}Ag		148006	16	148007.5	0.4	0.1	U			R12	1.5	83De51	
^{109}Ag−^{133}Cs$_{.820}$		−17766	83	−17714.3	0.4	0.6	U			MA8	1.0	08Br.A	*
^{109}Cd−^{85}Rb$_{1.282}$		18072.9	1.9	18072.3	1.6	−0.3	1	72	72 ^{109}Cd	MA8	1.0	09Br09	
^{109}In−u		−92885.2	36.5	−92852.6	2.8	0.9	U			TR1	1.0	20Ho03	
^{109}Sn−u		−88747	30	−88707	9	1.3	U			GS2	1.0	05Li24	
^{109}Sb−^{87}Rb$_{1.253}$		31937.9	5.9	31938	6	0.0	1	92	92 ^{109}Sb	JY1	1.0	09El07	
^{109}Sb−^{133}Cs$_{.820}$		−4360	19	−4329	6	1.6	U			SH1	1.0	07Ma92	
^{109}Te−^{87}Rb$_{1.253}$		41101.6	6.4	41101	5	−0.1	1	54	54 ^{109}Te	JY1	1.0	09El07	
^{109}Te−^{133}Cs$_{.820}$		4835.6	8.3	4834	5	−0.2	1	32	32 ^{109}Te	SH1	1.0	07Ma92	
^{109}Tc−^{105}Ru$_{1.038}$		16014.3	10.0				2			JY1	1.0	07Ha20	
^{109}Ru−^{105}Ru$_{1.038}$		9083.9	9.2				2			JY1	1.0	07Ha20	
^{109}Ag−^{110}Cd$_{.991}$		875.974	0.048	875.97	0.05	−0.0	1	100	97 ^{109}Ag	JY2	1.0	25Ru03	
^{109}Ag−^{107}Ag		−335	28	−332.3	1.3	0.1	U			R12	1.5	83De51	
^{109}Te(α)^{105}Sn		3197.6	30.	3199	6	0.0	U					65Ma12	
		3197.6	15.			0.1	1	13	7 ^{109}Te			79Sc22	
		3225.6	4.			−6.8	C					91He21	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{109}I(α)^{105}Sb		3918.1	20.8				3			ORa		07Ma35	
^{109}Xe(α)^{105}Te		4217.0	7.3				3					06Li41	*
^{109}Ag(p,t)^{107}Ag		−7995	15	−7970.4	1.2	1.6	U			Min		75Ku14	*
^{108}Pd(n,γ)^{109}Pd		6153.8	0.3	6153.58	0.15	−0.7	–			ILn		80Ca02	Z
		6153.54	0.17			0.3	–			Bdn		06Fi.A	
^{108}Pd(d,p)^{109}Pd		3936	30	3929.02	0.15	−0.2	U			Pit		64Co11	
^{108}Pd(n,γ)^{109}Pd	ave.	6153.60	0.15	6153.58	0.15	−0.1	1	100	73 ^{109}Pd			average	
^{109}Ag(γ,n)^{108}Ag		−9196	26	−9180.8	1.2	0.6	U			Phi		60Ge01	
^{109}Ag(d,t)^{108}Ag		−2947	30	−2923.5	1.2	0.8	U			Pit		64Co11	
^{108}Cd(^{3}He,d)^{109}In−^{110}Cd()^{111}In		−806.5	2.6	−807.3	2.4	−0.3	1	89	87 ^{109}In			80Ta07	
^{109}Te(εp)^{108}Sn		7140	60	7065	7	−1.2	U					73Bo20	
^{109}I(p)^{108}Te		819	5	820	4	0.2	o					84Fa04	
		819.6	2.0			0.3	o					92He.A	
		820.2	4.0				2					95Ho26	
^{109}Tc(β^-)^{109}Ru		6315	70	6456	13	2.0	U			Bwg		89Gr23	
^{109}Ru(β^-)^{109}Rh		4160	65	4260	10	1.5	U			Bwg		89Gr23	
^{109}Rh(β^-)^{109}Pd		2577	50	2607	4	0.6	U					78Ka10	*
^{109}Pd(β^-)^{109}Ag		1116	2	1112.6	1.1	−1.7	1	29	27 ^{109}Pd			62Br15	*
^{109}Cd(ε)^{109}Ag		182	3	214.6	1.5	10.9	C					68Go.A	*
		214	3			0.2	1	26	25 ^{109}Cd			Averag	*
^{109}In(β^+)^{109}Cd		2015	8	2012.7	2.9	−0.3	–					62No06	*
		2030	15			−1.2	–					71Ba08	*
	ave.	2018	7			−0.8	1	17	13 ^{109}In			average	
^{109}Sn(β^+)^{109}In		4120	50	3861	8	−5.2	B					70Sh05	
^{109}Sb(β^+)^{109}Sn		6380	16	6379	9	−0.0	1	30	22 ^{109}Sn			82Jo03	*
*^{109}Ag−^{133}Cs$_{.820}$	D_M=–17719(78) μu for mixture gs+m at 88.0337 keV; $M-A$=–88723(73) keV											Nub269	**
*^{109}Xe(α)^{105}Te	Also E_α=3918(9) keV to 150(13) level											06Li41	**
*^{109}Ag(p,t)^{107}Ag	Recalibrated with (p,t) results on ^{104}Pd, ^{105}Pd, ^{106}Pd and ^{108}Pd											AHW	**
*^{109}Rh(β^-)^{109}Pd	E_{β^-}=2250(50) to 5/2$^+$ level at 326.869 keV											Ens062	**
*^{109}Pd(β^-)^{109}Ag	E_{β^-}=1028(2) to ^{109}Ag^m at 88.0337 keV											Nub269	**
*^{109}Cd(ε)^{109}Ag	IBE=68(3) gives 94(3) to ^{109}Ag^m at 88.0337 keV											Nub269	**
*^{109}Cd(ε)^{109}Ag	From average LM/K=0.2265(0.0026) → Q_{β^+}=126(3); recalculated Q											AHW	**
*	to ^{109}Ag^m at 88.0337 keV											Nub269	**
*	LMN/K=0.228(0.003)											65Le06	**
*	L/K=0.195(0.005) → LMN/K=0.258(0.006) → Q_{β^+}=109(5) not used											65Le06	**
*	LMN/K=0.226(0.003)											70Go39	**
*^{109}In(β^+)^{109}Cd	E_{β^+}=790(8) 805(15) respectively, to 7/2$^+$ level at 203.30 keV											Ens062	**
*^{109}Sb(β^+)^{109}Sn	E_{β^+}=4416(21) to 3/2$^+$ level at 925.6 keV, and other E_{β^+}											Ens062	**
^{110}Nb−u		−56157	360	−56030#	220#	0.1	D			GT3	2.5	16Kn03	*
^{110}Mo−^{97}Zr$_{1.134}$		31685	26				2			JY1	1.0	06Ha03	
^{110}Mo−u		−69544	268	−69282	26	0.7	U			GT1	1.5	04Ma.A	
^{110}Ru−u		−85897	78	−85961.7	0.6	−0.8	U			JY0	1.0	03Ko.A	
^{110}Rh−u		−88738	79	−88787.7	0.8	−0.6	U			JY0	1.0	03Ko.A	*
^{110}Rh−^{120}Sn$_{.917}$		912	14	892.7	1.2	−1.4	U			JY1	1.0	07Ha20	*
^{110}Rh^m−^{133}Cs$_{.827}$		−10552.6	2.0	−10553.8	0.7	−0.6	1	13	13 ^{110}Rh^m	JY2	1.0	23Hu01	
C$_8$ H$_{14}$−^{110}Pd		204389	9	204377.5	0.6	−0.5	U			M16	2.5	63Da10	
		204380	20			−0.1	U			R12	1.5	83De51	
C$_7$ ^{13}C H$_{13}$−^{110}Pd		199913	20	199907.3	0.6	−0.2	U			R12	1.5	83De51	
C$_6$ N O H$_8$−^{110}Pd		155418	17	155415.9	0.6	−0.1	U			R12	1.5	83De51	
C$_5$ ^{13}C N O H$_7$−^{110}Pd		150946	17	150945.7	0.6	−0.0	U			R12	1.5	83De51	
C$_6$ O$_2$ H$_6$−^{110}Pd		131609	18	131606.5	0.6	−0.1	U			R12	1.5	83De51	
^{110}Pd−u		−94829.7	1.5	−94827.0	0.6	1.8	–			MA8	1.0	12Fi01	
		−94829.5	1.7			1.0	–			TG1	1.5	12Sm01	
		−94830.9	3.0			0.9	–			TG1	1.5	12Sm01	
	ave.	−94829.7	1.2			2.2	1	25	25 ^{110}Pd			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$C_8\ H_{14}-{}^{110}Cd$		206548.4	4.6	206543.0	0.4	−0.5	U			M16	2.5	63Da10	
		206550	45			−0.1	U			R12	1.5	83De51	
		206569	13			−1.3	U			R12	1.5	83De51	
$C_7\ {}^{13}C\ H_{13}-{}^{110}Cd$		202093	14	202072.8	0.4	−1.0	U			R12	1.5	83De51	
		202053	28			0.5	U			R12	1.5	83De51	
$C_7\ O\ H_{10}-{}^{110}Cd$		170156	16	170157.5	0.4	0.1	U			R12	1.5	83De51	
$C_6\ N\ O\ H_8-{}^{110}Cd$		157614	17	157581.4	0.4	−1.3	U			R12	1.5	83De51	
$C_5\ {}^{13}C\ N\ O\ H_7-{}^{110}Cd$		153131	17	153111.3	0.4	−0.8	U			R12	1.5	83De51	
$C_6\ O_2\ H_6-{}^{110}Cd$		133801	18	133772.0	0.4	−1.1	U			R12	1.5	83De51	
$C_9\ H_2-{}^{110}Cd$		112661	19	112642.6	0.4	−0.6	U			R12	1.5	83De51	
$^{110}Cd-u$		−96993.6	1.5	−96992.6	0.4	0.7	–			MA8	1.0	12Fi01	
		−96997.0	1.5			2.0	–			TG1	1.5	12Sm01	
		−96992.2	2.4			−0.1	–			TG1	1.5	12Sm01	
	ave.	−96994.4	1.2			1.5	1	11	11 ^{110}Cd			average	
$^{110}In-u$		−92898	36	−92829	12	1.9	U			GS2	1.0	05Li24	*
$^{110}Sn-u$		−92189	30	−92155	15	1.1	2			GS2	1.0	05Li24	
$^{110}Sb-{}^{87}Rb_{1.264}$		31650.1	6.4				2			JY1	1.0	09El07	
$^{110}Te-{}^{133}Cs_{.827}$		643.8	7.7	650	7	0.8	1	84	84 ^{110}Te	SH1	1.0	07Ma92	
$^{110}Tc-{}^{105}Ru_{1.048}$		20424.0	9.8				2			JY1	1.0	07Ha20	
$^{110}Ru-{}^{105}Ru_{1.048}$		10719.5	9.3	10721.9	1.5	0.3	U			JY1	1.0	07Ha20	
$^{110}Cd\ {}^{35}Cl-{}^{108}Cd\ {}^{37}Cl$		1764	5	1773.8	1.2	0.5	U			H11	4.0	63Bi12	
$^{110}Ru-{}^{110}Pd$		8865.389	0.043				2			CP2	1.0	24Po14	
$^{110}Rh-{}^{110}Rh^m$		−40.8	1.7	−42.6	0.6	−1.1	1	14	9 ^{110}Rh	JY2	1.0	23Hu01	
$^{110}Rh-{}^{110}Pd$		6039.19	0.54	6039.4	0.5	0.3	1	91	91 ^{110}Rh	CP2	1.0	25Li19	
$^{110}Rh^m-{}^{110}Pd$		6082.07	0.43	6082.0	0.4	−0.1	1	91	82 $^{110}Rh^m$	CP2	1.0	25Li19	
$^{110}Pd-{}^{110}Cd$		2166.24	0.69	2165.5	0.6	−1.0	–			MA8	1.0	12Fi01	
		2166.2	1.3			−0.3	–			TG1	1.5	12Sm01	
	ave.	2166.2	0.7			−1.1	1	75	65 ^{110}Pd			average	
$^{110}Pd-{}^{108}Pd$		1288	35	1281.0	1.3	−0.1	U			R12	1.5	83De51	
$^{110}Cd-{}^{108}Cd$		−1219	34	−1176.3	1.2	0.8	U			R12	1.5	83De51	
$^{110}Te(\alpha)^{106}Sn$		2723.2	15.6	2698	8	−1.6	1	24	16 ^{110}Te			81Sc17	
$^{110}I(\alpha)^{106}Sb$		3574.2	10.	3580	60	0.1	3					81Sc17	
		3586.7	5.			−0.1	3					91He21	
$^{110}Xe(\alpha)^{106}Te$		3878.3	30.	3872	9	−0.2	4					81Sc17	
		3886.6	15.			−0.9	4					92He.A	
		3871.0	30.			0.0	4					02Ma19	
		3857.5	19.7			0.7	4			Jya		07Sa36	
		3860.7	20.8			0.6	4					16Ca33	
$^{110}Pd(p,t)^{108}Pd$		−6495	15	−6467.6	1.2	1.8	U			Min		75Ku14	*
$^{110}Pd(d,{}^3He)^{109}Rh$		−5134	5	−5127	4	1.4	1	65	64 ^{109}Rh	VUn		87Ka29	
$^{110}Pd(t,\alpha)^{109}Rh$		9206	25	9193	4	−0.5	U			LAl		82Fl09	
$^{109}Ag(n,\gamma)^{110}Ag$		6809.2	0.1				2					81Bo.B	
		6808.20	0.16	6809.20	0.10	6.2	C			Bdn		06Fi.A	
$^{109}Ag(d,p)^{110}Ag$		4590	5	4584.63	0.10	−1.1	U			MIT		64Sp12	
$^{110}Cd(d,t)^{109}Cd$		−3664	30	−3657.8	1.5	0.2	U			Pit		64Ro17	
$^{110}Tc(\beta^-)^{110}Ru$		6680	120	9037	9	19.6	C			Jyv		89Jo.A	
		9021	55			0.3	U			Jyv		00Kr.A	
$^{110}Ru(\beta^-)^{110}Rh$		2810	50	2632.4	0.5	−3.6	B			Jyv		91Jo11	*
$^{110}Rh(\beta^-)^{110}Pd$		5500	500	5625.6	0.5	0.3	U					63Ka21	
		5400	100			2.3	U					70Pi01	*
		5510	19			6.1	B			Bwg		00Kr.A	
$^{110}Ag(\beta^-)^{110}Cd$		2891.4	3.0	2891.21	0.11	−0.1	U					63Da03	*
		2892.9	2.0			−0.8	U					67Mo12	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{110}In(β^+)^{110}Cd	3928	20	3878	12	−2.5	2						51Mc11	*
	3868	20			0.5	2						53Bl44	*
	3838	20			2.0	2						62Ka08	*
^{110}Sb(β^+)^{110}Sn	8750	200	8392	15	−1.8	U						72Mi26	*
	9085	100			−6.9	B						72Si28	*
*^{110}Nb−u	Trends in Mass Surface TMS suggest ^{110}Nb 120 keV less bound											HWJ266	**
*^{110}Rh−u	$M-A$=–82639(73) keV for mixture gs+m at 39.7(0.6) keV											Nub269	**
*^{110}Rh−^{120}Sn$_{.917}$	D_M=933.3(7.2) μu for mixture gs+m at 39.7(0.6) keV; $M-A$=–82668.1(6.8) keV											Nub269	**
*^{110}In−u	$M-A$=–86503(28) keV for mixture gs+m at 62.08 keV											Nub269	**
*^{110}Pd(p,t)^{108}Pd	Recalibrated with (p,t) results on ^{104}Pd, ^{105}Pd, ^{106}Pd and ^{108}Pd											AHW	**
*^{110}Ru(β^-)^{110}Rh	E_{β^-}=2700(50) to 1^+ level at 112.19 keV											Ens126	**
*^{110}Rh(β^-)^{110}Pd	E_{β^-}=2600(100) to (4^+) levels at 2790.64 and 2805.03 keV											Ens126	**
*^{110}Ag(β^-)^{110}Cd	E_{β^-}=529(3) from ^{110}Ag^n at 117.59 to 6^+ level at 2479.9339 keV											Ens126	**
*^{110}Ag(β^-)^{110}Cd	E_{β^-}=2891(4); and 531(2) from ^{110}Ag^n at 117.59 to 6^+ level at 2479.9339 keV											Ens126	**
*^{110}In(β^+)^{110}Cd	E_{β^+}=2310(20) 2250(20) 2220(20) respectively, from ^{110}In^m at 62.08(0.04) keV											Nub266	**
*	to 2^+ level at 657.7623 keV											Ens126	**
*^{110}Sb(β^+)^{110}Sn	E_{β^+}=6500(200) 6850(100) respectively, to 2^+ level at 1211.72; and other E_{β^+}											Ens126	**
^{111}Mo−u	−64348	279	−64340	5	0.0	U				GT1	1.5	04Ma.A	
^{111}Mo−^{133}Cs$_{.835}$	14599.7	9.3	14607	5	0.8	1	32	32	^{111}Mo	JY2	1.0	24Hu16	
^{111}Ru−u	−82302	79	−82425	5	−1.6	U				JY0	1.0	03Ko.A	
^{111}Ru−^{111}Tc	−8335.4	5.0	−8334	4	0.3	1	73	37	^{111}Tc	RI1	1.0	23Ho11	
^{111}Rh−u	−88282	79	−88354	5	−0.9	U				JY0	1.0	03Ko.A	
^{111}Rh−^{120}Sn$_{.925}$	2105.8	7.3	2109	5	0.4	1	39	38	^{111}Rh	JY1	1.0	07Ha20	
^{111}Rh−^{111}Tc	−14269.5	8.1	−14263	5	0.9	1	45	23	^{111}Tc	RI1	1.0	23Ho11	
^{111}Rh ^{40}Ar−^{151}Pr	−54274.9	7.1	−54283	5	−1.1	1	41	40	^{111}Rh	RI1	1.0	24Ki16	*
^{111}Ag−u	−94741	51	−94703.2	1.6	0.7	U				GS2	1.0	05Li24	*
C_8 H_{15}−^{111}Cd	213184.4	3.9	213191.7	0.4	0.8	U				M16	2.5	63Da10	
	213197	40			−0.1	U				R12	1.5	83De51	
C_7 ^{13}C H_{14}−^{111}Cd	208719	19	208721.5	0.4	0.1	U				R12	1.5	83De51	
C_7 O H_{11}−^{111}Cd	176814	16	176806.2	0.4	−0.3	U				R12	1.5	83De51	
C_9 H_3−^{111}Cd	119317	18	119291.4	0.4	−0.9	U				R12	1.5	83De51	
C_8 N H−^{111}Cd	106723	17	106715.3	0.4	−0.3	U				R12	1.5	83De51	
^{111}Cd−u	−95774	30	−95816.3	0.4	−1.4	U				GS2	1.0	05Li24	*
^{111}Sb−u	−86837	30	−86782	10	1.8	U				GS2	1.0	05Li24	
^{111}Sb−^{133}Cs$_{.835}$	−7834.2	9.5				2				SH1	1.0	07Ma92	
^{111}Te−^{133}Cs$_{.835}$	−51.8	6.9				2				SH1	1.0	07Ma92	
^{111}I−^{87}Rb$_{1.276}$	46150.4	6.1	46155	5	0.7	1	70	70	^{111}I	JY1	1.0	09El07	
^{111}I−^{133}Cs$_{.835}$	9197	19	9217	5	1.0	U				SH1	1.0	07Ma92	
^{111}Tc−^{105}Ru$_{1.057}$	23412	11	23422	5	0.9	1	19	18	^{111}Tc	JY1	1.0	07Ha20	
^{111}Ru−^{105}Ru$_{1.057}$	15080.6	10.0	15088	5	0.8	1	22	21	^{111}Ru	JY1	1.0	07Ha20	
^{110}Cd H−^{111}Cd	6638	18	6648.73	0.18	0.4	U				R12	1.5	83De51	
^{111}Mo−^{111}Tc	9753.0	7.3	9752	3	−0.2	–				JY1	1.0	11Ha48	*
	9752.6	4.1			−0.3	–				RI1	1.0	23Ho11	
ave.	9753	4			−0.3	1	90	68	^{111}Mo			average	
^{111}Pd−^{111}Tc	−18238	12	−18218	5	1.7	U				RI1	1.0	23Ho11	
^{111}Ag−^{111}Tc	−20579	13	−20612	5	−2.5	U				RI1	1.0	23Ho11	
^{111}In−^{111}Cd	920.71	0.19	920.71	0.19	0.0	1	100	100	^{111}In	JY2	1.0	22Ge11	
^{111}Cd−^{110}Cd	1180	11	1176.31	0.18	−0.1	U				M16	2.5	63Da10	
	1208	34			−0.6	U				R12	1.5	83De51	
^{111}Cd H−^{110}Cd	8994	35	9001.34	0.18	0.1	U				R12	1.5	83De51	
^{111}I(α)^{107}Sb	3270.1	10.	3275	5	0.4	–						79Sc22	
	3293.0	10.			−1.8	–						92He.A	
ave.	3281	7			−0.9	1	50	30	^{111}I			average	
^{111}Xe(α)^{107}Te	3693.3	25.9	3710	8	0.6	o						79Sc22	*
	3707	31			0.1	2						81Sc17	*
	3716.1	10.4			−0.6	2						91He21	
	3694.2	16.6			0.9	2						20Ca01	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{110}Pd(n,γ)^{111}Pd		5726.3	0.4				2			Bdn		06Fi.A	
^{110}Cd(n,γ)^{111}Cd		6980	100	6975.60	0.17	−0.0	U					61Ja21	
		6975.5	0.5			0.2	–					86Ba72	
		6975.9	0.2			−1.5	–					90Ne.B	
		6975.1	0.4			1.2	–			Bdn		06Fi.A	
^{111}Cd(γ,n)^{110}Cd		−6975	3	−6975.60	0.17	−0.2	U			McM		79Ba06	
^{110}Cd(d,p)^{111}Cd		4740	30	4751.03	0.17	0.4	U			Pit		64Ro17	
		4750.68	0.88			0.4	U			Rez		90Pi05	*
^{111}Cd(d,t)^{110}Cd		−745	30	−718.37	0.17	0.9	U			Pit		64Co11	
^{110}Cd(n,γ)^{111}Cd	ave.	6975.71	0.17	6975.60	0.17	−0.7	1	96	75 ^{110}Cd			average	
^{111}Te(εp)^{110}Sn		5070	70	4966	15	−1.5	U					68Ba53	
^{111}Tc(β^-)^{111}Ru		7480	80	7763	4	3.5	B			Jyv		96Kl.A	
		7449	80			3.9	B			Jyv		00Kr.A	
^{111}Ru(β^-)^{111}Rh		5039	50	5523	5	9.7	C			Jyv		00Kr.A	
^{111}Rh(β^-)^{111}Pd		3640	50	3684	4	0.9	U			Jyv		00Kr.A	
		3650	33			1.0	U			Bwg		00Kr.A	
^{111}Pd(β^-)^{111}Ag		2210	100	2229.7	1.6	0.2	U					52Mc34	*
		2190	50			0.8	U					57Kn.A	*
		2160	100			0.7	U					60Pr07	*
^{111}Ag(β^-)^{111}Cd		1028	3	1036.8	1.4	2.9	B					67Le06	
		1035	2			0.9	2					71Na02	
		1038.6	2.			−0.9	2					77Re12	
^{111}Cd(p,n)^{111}In		−1635	20	−1639.99	0.18	−0.2	U			Oak		74Ki02	
^{111}Sn(β^+)^{111}In		2530	30	2456	5	−2.5	U					51Mc11	
^{111}Sb(β^+)^{111}Sn		4470	50	5102	10	12.6	B					72Si28	*
*^{111}Rh ^{40}Ar−^{151}Pr	Reported $M-A$=–117336(24) keV, author confirmed mistake in the unc.											24Ki16	**
*^{111}Ag−u	$M-A$=–88221(44) keV for mixture gs+m at 59.82 keV											Nub269	**
*^{111}Cd−u	$M-A$=–88817(28) keV for ^{111}Cd^m 11/2$^-$ at 396.214 keV											Nub269	**
*^{111}Mo−^{111}Tc	Taken as low-spin isomer (see also ^{102}Y and ^{114}Tc doublets in same work)											GAu	**
*^{111}Xe(α)^{107}Te	other E_α=3463(50) to level at 90.3 keV; doublet											79Sc22	**
*^{111}Xe(α)^{107}Te	E_α=3580(30), 3480(30) to ground state, 90.3 keV level											81Sc17	**
*^{111}Xe(α)^{107}Te	E_α=3562(16), 3474(16) to ground state, 90.3 keV level											20Ca01	**
*^{110}Cd(d,p)^{111}Cd	Estimated systematic error 0.5 added to statistical error 0.73 keV											AHW	**
*^{111}Pd(β^-)^{111}Ag	Q_{β^-}=2150(100) 2130(50) 2100(100) respectively, to ^{111}Ag^m at 59.82 keV											Nub269	**
*^{111}Sb(β^+)^{111}Sn	E_{β^+}=3290(50) to 5/2$^+$ level at 154.48 keV											Ens095	**
^{112}Mo−u		−60713	850	−61693	8	−1.2	U			MT1	1.0	24Wa19	
^{112}Mo−^{112}Tc		8362.0	6.8				2			RI1	1.0	23Ho11	
^{112}Mo−^{133}Cs$_{.842}$		17931	133	17916	8	−0.1	U			JY2	1.0	24Hu16	
^{112}Tc−^{102}Ru$_{1.098}$		34976.0	5.9	34979	5	0.5	1	68	67 ^{112}Tc	JY1	1.0	07Ha20	
^{112}Ru−u		−81033	78	−81183	5	−1.9	U			JY0	1.0	03Ko.A	
^{112}Ru ^{40}Ar−^{152}Pr		−50325.3	7.7	−50337	5	−1.5	1	45	37 ^{112}Ru	RI1	1.0	24Ki16	*
^{112}Ru−^{112}Tc		−11130.6	5.6	−11128	5	0.5	1	71	38 ^{112}Ru	RI1	1.0	23Ho11	
^{112}Rh−u		−85346	41	−85457.3	1.6	−2.7	U			JY0	1.0	03Ko.A	*
^{112}Rh−^{85}Rb$_{1.318}$		30803.1	1.9	30803.8	1.6	0.4	1	73	73 ^{112}Rh	JY2	1.0	23Hu01	
^{112}Rh ^{40}Ar−^{152}Pr		−54575.8	20.8	−54611	3	−1.7	U			RI1	1.0	24Ki16	*
^{112}Rh^m−^{133}Cs$_{.842}$		−5807.3	2.1	−5805.0	0.7	1.1	1	11	11 ^{112}Rh^m	JY2	1.0	23Hu01	
^{112}Ag−^{133}Cs$_{.842}$		−13342.0	2.6	−13343.1	0.3	−0.4	U			MA8	1.0	10Br02	
C$_8$ H$_{16}$−^{112}Cd		222445.3	3.9	222436.67	0.25	−0.9	U			M16	2.5	63Da10	
C$_7$ O H$_{12}$−^{112}Cd		186063	16	186051.16	0.25	−0.5	U			R12	1.5	83De51	
C$_9$ H$_4$−^{112}Cd		128541	19	128536.29	0.25	−0.2	U			R12	1.5	83De51	
		128550	10			−0.9	U			R12	1.5	83De51	
C$_8$ N H$_2$−^{112}Cd		115979	14	115960.23	0.25	−0.9	U			R12	1.5	83De51	
^{112}In−u		−94366	58	−94461	5	−1.6	U			GS2	1.0	05Li24	*
C$_8$ H$_{16}$−^{112}Sn		220384	9	220375.67	0.28	−0.4	U			M16	2.5	63Da10	
		220385	8			−0.8	U			R13	1.5	83De51	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{112}Sn−^{86}Kr$_{1.302}$		21210.3	2.5	21209.82	0.28	−0.2	U			JY1	1.0	11Ha48	
^{112}Sn−^{112}Tc		−25130	15	−25120	5	0.7	U			RI1	1.0	23Ho11	
^{112}Sb−u		−87597	30	−87600	19	−0.1	2			GS2	1.0	05Li24	
^{112}Te−^{133}Cs$_{.842}$		−3662.7	9.0				2			SH1	1.0	07Ma92	
^{112}I−^{133}Cs$_{.842}$		7614	11				2			SH1	1.0	07Ma92	
^{112}Rh−^{120}Sn$_{.933}$		5801	14	5787.8	1.8	−0.9	U			JY1	1.0	07Ha20	*
^{112}Rh−^{112}Tc		−15394	13	−15402	5	−0.6	U			RI1	1.0	23Ho11	
^{112}Pd−^{120}Sn$_{.933}$		−1423.7	7.4	−1425	7	−0.1	1	89	89 ^{112}Pd	JY1	1.0	07Ha20	
^{112}Sn−^{120}Sn$_{.933}$		−3930.2	1.9	−3930.0	0.9	0.1	1	23	22 ^{120}Sn	JY1	1.0	11Ha48	
^{112}Ru−^{105}Ru$_{1.067}$		17242.5	9.9	17253	5	1.1	1	27	25 ^{112}Ru	JY1	1.0	07Ha20	
^{112}Cd ^{35}Cl−^{110}Cd ^{37}Cl		2701	2	2706.5	0.3	0.7	U			H11	4.0	63Bi12	
^{111}Cd H−^{112}Cd		9255	20	9244.9	0.3	−0.3	U			R12	1.5	83De51	
^{112}Rh^{m}−^{112}Sn		9761.10	0.72	9760.7	0.7	−0.5	1	87	85 ^{112}Rh^{m}	CP2	1.0	25Li19	
^{112}Rh^{m}−^{112}Rh		41.2	3.0	42.9	1.7	0.6	1	31	27 ^{112}Rh	JY2	1.0	23Hu01	
^{112}Pd−^{112}Tc		−22613	25	−22615	9	−0.1	U			RI1	1.0	23Ho11	
^{112}Ag−^{112}Cd		4283.61	0.24				2			CP2	1.0	22Ga27	
^{112}Sn−^{112}Cd		2061.01	0.17	2061.00	0.14	−0.0	–			JY1	1.0	09Ra11	
		2061.03	0.27			−0.1	–			CP2	1.0	22Ga27	
	ave.	2061.02	0.14			−0.1	1	97	86 ^{112}Sn			average	
^{112}Cd−^{110}Cd H		−8060	40	−8068.6	0.3	−0.1	U			R12	1.5	83De51	
^{112}Cd−^{111}Cd		−1419	11	−1419.9	0.3	−0.0	U			M16	2.5	63Da10	
		−1410	42			−0.2	U			R12	1.5	83De51	
^{112}Cd−^{110}Cd		−238	39	−243.6	0.3	−0.1	U			R12	1.5	83De51	
^{112}Cd H−^{111}Cd		6402	35	6405.1	0.3	0.1	U			R12	1.5	83De51	
^{112}I(α)^{108}Sb		2986.9	31.1	2957	12	−0.5	U					81Sc17	
^{112}Xe(α)^{108}Te		3329.1	20.	3330	6	0.1	2					81Sc17	
		3308.4	15.6			1.4	2					92He.A	
		3335.4	7.			−0.7	2					94Pa11	
^{112}Sn(^{3}He,^{6}He)^{109}Sn		−8686	9	−8686	8	0.0	1	78	78 ^{109}Sn	MSU		78Pa11	
^{110}Pd(t,p)^{112}Pd		5659	20	5651	7	−0.4	1	11	11 ^{112}Pd	LAl		72Ca10	
^{112}Cd(^{14}C,^{16}O)^{110}Pd		5543	29	5512.8	0.6	−1.0	U			LAl		84Co19	
^{110}Cd(t,p)^{112}Cd		7888	20	7887.7	0.3	−0.0	U			Ald		67Hi01	
^{112}Cd(p,t)^{110}Cd		−7891	5	−7887.7	0.3	0.7	U			Min		73Oo01	
^{112}Sn(p,t)^{110}Sn		−10485	15	−10474	14	0.7	R			Roc		70Fl08	
^{111}Cd(n,γ)^{112}Cd		9460	50	9393.94	0.28	−1.3	U					61Ja21	
		9394.3	0.3			−1.2	1	88	79 ^{111}Cd	ILn		97Dr03	
^{112}Cd(γ,n)^{111}Cd		−9403	5	−9393.94	0.28	1.8	U			McM		79Ba06	
^{111}Cd(d,p)^{112}Cd		7183	30	7169.38	0.28	−0.5	U			Pit		64Co11	
		7170	10			−0.1	U			Yal		67Ba15	
		7171	5			−0.3	U			MIT		67Sp09	
^{112}Cd(d,t)^{111}Cd		−3129	30	−3136.71	0.28	−0.3	U			Pit		64Ro17	
^{112}Sn(d,^{3}He)^{111}In		−2050	50	−2055.9	0.4	−0.1	U			Sac		69Co03	
^{112}Sn(p,d)^{111}Sn		−8574	15	−8563	5	0.7	2			Har		70Ca01	
^{112}Sn(d,t)^{111}Sn		−4529.0	5.7	−4531	5	−0.3	2			SPa		75Be09	
^{112}Cs(p)^{111}Xe		814.3	7.	816	4	0.3	3					94Pa12	
		817.3	5.			−0.2	3					12Wa10	
^{112}Tc(β^-)^{112}Ru		6060	130	10366	4	33.1	C			Jyv		89Jo.A	
		9484	100			8.8	C			Jyv		00Kr.A	
^{112}Ru(β^-)^{112}Rh		4520	80	3981	5	−6.7	B			Jyv		91Jo11	*
^{112}Rh(β^-)^{112}Pd		6200	500	6718	7	1.0	U			Jyv		88Ay02	
		6573	54			2.7	U			Bwg		00Kr.A	
^{112}Rh^{m}(β^-)^{112}Pd		6929	56	6758	7	−3.0	B			Bwg		00Kr.A	
^{112}Pd(β^-)^{112}Ag		299	20	263	7	−1.8	U					55Nu11	*
^{112}Ag(β^-)^{112}Cd		4057	20	3990.16	0.22	−3.3	C					57Je.A	*
		3940	40			1.3	U					62In01	*
^{112}In(β^+)^{112}Cd		2582	20	2585	4	0.1	U					62Ru05	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{112}Cd(p,n)^{112}In	−3399.3	20.	−3367	4	1.6	U			Oak		64Jo11	
	−3376	6			1.5	1	50	50 ^{112}In	Tky		80Ad04	
^{112}In(β^-)^{112}Sn	656	6	665	4	1.5	1	50	50 ^{112}In			53Bl44	
^{112}Sb(β^+)^{112}Sn	7530	100	7056	18	−4.7	B					72Mi27	*
	7029	50			0.5	R					72Si28	*
	7062	26			−0.2	R					82Jo03	*
^{112}Sn(p,n)^{112}Sb	−7995	55	−7838	18	2.8	U			VUn		76Ka19	
*^{112}Ru ^{40}Ar−^{152}Pr	Reported $M-A$=–110660(21) keV, author confirmed mistake in the unc.										24Ki16	**
*^{112}Rh−u	Average of 2 values; $M-A$=–79479(36) keV for mixture gs+m at 40(2) keV										Nub269	**
*^{112}Rh ^{40}Ar−^{152}Pr	D_M=–54555.2(8.4) μu, $M-A$=–114600.0(12.5) keV for mixture of ground state and m										Nub269	**
*^{112}Rh ^{40}Ar−^{152}Pr	at 40.0(1.6) keV; reported $M-A$=–114600(21) keV, author confirmed										Nub269	**
*^{112}Rh ^{40}Ar−^{152}Pr	mistake in the uncertainty										Nub269	**
*^{112}In−u	$M-A$=–87823(30) keV for mixture gs+m at 156.592 keV										Nub269	**
*^{112}Rh−^{120}Sn$_{.933}$	D_M=5822.6(7.4) μu for mixture gs+m at 40.0(1.6) keV; $M-A$=–79571.2(7.0) keV										Nub269	**
*^{112}Rh−^{120}Sn$_{.933}$	D_M=–15374(19) μu for mixture gs+m at 40.0(1.6) keV; $M-A$=–79579(19) keV										Nub269	**
*^{112}Ru(β^-)^{112}Rh	E_{β^-}=4190(80) to 1^+ level at 327.03 keV										Ens14c	**
*^{112}Pd(β^-)^{112}Ag	E_{β^-}=280(20) to (1^+) level at 18.5 keV										Ens14c	**
*^{112}Ag(β^-)^{112}Cd	E_{β^-}=3440(20) to 2^+ level at 617.518 keV										Ens14c	**
*^{112}Ag(β^-)^{112}Cd	E_{β^-}=3350(20) to 2^+ level at 617.518 keV; error increased by evaluator										Ens14c	**
*^{112}Sb(β^+)^{112}Sn	E_{β^+}=5200(100) 4750(50) 4783(26) respectively, to 2^+ level at 1256.69 keV										Ens14c	**
^{113}Mo−u	−57317	337	−56420#	220#	1.1	D			GT3	2.5	16Kn03	*
^{113}Tc−u	−67633	268	−67430	4	0.5	o			GT1	1.5	04Ma.A	
	−67502	106			0.3	U			GT2	2.5	08Kn.A	
^{113}Tc−^{129}Xe$_{.876}$	15981.0	3.6	15982	4	0.2	1	96	96 ^{113}Tc	JY1	1.0	11Ha48	
^{113}Ru−u	−77015	88	−77160.5	1.6	−1.7	U			JY0	1.0	03Ko.A	*
	−77110	100			−0.2	o			GT2	2.5	08Kn.A	
	−77170	130			0.0	U			GT2	2.5	08Su19	
^{113}Ru−^{113}Tc	−9774	32	−9730	4	1.4	U			RI1	1.0	23Ho11	*
^{113}Ru−^{133}Cs$_{.850}$	3205.3	1.6				2			JY2	1.0	23Hu25	
^{113}Ru ^{40}Ar−^{153}Nd	−42462	178	−42500.6	2.5	−0.2	U			RI1	1.0	24Ki16	*
^{113}Rh−u	−84464	83	−84563	7	−1.2	U			JY0	1.0	03Ko.A	
^{113}Rh ^{40}Ar−^{153}Nd	−49866.8	7.3	−49903	7	−4.9	B			RI1	1.0	24Ki16	*
^{113}Pd ^{40}Ar−^{153}Nd	−55081.1	22.9	−55079	7	0.1	1	10	10 ^{113}Pd	RI1	1.0	24Ki16	*
^{113}Ag^m−^{133}Cs$_{.850}$	−12944.7	3.6	−12943.2	2.9	0.4	1	66	66 ^{113}Ag^m	JY2	1.0	24De01	
C$_9$ H$_5$−^{113}Cd	134721.1	3.9	134717.13	0.24	−0.4	U			M16	2.5	63Da10	
	134727	19			−0.3	U			R12	1.5	83De51	
	134728	5			−1.4	U			R12	1.5	83De51	
C$_7$ O H$_{13}$−^{113}Cd	192250	16	192232.01	0.24	−0.7	U			R12	1.5	83De51	
C$_6$ ^{13}C O H$_{12}$−^{113}Cd	187772	17	187761.81	0.24	−0.4	U			R12	1.5	83De51	
C$_8$ N H$_3$−^{113}Cd	122161	19	122141.07	0.24	−0.7	U			R12	1.5	83De51	
^{113}Cd−u	−95506	93	−95591.97	0.24	−0.9	U			GS2	1.0	05Li24	*
C$_9$ H$_5$−^{113}In	135015	9	135064.74	0.20	2.2	U			M16	2.5	63Da10	
	135087	6			−2.5	U			R12	1.5	83De51	
C$_8$ N H$_3$−^{113}In	122506	14	122488.68	0.20	−0.8	U			R12	1.5	83De51	
^{113}In−u	−95969	126	−95939.59	0.20	0.2	U			GS2	1.0	05Li24	*
^{113}Sn−u	−94796	39	−94823.90	0.20	−0.7	U			GS2	1.0	05Li24	*
^{113}Sb−u	−90635	30	−90625	18	0.3	R			GS2	1.0	05Li24	
^{113}Te−u	−84109	30				2			GS2	1.0	05Li24	
^{113}I−^{133}Cs$_{.850}$	4015.9	8.6				2			SH1	1.0	07Ma92	
^{113}Xe−^{133}Cs$_{.850}$	13585.5	8.1	13587	7	0.2	1	82	82 ^{113}Xe	SH1	1.0	07Ma92	
^{113}Ru−^{105}Ru$_{1.076}$	22103	34	22106.1	2.1	0.1	U			JY1	1.0	07Ha20	*
^{113}Rh−^{120}Sn$_{.942}$	7565.4	7.6	7563	7	−0.4	1	84	84 ^{113}Rh	JY1	1.0	07Ha20	
^{113}Pd−^{120}Sn$_{.942}$	2387.1	7.4	2387	7	−0.0	1	91	90 ^{113}Pd	JY1	1.0	07Ha20	
^{113}Cd ^{35}Cl−^{111}Cd ^{37}Cl	3174	2	3174.4	0.4	0.1	U			H11	4.0	63Bi12	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{112}Cd H$-^{113}$Cd	6164	20	6180.84	0.23	0.6	U			R12	1.5	83De51	
^{113}Cd$-^{112}$Cd$_{1.009}$	2519.36	0.29	2519.31	0.23	−0.2	1	62	33 ^{112}Cd	MS1	1.0	16Ga33	
^{113}In$-^{112}$Cd$_{1.009}$	2171.26	0.32	2171.70	0.25	1.4	1	61	42 ^{112}Cd	MS1	1.0	16Ga33	
^{113}Ru$^{m}-^{113}$Ru	107.4	1.1	107.8	0.9	0.4	3			JY2	1.0	23Hu25	
	108.7	1.6			−0.6	3			JY2	1.0	23Hu25	*
^{113}Rh$-^{113}$Tc	−17146	17	−17132	8	0.8	1	20	16 ^{113}Rh	RI1	1.0	23Ho11	
^{113}Pd$-^{113}$Tc	−22291	51	−22308	8	−0.3	U			RI1	1.0	23Ho11	
^{113}Ag$-^{113}$Tc	−25962	43	−25925	5	0.9	U			RI1	1.0	23Ho11	
^{113}Ag$-^{113}$Ag^{m}	−49.2	4.2	−46.70	0.11	0.6	U			JY2	1.0	24De01	
^{113}Ag$-^{113}$Cd	2239.0	5.0	2236.2	2.9	−0.6	1	34	34 ^{113}Ag	CP2	1.0	22Ga27	
^{113}Sn$^{m}-^{113}$In	1198.762	0.020				2			JY2	1.0	26Ge.A	
^{113}Cd$-^{111}$Cd H	−7623	42	−7600.7	0.4	0.4	U			R12	1.5	83De51	
^{113}Cd$-^{112}$Cd	1642	11	1644.19	0.23	0.1	U			M16	2.5	63Da10	
	1620	40			0.4	U			R12	1.5	83De51	
^{113}In$-^{112}$Cd	1297	45	1296.58	0.25	−0.0	U			R12	1.5	83De51	
^{113}Cd$-^{110}$Cd H	−6412	32	−6424.4	0.4	−0.3	U			R12	1.5	83De51	
^{113}Cd$-^{111}$Cd	242	35	224.3	0.4	−0.3	U			R12	1.5	83De51	
^{113}Cd H$-^{112}$Cd	9467	35	9469.22	0.23	0.0	U			R12	1.5	83De51	
^{113}I(α)^{109}Sb	2706.0	41.5	2707	10	0.0	U					81Sc17	
^{113}Xe(α)^{109}Te	3094.8	15.	3087	8	−0.5	1	24	18 ^{113}Xe			79Sc22	
^{111}Cd(t,p)^{113}Cd	7456	20	7451.9	0.3	−0.2	U			Ald		67Hi01	
^{113}Cd(p,t)^{111}Cd	−7456	5	−7451.9	0.3	0.8	U			Min		73Oo01	
^{113}In(p,t)^{111}In$-^{115}$In()^{113}In	−810	10	−803.3	0.5	0.7	U			Roc		74Ma09	
^{113}In(p,t)^{111}In$-^{112}$Cd()^{110}Cd	−746.3	4.1	−745.6	0.3	0.2	U			SPa		80Ta07	
^{112}Cd(n,γ)^{113}Cd	6550	100	6539.77	0.21	−0.1	U					61Ja21	
	6542.0	0.2			−11.2	C					90Ne.A	
^{112}Cd(d,p)^{113}Cd	4318	30	4315.20	0.21	−0.1	U			Pit		64Ro17	
	4315.56	0.64			−0.6	1	11	6 ^{112}Cd	Rez		90Pi05	*
^{113}Cd(d,t)^{112}Cd	−254	30	−282.54	0.21	−1.0	U			Pit		64Co11	
^{113}In(d,t)^{112}In	−3180	25	−3191	4	−0.4	U			Pit		67Hj03	
^{112}Sn(n,γ)^{113}Sn	7741.9	2.3	7744.13	0.26	1.0	U			ORn		75Sl.A	
^{112}Sn(d,p)^{113}Sn	5504	25	5519.56	0.26	0.6	U			Pit		64Co11	
	5518.2	3.2			0.4	U			SPa		75Be09	
^{112}Sn(^{3}He,d)^{113}Sb	−2400	40	−2443	17	−1.1	R			Sac		68Co22	
^{113}Xe(εp)^{112}Te	7920	150	8075	11	1.0	o					82Pl05	
	8300	150			−1.5	U					05Ja10	
^{113}Cs(p)^{112}Xe	967	4	972.8	2.2	1.5	o					84Fa04	*
	982.7	4.			−2.5	3					92He.A	
	967.6	6.			0.9	3					94Pa12	
	968.6	3.			1.4	3					95Ho26	
^{113}Ru(β^{-})^{113}Rh	6480	50	6895	7	8.3	C			Jyv		00Kr.A	
^{113}Rh(β^{-})^{113}Pd	5008	50	4821	9	−3.7	C			Jyv		00Kr.A	
^{113}Pd(β^{-})^{113}Ag	3360	150	3369	7	0.1	U					75Br.A	
	3340	35			0.8	U			Stu		90Fo07	
^{113}Ag^{m}(IT)^{113}Ag	43.50	0.10	43.50	0.10	−0.0	1	100	66 ^{113}Ag			FGK243	*
^{113}Ag(β^{-})^{113}Cd	2010	20	2083.0	2.7	3.6	B					57Je.A	
	2070	150			0.1	U					70Ma47	*
	2031	30			1.7	U			Stu		90Fo07	*
^{113}Cd(β^{-})^{113}In	326.4	15.	323.80	0.26	−0.2	U					51Ca43	*
	316.4	30.			0.2	U					54De13	*
	320	10			0.4	U			CIT		88Mi13	
	344.9	21.0			−1.0	U					07Be61	
	322.2	0.9			1.8	U					09Da03	
	322.8	1.1			0.9	U					24Ba40	
^{113}Sn(β^{+})^{113}In	1034.6	5.0	1039.251	0.027	0.9	U					93Li10	*
^{113}In(p,n)^{113}Sn	−1809	6	−1821.598	0.027	−2.1	U			Oak		73Ra13	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$^{113}Sn^{m}$(IT)^{113}Sn	77.389	0.019				3					Ens095	
$^{113}Sb(\beta^{+})^{113}Sn$	3934	30	3911	17	−0.8	4					61Se08	*
	3945	50			−0.7	4					69Ki16	*
$^{113}Te(\beta^{+})^{113}Sb$	5520	300	6070	30	1.8	U					74Bu21	
	5720	200			1.7	U					74Ch17	
$*^{113}Mo-u$	Trends in Mass Surface TMS suggest ^{113}Mo 840 keV less bound										GAu	**
$*^{113}Ru-u$	$M-A$=–71689(77) keV for mixture gs+m at 100.4(0.8) keV										Nub269	**
$*^{113}Ru-u$	$M-A$=–71882(93) keV for mixture gs+m at 100.4(0.8) keV										Nub269	**
$*^{113}Ru-u$	$M-A$=–71931(120) keV for mixture gs+m at 100.4(0.8) keV										Nub269	**
$*^{113}Ru-^{113}Tc$	D_M=–9720.7(6.3) μu for mixture gs+m at 100.4(0.8) keV; $M-A$=–71866.4(6.8) keV										Nub269	**
$*^{113}Ru\ ^{40}Ar-^{153}Nd$	D_M=–42408.1(30.6) μu, $M-A$=–106833.3(32.7) keV for mixture of ground state and m										Nub269	**
$*^{113}Ru\ ^{40}Ar-^{153}Nd$	at 100.4(0.8) keV; reported $M-A$=–106833(44) keV, author confirmed										Nub269	**
$*^{113}Ru\ ^{40}Ar-^{153}Nd$	mistake in the uncertainty										Nub269	**
$*^{113}Rh\ ^{40}Ar-^{153}Nd$	Reported $M-A$=–113781(34) keV, author confirmed mistake in the unc.										24Ki16	**
$*^{113}Pd\ ^{40}Ar-^{153}Nd$	Reported $M-A$=–118639(39) keV, author confirmed mistake in the unc.										24Ki16	**
$*^{113}Cd-u$	$M-A$=–88832(41) keV for mixture gs+m at 263.361 keV										Nub269	**
$*^{113}In-u$	$M-A$=–89199(30) keV for mixture gs+m at 391.699 keV										Nub269	**
$*^{113}Sn-u$	$M-A$=–88263(29) keV for mixture gs+m at 77.389 keV										Nub269	**
$*^{113}Ru-^{105}Ru_{1.076}$	D_M=22157(12) μu for mixture gs+m at 100.4(0.8) keV; $M-A$=–71822(12) keV										Nub269	**
$*^{113}Ru^{m}-^{113}Ru$	measured with 2+ charge state; systematic errors are negligible										23Hu25	**
$*^{112}Cd(d,p)^{113}Cd$	Estimated systematic error 0.5 added to statistical error 0.40										AHW	**
$*^{113}Cs(p)^{112}Xe$	E_p from another GSI work; superseded by 95Ho26										87Gi07	**
$*^{113}Ag^{m}$(IT)^{113}Ag	From a least-squares fit to Eg										Ens10a	**
$*^{113}Ag(\beta^{-})^{113}Cd$	E_{β^-}=1530(150) from $^{113}Ag^{m}$ at 43.50 to 5/2$^+$ level at 583.962 keV										Ens106	**
$*^{113}Ag(\beta^{-})^{113}Cd$	Q_{β^-}=2075(30) from $^{113}Ag^{m}$ at 43.50 keV										Nub269	**
$*^{113}Cd(\beta^{-})^{113}In$	Q_{β^-}=590(15) 580(30) respectively, from $^{113}Cd^{m}$ at 263.361 keV										Nub269	**
$*^{113}Sn(\beta^{+})^{113}In$	Q_{β^+}=642.9(5.0) to 1/2$^-$ level at 391.699 keV										Ens106	**
$*^{113}Sb(\beta^{+})^{113}Sn$	E_{β^+}=2420(20) 2430(50) respectively, to 3/2$^+$ level at 498.06 keV,										Ens106	**
*	plus 6% to 5/2$^+$ at 409.83 keV										Ens106	**
$^{114}Tc-u$	−62459	365	−62910	470	−0.8	o			GT1	1.5	04Ma.A	
	−62910	186				2			GT3	2.5	16Kn03	
$^{114}Ru-^{105}Ru_{1.086}$	24805	13	24804	4	−0.1	U			JY1	1.0	07Ha20	
$^{114}Ru-u$	−75642	236	−75386	4	0.7	U			GT1	1.5	04Ma.A	
$^{114}Rh-u$	−81160	70	−81225.5	0.7	−0.9	U			JY0	1.0	03Ko.A	*
$^{114}Rh-^{133}Cs_{.857}$	−199.5	2.8	−197.8	0.7	0.6	1	6	6 ^{114}Rh	JY2	1.0	23Hu01	
$^{114}Rh-^{85}Rb_{1.341}$	37062.9	2.0	37064.5	0.7	0.8	1	13	13 ^{114}Rh	JY2	1.0	25St02	
$^{114}Rh^{m}-^{85}Rb_{1.341}$	37162.5	2.0	37164.2	0.6	0.8	–			JY2	1.0	25St02	*
	37165.3	1.5			−0.7	–			JY2	1.0	25St02	*
ave.	37164.3	1.2			−0.1	1	26	26 $^{114}Rh^{m}$			average	
$^{114}Rh^{n}-^{85}Rb_{1.341}$	37187.6	1.8	37188.2	0.6	0.3	–			JY2	1.0	25St02	*
	37189.8	1.5			−1.1	–			JY2	1.0	25St02	
ave.	37188.9	1.2			−0.6	1	28	28 $^{114}Rh^{n}$			average	
$^{114}Ag-^{133}Cs_{.857}$	−10149.3	4.9	−10141.3	1.3	1.6	U			MA8	1.0	10Br02	
	−10141.3	1.3				2			JY2	1.0	25Va02	
$C_8\ H_{18}-^{114}Cd$	237487.6	4.	237485.67	0.27	−0.2	U			M16	2.5	63Da10	
$C_6\ O_2\ H_{10}-^{114}Cd$	164713	15	164714.65	0.27	0.1	U			R12	1.5	83De51	
	164711	15			0.2	U			R12	1.5	83De51	
$C_9\ H_6-^{114}Cd$	143591	5	143585.29	0.27	−0.8	U			R12	1.5	83De51	
	143586	8			−0.1	U			R12	1.5	83De51	
$C_8\ ^{13}C\ H_5-^{114}Cd$	139117	17	139115.09	0.27	−0.1	U			R12	1.5	83De51	
$C_8\ N\ H_4-^{114}Cd$	131017	12	131009.23	0.27	−0.4	U			R12	1.5	83De51	
	131009	20			0.0	U			R12	1.5	83De51	
$^{114}Cd-^{133}Cs_{.857}$	−15611.4	4.2	−15607.43	0.27	0.9	U			MA8	1.0	10Br02	
$^{114}In-u$	−94986	68	−95083.6	0.3	−1.4	U			GS2	1.0	05Li24	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$C_8\ H_{18}-{}^{114}Sn$		238092	10	238070.45	0.03	−0.9	U			M16	2.5	63Da10	
		238066	8			0.4	U			R13	1.5	83De51	
$^{114}Sb-u$		−90731	30	−90711	21	0.7	–			GS2	1.0	05Li24	
		−90711	50			0.0	–			GR1	1.0	19An10	
	ave.	−90726	26			0.6	1	68	68 ^{114}Sb			average	
$^{114}Te-u$		−87911	30	−87912	26	−0.0	2			GS2	1.0	05Li24	
		−87916	54			0.1	2			GR1	1.0	19An10	
$^{114}I-u$		−77981.1	21.5				2			GR1	1.0	20Ay05	
$^{114}Xe-{}^{133}Cs_{.857}$		9008	12				2			MA6	1.0	04Di18	
$^{114}Ru-{}^{120}Sn_{.950}$		17522.0	3.7				2			JY1	1.0	11Ha48	
$^{114}Rh-{}^{120}Sn_{.950}$		11603	28	11682.2	1.1	2.8	U			JY1	1.0	07Ha20	*
$^{114}Pd-{}^{120}Sn_{.950}$		3277.0	7.4				2			JY1	1.0	07Ha20	
$^{114}Rh-{}^{112}Sn_{1.018}$		15663.14	0.74	15662.8	0.7	−0.4	1	83	81 ^{114}Rh	CP2	1.0	25Li19	
$^{114}Rh^m-{}^{112}Sn_{1.018}$		15762.98	0.79	15762.5	0.6	−0.6	1	55	51 $^{114}Rh^m$	CP2	1.0	25Li19	
$^{114}Rh^n-{}^{112}Sn_{1.018}$		15785.75	0.79	15786.6	0.6	1.0	1	54	50 $^{114}Rh^n$	CP2	1.0	25Li19	
$^{114}Cd\ ^{35}Cl-{}^{112}Cd\ ^{37}Cl$		3546	3	3551.19	0.27	0.4	U			H11	4.0	63Bi12	
		3547	3			0.6	U			H20	2.5	66Ma05	
		3548.5	1.0			1.1	U			H26	2.5	73Me28	
$^{113}Cd\ H-{}^{114}Cd$		8859	18	8868.16	0.14	0.3	U			R12	1.5	83De51	
$^{114}Tc^m-{}^{114}Ru$		12651	13				3			JY1	1.0	11Ha48	*
$^{114}Rh^m-{}^{114}Rh$		99.6	8.1	99.7	0.9	0.0	U			JY2	1.0	23Hu01	*
$^{114}Rh^n-{}^{114}Rh^m$		24.8	1.0	24.0	0.7	−0.8	1	46	23 $^{114}Rh^m$	JY2	1.0	25St02	
$^{114}Cd-{}^{112}Cd\ H$		−7225	33	−7223.97	0.26	0.0	U			R12	1.5	83De51	
$^{114}Cd-{}^{113}Cd$		−1040	11	−1043.12	0.14	−0.1	U			M16	2.5	63Da10	
		−1032	33			−0.2	U			R12	1.5	83De51	
$^{114}Cd-{}^{113}In$		−679	45	−695.5	0.3	−0.2	U			R12	1.5	83De51	
$^{114}Cd-{}^{111}Cd\ H$		−8651	35	−8643.9	0.4	0.1	U			R12	1.5	83De51	
$^{114}Cd-{}^{112}Cd$		587	33	601.06	0.26	0.3	U			R12	1.5	83De51	
$^{114}Cd\ H-{}^{113}Cd$		6821	35	6781.91	0.14	−0.7	U			R12	1.5	83De51	
$^{114}Ba(\gamma,{}^{12}C)^{102}Sn$		18110	780	19029	23	1.2	U					95Gu01	
$^{114}Cs(\alpha)^{110}I$		3343.5	30.	3360	60	0.2	o			GSa		80Ro04	
		3357.0	30.				4			GSa		81Sc17	
$^{114}Ba(\alpha)^{110}Xe$		3534.2	40.	3592	19	1.4	5					02Ma19	
		3606.8	20.			−0.7	5					16Ca33	
$^{112}Cd(t,p)^{114}Cd$		7105	20	7100.95	0.25	−0.2	U			Ald		67Hi01	
$^{114}Cd(p,t)^{112}Cd$		−7106	5	−7100.95	0.25	1.0	U			Min		73Oo01	
$^{112}Sn(t,p)^{114}Sn$		9579	15	9565.48	0.26	−0.9	U			Ald		69Bj01	
$^{114}Sn(p,t)^{112}Sn$		−9582	15	−9565.48	0.26	1.1	U			Roc		70Fl08	
$^{113}Cd(n,\gamma)^{114}Cd$		9042.76	0.20	9042.98	0.13	1.1	–			ILn		79Br25	Z
		9043.18	0.19			−1.0	–			Bdn		06Fi.A	
$^{114}Cd(\gamma,n)^{113}Cd$		−9050	4	−9042.98	0.13	1.8	U			McM		79Ba06	
$^{113}Cd(d,p)^{114}Cd$		6817	30	6818.42	0.13	0.0	U			Pit		64Co11	
		6822	8			−0.4	U			MIT		67Co15	
$^{114}Cd(d,t)^{113}Cd$		−2801	30	−2785.75	0.13	0.5	U			Pit		64Ro17	
$^{113}Cd(n,\gamma)^{114}Cd$	ave.	9042.98	0.14	9042.98	0.13	0.0	1	95	80 ^{114}Cd			average	
$^{113}In(n,\gamma)^{114}In$		7274.0	1.2	7274.00	0.25	−0.0	U					75Ra07	Z
		7273.83	0.27			0.6	1	88	82 ^{114}In	Bdn		06Fi.A	
$^{113}In(d,p)^{114}In$		5082	25	5049.43	0.25	−1.3	U			Pit		67Hj03	
$^{114}Sn(d,{}^3He)^{113}In$		−2980	50	−2988.08	0.19	−0.2	U			Sac		69Co03	
$^{114}Sn(p,d)^{113}Sn$		−8101	15	−8078.58	0.19	1.5	U			Har		70Ca01	
$^{114}Sn(d,t)^{113}Sn$		−4052	20	−4045.92	0.19	0.3	U			Pit		64Co11	
		−4043.7	4.2			−0.5	U			SPa		75Be09	
$^{114}Cs(\varepsilon p)^{113}I$		8730	150	9140	90	2.8	B					82Pl05	
$^{114}Ru(\beta^-)^{114}Rh$		6100	200	5440	4	−3.3	B			Jyv		92Jo05	*
		6120	200			−3.4	C			Jyv		94Jo.A	
$^{114}Rh(\beta^-)^{114}Pd$		6500	500	7829	7	2.7	U			Jyv		88Ay02	
		7392	53			8.3	C			Jyv		00Kr.A	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{114}Pd(β^-)^{114}Ag		1450	100	1433	7	−0.2	U					75Br.A	
		1450	100			−0.2	o			Jyv		89Ay.A	
		1450	100			−0.2	o			Jyv		89Ko22	
		1414	30			0.6	U			Stu		90Fo07	
		1451	25			−0.7	U			Jyv		94Jo.A	
^{114}Ag(β^-)^{114}Cd		4850	150	5091.7	1.2	1.6	U					71Ro19	
		4900	260			0.7	U					72Wa06	
		5160	110			−0.6	o			Stu		84Lu02	
		5018	35			2.1	U			Stu		90Fo07	
^{114}In(β^+)^{114}Cd		1422	25	1445.2	0.4	0.9	U					56Gr35	
		1417	20			1.4	U					57Dz64	
^{114}In(β^-)^{114}Sn		1987	2	1989.9	0.3	1.4	U					61Da01	
		1989	1			0.9	–					61Ni02	
		1980	2			4.9	B					64An12	
		1988.5	1.0			1.4	–					68Ze04	
	ave.	1988.7	0.7			1.6	1	18	18 ^{114}In			average	
^{114}Sb(β^+)^{114}Sn		5690	100	6063	20	3.7	C					69Bu.A	*
		6370	100			−3.1	B					72Mi27	*
^{114}Sn(p,n)^{114}Sb		−6875	35	−6845	20	0.8	1	32	32 ^{114}Sb	VUn		76Ka19	
*^{114}Rh−u	Average of 2 values; $M-A$=–75531(60) keV for mixture gs+m+n at											Nub269	**
*^{114}Rh−u	92.9(0.8) 115.3(0.8) keV											Nub269	**
*^{114}Rh^m−^{85}Rb$_{1.341}$	from frequency ratio ^{114}Rhm$^+$/(^{85}Rb)$^+$ =1.341616521(23)											25St02	**
*^{114}Rh^m−^{85}Rb$_{1.341}$	from frequency ratio ^{114}Rhm^{++}/(^{85}Rb)$^+$ =0.670805047(9)											25St02	**
*^{114}Rh^n−^{85}Rb$_{1.341}$	from frequency ratio ^{114}Rhn$^+$/(^{85}Rb)$^+$ =1.341616817(21)											25St02	**
*^{114}Rh^n−^{85}Rb$_{1.341}$	from frequency ratio ^{114}Rhn^{++}/(^{85}Rb)$^+$ =0.670805191(9)											25St02	**
*^{114}In−u	$M-A$=–88384(31) keV for mixture gs+m at 190.2682 keV											Nub269	**
*^{114}Rh−^{120}Sn$_{.950}$	D_M=11678.0(7.8) μu for mixture gs+m at 92.9(0.8) 115.3(0.8) keV;											Nub269	**
*^{114}Rh−^{120}Sn$_{.950}$	$M-A$=–75665.6(7.3) keV											Nub269	**
*^{114}Tc^m−^{114}Ru	Mixture of two isomeric states with stronger component of low-spin state											11Ri01	**
*	however, estimates from TMS suggest this is ^{114}Tc^m											GAu	**
*^{114}Rh^m−^{114}Rh	D_M=119.0(3.4) and $M-A$=–75551.8(4.1) for a mixture of m and n with a											23Hu01	**
*^{114}Rh^m−^{114}Rh	energy difference of 22.4(0.9) keV as per 25St02											23Hu01	**
*^{114}Ru(β^-)^{114}Rh	E_{β^-}=5910(120) doublet to (2)$^+$ level at 127.0, 1$^+$ at 255.2 keV											Ens123	**
*^{114}Sb(β^+)^{114}Sn	E_{β^+}=3365(50) to 2$^+$ at 1299.907, original error doubled see ^{114}Sn(p,n)											Ens123	**
*^{114}Sb(β^+)^{114}Sn	E_{β^+}=4050(100) to 2$^+$ at 1299.907 level, see ^{112}Sb(β^+)											Ens123	**
^{115}Tc−u		−60462	339	−59900#	220#	0.7	D			GT3	2.5	16Kn03	*
		−59151	940			−0.8	D			MT1	1.0	24Wa19	*
^{115}Ru−^{133}Cs$_{.865}$		10871.4	3.1				2			JY2	1.0	23Hu25	
^{115}Rh−u		−79664	85	−79678	6	−0.2	U			JY0	1.0	03Ko.A	
^{115}Rh ^{40}Ar−^{155}Pm		−45424.0	7.7	−45434	6	−1.4	1	53	49 ^{115}Rh	RI1	1.0	24Ki16	*
^{115}Pd−^{133}Cs$_{.865}$		−4550.9	1.9	−4551.8	1.5	−0.5	1	63	63 ^{115}Pd	JY2	1.0	24Ja10	
^{115}Pd ^{40}Ar−^{155}Pm		−52040.1	81.5	−52092.5	2.5	−0.6	U			RI1	1.0	24Ki16	*
^{115}Pd^m−^{133}Cs$_{.865}$		−4457.7	2.5	−4456.1	1.5	0.6	1	37	37 ^{115}Pd^m	JY2	1.0	24Ja10	
^{115}Ag−^{133}Cs$_{.865}$		−9439	24	−9412.8	2.7	1.1	U			MA8	1.0	10Br02	*
^{115}Ag^m−^{133}Cs$_{.865}$		−9368.7	2.7				2			JY2	1.0	24De01	
^{115}Cd ^{40}Ar−^{155}Pm		−60228.4	95.6	−60319.7	2.0	−1.0	U			RI1	1.0	24Ki16	*
C_9 H_7−^{115}In		150910	8	150896.453	0.013	−0.7	U			M16	2.5	63Da10	
		150932	16			−1.5	U			R12	1.5	83De51	
C_6 O_2 H_{11}−^{115}In		172055	16	172025.818	0.012	−1.2	U			R12	1.5	83De51	
C_8 N H_5−^{115}In		138355	13	138320.393	0.013	−1.8	U			R12	1.5	83De51	
^{115}In−u		−96095	30	−96121.229	0.013	−0.9	U			GS2	1.0	05Li24	
C_9 H_7−^{115}Sn		151411	8	151430.529	0.016	1.0	U			M16	2.5	63Da10	
		151440	8			−0.8	U			R13	1.5	83De51	
^{115}Sb−u		−93402	30	−93402	17	0.0	2			GS2	1.0	05Li24	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{115}Te−u	−88098	30				2			GS2	1.0	05Li24	*
^{115}I−u	−81952	31				2			GS2	1.0	05Li24	
^{115}Xe−^{133}Cs$_{.865}$	8078	13				2			MA6	1.0	04Di18	
^{115}Ru−^{120}Sn$_{.958}$	22699	41	22777	3	1.9	U			JY1	1.0	07Ha20	*
^{115}Rh−^{120}Sn$_{.958}$	14001.6	7.8	14012	6	1.4	1	51	51 ^{115}Rh	JY1	1.0	07Ha20	
^{115}Pd−^{120}Sn$_{.958}$	7347	15	7354.2	1.8	0.5	U			JY1	1.0	07Ha20	*
^{115}Sn−^{120}Sn$_{.958}$	−2963.9	2.0	−2965.2	0.9	−0.7	1	21	21 ^{120}Sn	JY1	1.0	11Ha48	
^{113}Cd−^{115}In$_{.983}$	−1104.76	0.34	−1104.81	0.24	−0.1	1	51	51 ^{113}Cd	MS1	1.0	16Ga33	
^{113}In−^{115}In$_{.983}$	−1452.08	0.23	−1452.42	0.20	−1.5	1	75	75 ^{113}In	MS1	1.0	16Ga33	
^{115}Ru^{m}−^{115}Ru	138.6	5.4				3			JY2	1.0	23Hu25	
^{115}Ag−^{115}Ag^{m}	−39.3	1.5	−44.06	0.13	−3.2	B			JY2	1.0	24De01	
^{115}Cd−^{115}In	1558.07	0.36	1558.2	0.3	0.3	1	79	79 ^{115}Cd	CP2	1.0	22Ga27	
^{115}In−^{115}Sn	534.0768	0.0104	534.076	0.010	−0.0	1	100	100 ^{115}Sn	FS1	1.0	09Mo23	
	534.28	0.18			−1.1	U			JY1	1.0	09Wi10	
	536.29	0.81			−2.7	U			CP2	1.0	22Ga27	
^{115}In−^{114}Cd	483	45	513.87	0.27	0.5	U			R12	1.5	83De51	
^{115}Sn−^{114}Sn	573	11	564.566	0.027	−0.3	U			M16	2.5	63Da10	
^{115}In−^{113}In	−200	28	−181.64	0.20	0.4	U			R12	1.5	83De51	
^{115}In−^{129}Xe	−902.0845	0.0111	−902.085	0.011	−0.0	1	100	100 ^{115}In	FS1	1.0	09Mo23	
^{115}Sn−^{129}Xe	−1436.1613	0.0130	−1436.161	0.015	−0.0	o			FS1	1.0	09Mo23	*
^{113}Cd(t,p)^{115}Cd	6702	20	6702.4	0.3	0.0	U			Ald		67Hi01	
^{114}Cd(n,γ)^{115}Cd	6160	100	6141.2	0.3	−0.2	U					61Ja21	
^{114}Cd(d,p)^{115}Cd	3923	30	3916.6	0.3	−0.2	U			Pit		64Ro17	
	3929	20			−0.6	U			Oak		64Si18	
	3916.30	0.59			0.6	1	35	21 ^{115}Cd	Rez		90Pi05	*
^{114}Cd(^{3}He,d)^{115}In	1320	15	1316.83	0.25	−0.2	U					70Th.A	
^{115}In(γ,n)^{114}In	−9025	29	−9037.84	0.30	−0.4	U			Phi		60Ge01	
	−9039	5			0.2	U			McM		79Ba06	
^{115}In(d,t)^{114}In	−2789	30	−2780.61	0.30	0.3	U			Pit		64Co11	
	−2766	25			−0.6	U			Pit		67Hj03	
^{114}Sn(n,γ)^{115}Sn	7545.5	2.0	7545.428	0.025	−0.0	U			ORn		78Ra16	Z
	7545.427	0.025			0.1	1	100	100 ^{114}Sn			16Ur03	
^{114}Sn(d,p)^{115}Sn	5329	25	5320.862	0.025	−0.3	U			Pit		64Co11	
	5320.6	3.4			0.1	U			SPa		75Be09	
^{115}Sn(d,t)^{114}Sn	−1304	30	−1288.198	0.025	0.5	U			Pit		64Co11	
^{115}Xe(εp)^{114}Te	6200	130	5944	27	−2.0	U					72Ho18	
^{115}Ru(β^-)^{115}Rh	7780	100	8165	6	3.8	C			Jyv		00Kr.A	
^{115}Rh(β^-)^{115}Pd	6000	500	6202	5	0.4	U			Jyv		88Ay01	
	6566	50			−7.3	C			Jyv		00Kr.A	
^{115}Pd^{m}(IT)^{115}Pd	89.21	0.16	89.20	0.16	−0.0	1	100	63 ^{115}Pd^{m}			Ens12a	
^{115}Pd(β^-)^{115}Ag	4584	50	4527.9	2.9	−1.1	U			Stu		90Fo07	
^{115}Ag^{m}(IT)^{115}Ag	41.1	0.2	41.04	0.12	−0.3	3					90Fo07	
	41.00	0.16			0.2	3					25Ur02	
^{115}Ag(β^-)^{115}Cd	3180	100	3135.6	2.5	−0.4	U					64Ba36	*
	3118	100			0.2	U					78Ma18	*
	3091	40			1.1	U					90Fo07	*
^{115}Cd(β^-)^{115}In	1460	4	1451.44	0.30	−2.1	U					74Bo26	*
	1431	5			4.1	B					75Bo29	*
	1440	2			5.7	B					76Ra16	*
^{115}In(β^-)^{115}Sn	494	20	497.489	0.010	0.2	U					49Be53	*
	630	30			−4.4	B					50Ma76	
	625	70			−1.8	U					61Be15	
	494	30			0.1	U					62Se03	*
	480	30			0.6	U					62Wa15	
	495	20			0.1	U					72Mu02	
	482	15			1.0	U					78Pf01	*
^{115}Sb(β^+)^{115}Sn	3030	20	3030	16	0.0	R					61Se08	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value	Adjusted value	v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{115}$Tc−u	Trends in Mass Surface TMS suggest ^{115}Tc 530 keV less bound								HWJ20a	**
$*^{115}$Tc−u	Trends in Mass Surface TMS suggest ^{115}Tc 690 keV more bound								HWJ245	**
$*^{115}$Rh ^{40}Ar−^{155}Pm	Reported $M-A$=–109252(16) keV, author confirmed mistake in the unc.								24Ki16	**
$*^{115}$Pd ^{40}Ar−^{155}Pm	D_M=–51992.1(4.0) μu, $M-A$=–115370.4(8.0) keV for a mixture of ground state and m								24Ki16	**
$*^{115}$Pd ^{40}Ar−^{155}Pm	at 89.21(0.16) keV; reported $M-A$=–115370(15) keV, author confirmed								24Ki16	**
$*^{115}$Pd ^{40}Ar−^{155}Pm	mistake in the uncertainty								24Ki16	**
$*^{115}$Ag−^{133}Cs$_{.865}$	D_M=–9416.7(9.2) μu for ground state or ^{115}Ag^m at 41.04 keV; $M-A$=–84952.9(8.6) keV								Nub269	**
$*^{115}$Cd ^{40}Ar−^{155}Pm	D_M=–60131.2(10.8) μu, $M-A$=–122951.9(12.3) keV for a mixture of ground state								24Ki16	**
$*^{115}$Cd ^{40}Ar−^{155}Pm	and m at 181.0(0.5) keV; reported $M-A$=–122952(17) keV, author								24Ki16	**
$*^{115}$Cd ^{40}Ar−^{155}Pm	confirmed mistake in the uncertainty								24Ki16	**
$*^{115}$Te−u	$M-A$=–82058(28) keV for mixture gs+m at 10(6) keV								Nub269	**
$*^{115}$Ru−^{120}Sn$_{.958}$	D_M=22767.3(7.3) μu for mixture gs+m at 129(5) keV; $M-A$=–66064.8(6.9) keV								Nub269	**
$*^{115}$Pd−^{120}Sn$_{.958}$	D_M=7348(15), 7442(15) μu for ground state, 89.20 isomer								Nub269	**
$*^{115}$Sn−^{129}Xe	Used are the equations for the ^{115}In−^{129}Xe and ^{115}In−^{115}Sn doublets								GAu	**
$*^{114}$Cd(d,p)^{115}Cd	Estimated systematic error 0.5 added to statistical error 0.32 keV								AHW	**
$*^{115}$Ag(β^-)^{115}Cd	E_{β^-}=2950(100) to (3/2)$^+$ level at 229.1 keV, and other E_{β^-}								Ens12a	**
$*^{115}$Ag(β^-)^{115}Cd	E_{β^-}=721(100) to (23/2$^-$) level at 2397.2 keV, and other E_{β^-}								Ens12a	**
$*^{115}$Ag(β^-)^{115}Cd	Q_{β^-}=3132(40) from ^{115}Ag^m at 41.04 keV								Nub269	**
$*^{115}$Cd(β^-)^{115}In	E_{β^-}=593(2), 636(2) to 1/2$^+$ level at 864.139, 3/2$^+$ at 828.588 keV								Ens12a	**
$*^{115}$Cd(β^-)^{115}In	E_{β^-}=320(5), 679(6) from ^{115}Cd^m 181.0 to 13/2$^+$ 1290.592, 7/2$^+$ 933.780								Ens12a	**
$*^{115}$Cd(β^-)^{115}In	Q_{β^-}=1621(2) from ^{115}Cd^m at 181.0 keV								Nub269	**
$*^{115}$In(β^-)^{115}Sn	Q_{β^-}=830(20) from ^{115}In^m at 336.244 keV								Nub269	**
$*^{115}$In(β^-)^{115}Sn	Q_{β^-}=830(30) from ^{115}In^m at 336.244 keV								Nub269	**
$*^{115}$In(β^-)^{115}Sn	Q_{β^-} is larger than first excitation energy 497.334(0.023) in ^{115}Sn								05Ca03	**
$*^{115}$Sb(β^+)^{115}Sn	E_{β^+}=1510(20) to 3/2$^+$ level at 497.334 keV								Ens12a	**

^{116}Ru−^{129}Xe$_{.899}$		16821.2	4.0				2				JY1	1.0	11Ha48	
^{116}Rh−u		−75890	110	−75931.9	1.9	−0.4	U				JY0	1.0	03Ko.A	*
		−75920	110			−0.0	U				GT2	2.5	08Kn.A	*
^{116}Rh−^{133}Cs$_{.872}$		6515.1	2.3	6514.0	1.9	−0.5	1	71	71	^{116}Rh	JY2	1.0	23Hu01	
^{116}Ag−^{133}Cs$_{.872}$		−6167.3	3.5	−6168.3	2.4	−0.3	1	47	47	^{116}Ag	MA8	1.0	10Br02	*
^{116}Ag^n−^{133}Cs$_{.872}$		−6029.8	3.3	−6028.9	2.4	0.3	1	53	53	^{116}Ag^n	JY2	1.0	25Va02	
C$_9$ H$_8$−^{116}Cd		157837.4	2.9	157837.01	0.17	−0.1	U				M16	2.5	63Da10	
		157851	5			−1.9	U				R12	1.5	83De51	
		157846	22			−0.3	U				R12	1.5	83De51	
C$_6$ O$_2$ H$_{12}$−^{116}Cd		178982	15	178966.38	0.17	−0.7	U				R12	1.5	83De51	
C$_8$ ^{13}C H$_7$−^{116}Cd		153376	8	153366.81	0.17	−0.8	U				R12	1.5	83De51	
		153382	22			−0.5	U				R12	1.5	83De51	
C$_8$ N H$_6$−^{116}Cd		145262	17	145260.95	0.17	−0.0	U				R12	1.5	83De51	
C$_9$ H$_8$−^{116}Sn		160861	8	160857.42	0.10	−0.2	U				M16	2.5	63Da10	
		160857	8			0.0	U				R13	1.5	83De51	
^{116}Sb−u		−93128	129	−93207	6	−0.6	U				GS2	1.0	05Li24	*
^{116}Te−u		−91540	30	−91534	26	0.2	–				GS2	1.0	05Li24	
		−91539	55			0.1	–				GR1	1.0	19An10	
	ave.	−91540	26			0.2	1	97	97	^{116}Te			average	
^{116}I−u		−83005	126	−83110	80	−0.9	1	41	41	^{116}I	GR1	1.0	20Ay05	
^{116}Xe−^{133}Cs$_{.872}$		4027	14	4027	14	−0.0	1	100	100	^{116}Xe	MA6	1.0	04Di18	
^{116}Rh−^{120}Sn$_{.967}$		18674	38	18638.3	2.2	−0.9	U				JY1	1.0	07Ha20	*
^{116}Pd−^{120}Sn$_{.967}$		8868.0	7.6	8866.0	0.9	−0.3	U				JY1	1.0	07Ha20	
^{116}Cd ^{35}Cl−^{114}Cd ^{37}Cl		4353	3	4348.5	0.3	−0.4	U				H11	4.0	63Bi12	
		4344	2			0.9	U				H20	2.5	66Ma05	
		4348.7	1.2			−0.1	o				H26	2.5	73Me28	
		4347.46	0.44			0.9	1	8	6	^{114}Cd	H49	2.5	10Mc04	
^{116}Rh^m−^{116}Rh		129.7	2.1	129.2	1.9	−0.2	1	84	69	^{116}Rh^m	JY2	1.0	23Hu01	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{116}Rh−^{116}Cd		19301.8	5.4	19304.8	1.9	0.6	1	13	13 ^{116}Rh	CP2	1.0	25Li19	
^{116}Rh^{m}−^{116}Cd		19431.9	4.4	19434.0	2.5	0.5	1	32	31 ^{116}Rh^{m}	CP2	1.0	25Li19	
^{116}Pd−^{116}Cd		9532.479	0.086				2			CP2	1.0	24Po14	
^{116}Ag−^{116}Ag^{n}		−146.0	4.4	−139.34	0.24	1.5	U			JY2	1.0	25Va02	
^{116}Ag^{m}−^{116}Ag^{n}		−93.0	3.1	−87.92	0.26	1.6	U			JY2	1.0	25Va02	
^{116}Cd−^{116}Sn		3020.42	0.14	3020.41	0.14	−0.1	1	98	98 ^{116}Cd	JY1	1.0	11Ra24	
^{116}Cd−^{114}Cd H		−6452	32	−6426.7	0.3	0.5	U			R12	1.5	83De51	
^{116}Sn−^{115}Sn		−1602	11	−1601.86	0.10	0.0	U			M16	2.5	63Da10	
^{116}Cd−^{113}Cd H		−7458	32	−7469.82	0.29	−0.2	U			R12	1.5	83De51	
^{116}Cd−^{114}Cd		1370	32	1398.3	0.3	0.6	U			R12	1.5	83De51	
^{116}Cs($\varepsilon\alpha$)^{112}Te		12300	400	13100#	100#	2.0	D					77Bo28	
		12400	900			0.8	D					76Jo.A	*
^{116}Cd(^{14}C,^{16}O)^{114}Pd		2497	29	2535	7	1.3	U			LAl		84Co19	
^{114}Cd(t,p)^{116}Cd		6362	20	6358.29	0.29	−0.2	U			Ald		67Hi01	
^{116}Cd(p,t)^{114}Cd		−6363	5	−6358.29	0.29	0.9	U			Min		73Oo01	
^{116}Sn(p,t)^{114}Sn		−8619	15	−8627.07	0.10	−0.5	U			Roc		70Fl08	
^{116}Cd(γ,n)^{115}Cd		−8702	4	−8698.9	0.3	0.8	U			McM		79Ba06	
^{116}Cd(d,t)^{115}Cd		−2458	30	−2441.7	0.3	0.5	U			Pit		64Ro17	
^{115}In(n,γ)^{116}In		6783.8	1.2	6784.72	0.22	0.8	U					72Ra39	Z
		6784.4	1.1			0.3	U					74Co35	
		6784.72	0.22				2			Bdn		06Fi.A	
^{115}In(d,p)^{116}In		4494	25	4560.15	0.22	2.6	U			Pit		64Co11	
		4580	30			−0.7	U			Pit		67Hj03	
^{116}Sn(d,^{3}He)^{115}In		−3740	50	−3785.11	0.09	−0.9	U			Sac		69Co03	
^{115}Sn(n,γ)^{116}Sn		9562.2	1.5	9563.44	0.09	0.8	U					72Mc08	
		9563.5	0.5			−0.1	U					84Ga.B	
		9563.41	0.11			0.3	–			ORn		91Ra01	Z
		9563.55	0.19			−0.6	–			Bdn		06Fi.A	
^{115}Sn(d,p)^{116}Sn		7358	30	7338.87	0.09	−0.6	U			Pit		64Co11	
^{116}Sn(p,d)^{115}Sn		−7344	15	−7338.87	0.09	0.3	U			Har		70Ca01	
^{116}Sn(d,t)^{115}Sn		−3309	20	−3306.21	0.09	0.1	U			Pit		64Co11	
		−3305.0	2.5			−0.5	U			SPa		75Be09	
^{115}Sn(n,γ)^{116}Sn	ave.	9563.45	0.10	9563.44	0.09	−0.1	1	96	96 ^{116}Sn			average	
^{115}Sn(^{3}He,d)^{116}Sb−^{120}Sn()^{121}Sb		−1722	10	−1714	5	0.8	1	29	25 ^{116}Sb	VUn		78Ka12	
^{116}Cs(εp)^{115}I		6350	300	7010#	100#	2.2	U					78Da07	*
^{116}La(p)^{115}Ba		724.2	9.1				3			Jyp		22Zh76	
^{116}Rh(β^-)^{116}Pd		8000	500	9102.9	1.8	2.2	U			Jyv		88Ay02	
^{116}Pd(β^-)^{116}Ag		2615	100	2710.5	2.2	1.0	U					75Br.A	
		2620	100			0.9	o			Jyv		89Ay.A	
		2607	30			3.5	B			Stu		90Fo07	
		2620	100			0.9	U			Jyv		94Jo.A	
^{116}Ag^{m}(IT)^{116}Ag		47.90	0.10				2					05Ba94	
^{116}Ag^{n}(IT)^{116}Ag		129.80	0.22	129.80	0.22	−0.0	1	100	53 ^{116}Ag			05Ba94	
^{116}Ag(β^-)^{116}Cd		6062	130	6168.9	2.2	0.8	o			Stu		82Al29	*
		5800	200			1.8	U					82Br10	
		6194	50			−0.5	U			Stu		90Fo07	*
^{116}In(β^-)^{116}Sn		3290	60	3276.21	0.24	−0.2	U					54Bo39	
^{116}Sb(β^+)^{116}Sn		4586	100	4704	5	1.2	U					61Fi05	*
		4606	50			2.0	U					68Ki07	*
^{116}Sn(p,n)^{116}Sb		−5515	40	−5486	5	0.7	U			VUn		76Ka19	
		−5483.2	6.			−0.5	1	75	75 ^{116}Sb	Oak		77Jo03	
^{116}Sb^{n}(β^+)^{116}Sn		5090	40				2					60Je03	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{116}Te(β^+)^{116}Sb		1554	100	1558	25	0.0	U					61Fi05	*
^{116}I(β^+)^{116}Te		7760	130	7840	80	0.6	–					70Be.A	
		7710	200			0.7	–					76Go02	
	ave.	7750	110			0.9	1	48	45 ^{116}I			average	
^{116}Xe(β^+)^{116}I		4340	200	4370	80	0.2	1	14	14 ^{116}I			76Go02	
*^{116}Rh−u	$M-A$=–70634(96) keV for mixture gs+m at 120.4(1.8) keV											Nub269	**
*^{116}Rh−u	$M-A$=–70662(93) keV for mixture gs+m at 120.4(1.8) keV											Nub269	**
*^{116}Ag−^{133}Cs$_{.872}$	maybe contaminated by the 3+ isomer at 48 keV											HWJ258	**
*^{116}Sb−u	$M-A$=–86553(34) keV for mixture gs+n at 390(40) keV											Nub269	**
*^{116}Rh−^{120}Sn$_{.967}$	D_M=18740.7(8.4) μu for mixture gs+m at 120.4(1.8) keV ; $M-A$=–70635.5(7.9) keV											Nub269	**
*^{116}Cs($\varepsilon\alpha$)^{112}Te	Q =12500(900) from ^{116}Cs^m estimated at 100#60 keV											Nub269	**
*^{116}Cs($\varepsilon\alpha$)^{112}Te	Trends in Mass Surface TMS suggest ^{116}Cs 780 keV less bound											GAu	**
*^{116}Cs(εp)^{115}I	Q =6450(300) from ^{116}Cs^m at estimated 100#60 keV											Nub269	**
*^{116}Ag(β^-)^{116}Cd	Q_{β^-}=6110(130) from ^{116}Ag^m at 47.90 keV											Nub269	**
*^{116}Ag(β^-)^{116}Cd	Q_{β^-}=6199(100); and 6241(50) from ^{116}Ag^m at 47.90 keV											Nub269	**
*^{116}Sb(β^+)^{116}Sn	E_{β^+}=2270(100) 2290(50) respectively, to 2^+ level at 1293.560 keV											Ens104	**
*^{116}Sb^n(β^+)^{116}Sn	E_{β^+}=1160(40) to 7^- level at 2908.85 keV											Ens104	**
*^{116}Te(β^+)^{116}Sb	E_{β^+}=440(100) to 1^+ level at 93.99 keV											Ens104	**
^{117}Ru−u		−63897	419	−63900	70	−0.0	o			GT1	1.5	04Ma.A	
		−63865	186			−0.1	U			GT3	2.5	16Kn03	
^{117}Ru−^{133}Cs$_{.880}$		19298	69				2			JY2	1.0	23Hu25	
^{117}Rh−u		−73903	408	−73964	10	−0.1	U			GT1	1.5	04Ma.A	
^{117}Ag−^{133}Cs$_{.880}$		−5029	16	−5029.7	2.3	−0.0	U			MA8	1.0	10Br02	*
^{117}Ag^m−^{133}Cs$_{.880}$		−4998.6	2.3				2			JY2	1.0	24De01	
C$_9$ H$_9$−^{117}Sn		167486	12	167470.94	0.15	−0.5	U			M16	2.5	63Da10	
		167471	8			−0.0	U			R13	1.5	83De51	
C ^{35}Cl$_3$−^{117}Sn		3596	2	3603.75	0.19	1.0	U			H14	4.0	62Ba24	
^{117}Te−u		−91318	30	−91354	14	−1.2	–			GS2	1.0	05Li24	
		−91359	30			0.2	–			GS2	1.0	05Li24	*
	ave.	−91339	21			−0.7	1	46	46 ^{117}Te			average	
^{117}I−u		−86350	30	−86354	27	−0.1	–			GS2	1.0	05Li24	
		−86407	50			0.4	–			Gr1	2.5	19An10	
	ave.	−86353	29			−0.0	1	88	88 ^{117}I			average	
^{117}Xe−u		−79647	30	−79641	11	0.2	R			GS2	1.0	05Li24	
^{117}Xe−^{133}Cs$_{.880}$		3562	12	3561	11	−0.1	2			MA6	1.0	04Di18	
^{117}Cs−^{133}Cs$_{.880}$		11846	46				2			MA4	1.0	99Am05	*
^{117}Rh−^{120}Sn$_{.975}$		21388.8	9.5				2			JY1	1.0	07Ha20	
^{117}Pd−^{120}Sn$_{.975}$		13309.4	7.9				2			JY1	1.0	07Ha20	
^{117}Ag−^{117}Ag^m		−32.0	3.3	−31.08	0.15	0.3	U			JY2	1.0	24De01	
^{117}In−^{117}Sn^m		1224.1	2.8	1222.4	2.3	−0.6	1	70	70 ^{117}In	JY2	1.0	23Ja11	
^{117}Sn−^{116}Sn		1219	11	1211.51	0.17	−0.3	U			M16	2.5	63Da10	
^{116}Cd(d,p)^{117}Cd		3538	30	3552.7	1.0	0.5	U			Pit		64Ro17	
		3550	20			0.1	U			Oak		64Si18	
		3552.66	1.0				2			Rez		90Pi05	*
^{116}Sn(n,γ)^{117}Sn		6943.5	2.0	6942.80	0.16	−0.3	U					75Bh01	Z
		6943.3	1.5			−0.3	U			ORn		78Ra16	Z
		6942.9	0.5			−0.2	–			Bdn		06Fi.A	
^{116}Sn(d,p)^{117}Sn		4721.0	1.8	4718.24	0.16	−1.5	–			SPa		75Be09	
^{116}Sn(n,γ)^{117}Sn	ave.	6943.1	0.5	6942.80	0.16	−0.6	1	11	8 ^{117}Sn			average	
^{116}Sn(^{3}He,d)^{117}Sb		−1091	10	−1091	8	0.0	1	71	71 ^{117}Sb	VUn		78Ka12	*
^{117}Xe(εp)^{116}Te		4100	200	3789	26	−1.6	U					72Ho18	
^{117}Ba(εp)^{116}Xe		7900	300	8300	250	1.3	F					78Bo20	*
		8300	250				2			GSI		05Ja06	
^{117}La(p)^{116}Ba		789.8	6.	818.7	2.6	4.8	B					01So02	*
		813.0	5.			1.1	o			Arp		01Ma69	
		820.1	3.			−0.5	3			Arp		11Li28	
		815.0	5.0			0.7	3			Jyp		22Zh76	
^{117}Pd(β^-)^{117}Ag		5735	32	5765	8	0.9	U			Jyv		00Kr.A	
^{117}Ag^m(IT)^{117}Ag		28.6	0.2	28.95	0.14	1.8	3					90Fo07	*
		29.3	0.2			−1.7	3					25Ur02	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{117}Ag(β^-)^{117}Cd		4160	50	4230.8	2.4	1.4	U				Stu		82Al29	*
^{117}Cd(β^-)^{117}In		2535	20	2525.8	2.4	−0.5	U						75Ta06	*
^{117}In(β^-)^{117}Sn		1456.6	5.	1453.3	2.2	−0.7	1	19	19	^{117}In			55Mc17	*
^{117}Sn^m(IT)^{117}Sn		314.58	0.04	314.58	0.04	−0.0	1	100	100	^{117}Sn^m			Ens093	
^{117}Sb(β^+)^{117}Sn		1751	40	1758	8	0.2	U						64Ba46	*
^{117}Sn(p,n)^{117}Sb		−2525	20	−2540	8	−0.8	1	18	18	^{117}Sb	Oak		71Ke21	
^{117}Te(β^+)^{117}Sb		3552	20	3544	13	−0.4	–						62Kh05	*
		3492	30			1.7	–						67Be46	*
	ave.	3534	17			0.6	1	62	51	^{117}Te			average	
^{117}I(β^+)^{117}Te		4680	100	4657	28	−0.2	–						69La33	*
		4610	110			0.4	–						70Be.A	*
	ave.	4650	70			0.1	1	14	12	^{117}I			average	
^{117}Xe(β^+)^{117}I		6270	300	6253	28	−0.1	U						85Le10	*
^{117}Cs^x(IT)^{117}Cs		50	50				3						AHW	
*^{117}Ag−^{133}Cs$_{.880}$	D_M=–5013.3(4.0) μu for ground state or ^{117}Ag^m at 28.9 keV; $M-A$=–82172.3(3.7) keV												Nub269	**
*^{117}Te−u	$M-A$=–84804(28) keV for ^{117}Te^m 11/2$^-$ at 296.1 keV												Nub269	**
*^{117}Cs−^{133}Cs$_{.880}$	D_M=11900(21) μu for mixture gs+m at 100#50 keV; $M-A$=–66418(19) keV												Nub269	**
*^{116}Cd(d,p)^{117}Cd	Estimated systematic error 0.5 added to statistical error 0.85 keV												AHW	**
*^{116}Sn(^{3}He,d)^{117}Sb	$Q-Q(^{120}$Sn(^{3}He,d))= 1373(10,Ka), Q(120)=282.1(2.0) keV												AHW	**
*^{117}Ba(εp)^{116}Xe	F : disagrees with next and with TMS												GAu	**
*^{117}La(p)^{116}Ba	Reports also an isomer ^{117}La^m E_p=933(10) Q_p=941.1 keV T=10(5) ms,												01So02	**
*	not observed in reference using similar set-up and far greater statistics												11Li28	**
*^{117}Ag^m(IT)^{117}Ag	E_3 ce transition (complex peak) to ground state												90Fo07	**
*^{117}Ag(β^-)^{117}Cd	Q_{β^-}=4260(110); and 4170(50) from ^{117}Ag^m at 28.9 keV												Nub269	**
*^{117}Cd(β^-)^{117}In	Q_{β^-}=2220(20) to ^{117}In^m at 315.303 keV												Nub269	**
*^{117}In(β^-)^{117}Sn	E_{β^-}=740(10) to 7/2$^+$ level at 711.54; and 1772(5), 1616(5) from												Ens111	**
*	^{117}In^m at 315.303 to 1/2$^+$ ground state, 3/2$^+$ level at 158.56 keV												Nub269	**
*^{117}Sb(β^+)^{117}Sn	E_{β^+}=570(40) to 3/2$^+$ level at 158.562 keV												Ens111	**
*^{117}Te(β^+)^{117}Sb	E_{β^+}=1810(20) 1750(30) respectively, to 1/2$^+$ level at 719.7 keV												Ens111	**
*^{117}I(β^+)^{117}Te	E_{β^+}=3500(50), 3250(50) to 5/2$^+$ level at 274.4, (3/2)$^+$ at 325.9 keV												Ens111	**
*^{117}I(β^+)^{117}Te	Q_{β^+}=4310(100) assumed to 5/2$^+$ level at 274.4, (3/2)$^+$ at 325.9 keV												Ens111	**
*^{117}Xe(β^+)^{117}I	May be lower limit												AHW	**
^{118}Ru−u		−61879	196	−61390#	220#	1.0	D				GT3	2.5	16Kn03	*
^{118}Rh−u		−69598	290	−69774	5	−0.4	U				GT1	1.5	04Ma.A	
^{118}Rh−^{133}Cs$_{.887}$		14090.2	5.3				2				JY2	1.0	23Hu01	
^{118}Pd−^{129}Xe$_{.915}$		6193.6	4.3	6192.7	2.7	−0.2	1	39	39	^{118}Pd	JY1	1.0	11Ha48	
^{118}Ag−^{133}Cs$_{.887}$		−1527.6	3.2				2				JY2	1.0	25Va02	*
^{118}Ag^p−u		−85267.6	2.7	−85255	3	4.8	B				MA8	1.0	10Br02	
^{118}Cd−^{133}Cs$_{.887}$		−9201.9	2.1				2				JY2	1.0	23Ja11	
C_9 H_{10}−^{118}Sn		176645	7	176643.35	0.06	−0.1	U				M16	2.5	63Da10	
		176637	8			0.5	U				R13	1.5	83De51	
^{118}Sn−u		−98393.03	0.06	−98393.03	0.06	−0.1	1	99	99	^{118}Sn	Hea	1.0	23Mo.1	
^{118}Te−u		−94162	30	−94140	20	0.7	R				GS2	1.0	05Li24	
^{118}I−u		−86932	30	−86926	21	0.2	2				GS2	1.0	05Li24	
		−86920	30			−0.2	2				GS2	1.0	05Li24	*
^{118}Xe−u		−83785	30	−83821	11	−1.2	R				GS2	1.0	05Li24	
^{118}Xe−^{133}Cs$_{.887}$		37	12	43	11	0.5	2				MA6	1.0	04Di18	
^{118}Cs^x−^{133}Cs$_{.887}$		10424	16	10429	13	0.3	o				MA1	1.0	90St25	
		10429	13				2				MA1	1.0	99Am05	
^{118}Rh−^{120}Sn$_{.983}$		26476	26	26361	5	−4.4	B				JY1	1.0	07Ha20	
^{118}Pd−^{120}Sn$_{.983}$		15199.7	7.9	15202.2	2.6	0.3	–				JY1	1.0	07Ha20	
		15202.1	3.6			0.0	–				JY1	1.0	11Ha48	
	ave.	15202	3			0.2	1	64	61	^{118}Pd			average	
^{118}Sn ^{35}Cl−^{116}Sn ^{37}Cl		2814	2	2814.26	0.14	0.0	U				H15	4.0	62Ba23	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$^{118}Rh^{m}-^{118}Rh$	202.8	6.5				3			JY2	1.0	23Hu01	
$^{118}Ag-^{118}Ag^{p}$	−132.5	1.3	−137.02	0.11	−3.5	B			JY2	1.0	25Va02	*
$^{118}In-^{118}Sn$	4697.5	2.0				2			JY2	1.0	23Ja11	
$^{118}In^{m}-^{118}Sn$	4740.8	1.5	4740.9	1.1	0.1	1	53	53 $^{118}In^{m}$	JY2	1.0	23Ja11	
$^{118}In^{n}-^{118}Sn$	4889.4	1.5	4889.3	1.1	−0.1	1	53	53 $^{118}In^{n}$	JY2	1.0	23Ja11	
$^{118}Sn-^{117}Sn$	−1338	11	−1347.38	0.13	−0.3	U			M16	2.5	63Da10	
$^{117}Cs^{x}-^{118}Cs^{x}_{496}$ $^{116}Cs_{.504}$	−1160	400	−1220#	80#	−0.1	U			P32	2.5	86Au02	
$^{118}Cs(\varepsilon\alpha)^{114}Te$	10600	200	11055	28	2.3	U					77Bo28	
	10750	200			1.5	U					78Da07	*
$^{116}Cd(t,p)^{118}Cd$	5650	20	5638.8	2.0	−0.6	U			Ald		67Hi01	
$^{116}Sn(t,p)^{118}Sn$	7769	15	7787.40	0.11	1.2	U			Ald		68Bj02	
$^{118}Sn(p,t)^{116}Sn$	−7790	10	−7787.40	0.11	0.3	U			Roc		70Fl08	
$^{118}Sn(d,^{3}He)^{117}In$	−4440	40	−4503.8	2.2	−1.6	U			Sac		69Co03	
	−4481	15			−1.5	U			MSU		71We01	
$^{117}Sn(n,\gamma)^{118}Sn$	9326.5	2.	9326.39	0.13	−0.1	U					70Or.A	
	9324.8	2.1			0.8	U			ORn		75Sl.A	
	9326.42	0.13			−0.2	1	93	92 ^{117}Sn			02Bo11	
	9327.9	1.1			−1.4	U			Bdn		06Fi.A	
$^{117}Sn(d,p)^{118}Sn$	7090	12	7101.83	0.13	1.0	U			Tal		64No06	
$^{118}Sn(p,d)^{117}Sn$	−7097	15	−7101.83	0.13	−0.3	U			Har		70Ca01	
$^{118}Cs(\varepsilon p)^{117}I$	4700	300	4740	29	0.1	U					78Da07	
$^{118}Pd(\beta^{-})^{118}Ag$	4100	200	4153	4	0.3	U			Jyv		89Ko22	*
$^{118}Ag^{p}(IT)^{118}Ag$	127.63	0.10				3					Ens932	
$^{118}Ag(\beta^{-})^{118}Cd$	7122	100	7149	4	0.3	U			Stu		82Al29	*
	7110	470			0.1	U			Stu		82Al29	*
	7155	76			−0.1	U					95Ap.A	
$^{118}In^{n}(IT)^{118}In^{m}$	138.2	0.5	138.2	0.5	0.0	1	94	47 $^{118}In^{m}$			69Ha08	
$^{118}In(\beta^{-})^{118}Sn$	4200	400	4375.7	1.9	0.4	U					61Gl02	
	4200	300			0.6	U					64Ka10	
	4310	100			0.7	U					87Ga.A	
$^{118}In^{m}(\beta^{-})^{118}Sn$	4270	100	4416.1	1.0	1.5	U					64Ka10	*
$^{118}Sb(\beta^{+})^{118}Sn$	3610	50	3656.6	3.0	0.9	U					61Fi05	
$^{118}Sn(p,n)^{118}Sb$	−4477.7	5.7	−4439.0	3.0	6.8	F			Tkm		63Ok01	*
	−4439.0	3.				2			Oak		77Jo03	
$^{118}Sb^{n}(\beta^{+})^{118}Sn$	3907	5				2					61Bo13	*
$^{118}I(\beta^{+})^{118}Te$	7080	150	6720	27	−2.4	U					68La18	*
	7068	100			−3.5	C					70Be.A	*
$^{118}Xe(\beta^{+})^{118}I$	3720	110	2892	22	−7.5	F					70Be.A	*
$^{118}Cs(\beta^{+})^{118}Xe$	9300	1000	9670	16	0.4	U					76Da.C	
$^{118}Cs^{x}(IT)^{118}Cs$	5	4				3					82Au01	
$*^{118}Ru-u$	Trends in Mass Surface TMS suggest ^{118}Ru 460 keV less bound										HWJ261	**
$*^{118}Ag-^{133}Cs_{.887}$	re-calculated from two frequency ratios between $^{133}Cs^{+}/^{118}Agx^{+}$ and										HWJ258	**
$*^{118}Ag-^{133}Cs_{.887}$	$^{118}Agx^{+}/^{118}Ag+$										HWJ258	**
$*^{118}Sn-u$	M=117.901606974(56,exp)(5,theo) u, the first uncertainty is from										23Mo.1	*W
$*^{118}Sn-u$	measurement and the second from theoretical electron-binding energies										23Mo.1	*W
$*^{118}I-u$	$M-A$=–80775(28) keV for $^{118}I^{m}$ 7^{-} at 188.8 keV										Nub269	**
$*^{118}Ag-^{118}Ag^{p}$	authors suspects $^{118}Ag^{n}$ is a mixture										25Va02	**
$*^{118}Cs(\varepsilon\alpha)^{114}Te$	As read from Fig. 2 (p.401)										GAu	**
$*^{118}Pd(\beta^{-})^{118}Ag$	Original value 4000(200) corrected for new branching ratios										93Ja03	**
$*^{118}Ag(\beta^{-})^{118}Cd$	$E_{\beta^{-}}$=4330(240), 3960(170), 3810(150) reinterpreted as feeding										95Ap.A	**
*	(1) level at 2788.72, (1) at 3224.32, (2,3,4) at 3265.77 keV										Ens959	**
$*^{118}Ag(\beta^{-})^{118}Cd$	$E_{\beta^{-}}$=3990(720), 3910(630) reinterpreted as $^{118}Ag^{n}$ at 127.63(0.10) keV										95Ap.A	**
*	to (2,3,4) level at 3181.73, 3381.8 keV										Ens959	**
$*^{118}In^{m}(\beta^{-})^{118}Sn$	$E_{\beta^{-}}$=2000(100) to 4^{+} level at 2280.342 level, and other $E_{\beta^{-}}$										Ens959	**
$*^{118}Sn(p,n)^{118}Sb$	F : see note added in proof to reference										77Jo03	**
$*^{118}Sb^{n}(\beta^{+})^{118}Sn$	$p^{+}=16(1)\times10^{-4}$ to 7^{-} level at 2574.91 keV, recalculated Q										Ens959	**
$*^{118}I(\beta^{+})^{118}Te$	$E_{\beta^{+}}$=5450(150) to 2^{+} level at 605.70 keV										Ens959	**
$*^{118}I(\beta^{+})^{118}Te$	$E_{\beta^{+}}$=5440(100) to 2^{+} level at 605.70 keV										Ens959	**
$*^{118}Xe(\beta^{+})^{118}I$	F : probably contaminated by isobars										GAu	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{119}Rh−u		−67698	268	−67443	10	0.6	U			GT1	1.5	04Ma.A	
^{119}Rh−^{129}Xe$_{.922}$		20349	10				2			JY1	1.0	11Ha48	
^{119}Pd−u		−76844	208	−76876	4	−0.1	U			GT1	1.5	04Ma.A	
^{119}Pd^m−^{120}Sn$_{.992}$		20356.2	8.8	20353	3	−0.4	1	12	11 ^{119}Pd^m	JY1	1.0	07Ha20	*
^{119}Pd^m−^{133}Cs$_{.895}$		7958.0	3.1	7958.4	2.9	0.1	1	89	89 ^{119}Pd^m	JY2	1.0	24Ja10	
^{119}Ag−^{133}Cs$_{.895}$		181	12	185.3	2.3	0.4	U			MA8	1.0	10Br02	*
		197	19			−0.6	o			CP2	1.0	22Ra.A	
		184.1	4.0			0.3	1	32	32 ^{119}Ag	CP2	1.0	25Ri09	
^{119}Ag^m−^{133}Cs$_{.895}$		222.4	3.7	221.2	2.3	−0.3	–			JY2	1.0	24De01	
		218.0	4.1			0.8	o			CP2	1.0	22Ra.A	
		221.0	4.1			0.0	–			CP2	1.0	25Ri09	
	ave.	221.8	2.7			−0.2	1	68	68 ^{119}Ag^m			average	
^{119}Cd−^{133}Cs$_{.895}$		−5626.7	2.3	−5625.6	2.0	0.5	2			JY2	1.0	23Ja11	
		−5627.9	4.1			0.5	o			CP2	1.0	22Ra.A	
		−5621.8	4.4			−0.9	2			CP2	1.0	25Ri09	
^{119}Cd^m−^{133}Cs$_{.895}$		−5473.2	2.4	−5468.3	2.0	2.0	U			JY2	1.0	23Ja11	
		−5472.7	4.1			1.1	o			CP2	1.0	22Ra.A	
		−5470.0	4.3			0.4	U			CP2	1.0	25Ri09	
^{119}In−^{133}Cs$_{.895}$		−9542.5	2.3	−9540.7	2.2	0.8	1	90	90 ^{119}In	JY2	1.0	23Ja11	
C_9 H_{11}−^{119}Sn		182778	7	182763.8	0.6	−0.8	U			M16	2.5	63Da10	
		182762	8			0.1	U			R13	1.5	83De51	
^{119}Sn−u		−96726	39	−96688.4	0.6	1.0	U			GR1	1.0	19Mi18	*
^{119}Sb−u		−96063	18	−96056	8	0.4	1	17	17 ^{119}Sb	GR1	1.0	19Mi18	
^{119}Te−u		−93731	190	−93594	8	0.7	U			GR1	1.0	19Mi18	*
^{119}I−u		−89926	30	−89939	23	−0.4	2			GS2	1.0	05Li24	
		−89959	37			0.5	2			GR1	1.0	19An10	
^{119}Xe−u		−84601	30	−84589	11	0.4	1	14	14 ^{119}Xe	GS2	1.0	05Li24	
		−84613	61			0.4	U			GR1	1.0	19An10	
^{119}Xe−^{133}Cs$_{.895}$		33	12	31	11	−0.2	1	86	86 ^{119}Xe	MA6	1.0	04Di18	
^{119}Cs−u		−77528	53	−77623	15	−1.8	U			GS2	1.0	05Li24	*
^{119}Cs^x−^{133}Cs$_{.895}$		7011	16	7015	9	0.2	o			MA1	1.0	90St25	
		7018	13			−0.2	2			MA1	1.0	99Am05	
		7012	13			0.2	2			MA4	1.0	99Am05	
^{119}Sn ^{35}Cl−^{117}Sn ^{37}Cl		3306	2	3307.4	0.6	0.2	U			H15	4.0	62Ba23	
^{119}Pd−^{119}Pd^m		−213.7	3.2				2			JY2	1.0	24Ja10	
^{119}Ag−^{119}Ag^m		−35.0	8.2	−35.86	0.11	−0.1	U			JY2	1.0	24De01	
^{119}In^m−^{119}In		334.2	3.7	334.27	0.03	0.0	U			JY2	1.0	23Ja11	
^{119}Sn−^{118}Sn		1709	12	1704.6	0.6	−0.1	U			M16	2.5	63Da10	
^{119}I−^{118}I		−2747	155	−3010	30	−1.1	U			CR2	1.5	92Sh.A	*
^{119}I−^{117}I		−3570	155	−3580	40	−0.1	U			CR2	1.5	92Sh.A	*
^{118}Cs^x−^{119}Cs$^x_{.661}$ ^{116}Cs$_{.339}$		530	80	410#	40#	−0.6	U			P32	2.5	86Au02	
^{118}Cs^x−^{119}Cs$^x_{.496}$ ^{117}Cs$^x_{.504}$		870	50	930	40	0.4	U			P22	2.5	82Au01	
		980	40			−0.5	U			P32	2.5	86Au02	
^{119}Sn(t,α)^{118}In−^{118}Sn()^{117}In		−127	6	−79.5	2.9	7.9	B			McM		85Pi03	
^{118}Sn(n,γ)^{119}Sn		6484.6	1.5	6483.5	0.5	−0.7	–			ORn		78Ra16	
		6483.3	0.6			0.3	–			Bdn		06Fi.A	
^{118}Sn(d,p)^{119}Sn		4238	12	4258.9	0.5	1.7	U			MIT		67Sp09	
^{118}Sn(n,γ)^{119}Sn	ave.	6483.5	0.6	6483.5	0.5	0.0	1	96	96 ^{119}Sn			average	
^{118}Sn(^{3}He,d)^{119}Sb		−388	10	−382	7	0.6	1	49	49 ^{119}Sb	VUn		78Ka12	*
^{119}Ba(εp)^{118}Xe		6200	200				3					78Bo20	
^{119}Ag^m(IT)^{119}Ag		33.4	0.1	33.40	0.10	0.0	1	100	68 ^{119}Ag			22Ku09	*
^{119}Ag(β^-)^{119}Cd		5350	40	5412.9	2.8	1.6	U			Stu		82Al29	
^{119}Cd^m(IT)^{119}Cd		146.54	0.11				3					Ens08c	
^{119}Cd(β^-)^{119}In		3797	80	3646.9	2.8	−1.9	U			Stu		82Al29	*
^{119}In^m(IT)^{119}In		311.37	0.03				2					Ens08c	
^{119}In(β^-)^{119}Sn		2387	100	2354.1	2.1	−0.3	U					60Yu01	*
		2413	200			−0.3	U					61Gl06	*
^{119}Sb(ε)^{119}Sn		579	20	589	7	0.5	–					57Ol05	*
^{119}Sn(p,n)^{119}Sb		−1369	15	−1372	7	−0.2	–			Oak		71Ke21	
^{119}Sb(ε)^{119}Sn	ave.	584	12	589	7	0.5	1	34	34 ^{119}Sb			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{119}Te(β^+)^{119}Sb	2293	2				2					60Ko12	*
^{119}I(β^+)^{119}Te	3630	100	3405	23	−2.3	U					69La33	*
	3370	100			0.3	U					70Be.A	
^{119}Xe(β^+)^{119}I	4990	120	4983	24	−0.1	U					70Be.A	
^{119}Cs(β^+)^{119}Xe	6260	290	6489	17	0.8	U					83Pa.A	*
^{119}Cs^x(IT)^{119}Cs	16	11				3					82Au01	*
*^{119}Pd^m−^{120}Sn$_{.992}$	re-assigned to the gs.										24Ja10	**
*^{119}Ag−^{133}Cs$_{.895}$	D_M=198.4(5.7) μu mixture for gs+m at 33.40(0.10) keV; $M-A$=–78638.7(5.3)										Nub269	**
*^{119}Sn−u	could be mixture of gs+iso, not corrected										WgM209	**
*^{119}Te−u	could be mixture of gs+iso, not corrected										WgM209	**
*^{119}Cs−u	$M-A$=–72195(48) keV for mixture gs+m at 43.0(0.2) keV										Nub269	**
*^{119}I−^{118}I	From ^{118}I/^{119}I=0.99161584(117), originally D_M=–3039(139) μu, revised by										GAu	**
*	authors: –2849(139) μu for ^{118}I gs+m mixture at 188.8 keV										Nub269	**
*^{119}I−^{117}I	From ^{117}I/^{119}I=0.98321059(130)										GAu	**
*^{118}Sn(^{3}He,d)^{119}Sb	$Q-Q$(^{120}Sn(^{3}He,d)^{121}Sb)=–673(10), Q(120)=285.1(2.1) keV										AHW	**
*^{119}Ag^m(IT)^{119}Ag	lsq fit to Eg in the decay scheme of 22Ku09										22Ku09	**
*^{119}Cd(β^-)^{119}In	Q_{β^-}=3800(90); and 3940(80) from ^{119}Cd^m at 146.54 keV										Nub269	**
*^{119}In(β^-)^{119}Sn	E_{β^-}=1600(100) to 7/2$^+$ level at 787.01 keV										Ens09a	**
*^{119}In(β^-)^{119}Sn	E_{β^-}=2700(200) from ^{119}In^m at 311.37(0.03) to 3/2$^+$ level at 23.871 keV										Ens09a	**
*^{119}Sb(ε)^{119}Sn	IBE=526(20) to 3/2$^+$ level at 23.871 keV										Ens09a	**
*^{119}Te(β^+)^{119}Sb	E_{β^+}=627(2) to 1/2$^+$ level at 644.03 keV										Ens09a	**
*^{119}I(β^+)^{119}Te	E_{β^+}=2350(100) to 3/2$^+$ level at 257.484 keV										Ens09a	**
*^{119}Cs(β^+)^{119}Xe	E_{β^+}=4980(290) to 9/2$^+$ level at 257.84 keV										Ens09a	**
*^{119}Cs^x(IT)^{119}Cs	Original 33(22) corrected for new estimated IT=50(30)#										GAu	**
^{120}Rh−^{133}Cs$_{.902}$	22358	63				2			JY1	1.0	23Hu01	
^{120}Pd−^{129}Xe$_{.930}$	13107.0	4.4	13105.5	2.5	−0.4	1	31	31 ^{120}Pd	JY1	1.0	11Ha48	
^{120}Ag−^{133}Cs$_{.902}$	4067.1	4.8	4043.0	2.9	−5.0	B			MA8	1.0	10Br02	
	4086	12			−3.6	B			MA8	1.0	10Br02	
	4043.0	2.9				2			JY2	1.0	25Va02	
^{120}Cd−^{133}Cs$_{.902}$	−4849.6	4.0				2			MA8	1.0	10Br02	
^{120}In−^{133}Cs$_{.902}$	−6730.3	3.3				2			JY2	1.0	23Ne13	*
^{120}In^m−^{133}Cs$_{.902}$	−6637.5	3.7				2			JY2	1.0	23Ne13	*
C_9H_{12}−^{120}Sn	191709	11	191698.0	1.0	−0.4	U			M16	2.5	63Da10	
	191705	8			−0.6	U			R13	1.5	83De51	
^{13}C ^{35}Cl$_2$ ^{37}Cl−^{120}Sn	4758	3	4760.4	1.0	0.2	U			H14	4.0	62Ba24	
^{120}Sb−u	−94796	76	−94920	8	−1.6	U			GS2	1.0	05Li24	*
C_9H_{12}−^{120}Te	189879	9	189835.0	1.8	−2.0	U			M16	2.5	63Da10	
	189868	8			−2.7	U			R13	1.5	83De51	
^{120}I−u	−90121	52	−89907	16	4.1	C			GS2	1.0	05Li24	*
^{120}Xe−u	−88231	30	−88216	13	0.5	R			GS2	1.0	05Li24	
^{120}Xe−^{133}Cs$_{.902}$	−2930	14	−2933	13	−0.2	2			MA6	1.0	04Di18	
^{120}Cs−u	−79342	54	−79323	11	0.4	U			GS2	1.0	05Li24	*
^{120}Cs^x−^{133}Cs$_{.902}$	5947	16	5965	10	1.1	o			MA1	1.0	90St25	
	5956	12			0.7	2			MA1	1.0	99Am05	
	5983	17			−1.1	2			MA4	1.0	99Am05	
^{120}Sn ^{35}Cl−^{118}Sn ^{37}Cl	3546	2	3545.6	1.0	−0.1	U			H15	4.0	62Ba23	
^{120}Pd−^{120}Sn	22317.1	9.7	22349.2	2.4	3.3	B			JY1	1.0	07Ha20	
	22348.6	2.8			0.2	1	72	69 ^{120}Pd	JY1	1.0	11Ha48	
^{120}Ag^m−^{120}Ag	123.5	3.7				3			JY2	1.0	25Va02	
^{120}Ag^n−^{120}Ag	217.9	4.3	218.14	0.21	0.1	U			JY2	1.0	25Va02	
^{120}Te−^{120}Sn	1842.2	1.7	1862.9	1.8	12.2	B			CP1	1.0	09Sc19	
	1839.7	1.7			13.7	B			CP1	1.0	09Sc19	
^{120}Sn−^{119}Sn	−1113	11	−1109.2	1.1	0.1	U			M16	2.5	63Da10	
^{118}Cs^x−^{120}Cs$^x_{.328}$ ^{117}Cs$^x_{.672}$	460	120	460	50	0.0	U			P22	2.5	82Au01	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$^{119}Cs^{x}-{}^{120}Cs^{x}_{.661}$ $^{117}Cs^{x}_{.339}$	−940	50	−936	25	0.0	U			P22	2.5	82Au01	
$^{119}Cs^{x}-{}^{120}Cs^{x}_{.496}$ $^{118}Cs^{x}_{.504}$	−1220	30	−1167	11	0.7	U			P22	2.5	82Au01	
	−1180	60			0.1	U			P32	2.5	86Au02	
	−1200	30			0.4	U			P32	2.5	86Au02	
	−1270	50			0.8	F			P32	2.5	86Au02	*
$^{120}Cs(\varepsilon\alpha)^{116}Te$	9200	300	8950	26	−0.8	U					76Jo.A	
$^{118}Sn(t,p)^{120}Sn$	7107	15	7106.2	0.9	−0.1	U			Ald		68Bj02	
$^{120}Sn(p,t)^{118}Sn$	−7109	10	−7106.2	0.9	0.3	U			Roc		70Fl08	
$^{120}Te(p,t)^{118}Te$	−9343	24	−9332	18	0.4	2			Win		74De31	*
$^{120}Sn(d,^{3}He)^{119}In$	−5160	40	−5182.7	2.2	−0.6	U			Sac		69Co03	
	−5169	20			−0.7	U			MSU		71We01	
$^{120}Sn(t,\alpha)^{119}In-{}^{118}Sn()^{117}In$	−692	6	−678.9	2.9	2.2	1	23	11 ^{117}In	McM		85Pi03	
$^{119}Sn(d,p)^{120}Sn$	6890	12	6879.9	1.0	−0.8	U			Tal		64No06	
$^{120}Sn(p,d)^{119}Sn$	−6889	15	−6879.9	1.0	0.6	U			Har		70Ca01	
$^{120}Sn(d,t)^{119}Sn$	−2847.0	2.5	−2847.3	1.0	−0.1	1	16	12 ^{120}Sn	SPa		75Be09	
$^{120}Pd(\beta^{-})^{120}Ag$	5500	100	5394	4	−1.1	U			Jyv		94Jo.A	
$^{120}Ag^{n}(IT)^{120}Ag$	203	1	203.20	0.20	0.2	U					71Fo22	
	203.2	0.2				3					12Ba58	
$^{120}Ag(\beta^{-})^{120}Cd$	8200	100	8283	5	0.8	U			Stu		82Al29	
	8450	100			−1.7	U					95Ap.A	
$^{120}In^{n}(IT)^{120}In^{m}$	7.5	4.3				3					23Ne12	*
$^{120}In(\beta^{-})^{120}Sn$	5300	170	5389	3	0.5	U			Stu		78Al18	
	5370	40			0.5	U					87Ga.A	
$^{120}In^{m}(\beta^{-})^{120}Sn$	5280	200	5475	4	1.0	U					64Ka10	*
	5340	170			0.8	U			Stu		78Al18	*
$^{120}Sb(\beta^{+})^{120}Sn$	2720	20	2681	7	−2.0	U					50Bl92	
	2770	30			−3.0	B					69Ki15	
$^{120}Sn(p,n)^{120}Sb$	−3462.9	7.1				2			Tkm		63Ok01	
$^{120}I(\beta^{+})^{120}Te$	5615	15				2					70Ga32	*
	5608	150	5615	15	0.0	U					68La18	*
$^{120}Xe(\beta^{+})^{120}I$	1960	40	1575	19	−9.6	B					74Mu10	*
$^{120}Cs(\beta^{+})^{120}Xe$	7300	500	8284	15	2.0	U					76Ba.A	*
	7800	1000			0.5	U					76Da.C	*
	7380	230			3.9	C					83Pa.A	*
	8210	200			0.4	U			IRS		93Al03	
$^{120}Cs^{x}(IT)^{120}Cs$	5	4				3					82Au01	
$^{120}Ba(\beta^{+})^{120}Cs$	5000	300				4					92Xu04	
$*^{120}In-{}^{133}Cs_{.902}$	Reconstructed from frequency ratio $^{120}In^{+}/(^{120}In^{x})^{+}$ =0.999999193(23)										HWJ24a	**
$*^{120}In-{}^{133}Cs_{.902}$	$^{120}Inx^{+}/(^{133}Cs+)$=0.902205506(13)										HWJ24a	**
$*^{120}In^{m}-{}^{133}Cs_{.902}$	D_M=–6633.5(1.7) μu, $M-A$=–85619.0(1.6) keV mixture of m and n state										23Ne13	**
$*^{120}In^{m}-{}^{133}Cs_{.902}$	with energy difference below 15 keV										23Ne13	**
$*^{120}Sb-u$	$M-A$=–88302(50) keV for mixture gs+m at 0#100 keV										Nub269	**
$*^{120}I-u$	$M-A$=–83881(28) keV for mixture gs+n at 133#14 keV										Nub269	**
$*^{120}Cs-u$	$M-A$=–73856(29) keV for mixture gs+m at 100#60 keV										Nub269	**
$*^{119}Cs^{x}-{}^{120}Cs^{x}_{.496}$ ^{118}C	F : rejection based on line-shape analysis										86Au02	**
$*^{120}Te(p,t)^{118}Te$	Original error 12; added systematic error 21 keV										GAu	**
$*^{120}In^{n}(IT)^{120}In^{m}$	from 23Ne12<15 keV										FKo259	**
$*^{120}In^{m}(\beta^{-})^{120}Sn$	E_{β^-}=3100(200),2200(200) to 4^+ levels at 2194.299,3057.946 keV										Ens029	**
$*^{120}In^{m}(\beta^{-})^{120}Sn$	E_{β^-}=3100(170) to 4^+ level at 2194.299 keV, and other E_{β^-}										Ens029	**
$*^{120}I(\beta^{+})^{120}Te$	E_{β^+}=4595(15), 4030(20) to ground state, 2^+ level at 560.438 keV										Ens029	**
$*^{120}I(\beta^{+})^{120}Te$	E_{β^+}=3130(150) from $^{120}I^{n}$ at 320(15) to 6^+ level at 1776.23 keV										Ens029	**
$*^{120}Xe(\beta^{+})^{120}I$	p^{+} =0.07(0.01) to 1^+ level at 25.1 keV, recalculated Q										Ens029	**
$*^{120}Cs(\beta^{+})^{120}Xe$	E_{β^+}=6000(500) 6500(1000) 6040(230), respectively, to 2^+ level at 322.61 keV										Ens029	**
$^{121}Rh-u$	−60387	266				2			GT3	2.5	16Kn03	
$^{121}Pd-u$	−71820	311	−71049.1	2.7	1.7	U			GT1	1.5	04Ma.A	
$^{121}Pd-{}^{133}Cs_{.910}$	14990.3	4.0	14989.6	2.7	−0.2	1	45	45 ^{121}Pd	JY2	1.0	24Ja10	
$^{121}Ag-{}^{133}Cs_{.910}$	6164	13	6163	9	−0.1	2			MA8	1.0	10Br02	*
	6170	17			−0.4	o			MA8	1.0	10Br02	*
	6162.3	12.9			0.1	2			JY2	1.0	24De01	*
$^{121}Cd-{}^{130}Xe_{.931}$	2796.2	3.0	2796.5	2.1	0.1	2			JY1	1.0	12Ha25	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{121}Cd−^{130}Xe$_{.931}$		2796.7	2.9	2796.5	2.1	−0.1	2			JY1	1.0	13Ka08	*
^{121}In−^{133}Cs$_{.910}$		−6119.7	1.3				2			JY2	1.0	23Ne13	
C_9H_{13}−^{121}Sb		197910.5	3.7	197914.7	2.6	0.5	U			M16	2.5	63Da10	
		197910	8			0.4	U			R13	1.5	83De51	
^{121}Sb−C ^{35}Cl ^{37}Cl$_2$		3162	3	3152.8	2.6	−0.8	U			H14	4.0	62Ba24	
^{121}Sb−u		−96180	30	−96189.3	2.6	−0.3	U			GS2	1.0	05Li24	
^{121}I−u		−92609	30	−92589	5	0.7	U			GS2	1.0	05Li24	
^{121}Xe−u		−88562	30	−88547	11	0.5	R			GS2	1.0	05Li24	
^{121}Xe−^{133}Cs$_{.910}$		−2495	13	−2508	11	−1.0	–			MA6	1.0	04Di18	
	ave.	−2499	12			−0.7	1	85	85 ^{121}Xe			average	
^{121}Cs−^{133}Cs$_{.910}$		3247	25	3266	15	0.8	o			MA1	1.0	90St25	*
		3248	25			0.7	1	38	38 ^{121}Cs	MA1	1.0	99Am05	*
^{121}Cs−u		−82821	38	−82773	15	1.3	1	16	16 ^{121}Cs	GS2	1.0	05Li24	*
^{121}Pd−^{129}Xe$_{.938}$		18265.9	3.6	18266.5	2.7	0.2	1	55	55 ^{121}Pd	JY1	1.0	11Ha48	*
^{121}Sb ^{35}Cl−^{119}Sn ^{37}Cl		3452	2	3449.2	2.7	−0.3	U			H14	4.0	62Ba24	
^{121}In^m−^{121}In		337.03	0.86	336.75	0.08	−0.3	U			JY2	1.0	23Ne13	
^{119}Cs^x−^{121}Cs$^x_{.328}$ ^{118}Cs$^x_{.672}$		−1080	30	−1047	13	0.4	U			P22	2.5	82Au01	
^{120}Cs^x−^{121}Cs$^x_{.661}$ ^{118}Cs$^x_{.339}$		280	30	240	15	−0.5	U			P22	2.5	82Au01	
^{120}Cs^x−^{121}Cs$^x_{.496}$ ^{119}Cs$^x_{.504}$		790	30	770	13	−0.3	U			P22	2.5	82Au01	
		860	50			−0.7	U			P32	2.5	86Au02	
		813	14			−1.2	U			P32	2.5	86Au02	
^{120}Sn(n,γ)^{121}Sn		6170.3	2.	6170.2	0.3	−0.0	U					76Ca24	
		6170.5	0.7			−0.4	–					81Ba53	
		6170.1	0.4			0.3	–			Bdn		06Fi.A	
^{120}Sn(d,p)^{121}Sn		3946.2	1.7	3945.6	0.3	−0.3	–			SPa		75Be09	
^{120}Sn(n,γ)^{121}Sn	ave.	6170.2	0.3	6170.2	0.3	−0.0	1	99	95 ^{121}Sn			average	
^{121}Sb(γ,n)^{120}Sb		−9310	60	−9254	8	0.9	U			Phi		60Ge01	
		−9240	25			−0.6	U			McM		79Ba06	
^{120}Te(^{3}He,d)^{121}I		−1320.5	4.4	−1321	4	−0.1	1	99	99 ^{121}I	Hei		78Sz09	
^{121}Ba(εp)^{120}Xe		4200	300	4140	140	−0.2	R					78Bo20	
^{121}Pr(p)^{120}Ce		837	50	890	10	1.1	F					90Bo39	*
		889.6	10.				3			Arp		05Ro19	*
^{121}Ag(β^-)^{121}Cd		6400	120	6670	9	2.3	U			Stu		82Al29	
^{121}Cd(β^-)^{121}In		4780	80	4771.2	2.3	−0.1	U			Stu		82Al29	*
^{121}In^m(IT)^{121}In		313.68	0.07				3					Ens091	
^{121}In(β^-)^{121}Sn		3426	200	3351.7	1.5	−0.4	U					60Yu01	*
		3406	50			−1.1	R			Stu		78Al18	*
^{121}Sn(β^-)^{121}Sb		383	5	403.0	2.5	4.0	B					49Du15	
		383.4	3.			6.5	B					68Sn01	*
^{121}Te(β^+)^{121}Sb		1080	30	1056	26	−0.8	1	74	74 ^{121}Te			75Me23	*
^{121}I(β^+)^{121}Te		2364	50	2298	26	−1.3	1	27	26 ^{121}Te			53Fi.A	*
		2384	100			−0.9	U					65Bu03	*
^{121}Xe(β^+)^{121}I		3790	100	3765	11	−0.2	U					60Mo.A	
		4160	140			−2.8	U					70Be.A	
^{121}Cs(β^+)^{121}Xe		5650	490	5379	14	−0.6	U					75We23	
		5400	20			−1.1	–					81So06	
		5210	220			0.8	U					83Pa.A	*
		5300	100			0.8	U			IRS		93Al03	*
		5400	40			−0.5	–			JAE		96Os04	*
	ave.	5400	18			−1.2	1	61	46 ^{121}Cs			average	
^{121}Cs^x(IT)^{121}Cs		46	8				2					GAu	
^{121}Ba(β^+)^{121}Cs		6340	160	6360	140	0.1	2			JAE		96Os04	
*^{121}Ag−^{133}Cs$_{.910}$	D_M=6175.1(5.0) μu for ^{121}Ag^m at 20#20 keV; $M-A$=–74392.5(4.7) keV											Nub269	**
*^{121}Ag−^{133}Cs$_{.910}$	D_M=6180(12) μu for ^{121}Ag^m at 20#20 keV; $M-A$=–74388(11) keV											Nub269	**
*^{121}Ag−^{133}Cs$_{.910}$	D_M=6173.4(1.2) μu for ^{121}Ag^m at 20#20 keV; $M-A$=–74394.0(1.1) keV											Nub269	**
*^{121}Cd−^{130}Xe$_{.931}$	D_M=3027.4(2.9) μu for ^{121}Cd^m at 214.86(0.15) keV; $M-A$=–80858.7(2.7) keV											Nub269	**
*^{121}Cs−^{133}Cs$_{.910}$	D_M=3284(16) μu for mixture gs+m at 68.5 keV											Nub269	**
*^{121}Cs−^{133}Cs$_{.910}$	D_M=3285(13) μu for mixture gs+m at 68.5 keV; $M-A$=–77084(12) keV											Nub269	**
*^{121}Cs−u	$M-A$=–77113(29) keV for mixture gs+m at 68.5 keV											Nub269	**
*^{121}Pd−^{129}Xe$_{.938}$	Taken as low-spin isomer (see also ^{102}Y and ^{114}Tc doublets in same paper)											GAu	**
*^{121}Pr(p)^{120}Ce	F : misassigned according to reference											05Ro19	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value	Adjusted value	v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{121}$Pr(p)^{120}Ce	E_p=882(10); in publication Q_p=900(10) keV								WgM10c	**
$*^{121}$Pr(p)^{120}Ce	Trends in Mass Surface TMS suggest ^{121}Pr 100 keV less bound								GAu	**
$*^{121}$Cd(β^-)^{121}In	Q_{β^-}=4890(150); and 4960(80) from ^{121}Cd^m at 214.86 keV								Nub269	**
$*^{121}$In(β^-)^{121}Sn	E_{β^-}=3700(200) from ^{121}In^m at 313.68(0.07) to ground state and 1/2$^+$ level at 60.34 keV								Ens106	**
$*^{121}$In(β^-)^{121}Sn	E_{β^-}=2480(50) to 7/2$^+$ level at 925.59 keV								Ens106	**
$*^{121}$Sn(β^-)^{121}Sb	E_{β^-}=383(3); and 354(5) from ^{121}Sn^m at 6.31 to 7/2$^+$ level at 37.1298 keV								Ens106	**
$*^{121}$Te(β^+)^{121}Sb	p$^+$ =0.024(0.011) gives Q_{β^+}=315(30), recalculated Q_{β^+}								AHW	**
*	from ^{121}Te^m at 293.974 to 7/2$^+$ level at 37.1298 keV								Ens106	**
$*^{121}$I(β^+)^{121}Te	E_{β^+}=1130(50) 1150(100) respectively, to 3/2$^+$ level at 212.191 keV								Ens106	**
$*^{121}$Cs(β^+)^{121}Xe	E_{β^+}=3730(220) to 7/2$^+$ level at 459.59 keV								Ens106	**
$*^{121}$Cs(β^+)^{121}Xe	Q_{β^+}=5370(100) 5470(40) respectively, from ^{121}Cs^m at 68.5 keV								Ens106	**

^{122}Pd−u		−69308	397	−69368	21	−0.1	U			GT1	1.5	04Ma.A	
^{122}Ag−u		−76400	140	−76458	6	−0.2	o			GT2	2.5	08Kn.A	*
		−76450	160			−0.0	U			GT2	2.5	08Su19	*
^{122}Ag−^{133}Cs$_{.917}$		10245	96	10243	6	−0.0	U			MA8	1.0	10Br02	*
		10242.5	5.6				2			JY2	1.0	24Ja10	
^{122}Cd−^{133}Cs$_{.917}$		155.1	4.7	159.6	2.5	1.0	1	28	28 ^{122}Cd	MA8	1.0	10Br02	
^{122}In−^{133}Cs$_{.917}$		−2999.1	3.5				2			JY2	1.0	23Ne13	*
^{122}In^n−^{85}Rb$_{1.435}$		36969.8	1.4	36968.8	1.0	−0.7	1	46	46 ^{122}In^n	JY2	1.0	23Ne13	
^{122}In^n−^{133}Cs$_{.917}$		−2913.2	1.3	−2912.4	1.0	0.7	1	54	54 ^{122}In^n	JY2	1.0	23Ne13	*
C$_8$ H$_{12}$ N−^{122}Sn		193541	8	193529.3	1.4	−0.6	U			M16	2.5	63Da10	
		193558	8			−2.4	U			R13	1.5	83De51	
C$_8$ H$_{12}$ N−^{122}Te		193925	9	193930.4	1.4	0.2	U			M16	2.5	63Da10	
		193926	8			0.4	U			R13	1.5	83De51	
^{122}Xe−u		−91637	30	−91632	12	0.2	R			GS2	1.0	05Li24	
^{122}Xe−^{133}Cs$_{.917}$		−4931	13	−4932	12	−0.1	2			MA6	1.0	04Di18	
^{122}Cs−^{133}Cs$_{.917}$		2812	48	2810	40	−0.1	o			MA1	1.0	90St25	*
		2805	48			0.1	1	57	57 ^{122}Cs	MA1	1.0	99Am05	*
^{122}Cs−u		−83887	55	−83890	40	−0.1	1	43	43 ^{122}Cs	GS2	1.0	05Li24	*
^{122}Cs^n−^{133}Cs$_{.917}$		2952	16	2959	10	0.4	o			MA1	1.0	90St25	
		2961	12			−0.2	2			MA1	1.0	99Am05	
		2955	17			0.2	2			MA4	1.0	99Am05	
^{122}Ba−u		−80096	30				2			GS2	1.0	05Li24	
^{122}Cd−^{130}Xe$_{.938}$		3969.0	2.9	3967.3	2.5	−0.6	1	72	72 ^{122}Cd	JY1	1.0	12Ha25	
^{122}Pd−^{129}Xe$_{.946}$		20709	21				2			JY1	1.0	11Ha48	
^{122}Sn ^{35}Cl−^{120}Sn ^{37}Cl		4196	2	4192.8	1.3	−0.4	U			H15	4.0	62Ba23	
^{122}Ag^n−^{122}Ag		326.0	5.4				3			JY2	1.0	24Ja10	
^{122}Sn−^{122}Te		401.05	0.12	401.05	0.12	0.0	1	100	87 ^{122}Sn	JY2	1.0	25Ka40	
^{119}Cs^x−^{122}Cs$^x_{.244}$ ^{118}Cs$^x_{.756}$		−1600	80	−1511	15	0.4	U			P32	2.5	86Au02	
^{120}Cs^x−^{122}Cs$^x_{.492}$ ^{118}Cs$^x_{.508}$		−724	27	−694	20	0.4	U			P32	2.5	86Au02	
^{120}Cs^x−^{122}Cs$^x_{.328}$ ^{119}Cs$^x_{.672}$		350	50	321	16	−0.2	U			P22	2.5	82Au01	
		360	17			−0.9	U			P32	2.5	86Au02	
^{121}Cs^x−^{122}Cs$^x_{.496}$ ^{120}Cs$^x_{.504}$		−1100	40	−1066	24	0.3	U			P32	2.5	86Au02	
		−1169	15			2.7	U			P32	2.5	86Au02	
^{122}Sn(p,t)^{120}Sn		−6504	15	−6503.3	1.3	0.0	U			Roc		70Fl08	
^{122}Te(p,t)^{120}Te		−8560	24	−8612.2	1.1	−2.2	U			Win		74De31	*
^{122}Te(p,t)^{120}Te−^{132}Ba()^{130}Ba		227.0	0.2	227.00	0.20	−0.0	1	100	99 ^{120}Te			08Su14	
^{122}Te(p,t)^{120}Te−^{144}Sm()^{142}Sm		2032.6	0.4	2032.6	0.4	0.0	1	100	98 ^{142}Sm			09Bu.A	
^{122}Sn(d,^{3}He)^{121}In		−5910	50	−5890.8	1.8	0.4	U			Sac		69Co03	
		−5861	43			−0.7	U			MSU		71We01	
		−5945.6	50.			1.1	U					121Sn-0	
^{122}Sn(p,d)^{121}Sn		−6587	15	−6590.4	1.3	−0.2	U			Har		70Ca01	
^{122}Sn(d,t)^{121}Sn		−2558.8	3.0	−2557.7	1.3	0.4	1	18	13 ^{122}Sn	SPa		75Be09	
^{121}Sb(n,γ)^{122}Sb		6806.4	0.3	6806.37	0.13	−0.1	–					72Sh.A	Z
		6806.36	0.15			0.0	–			Bdn		06Fi.A	
	ave.	6806.37	0.13			−0.0	1	100	96 ^{121}Sb			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{122}In(β^-)^{122}Sn		6440	200	6386	4	−0.3	U					71Ta07	*
		6510	230			−0.5	U			Stu		78Al18	
^{122}Sn(t,^{3}He)^{122}In		−6350	50	−6367	4	−0.3	U			LAl		78Aj01	
^{122}In^n(β^-)^{122}Sn		6736	200	6466.5	1.6	−1.3	U					71Ta07	*
		6590	180			−0.7	U			Stu		78Al18	
^{122}Sb(β^+)^{122}Sn		1587	25	1605.5	2.1	0.7	U					58Pe17	
^{122}Sb(β^-)^{122}Te		1970	5	1979.1	2.1	1.8	–					55Fa33	
		1980	3			−0.3	–					68Hs02	
	ave.	1977.4	2.6			0.7	1	68	68 ^{122}Sb			average	
^{122}I(β^+)^{122}Te		4140	40	4234	5	2.3	U					54Ma75	
		4140	40			2.3	U					60Mo.A	
		4234	5				2					77Re.A	
^{122}Cs(β^+)^{122}Xe		7150	700	7210	40	0.1	U					75We23	*
		7050	180			0.9	U					83Pa.A	*
		7000	150			1.4	U			IRS		93Al03	
		7080	50			2.6	B			JAE		96Os04	
^{122}Cs^n(β^+)^{122}Xe		6950	250	7350	14	1.6	U					83Pa.A	*
		7300	150			0.3	U			IRS		93Al03	
^{122}Cs^x(IT)^{122}Cs		14	7				2					82Au01	*
*^{122}Ag−u	$M-A$=–71014(94) keV for mixture gs+n at 304(5) keV											Nub269	**
*^{122}Ag−u	$M-A$=–71065(120) keV for mixture gs+m at 304(5) keV											Nub269	**
*^{122}Ag−^{133}Cs$_{.917}$	D_M=10408(18) μu for mixture gs+m at 304(5) keV; $M-A$=–71066(17) keV											Nub269	**
*^{122}In−^{133}Cs$_{.917}$	D_M=–2994.8(1.3) for mixture gs+m at 8(4) keV; $M-A$=–83550.7(1.2) keV											Nub269	**
*^{122}In^n−^{133}Cs$_{.917}$	from frequency ratio ^{122}Inn$^+$/(^{133}Cs)$^+$ =0.917271181(10) constructed											HWJ24a	**
*^{122}In^n−^{133}Cs$_{.917}$	from ^{122}Inn$^+$/(^{122}Inx$^+$)=1.0000006694(25) and											HWJ24a	**
*^{122}In^n−^{133}Cs$_{.917}$	and ^{122}Inx$^+$/(^{133}Cs+)=0.9172705671(99)											HWJ24a	**
*^{122}Cs−^{133}Cs$_{.917}$	D_M=2887(12) μu for mixture gs+n at 140(30) keV											Nub269	**
*^{122}Cs−^{133}Cs$_{.917}$	D_M=2880(12) μu for mixture gs+n at 140(30) keV; $M-A$=–78078(12) keV											Nub269	**
*^{122}Cs−u	$M-A$=–78070(28) keV for mixture gs+m at 140(30) keV											Nub269	**
*^{122}Te(p,t)^{120}Te	Original error 12; added systematic error 21 keV											GAu	**
*^{122}In(β^-)^{122}Sn	E_{β^-}=5300(200) to 2^+ level at 1140.51 keV											Ens074	**
*^{122}In^n(β^-)^{122}Sn	E_{β^-}=4400(200) to 4^+ level at 2331.09 keV											Ens074	**
*^{122}Cs(β^+)^{122}Xe	E_{β^+}=5800(700) 5690(180) respectively, to 2^+ level at 331.28 keV											Ens074	**
*^{122}Cs^n(β^+)^{122}Xe	E_{β^+}=3710(250) to 8^+ level at 2217.69 keV											Ens074	**
*^{122}Cs^x(IT)^{122}Cs	Original was 45(33); revised using ^{122}Cs^n=140(30) keV											Nub269	**
^{123}Pd−u		−64423	290	−64730	270	−0.7	o			GT1	1.5	04Ma.A	
		−64874	339			0.2	2			GT3	2.5	16Kn03	
		−64715	285			−0.1	o			RR3	1.0	22Li20	
		−64715	285			−0.1	2			RR3	1.0	23Na12	
^{123}Ag−u		−74729	215	−74722	4	0.0	o			GT1	1.5	04Ma.A	
		−74479	130			−0.7	U			GT2	2.5	08Su19	*
^{123}Ag−^{133}Cs$_{.925}$		12700	120	12735	4	0.3	U			MA8	1.0	10Br02	
		12772	35			−1.1	2			MA8	1.0	10Br02	*
		12734.1	3.9			0.1	2			JY2	1.0	24De01	
^{123}Cd−^{133}Cs$_{.925}$		4491	52	4349.3	2.9	−2.7	U			MA8	1.0	10Br02	*
^{123}In−^{133}Cs$_{.925}$		−2075.1	1.2				2			JY2	1.0	23Ne13	
C_8 H_{13} N−^{123}Sb		200580.0	3.3	200584.8	1.4	0.6	U			M16	2.5	63Da10	
		200615	8			−2.5	U			R13	1.5	83De51	
C_8 H_{13} N−^{123}Te		200538	16	200529.1	1.4	−0.2	U			M16	2.5	63Da10	
		200515	8			1.2	U			R13	1.5	83De51	
^{123}Te−u		−95615	83	−95729.7	1.4	−1.4	U			GS2	1.0	05Li24	*
^{123}I−u		−94444	30	−94411	4	1.1	U			GS2	1.0	05Li24	
^{123}Xe−^{133}Cs$_{.925}$		−4048	13	−4061	10	−1.0	1	62	62 ^{123}Xe	MA6	1.0	04Di18	
^{123}Cs−u		−87007	57	−87004	13	0.1	U			GS2	1.0	05Li24	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{123}Cs−^{133}Cs$_{.925}$		447	17	453	13	0.4	o			MA1	1.0	90St25	
		453	13				2			MA1	1.0	99Am05	
^{123}Ba−^{133}Cs$_{.925}$		6238	13				2			MA5	1.0	00Be42	
^{123}Ba−u		−81327	30	−81219	13	3.6	C			GS2	1.0	05Li24	
^{123}Cd−^{130}Xe$_{.946}$		8172.5	2.9				2			JY1	1.0	12Ha25	
^{123}Cd^m−^{130}Xe$_{.946}$		8326.5	3.3				2			JY1	1.0	13Ka08	
^{123}Sb ^{35}Cl−^{121}Sb ^{37}Cl		3343	2	3354.0	2.3	1.4	U			H14	4.0	62Ba24	
^{123}Ag^m−^{123}Ag		61.3	9.5	63.7	0.8	0.2	U			JY2	1.0	24De01	
^{123}In^m−^{123}In		351.04	0.43	351.27	0.04	0.5	U			JY2	1.0	23Ne13	
^{123}Te−^{123}Sb		55.729	0.071	55.73	0.07	0.0	1	100	100 ^{123}Sb	SH2	1.0	16Fi07	
^{119}Cs^x−^{123}Cs$^x_{.193}$ ^{118}Cs$^x_{.807}$		−1480	60	−1447	13	0.2	U			P32	2.5	86Au02	
^{123}Te(n,α)^{120}Sn		7564	30	7572.6	1.2	0.3	U			ILL		75Em04	
^{121}Sb(t,p)^{123}Sb		7295	20	7284.6	2.1	−0.5	U			Ald		67Hi01	
^{122}Sn(n,γ)^{123}Sn		5948	3	5945.9	1.1	−0.7	–					75Bh01	
		5945.8	1.5			0.1	–					77Ca09	
^{122}Sn(d,p)^{123}Sn		3726	12	3721.4	1.1	−0.4	U			Tal		64Ne10	
		3716	11			0.5	U					72Ca33	
		3721.8	2.6			−0.2	–			SPa		75Be09	
^{122}Sn(n,γ)^{123}Sn	ave.	5946.3	1.2	5945.9	1.1	−0.3	1	91	90 ^{123}Sn			average	
^{123}Sb(γ,n)^{122}Sb		−8980	50	−8960.0	2.1	0.4	U			Phi		60Ge01	
		−8966	4			1.5	1	28	28 ^{122}Sb	McM		79Ba06	
^{122}Te(n,γ)^{123}Te		6937	5	6929.01	0.08	−1.6	U					68Ch.A	
		6929.1	0.5			−0.2	U					91Ho08	
		6928.97	0.09			0.5	–					00Bo24	
		6929.16	0.17			−0.9	–			Bdn		06Fi.A	
^{122}Te(d,p)^{123}Te		4706	6	4704.45	0.08	−0.3	U			MIT		75Li22	
^{122}Te(n,γ)^{123}Te	ave.	6929.01	0.08	6929.01	0.08	0.0	1	100	86 ^{122}Te			average	
^{122}Te(^{3}He,d)^{123}I		−574.2	3.5	−575	3	−0.3	1	97	96 ^{123}I	Hei		78Sz04	
^{123}Ag^m(IT)^{123}Ag		59.3	0.7				3					19Ch24	*
^{123}Cd(β^-)^{123}In		6033	35	5984.3	2.9	−1.4	U			Stu		87Sp09	*
^{123}In^m(IT)^{123}In		327.21	0.04				3					Ens214	
^{123}In(β^-)^{123}Sn		4400	30	4416.4	2.0	0.5	U			Stu		87Sp09	*
^{123}Sn(β^-)^{123}Sb		1395	10	1408.6	1.1	1.4	U					49Du15	*
		1420	10			−1.1	U					50Ke11	
		1399	20			0.5	U					66Au04	*
^{123}I(β^+)^{123}Te		1260	7	1228	3	−4.5	C					86Ag.A	
^{123}Xe(β^+)^{123}I		2720	100	2695	10	−0.3	U					54Ma75	
		2676	15			1.2	1	42	38 ^{123}Xe			60Mo.A	*
^{123}Cs(β^+)^{123}Xe		4110	310	4205	15	0.3	U					75We23	*
		4000	140			1.5	U					81So06	*
		4050	180			0.9	U					83Pa.A	*
		4200	100			0.0	U			IRS		93Al03	
		4110	30			3.2	B			JAE		96Os04	
^{123}Cs^x(IT)^{123}Cs		7	4				3					82Au01	*
^{123}Ba(β^+)^{123}Cs		5330	100	5389	17	0.6	U			JAE		96Os04	
*^{123}Ag−u	Isomer should be expected, but ^{123}Ag^m at 20#(20#) keV, correction negligible											GAu	**
*^{123}Ag−^{133}Cs$_{.925}$	D_M=12805(30) μu for mixture gs+m at 59.3(0.7) keV; $M-A$=–69538(28)											Nub269	**
*^{123}Cd−^{133}Cs$_{.925}$	D_M=4568(26) μu for mixture gs+m at 143(4) keV; $M-A$=–77210(25)											Nub269	**
*^{123}Te−u	$M-A$=–88941(30) keV for mixture gs+m at 247.45 keV											Nub269	**
*^{123}Cs−u	$M-A$=–80968(28) keV for mixture gs+m at 156.27 keV											Nub269	**
*^{123}Ag^m(IT)^{123}Ag	1009.2(0.3)+27.2(0.4)-594.0(0.3)-383.1(0.3) in 19Ch24											FGK259	**
*^{123}Cd(β^-)^{123}In	Q =3590(51) 3464(41) 3547(36) from ground state to 2393 2529 2541 levels											89Hu03	**
*^{123}Cd(β^-)^{123}In	Q =3624(21) 3710(37) 3513(59) from ^{123}Cd^m at 143(4) to 2529 2461 2602 levels											Nub269	**
*^{123}In(β^-)^{123}Sn	Q_{β^-}=4410(31); and 4645(72) from ^{123}In^m at 327.21 keV											Nub269	**
*^{123}Sn(β^-)^{123}Sb	E_{β^-}=1260(10) from ^{123}Sn^m at 24.6 to 5/2$^+$ level at 160.33 keV											Ens04a	**
*^{123}Sn(β^-)^{123}Sb	E_{β^-}=310(20) to 9/2$^+$ level at 1088.64 keV											Ens04a	**
*^{123}Xe(β^+)^{123}I	E_{β^+}=1505(15) to 1/2$^+$ level at 148.92 keV											Ens04a	**
*^{123}Cs(β^+)^{123}Xe	E_{β^+}=2990(310) to 3/2$^+$ level at 97.30 keV											Ens04a	**
*^{123}Cs(β^+)^{123}Xe	E_{β^+}=2370(140) to (1/2,3/2)$^+$ level at 596.65 keV, and other E_{β^+}											Ens04a	**
*^{123}Cs(β^+)^{123}Xe	E_{β^+}=2930(180) to 3/2$^+$ level at 97.30 keV											Ens04a	**
*^{123}Cs^x(IT)^{123}Cs	Based on ^{123}Cs^m(IT)=156.27 and isomeric ratio R<0.1											Nub269	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{124}Pd−u		−64617	399	−62580#	320#	2.0	D			GT3	2.5	16Kn03	*
^{124}Ag−^{133}Cs$_{.932}$		16950	270	17051	7	0.4	U			MA8	1.0	10Br02	*
		17050.6	7.4				2			JY2	1.0	25Ru02	
^{124}Ag−u		−71101	273	−71068	7	0.1	U			RR3	1.0	23Na12	
^{124}Cd−^{133}Cs$_{.932}$		5781	10	5778.6	2.8	−0.2	–			MA8	1.0	10Br02	
		5786.1	5.7			−1.3	–			MA8	1.0	20Ma09	
	ave.	5785	5			−1.3	1	32	32 ^{124}Cd			average	
^{124}In−^{133}Cs$_{.932}$		1257.8	3.5				2			JY2	1.0	23Ne13	
C_7 ^{13}C H_{13} N−^{124}Sn		202886	8	202874.4	1.4	−0.6	U			M16	2.5	63Da10	
		202891	8			−1.4	U			R13	1.5	83De51	
^{124}Sn−^{13}C $^{37}Cl_3$		4210.47	0.71	4217.3	1.4	2.4	U			H39	4.0	84Ha20	
^{124}Sn−^{133}Cs$_{.932}$		−6598	21	−6601.4	1.4	−0.2	U			MA8	1.0	05Si34	
^{124}Te−^{13}C $^{37}Cl_3$		1754.63	1.26	1755.1	1.4	0.1	U			H39	4.0	84Ha20	
^{124}Te−^{54}Fe $^{35}Cl_2$		25501.65	2.56	25503.4	1.4	0.2	U			H39	4.0	84Ha20	
C_7 ^{13}C H_{13} N−^{124}Te		205336	13	205336.6	1.4	0.0	U			M16	2.5	63Da10	
		205325	8			1.0	U			R13	1.5	83De51	
^{124}I−u		−93786	30	−93790.4	2.4	−0.1	U			GS2	1.0	05Li24	
^{124}Xe−^{13}C $^{37}Cl_3$		4831.15	1.58	4821.9	1.4	−1.5	U			H39	4.0	84Ha20	
^{124}Xe−^{54}Fe $^{35}Cl_2$		28575.78	0.99	28570.2	1.4	−1.4	U			H39	4.0	84Ha20	
^{124}Xe−^{133}Cs$_{.932}$		−5986	13	−5996.8	1.4	−0.8	U			MA6	1.0	04Di18	
^{124}Cs−^{133}Cs$_{.932}$		361	16	366	10	0.3	o			MA1	1.0	90St25	
		370	13			−0.3	2			MA1	1.0	99Am05	
		361	15			0.3	2			MA8	1.0	05Gu37	
^{124}Cs−u		−87696	30	−87753	10	−1.9	U			GS2	1.0	05Li24	
		−87693	30			−2.0	U			GS2	1.0	05Li24	*
		−87708	42			−1.1	U			GR1	1.0	19An10	*
^{124}Ba−^{133}Cs$_{.932}$		3212	15	3212	13	0.0	2			MA1	1.0	99Am05	
^{124}Ba−u		−84905	30	−84906	13	−0.0	R			GS2	1.0	05Li24	
^{124}La−u		−75464	71	−75430	60	0.5	2			GS2	1.0	05Li24	*
^{124}Cd−^{130}Xe$_{.954}$		9708.9	3.4	9711.9	2.8	0.9	1	68	68 ^{124}Cd	JY1	1.0	12Ha25	
^{124}Sn−^{129}Xe$_{.961}$		−3214.3	2.1	−3214.5	1.4	−0.1	1	42	42 ^{124}Sn	JY1	1.0	11Ha48	
^{124}Sn−^{120}Sn$_{1.033}$		6305.1	2.1	6304.8	1.3	−0.2	1	41	30 ^{124}Sn	JY1	1.0	11Ha48	
^{124}Sn ^{35}Cl−^{122}Sn ^{37}Cl		4784	2	4784.93	0.21	0.1	U			H15	4.0	62Ba23	
^{124}Te ^{35}Cl−^{122}Te ^{37}Cl		2728	2	2723.76	0.15	−0.5	U			H16	4.0	63Ba47	
^{124}Ag^m−^{124}Ag		202.1	2.7				3			JY2	1.0	25Ru02	
^{124}In^m−^{124}In		25.9	2.9				3			JY2	1.0	23Ne13	
^{124}Sn−^{124}Te		2458.51	0.89	2462.22	0.09	1.0	U			H39	4.0	84Ha20	
		2461.26	0.42			2.3	U			SH1	1.0	12Ne10	
		2462.221	0.091			−0.0	1	100	73 ^{124}Te	JY2	1.0	25Ka40	
^{124}Xe−^{124}Te		3076.00	1.78	3066.83	0.14	−1.3	U			H39	4.0	84Ha20	
		3066.83	0.14			0.0	1	100	99 ^{124}Xe	SH1	1.0	12Ne10	
^{124}Sn−^{122}Sn		1838	22	1834.80	0.20	−0.1	U			M16	2.5	63Da10	
^{120}Cs^x−^{124}Cs$^x_{.194}$ ^{119}Cs$^x_{.807}$		310	30	306	12	−0.1	U			P22	2.5	82Au01	
^{121}Cs^x−^{124}Cs$^x_{.244}$ ^{120}Cs$^x_{.756}$		−1360	30	−1263	19	1.3	U			P22	2.5	82Au01	
^{123}Cs^x−^{124}Cs$^x_{.744}$ ^{120}Cs$^x_{.256}$		−1390	30	−1329	21	0.8	U			P22	2.5	82Au01	
^{124}Sn(d,^{6}Li)^{120}Cd		−5216	24	−5225	4	−0.4	U					79Ja21	
^{124}Sn(^{3}He,^{7}Be)^{120}Cd		−5098	30	−5112	4	−0.5	U			MSU		76St11	
^{124}Sn(^{18}O,^{20}Ne)^{122}Cd		−1266	39	−1359.7	2.6	−2.4	U					97Gu32	*
^{122}Sn(t,p)^{124}Sn		5931	15	5951.73	0.19	1.4	U			Roc		70Fl05	
^{124}Sn(p,t)^{122}Sn		−5956	10	−5951.73	0.19	0.4	U					64Al29	
^{124}Sn(d,^{3}He)^{123}In		−6610	50	−6628.2	1.7	−0.4	U			Sac		69Co03	
		−6572	66			−0.9	U			MSU		71We01	
^{124}Sn(p,d)^{123}Sn		−6279	15	−6263.0	1.1	1.1	U			Har		70Ca01	
^{124}Sn(d,t)^{123}Sn		−2260	35	−2230.4	1.1	0.8	U			Pit		64Co11	
		−2233.4	3.7			0.8	1	10	10 ^{123}Sn	SPa		75Be09	
^{123}Sb(n,γ)^{124}Sb		6467.55	0.10	6467.50	0.06	−0.5	2					73Sh.A	Z
		6467.40	0.10			1.0	2					81Su.A	Z
		6467.58	0.14			−0.6	2			Bdn		06Fi.A	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{123}Te(n,γ)^{124}Te		9425	2	9424.48	0.09	−0.3	U					69Bu05	
		9423.7	1.5			0.5	U					70Or.A	
		9424.05	0.30			1.4	–			Ltn		95Ge06	Z
		9423.89	0.20			3.0	C			Bdn		06Fi.A	
		9424.53	0.10			−0.5	–					06Vo09	
	ave.	9424.48	0.09			0.0	1	100	86 ^{123}Te			average	
^{124}Cd(β^-)^{124}In		4166	39	4211	4	1.2	U			Stu		87Sp09	
^{124}In(β^-)^{124}Sn		7180	50	7321	3	2.8	B			Stu		78Al18	
		7360	49			−0.8	U			Stu		87Sp09	
^{124}Sn(t,^{3}He)^{124}In		−7590	50	−7302	3	5.8	B			LAl		78Aj01	
^{124}In^m(β^-)^{124}Sn		7370	210	7345	4	−0.1	o			Stu		78Al18	
		7341	51			0.1	U			Stu		87Sp09	
^{124}Sb(β^-)^{124}Te		2907.7	5.	2905.07	0.13	−0.5	U					65Hs02	*
		2903.7	4.			0.3	U					66Ca10	*
		2904.7	2.			0.2	U					69Na05	*
^{124}I(β^+)^{124}Te		3157	4	3159.6	1.9	0.6	2					71Bo01	*
		3160.3	2.1			−0.3	2					92Wo03	
^{124}Cs(β^+)^{124}Xe		5920	460	5927	9	0.0	U					75We23	
		5900	90			0.3	U			IRS		93Al03	
		5910	30			0.6	U			JAE		96Os04	
^{124}Cs^x(IT)^{124}Cs		30	20				3					AHW	*
^{124}La(β^+)^{124}Ba		8930	110	8830	60	−0.9	R			JAE		98Ko66	
*^{124}Pd−u	Trends in Mass Surface TMS suggest ^{124}Pd 1900 keV less bound											HWJ268	**
*^{124}Ag−^{133}Cs$_{.932}$	D_M=17050(270) μu for mixture at gs+m at 188.3(2.5) keV; $M-A$=–66200(250)											Nub269	**
*^{124}Cs−u	$M-A$=–81223(28) keV for ^{124}Cs^m at 462.63 keV											Nub269	**
*^{124}La−u	$M-A$=–70244(32) keV for mixture gs+m at 100#100 keV											Nub269	**
*^{124}Sn(^{18}O,^{20}Ne)^{122}Cd	Original Q =–1250(39) calibrated with ^{120}Cd=–83973(19) keV											GAu	**
*^{124}Sb(β^-)^{124}Te	E_{β^-}=2305(5) 2301(4) 2302(2) respectively, to 2$^+$ level at 602.7271 keV											Ens087	**
*^{124}I(β^+)^{124}Te	Original error increased, see ^{84}Rb(β^+)											AHW	**
*^{124}Cs^x(IT)^{124}Cs	Based on ^{124}Cs^m(IT)=462.63 keV											Nub269	**
*^{124}Cs^x(IT)^{124}Cs	Isomeric ratio assumed <0.1 as for ^{118}Cs, ^{120}Cs, ^{122}Cs											AHW	**
^{125}Ag−u		−68954	429	−68994	4	−0.1	o			GT1	1.5	04Ma.A	
		−69265	186			0.6	U			GT3	2.5	16Kn03	
^{125}Cd−u		−78770	120	−78742	3	0.1	o			GT2	2.5	08Kn.A	*
		−78780	140			0.1	U			GT2	2.5	08Su19	*
		−78623	344			−0.3	o			RR3	1.0	22Li20	*
		−78623	344			−0.3	U			RR3	1.0	23Na12	
^{125}In−u		−91274	120	−86326.2	1.9	41.2	U			TR1	1.0	21Iz01	*
^{125}In N$_2$−^{153}Nd		−7900.2	27.5	−7901.3	2.7	−0.0	U			RI1	1.0	24Ki16	*
^{125}Sn N$_2$−^{153}Nd		−13712	175	−13785.6	2.3	−0.4	U			RI1	1.0	24Ki16	*
C$_7$ H$_6$ ^{35}Cl−^{125}Te		111363	6	111372.4	1.4	0.6	U			M16	2.5	63Da10	
		111368	8			0.4	U			R13	1.5	83De51	
^{125}I−u		−95374	30	−95370.1	1.4	0.1	U			GS2	1.0	05Li24	
^{125}Cs−u		−90280	30	−90274	8	0.2	U			GS2	1.0	05Li24	
		−90221	45			−1.2	U			GR1	1.0	19An10	
^{125}Ba−u		−85569	30	−85528	12	1.4	R			GS2	1.0	05Li24	
^{125}La−u		−79191	30	−79184	28	0.2	1	87	87 ^{125}La	GS2	1.0	05Li24	
^{125}Ag−^{133}Cs$_{.940}$		19878.1	4.4	19881	4	0.6	1	79	79 ^{125}Ag	JY2	1.0	25Ru02	
^{125}Ag^m−^{133}Cs$_{.940}$		19995.2	8.5	19985	4	−1.2	1	21	21 ^{125}Ag^m	JY2	1.0	25Ru02	
^{125}Cd−^{133}Cs$_{.940}$		10133	29	10133	3	−0.0	U			TT1	1.0	17La16	
^{125}Cd^m−^{133}Cs$_{.940}$		10337	11	10333	3	−0.4	U			TT1	1.0	17La16	
^{125}In−^{133}Cs$_{.940}$		2549.0	1.9				2			TT1	1.0	18Ba08	
^{125}In^m−^{133}Cs$_{.940}$		2927	13	2935.6	1.9	0.7	U			TT1	1.0	18Ba08	
^{125}Cs−^{133}Cs$_{.940}$		−1392	17	−1399	8	−0.4	o			MA1	1.0	90St25	
		−1382	14			−1.2	–			MA1	1.0	99Am05	
		−1386	14			−0.9	–			MA4	1.0	99Am05	
	ave.	−1384	10			−1.5	1	70	70 ^{125}Cs			average	
^{125}Ba−^{133}Cs$_{.940}$		3356	13	3347	12	−0.7	–			MA5	1.0	00Be42	
	ave.	3348	12			−0.1	1	98	98 ^{125}Ba			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{125}Cd−^{130}Xe$_{,962}$	14081.6	3.1				2			JY1	1.0	12Ha25	
^{125}Cd^{m}−^{130}Xe$_{,962}$	14281.6	3.4				2			JY1	1.0	13Ka08	
^{125}Te ^{35}Cl−^{123}Te ^{37}Cl	3090	2	3110.28	0.13	2.5	U			H16	4.0	63Ba47	
^{122}Cs^{x}−^{125}Cs$_{,244}$ ^{121}Cs$^{x}_{,756}$	715	23	640	40	−1.3	U			P32	2.5	86Au02	
^{123}Sb(t,p)^{125}Sb	6696	20	6693.0	2.1	−0.1	U			Ald		67Hi01	
^{124}Sn(n,γ)^{125}Sn	5733.1	1.5	5733.50	0.20	0.3	U					77Ca09	Z
	5733.1	0.6			0.7	U					81Ba53	
	5733.5	0.2				2					11To04	
^{124}Sn(d,p)^{125}Sn	3530	30	3508.93	0.20	−0.7	U			Pit		64Co11	
	3506	12			0.2	U			Tal		64Ne10	
	3515	11			−0.6	U					72Ca33	
	3509.4	3.6			−0.1	U			SPa		75Be09	
^{124}Te(n,γ)^{125}Te	6569.0	1.0	6568.970	0.030	−0.0	U					71Gr.A	
	6568.97	0.03			0.0	1	100	88 ^{125}Te	Prn		99Ho01	
	6569.39	0.19			−2.2	U			Bdn		06Fi.A	
^{125}Te(γ,n)^{124}Te	−6560	60	−6568.970	0.030	−0.1	U			Phi		60Ge01	
^{124}Te(d,p)^{125}Te	4344	8	4344.404	0.030	0.1	U			MIT		69Gr24	
^{124}Te(^{3}He,d)^{125}I	115.1	3.0	107.38	0.07	−2.6	U			Hei		78Sz04	
^{124}Te(α,t)^{125}I	−14203	7	−14213.01	0.07	−1.4	U			Hei		78Sz04	
^{124}Xe(n,γ)^{125}Xe	7603.3	0.4	7603.3	0.4	−0.1	1	100	99 ^{125}Xe			82Ka.A	
^{125}Ag^{m}(IT)^{125}Ag	97.1	0.5	97.1	0.5	0.1	1	100	79 ^{125}Ag^{m}			19Ch24	*
^{125}Cd(β^-)^{125}In	7122	62	7064	3	−0.9	U			Stu		87Sp09	*
^{125}Cd^{m}(β^-)^{125}In	7172	35	7251	4	2.2	B			Stu		87Sp09	*
^{125}In^{m}(IT)^{125}In	360.12	0.09				3					Ens101	
^{125}In(β^-)^{125}Sn	5418	30	5481.1	2.2	2.1	U			Stu		87Sp09	*
^{125}Sn(β^-)^{125}Sb	2330	10	2362.3	2.1	3.2	B					50Ha58	
	2370	20			−0.4	U					50Ke11	
	2335	40			0.7	U					64De02	*
^{125}Sb(β^-)^{125}Te	767.7	3.	766.7	2.1	−0.3	2					64Ma30	*
	765.7	3.			0.3	2					66Ma49	*
^{125}I(ε)^{125}Te	184	7	185.77	0.06	0.3	U					64Le05	*
	185	8			0.1	U					66Sm05	*
	177.2	2.			4.3	C					68Go.A	*
	186.1	0.3			−1.1	U					86Bo46	
	179.3	2.0			3.2	B					90Li14	*
	185.77	0.06				2					94Hi04	
^{125}Xe(ε)^{125}I	1735	40	1636.7	0.4	−2.5	U					69Lu09	*
^{125}Cs(β^+)^{125}Xe	3072	20	3110	8	1.9	–					54Ma54	
	3082	20			1.4	–					75We23	
	3100	100			0.1	U			IRS		93Al03	
ave.	3077	14			2.3	1	30	30 ^{125}Cs			average	
^{125}Ba(β^+)^{125}Cs	4560	250	4421	13	−0.6	U					68Da09	*
	4380	50			0.8	U			JAE		96Os04	
^{125}La(β^+)^{125}Ba	5950	70	5909	28	−0.6	1	16	13 ^{125}La	JAE		98Ko66	
*^{125}Cd−u	$M-A$=–73274(93) keV for mixture gs+m at 186(4) keV										Nub269	**
*^{125}Cd−u	$M-A$=–73287(120) keV for mixture gs+m at 186(4) keV										Nub269	**
*^{125}Cd−u	mixture gs+m at 186(4), not corrected										Nub269	**
*^{125}In−u	mixture gs+m is corrected by the author										21Iz01	**
*^{125}In N$_2$−^{153}Nd	Reported $M-A$=–74690(42) keV, author confirmed mistake in the unc.										24Ki16	**
*^{125}Sn N$_2$−^{153}Nd	D_M=–13697.6(24.5) μu, $M-A$=–80089.7(27.8) keV for mixture of ground state and m										24Ki16	**
*^{125}Sn N$_2$−^{153}Nd	at 27.50(0.14) keV; reported $M-A$=–80089(40) keV, author confirmed										24Ki16	**
*^{125}Sn N$_2$−^{153}Nd	mistake in the uncertainty										24Ki16	**
*^{125}Ag^{m}(IT)^{125}Ag	From 1118.5(0.3)-606.3(0.3)-415.1(0.3) keV in 19Ch24										FGK259	**
*^{125}Cd(β^-)^{125}In	E_{β^-}=4625(62) to (1/2$^+$,3/2$^+$) level at 2497.43 keV										Ens112	**
*^{125}Cd^{m}(β^-)^{125}In	E_{β^-}=5009(109), 4581(126), 4533(39) to 2101.40, 2640.29, 2641.26 levels										Ens112	**
*^{125}In(β^-)^{125}Sn	Q_{β^-}=5443(31); and 5730(43) from ^{125}In^{m} at 360.12 keV										Nub269	**
*^{125}Sn(β^-)^{125}Sb	E_{β^-}=2030(40) from ^{125}Sn^{m} at 27.50(0.14) to 5/2$^+$ level at 332.06 keV										Ens112	**
*^{125}Sb(β^-)^{125}Te	E_{β^-}=623(3) 621(3) respectively, to 1/2$^-$ level at 144.775 keV										Ens112	**
*^{125}I(ε)^{125}Te	LMK=0.254(0.003) 0.253(0.005) IBE=110(2) 150.6(0.3) respectively, all to										AHW	**
*	3/2$^+$ level at 35.4925 keV. Q(LMK) recalculated, error mainly theory										Ens112	**
*^{125}I(ε)^{125}Te	IBE=112.0(2.0)(1s)+31.8 to 3/2$^+$ level at 35.4925 keV										Ens112	**
*^{125}Xe(ε)^{125}I	E_{β^+}=470(40) to 3/2$^+$ at 188.416 and 1/2$^+$ at 243.382 keV, ratio 1:2										Ens112	**
*^{125}Ba(β^+)^{125}Cs	E_{β^+}=3450(250) to (5/2$^+$) level at 84.82 level										Ens112	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{126}Ag−u	−65926	329	−65190#	220#	0.9	D			GT3	2.5	16Kn03	*
^{126}In−u	−83420	290	−83532	5	−0.4	o			RR3	1.0	22Li20	
	−83506	47			−0.5	U			TR1	1.0	21Iz01	*
	−83422	289			−0.4	U			RR3	1.0	23Na12	
^{126}Sn ^{14}N$_2$−^{154}Nd	−15780.4	8.6	−15784	7	−0.4	1	64	64 ^{126}Sn	RI1	1.0	24Ki16	*
C$_{10}$ H$_6$−^{126}Te	143623	9	143638.7	1.4	0.7	U			M16	2.5	63Da10	
	143640	8			−0.1	U			R13	1.5	83De51	
^{126}Xe−u	−95647	30	−95702.580	0.012	−1.9	U			GS2	1.0	05Li24	
^{126}Cs−u	−90543	49	−90554	11	−0.2	U			GR1	1.0	19An10	
^{126}Ba−u	−88745	30	−88750	13	−0.2	R			GS2	1.0	05Li24	
^{126}La−u	−80503	232	−80490	100	0.1	2			GS2	1.0	05Li24	*
^{126}Ce−u	−76029	30				2			GS2	1.0	05Li24	
^{126}Xe−^{134}Xe$_{.940}$	−6772.8	2.9	−6772.028	0.014	0.3	o			MA8	1.0	05He.A	
	−6773.2	3.8			0.3	U			MA8	1.0	06He29	
^{126}Cd−^{133}Cs$_{.947}$	11966.5	4.5	11967.3	2.5	0.2	–			MA8	1.0	10Br02	
	11956	15			0.8	U			MA8	1.0	10Br02	
	11962	11			0.5	U			TT1	1.0	17La16	
	11973.7	6.7			−1.0	–			MA8	1.0	20Ma09	
ave.	11969	4			−0.4	1	44	44 ^{126}Cd			average	
^{126}In−^{133}Cs$_{.947}$	6005.2	4.5				2			TT1	1.0	18Ba08	
^{126}In^m−^{133}Cs$_{.947}$	6101.7	5.5				2			TT1	1.0	18Ba08	
^{126}Cs−^{133}Cs$_{.947}$	−1027	17	−1017	11	0.6	o			MA1	1.0	90St25	
	−1011	13			−0.5	1	73	73 ^{126}Cs	MA1	1.0	99Am05	
^{126}Ba−^{133}Cs$_{.947}$	786	15	787	13	0.1	2			MA1	1.0	99Am05	
^{126}Cd−^{130}Xe$_{.969}$	15928.6	3.3	15929.7	2.5	0.3	1	56	56 ^{126}Cd	JY1	1.0	12Ha25	
^{126}Te ^{35}Cl−^{124}Te ^{37}Cl	3432	2	3443.93	0.11	1.5	U			H16	4.0	63Ba47	
	3441.28	1.54			1.1	U			H43	1.5	90Dy04	
^{126}Xe−^{128}Xe$_{.984}$	−776.8366	0.0079	−776.837	0.008	0.0	1	100	100 ^{126}Xe	Hep	1.0	20Ri04	
^{123}Cs^x−^{126}Cs$_{.390}$ ^{121}Cs$^x_{.610}$	−1160	30	−1136	17	0.3	U			P22	2.5	82Au01	
^{124}Cs^x−^{126}Cs$_{.590}$ ^{121}Cs$^x_{.410}$	−340	30	−351	24	−0.1	U			P22	2.5	82Au01	
^{124}Cs^x−^{126}Cs$_{.492}$ ^{122}Cs$^x_{.508}$	−570	30	−520	29	0.7	U			P22	2.5	82Au01	
^{124}Cs^x−^{126}Cs$_{.328}$ ^{123}Cs$^x_{.672}$	390	30	413	24	0.3	U			P22	2.5	82Au01	
^{125}Cs−^{126}Cs$_{.496}$ ^{124}Cs$^x_{.504}$	−1130	30	−1070	14	0.8	U			P22	2.5	82Au01	
^{124}Sn(t,p)^{126}Sn	5445	15	5439	6	−0.4	–			Ald		69Bj01	
	5444	15			−0.3	–			Roc		70Fl05	
ave.	5445	11			−0.5	1	37	36 ^{126}Sn			average	
^{125}Te(n,γ)^{126}Te	9113.7	0.4	9113.69	0.08	−0.0	U					77Ko.A	
	9113.69	0.08			0.0	1	100	88 ^{126}Te			03Vo03	
^{126}Te(γ,n)^{125}Te	−8840	120	−9113.69	0.08	−2.3	U			Phi		60Ge01	
^{125}Te(d,p)^{126}Te	6892	6	6889.13	0.08	−0.5	U			MIT		71Gr01	
^{126}Cd(β^-)^{126}In	5486	36	5554	5	1.9	U			Stu		87Sp09	
^{126}In(β^-)^{126}Sn	8207	39	8200	8	−0.2	U			Stu		87Sp09	
^{126}In^m(β^-)^{126}Sn	8309	51	8290	8	−0.4	U			Stu		87Sp09	
^{126}Sn(β^-)^{126}Sb	378	30				2					71Or04	*
^{126}Sb(β^-)^{126}Te	3667	150	3680	30	0.1	U					71Or04	*
^{126}I(β^+)^{126}Te	2151	5	2154	4	0.6	1	54	52 ^{126}I			59Ha27	
^{126}I(β^-)^{126}Xe	1258	5	1235	4	−4.5	B					55Ko14	*
^{126}Cs(β^+)^{126}Xe	4670	140	4796	10	0.9	U					75We23	*
	4810	100			−0.1	U					76Pa11	*
	4830	40			−0.9	U			JAE		92Os07	
	4730	100			0.7	U			IRS		93Al03	
	4780	20			0.8	1	27	27 ^{126}Cs	JAE		96Os04	
^{126}La(β^+)^{126}Ba	7700	100	7700	90	−0.0	R			JAE		98Ko66	
^{126}La^m(β^+)^{126}Ba	7910	400				3			JAE		98Ko66	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
$*^{126}$Ag−u		Trends in Mass Surface TMS suggest ^{126}Ag 690 keV less bound											GAu	**
$*^{126}$In−u		mixture gs+m is corrected by the author											21Iz01	**
$*^{126}$Sn ^{14}N$_2$−^{154}Nd		Reported $M-A$=–80279(28) keV, author confirmed mistake in the unc.											24Ki16	**
$*^{126}$La−u		$M-A$=–74883(28) keV for mixture gs+m at 210(410) keV											Nub269	**
$*^{126}$Sn(β^-)^{126}Sb		E_{β^-}=250(30) to 2^+ level at 127.9 keV											Ens031	**
$*^{126}$Sb(β^-)^{126}Te		E_{β^-}=1900(150) from mixture ground state and ^{126}Sb^m at 17.7 to 6^+ level at 1776.19 keV											Ens031	**
$*^{126}$I(β^-)^{126}Xe		E_{β^-}=865(5) to 2^+ level at 388.631 keV, and other E_{β^-}											Ens031	**
$*^{126}$Cs(β^+)^{126}Xe		E_{β^+}=3260(140) 3400(100) respectively, to 2^+ level at 388.631 keV											Ens031	**
^{127}In−u		−82527	40	−82546.2	1.3	−0.5	U				TR1	1.0	21Iz01	
^{127}In^n−u		−80651	39	−80690.4	1.8	−1.0	U				TR1	1.0	21Iz01	
^{127}Sn N$_2$−^{155}Pm		−11591.4	76.8	−11600	10	−0.1	U				RI1	1.0	24Ki16	*
^{127}Sb N$_2$−^{155}Pm		−15028.1	12.9	−15067	6	−3.0	B				RI1	1.0	24Ki16	*
C$_{10}$ H$_7$−^{127}I		150297	6	150303	4	0.4	U				M16	2.5	63Da10	
		150305.3	3.4			−0.3	1	21	21	^{127}I	M16	2.5	63Da10	
		150322	8			−1.6	U				R13	1.5	83De51	
^{127}Cs−u		−92571	30	−92583	6	−0.4	U				GS2	1.0	05Li24	
^{127}Ba−u		−88923	39	−88909	12	0.4	R				GS2	1.0	05Li24	*
^{127}La−u		−83640	30	−83625	28	0.5	1	87	87	^{127}La	GS2	1.0	05Li24	*
^{127}Ce−u		−77273	31				2				GS2	1.0	05Li24	*
^{127}Cd−^{133}Cs$_{.955}$		16494.3	8.0	16497	7	0.3	2				TT1	1.0	17La16	
		16502	12			−0.4	2				IS1	1.0	20Ma09	
^{127}Cd^m−^{133}Cs$_{.955}$		16798.7	8.0	16803	5	0.5	2				TT1	1.0	17La16	
		16805.2	5.8			−0.4	2				IS1	1.0	20Ma09	
^{127}In−^{133}Cs$_{.955}$		7763	12	7747.2	1.3	−1.3	U				TT1	1.0	18Ba08	
		7747.2	1.3			0.0	1	100	100	^{127}In	JY2	1.0	24Ja10	
^{127}In^m−^{133}Cs$_{.955}$		8182	16	8185.3	1.4	0.2	U				TT1	1.0	18Ba08	
^{127}In^n−^{133}Cs$_{.955}$		9585	52	9603.0	1.8	0.3	U				TT1	1.0	18Ba08	
^{127}Sn ^{34}S−^{133}Cs$_{1.211}$		−7237	12	−7244	10	−0.6	1	66	66	^{127}Sn	MA8	1.0	08Dw01	*
^{127}Cs−^{133}Cs$_{.955}$		−2303	17	−2289	6	0.8	o				MA1	1.0	90St25	
		−2287	13			−0.2	–				MA1	1.0	99Am05	
		−2293.3	7.7			0.5	–				MA8	1.0	05Gu37	
	ave.	−2292	7			0.4	1	82	82	^{127}Cs			average	
^{127}Ba−^{133}Cs$_{.955}$		1389	13	1385	12	−0.3	–				MA5	1.0	00Be42	
	ave.	1387	12			−0.2	1	98	98	^{127}Ba			average	
^{127}Cd−^{130}Xe$_{.977}$		20598	89	20475	7	−1.4	U				JY1	1.0	12Ha25	*
^{127}In^m−^{127}In		437.9	5.3	438.1	0.5	0.0	U				JY2	1.0	24Ja10	*
^{127}In^n−^{127}In		1855.8	1.3				2				JY2	1.0	24Ja10	
^{125}Cs−^{127}Cs$_{.591}$ ^{122}Cs$^x_{.410}$		−1098	18	−1088	16	0.2	U				P32	2.5	86Au02	
^{126}Te(n,γ)^{127}Te		6289	3	6287.65	0.18	−0.5	U						72Mu.A	
		6287.8	0.4			−0.4	–				Bdn		06Fi.A	
		6287.6	0.2			0.2	–				Prn		05Ho15	
^{126}Te(d,p)^{127}Te		4044	8	4063.08	0.18	2.4	U				MIT		68Gr16	
^{126}Te(n,γ)^{127}Te	ave.	6287.64	0.18	6287.65	0.18	0.0	1	100	98	^{127}Te			average	
^{127}I(γ,n)^{126}I		−9135	22	−9144.0	2.7	−0.4	U				Phi		60Ge01	
		−9145	3			0.3	1	83	48	^{126}I	MMn		86Ts04	
^{127}Cd^m(β^-)^{127}In		8468	63	8435	5	−0.5	U				Stu		87Sp09	*
^{127}In^m(IT)^{127}In		408.1	0.5				2						19Lo04	
^{127}In(β^-)^{127}Sn		6514	31	6578	9	2.1	o				Stu		87Sp09	*
		6579	20			−0.0	1	21	20	^{127}Sn	Stu		04Ga24	*
^{127}In^n(β^-)^{127}Sn		8442	56	8307	9	−2.4	U				Stu		04Ga24	
^{127}Sn(β^-)^{127}Sb		3201	24	3229	10	1.2	1	17	14	^{127}Sn	Stu		77Lu06	*
^{127}Sb(β^-)^{127}Te		1581	5	1582	5	0.2	1	96	96	^{127}Sb			67Ra13	*
^{127}Te(β^-)^{127}I		683	10	702	4	1.9	–						55Da37	
		695	10			0.7	–						56Kn20	
	ave.	689	7			1.9	1	25	24	^{127}I			average	
^{127}Xe(ε)^{127}I		663.3	2.2	662.3	2.0	−0.4	–						68Sc14	
^{127}I(^{3}He,t)^{127}Xe		−676	6	−680.9	2.0	−0.8	–				Pri		89Ch01	
^{127}Xe(ε)^{127}I	ave.	662.6	2.1	662.3	2.0	−0.1	1	98	91	^{127}Xe			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{127}Cs(β^+)^{127}Xe		2115	25	2081	6	−1.4	–					54Ma54	*
		2076	20			0.3	–					67Sp08	*
		2089	20			−0.4	–					75We23	*
	ave.	2090	12			−0.7	1	27	18 ^{127}Cs			average	
^{127}Ba(β^+)^{127}Cs		3450	100	3422	13	−0.3	U					76Be11	*
^{127}La(β^+)^{127}Ba		5010	70	4922	28	−1.3	1	16	13 ^{127}La	JAE		98Ko66	
*^{127}Sn N$_2$−^{155}Pm	D_M=–11588.7(4.8) μu, $M-A$=–77734.8(8.4) keV for a mixture of ground state and m											24Ki16	**
*^{127}Sn N$_2$−^{155}Pm	at 5.07(0.06) keV; reported $M-A$=–77735(15) keV, author confirmed											24Ki16	**
*^{127}Sn N$_2$−^{155}Pm	mistake in the uncertainty											24Ki16	**
*^{127}Sb N$_2$−^{155}Pm	Reported $M-A$=–80939(19) keV, author confirmed mistake in the unc.											24Ki16	**
*^{127}Ba−u	$M-A$=–82791(28) keV for mixture gs+m at 80.32 keV											Nub269	**
*^{127}La−u	$M-A$=–77903(28) keV for mixture gs+m at 14.2 keV											Nub269	**
*^{127}Ce−u	$M-A$=–71976(29) keV for mixture gs+m at 7.3(1.1) keV											Nub269	**
*^{127}Sn ^{34}S−^{133}Cs$_{1.211}$	D_M=–7234.3(11.6) μu for mixture gs+m at 5.07(0.06) keV											Nub269	**
*^{127}Cd−^{130}Xe$_{.977}$	D_M=20741(14) μu for mixture of gs+m at 283 keV											WgM204	**
*^{127}In^m−^{127}In	Using corrected r=1.000003450(42), supersedes the published value											24Ja10	**
*^{127}Cd^m(β^-)^{127}In	Also E_{β^-}=7910(200) to ^{127}In^m at 408.1(0.5) keV											Nub269	**
*^{127}Cd^m(β^-)^{127}In	Re-assigned to ^{127}Cd^m by the evaluator											GAu	**
*^{127}In(β^-)^{127}Sn	Also E_{β^-}=6976(64) from ^{127}In^m at 408.1(0.5) keV											Nub269	**
*^{127}In(β^-)^{127}Sn	Also E_{β^-}=6999(63) from ^{127}In^m at 408.1(0.5) keV											Nub269	**
*^{127}Sn(β^-)^{127}Sb	Q_{β^-}=3206(24) from ^{127}Sn^m at 5.07 keV											Nub269	**
*^{127}Sb(β^-)^{127}Te	E_{β^-}=1493(5) to 11/2$^-$ level at 88.23 keV, and other E_{β^-}											Ens118	**
*^{127}Cs(β^+)^{127}Xe	E_{β^+}=1063(10), 685(25) to mixture gs+124.751 and 1/2$^+$ level at 411.965 keV											Ens118	**
*^{127}Cs(β^+)^{127}Xe	E_{β^+}=1068(20), 910(30), 650(30) to ground state, 3/2$^+$ at 124.751, 1/2$^+$ at 411.965											Ens118	**
*^{127}Cs(β^+)^{127}Xe	E_{β^+}=1040(20), 650(20) to mixture gs+124.751 and 1/2$^+$ level at 411.965 keV											Ens118	**
*^{127}Ba(β^+)^{127}Cs	E_{β^+}=2230(100) to 3/2$^+$ level at 180.92 keV, and other E_{β^+}											Ens118	**
^{128}In−u		−79639	41	−79646.4	1.4	−0.2	U			TR1	1.0	21Iz01	
^{128}In^n−u		−79361	47	−79340.3	2.1	0.4	U			TR1	1.0	21Iz01	
^{128}In^p−u		−77692	42	−77716.6	1.5	−0.6	U			TR1	1.0	21Iz01	
^{128}Sn−u		−89512	25	−89447	12	2.6	1	24	24 ^{128}Sn	GS3	1.0	12Ch19	
C$_{10}$ H$_8$−^{128}Te		158112	9	158139.0	0.8	1.2	U			M16	2.5	63Da10	
		158141.2	7.			−0.1	U			C3	2.5	70Ke05	
		158151	8			−1.0	U			R13	1.5	83De51	
C$_{10}$ H$_8$−^{128}Xe		159068.2	4.2	159069.504	0.010	0.1	U			M16	2.5	63Da10	
		159069.7	0.7			−0.1	U			C3	2.5	70Ke05	
^{128}Cs−u		−92181	30	−92252	6	−2.4	U			GS2	1.0	05Li24	
^{128}Ba−u		−91663	30	−91647.0	1.7	0.5	U			GS2	1.0	05Li24	
^{128}La−u		−84436	69	−84410	60	0.4	2			GS2	1.0	05Li24	*
^{128}Ce−u		−81089	30				2			GS2	1.0	05Li24	
^{128}Pr−u		−71209	32				2			GS2	1.0	05Li24	
^{128}Cd−^{133}Cs$_{.962}$		18759	11	18772	7	1.2	–			MA8	1.0	10Br02	
		18787	15			−1.0	–			MA8	1.0	20Ma09	
	ave.	18769	9			0.4	1	61	61 ^{128}Cd			average	
^{128}In−^{133}Cs$_{.962}$		11329	11	11308.9	1.4	−1.8	U			TT1	1.0	18Ba08	
^{128}In^n−^{133}Cs$_{.962}$		11610.9	9.8	11615.0	2.1	0.4	U			TT1	1.0	18Ba08	
^{128}Sn ^{34}S−^{133}Cs$_{1.218}$		−6396	14	−6420	12	−1.7	F			MA8	1.0	08Dw01	*
^{128}Sb−^{133}Cs$_{.962}$		124.8	2.7	125.0	2.3	0.1	o			CP2	1.0	22Ra.A	
		125.0	2.3				2			CP2	1.0	23Ho14	
^{128}Sb^m−^{133}Cs$_{.962}$		171.3	2.8	171.3	2.7	0.0	o			CP2	1.0	22Ra.A	
		172.1	2.7			−0.3	1	99	99 ^{128}Sb^m	CP2	1.0	23Ho14	
^{128}Cs−^{133}Cs$_{.962}$		−1306	17	−1296	6	0.6	o			MA1	1.0	90St25	
		−1293	13			−0.3	1	20	20 ^{128}Cs	MA1	1.0	99Am05	
^{128}Ba−^{133}Cs$_{.962}$		−720	13	−691.8	1.7	2.2	U			MA1	1.0	99Am05	
^{128}Cd−^{130}Xe$_{.985}$		22865	11	22860	7	−0.4	1	39	39 ^{128}Cd	JY1	1.0	12Ha25	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{128}Te ^{35}Cl−^{126}Te ^{37}Cl		4106	2	4099.9	1.5	−0.8	U			H16	4.0	63Ba47	
		4102.3	1.8			−0.5	1	12	9 ^{126}Te	C3	2.5	70Ke05	
^{128}Xe−^{129}Xe$_{.992}$		−2011.8587	0.0077	−2011.859	0.008	0.0	1	100	100 ^{128}Xe	Hep	1.0	20Ri04	
^{128}In−^{128}Te		15892.4	1.2				2			JY1	1.0	20Ne06	
^{128}In^{n}−^{128}Te		16198.5	2.0				2			JY1	1.0	20Ne06	
^{128}In^{p}−^{128}Te		17822.2	1.3				2			JY1	1.0	20Ne06	
^{128}Te−^{128}Xe		931.26	1.20	930.5	0.8	−0.4	–			H43	1.5	90Dy04	
		929.6	1.4			0.6	–			CP1	1.0	09Sc19	
	ave.	930.2	1.1			0.2	1	47	47 ^{128}Te			average	
^{128}Xe−^{126}Xe		−774	45	−766.669	0.008	0.1	U			M16	2.5	63Da10	
^{126}Cs−^{128}Cs$_{.656}$ ^{122}Cs$^{x}_{.344}$		−1130	30	−1102	16	0.4	U			P22	2.5	82Au01	
^{124}Cs^{x}−^{128}Cs$_{.323}$ ^{122}Cs$^{x}_{.678}$		−1070	30	−980	30	1.2	U			P22	2.5	82Au01	
^{126}Cs−^{128}Cs$_{.591}$ ^{123}Cs$^{x}_{.410}$		−350	30	−340	12	0.1	U			P22	2.5	82Au01	
^{124}Cs^{x}−^{128}Cs$_{.194}$ ^{123}Cs$^{x}_{.807}$		370	50	356	24	−0.1	U			P22	2.5	82Au01	
^{125}Cs−^{128}Cs$_{.244}$ ^{124}Cs$^{x}_{.756}$		−1440	30	−1349	18	1.2	U			P22	2.5	82Au01	
^{126}Cs−^{128}Cs$_{.492}$ ^{124}Cs$^{x}_{.508}$		−610	30	−563	15	0.6	U			P22	2.5	82Au01	
^{127}Cs−^{128}Cs$_{.661}$ ^{125}Cs$_{.339}$		−965	16	−933	7	0.8	U			P32	2.5	86Au02	
^{127}Cs−^{128}Cs$_{.496}$ ^{126}Cs$_{.504}$		−1160	30	−1105	8	0.7	U			P22	2.5	82Au01	
^{128}Te(γ,n)^{127}Te		−8410	120	−8784.0	1.4	−3.1	B			Phi		60Ge01	
^{127}I(n,γ)^{128}I		6825.7	0.5	6826.13	0.05	0.9	U					71Sc07	
		6826.12	0.05			0.2	–			MMn		90Is03	Z
		6826.22	0.14			−0.6	–			Bdn		06Fi.A	
	ave.	6826.13	0.05			0.0	1	100	87 ^{128}I			average	
^{128}Cd(β^-)^{128}In		7070	290	6952	7	−0.4	U			Stu		87Sp09	
^{128}In(β^-)^{128}Sn		9280	180	9129	11	−0.8	U			Stu		78Al18	*
		8984	37			3.9	F			Stu		87Sp09	*
		8950	103			1.7	F			Gsn		90St13	*
^{128}In^{n}(β^-)^{128}Sn		9390	220	9414	12	0.1	o			Stu		78Al18	*
		9306	30			3.6	F			Stu		87Sp09	*
		9230	90			2.0	U			Gsn		90St13	*
^{128}Sn(β^-)^{128}Sb^{m}		1265	30	1246	11	−0.6	–					76Nu01	*
		1290	40			−1.1	–			Stu		77Lu06	*
		1260	15			−1.0	–			Gsn		90St13	*
	ave.	1264	13			−1.4	1	77	76 ^{128}Sn			average	
^{128}Sb^{m}(IT)^{128}Sb		10	6	43	3	5.5	B					AHW	*
^{128}Sb(β^-)^{128}Te		4640	100	4386.0	2.3	−2.5	U					71Ki15	*
^{128}Sb^{m}(β^-)^{128}Te		4391	40	4429.1	2.6	1.0	U			Stu		77Lu06	*
		4395	30			1.1	U			Gsn		90St13	*
^{128}I(β^+)^{128}Te		1277	13	1255	4	−1.7	U					61La16	
^{128}I(β^-)^{128}Xe		2116	10	2122	4	0.6	1	13	13 ^{128}I			56Be18	*
^{128}Cs(β^+)^{128}Xe		3855	90	3929	5	0.8	U					75We23	*
		3928	6			0.1	1	80	80 ^{128}Cs			76Cr.B	
		3907	40			0.5	o			IRS		83Al06	
		3930	100			−0.0	U			IRS		93Al03	
^{128}La(β^+)^{128}Ba		6650	400	6740	50	0.2	U					66Li04	*
		6820	100			−0.8	R			JAE		98Ko66	*
*^{128}La−u	$M-A$=–78601(28) keV for mixture gs+m at 100#100 keV											Nub269	**
*^{128}Sn ^{34}S−^{133}Cs$_{1.218}$	F : authors say “possible contamination, measurement abandoned”											GAu	**
*^{128}In(β^-)^{128}Sn	E_{β^-}=4980(180) to (2)$^+$ level at 4297.70 keV											Ens15a	**
*^{128}In(β^-)^{128}Sn	E_{β^-}=5464(37) to (2)$^+$ at 3519.86; others 6986(170), 7857(109) to 2104.07,											Ens15a	**
*	1168.82; different equipment/method than in previous; low E_{β^-} not seen											FGK126	**
*^{128}In(β^-)^{128}Sn	E_{β^-}=4650(120), 5440(200) to 4297, 3520 levels and others											FGK126	**
*^{128}In(β^-)^{128}Sn	F : above 2 items conflict with 1st one and with trends in Ex of 8$^-$ isomer											GAu	**
*^{128}In^{n}(β^-)^{128}Sn	E_{β^-}=5430(220) to 3958 level											FGK126	**
*^{128}In^{n}(β^-)^{128}Sn	E_{β^-}=5239(40), 5350(44) to 4066, 3958 level											FGK126	**
*^{128}In^{n}(β^-)^{128}Sn	E_{β^-}=5160(170), 5250(130) to 4066, 3958 levels											FGK126	**
*^{128}Sn(β^-)^{128}Sb^{m}	E_{β^-}=630(30) 655(40) 625(15), to 1$^+$ level 635.2 above ^{128}Sb^{m} at 10(7) keV											Ens15a	**
*^{128}Sb^{m}(IT)^{128}Sb	From 3.6% IT for M3 transition											Ens15a	**
*^{128}Sb(β^-)^{128}Te	E_{β^-}=2300(100) to (7)$^-$ level at 2337.68 keV											Ens15a	**
*^{128}Sb^{m}(β^-)^{128}Te	E_{β^-}=2580(40) 2585(30) respectively, to 6$^+$ level at 1811.13 keV											Ens15a	**
*^{128}I(β^-)^{128}Xe	E_{β^-}=2120(10) and E_{β^-}=1665(15) to ground state and 2$^+$ level at 442.911 keV											Ens15a	**
*^{128}Cs(β^+)^{128}Xe	E_{β^+}=2390(90) to 2$^+$ level at 442.911 keV											Ens15a	**
*^{128}La(β^+)^{128}Ba	E_{β^+}=3200(400) 3370(100) respectively, to (4$^-$,5$^+$) level at 2425.45 keV											Ens15a	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{129}Cd−u		−67789	186	−67764	6	0.1	U			GT3	2.5	16Kn02	
^{129}In−u		−78203	40	−78191.5	2.1	0.3	U			TR1	1.0	21Iz01	
^{129}In^{m}−u		−77724	40	−77707.6	2.1	0.4	U			TR1	1.0	21Iz01	
^{129}In^{p}−u		−76432	86	−76424	4	0.1	U			TR1	1.0	21Iz01	*
^{129}Sn−u		−86521	31	−86520.1	1.9	0.0	U			MA8	1.0	05Si34	*
^{129}Xe−^{120}Sn$_{1.075}$		9913.3	2.4	9913.2	1.0	−0.0	1	18	18 ^{120}Sn	JY1	1.0	11Ha48	
C_{10} H_9−^{129}Xe		165643.6	3.6	165644.432	0.006	0.1	U			M16	2.5	63Da10	
^{129}Xe−u		−95228.7	5.4	−95219.145	0.006	0.7	U			ACC	2.5	90Me08	
^{129}Xe−C_2 $^{35}Cl_3$		−1777.98	0.68	−1777.23	0.11	0.7	U			H47	1.5	94Hy01	
^{129}Xe$_2$−^{86}Kr$_3$		77729.8547	0.0250	77729.840	0.014	−0.6	1	29	16 ^{86}Kr	FS1	1.0	05Sh38	*
^{129}La−u		−87300	30	−87304	23	−0.1	1	58	58 ^{129}La	GS2	1.0	05Li24	
^{129}Ce−u		−81898	30				2			GS2	1.0	05Li24	
^{129}Pr−u		−74905	32				2			GS2	1.0	05Li24	*
^{129}Xe−^{134}Xe$_{.963}$		−4114.7	3.8	−4112.633	0.009	0.5	o			MA8	1.0	05He.A	
		−4119.3	5.1			1.3	U			MA8	1.0	06He29	
^{129}Cd−^{133}Cs$_{.970}$		23831	107	23947	6	1.1	U			MA8	1.0	15At03	*
		23947.2	5.7				2			IS1	1.0	20Ma09	
^{129}Cd^{m}−^{133}Cs$_{.970}$		24315.5	6.0				2			IS1	1.0	20Ma09	
^{129}In−^{133}Cs$_{.970}$		13518.6	6.7	13520.1	2.1	0.2	1	10	10 ^{129}In	TT1	1.0	18Ba08	
^{129}In^{m}−^{133}Cs$_{.970}$		13995	15	14004.0	2.1	0.6	U			TT1	1.0	18Ba08	
^{129}Sn−^{133}Cs$_{.970}$		5191.2	2.7	5191.5	1.9	0.1	1	50	50 ^{129}Sn	CP2	1.0	25Va.A	
^{129}Sn^{m}−^{133}Cs$_{.970}$		5229.6	2.7	5229.3	1.9	−0.1	1	50	50 ^{129}Sn^{m}	CP2	1.0	25Va.A	
^{129}Cs−^{133}Cs$_{.970}$		−2234	18	−2222	5	0.6	o			MA1	1.0	90St25	
		−2216	14			−0.5	1	12	12 ^{129}Cs	MA1	1.0	99Am05	
^{129}In−^{130}Xe$_{.992}$		17523.9	2.9	17527.3	2.1	1.2	1	53	53 ^{129}In	JY1	1.0	12Ha25	
^{129}In^{m}−^{130}Xe$_{.992}$		18016.5	3.5	18011.2	2.1	−1.5	1	37	37 ^{129}In^{m}	JY1	1.0	13Ka08	
^{129}In^{m}−^{129}In		482.0	6.4	483.89	0.17	0.3	U			JY2	1.0	24Ja10	
^{129}In^{p}−^{129}In		1767.6	3.6				2			JY2	1.0	24Ja10	
^{129}Sb−^{129}Te^{m}		2475.4	1.3				3			JY2	1.0	26Wi.1	
C_{10} H_{10}−^{129}Xe		173469.4660	0.0147	173469.464	0.006	−0.2	1	16	16 ^{129}Xe	FS1	1.0	09Re03	
^{129}Xe−^{128}Xe		1247	12	1250.104	0.008	0.1	U			M16	2.5	63Da10	
C_3 O_6−^{129}Xe		64706.8420	0.0255	64706.859	0.006	0.7	o			FS1	1.0	05Sh38	*
		64706.8516	0.0181			0.4	1	10	9 ^{129}Xe	FS1	1.0	09Re03	
^{129}Xe$_2$−^{84}Kr$_3$		75068.5115	0.0405	75068.533	0.014	0.5	1	12	7 ^{84}Kr	FS1	1.0	05Sh38	*
^{128}Cs−^{129}Cs$_{.661}$ ^{126}Cs$_{.339}$		510	30	500	7	−0.1	U			P22	2.5	82Au01	
^{128}Te(n,γ)^{129}Te		6085	3	6082.41	0.08	−0.9	U					72Mu.A	
		6082.42	0.09			−0.1	–			Prn		03Wi02	
		6082.36	0.19			0.3	–			Bdn		06Fi.A	
^{128}Te(d,p)^{129}Te		3857	10	3857.84	0.08	0.1	U			MIT		67Mo22	
^{128}Te(n,γ)^{129}Te	ave.	6082.41	0.08	6082.41	0.08	−0.0	1	100	99 ^{129}Te			average	
^{129}Nd(εp)^{128}Ce		5300	300	5870#	200#	1.9	D					78Bo.A	*
^{129}In^{m}(IT)^{129}In		450.73	0.16	450.74	0.16	0.1	1	100	63 ^{129}In^{m}			FGK20c	*
^{129}In^{q}(IT)^{129}In^{p}		281.0	0.2				3					04Sc42	
^{129}In(β^-)^{129}Sn		7655	32	7758.1	2.7	3.2	B			Stu		87Sp09	
		7780	26			−0.8	U			Stu		04Ga24	*
^{129}In^{m}(β^-)^{129}Sn		8033	66	8208.8	2.7	2.7	B			Stu		87Sp09	
		8149	38			1.6	U			Stu		04Ga24	
^{129}In^{p}(β^-)^{129}Sn		9410	50	9405	4	−0.1	U			Stu		04Ga24	
^{129}Sn^{m}(IT)^{129}Sn		35.15	0.05	35.15	0.05	0.0	1	100	50 ^{129}Sn			Ens145	
^{129}Sn(β^-)^{129}Sb		3996	120	4000.6	2.3	0.0	U			Stu		77Lu06	*
^{129}Sb(β^-)^{129}Te		2345	30	2411.3	1.2	2.2	U					70Oh05	*
		2406	30			0.2	U					130Te-1	
^{129}Te^{m}(IT)^{129}Te		105.51	0.03				2					Ens145	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{129}Te(β^-)^{129}I		1453	28	1502	3	1.8	U					56Gr10	*
		1485	10			1.7	U					64De10	*
		1503	4			−0.2	1	61	60 ^{129}I			68Go34	*
^{129}I(β^-)^{129}Xe		190	5	189	3	−0.2	1	40	40 ^{129}I			54De17	*
^{129}Cs(β^+)^{129}Xe		1197	5	1197	5	0.0	1	83	83 ^{129}Cs			76Ma35	
^{129}Ba(β^+)^{129}Cs		2446	15	2438	11	−0.5	1	50	45 ^{129}Ba			61Ar05	*
^{129}La(β^+)^{129}Ba		3720	50	3737	22	0.3	–					79Br05	*
		3740	40			−0.1	–			JAE		98Ko66	
	ave.	3730	30			0.2	1	48	42 ^{129}La			average	
^{129}Ce(β^+)^{129}La		5600	200	5040	40	−2.8	U			IRS		93Al03	
*^{129}In^{p}−u	mixture at p+q is corrected by the author											21Iz01	**
*^{129}Sn−u	$M-A$=–80576(27) keV for mixture gs+m at 35.15 keV											Nub269	**
*^{129}Xe$_2$−^{86}Kr$_3$	Corrected in reference of same group											09Re03	**
*^{129}Pr−u	Isomer at 382.57 with estimated T=137# ms not considered											Nub269	**
*^{129}Cd−^{133}Cs$_{.970}$	D_M=24016(18) μu for mixture of gs+m at 343(8) keV; $M-A$=–63058(17)											WgM204	**
*C$_3$ O$_6$−^{129}Xe	Corrected in reference of same group											09Re03	**
*^{129}Xe$_2$−^{84}Kr$_3$	Corrected in reference of same group											09Re03	**
*^{129}Nd(εp)^{128}Ce	Trends in Mass Surface TMS suggest ^{129}Nd 570 keV less bound											GAu	**
*^{129}In^{m}(IT)^{129}In	from a least-squares fit to the level scheme in 15Ta13											FGK20c	**
*^{129}In(β^-)^{129}Sn	E_{β^-}=7780(26), 9410(50) from ground state, 1688.0(0.5) levels											03Ge04	**
*^{129}Sn(β^-)^{129}Sb	E_{β^-}=3350(120) to (5/2$^+$) level at 645.14 keV											Ens148	**
*^{129}Sb(β^-)^{129}Te	E_{β^-}=1800(30) to 5/2$^+$ level at 544.585 keV, and other E_{β^-}											Ens148	**
*^{129}Te(β^-)^{129}I	E_{β^-}=1453(5) to 5/2$^+$ level at 27.793 keV and 1530(5) from ^{129}Te^m											Ens148	**
*	at 105.51 to ground state (Birge=8.0: arithmetic average used)											Nub269	**
*^{129}Te(β^-)^{129}I	E_{β^-}=1452(10) to 5/2$^+$ level at 27.793 keV and 1595(10) from ^{129}Te^m at 105.51											Ens148	**
*^{129}Te(β^-)^{129}I	E_{β^-}=1476(4) to 5/2$^+$ level at 27.793 keV and 1607(7) from ^{129}Te^m at 105.51											Ens148	**
*^{129}I(β^-)^{129}Xe	E_{β^-}=150(5) to 3/2$^+$ level at 39.5774 keV											Ens148	**
*^{129}Ba(β^+)^{129}Cs	E_{β^+}=1425(15); and 1243(35), 975(60) from ^{129}Ba^m at 8.42(0.06) keV											Nub269	**
*	to 7/2$^+$ level at 188.91, (9/2)$^+$ at 426.47 keV											Ens148	**
*^{129}La(β^+)^{129}Ba	E_{β^+}=2420(50) to 1/2$^+$ level at 278.57 keV, and other E_{β^+}											Ens148	**
^{130}Cd−u		−66700	441	−65612	24	1.0	U			GT3	2.5	16Kn02	
^{130}In−u		−75033	46	−75047.7	1.9	−0.3	U			TR1	1.0	21Iz01	*
^{130}In^{n}−u		−74646	41	−74634.0	2.2	0.3	U			TR1	1.0	21Iz01	
C$_9$ H$_8$ N−^{130}Te		159446	10	159451.516	0.012	0.2	U			M16	2.5	63Da10	
^{13}C C$_8$ N H$_7$−^{130}Te		154990.6	7.	154981.319	0.012	−0.5	U			C3	2.5	70Ke05	
C$_9$ H$_8$ N−^{130}Te		159449	8	159451.516	0.012	0.2	U			R13	1.5	83De51	
C$_{10}$ H$_{10}$−^{130}Xe		174743.6	4.2	174740.973	0.010	−0.3	U			M16	2.5	63Da10	
^{13}C C$_8$ N H$_7$−^{130}Xe		157695.4	0.7	157694.717	0.010	−0.4	U			C3	2.5	70Ke05	
^{130}Xe−C ^{13}C ^{35}Cl$_3$		−6407.63	1.21	−6403.58	0.11	2.2	U			H47	1.5	94Hy01	
^{130}Cs−u		−93181	60	−93291	9	−1.8	U			GS2	1.0	05Li24	*
C$_{10}$ H$_{10}$−^{130}Ba		171926	68	171924.3	0.3	−0.0	U			R07	1.5	68De17	
^{130}Ba−^{85}Rb$_{1.529}$		41195.8	3.4	41199.6	0.3	1.1	U			MA8	1.0	05Gu37	
^{130}La−u		−87635	30	−87631	28	0.1	2			GS2	1.0	05Li24	
^{130}Ce−u		−85264	30				2			GS2	1.0	05Li24	
^{130}Pr−u		−76410	69				2			GS2	1.0	05Li24	*
^{130}Nd−u		−71494	30				2			GS2	1.0	05Li24	
^{130}Xe−^{134}Xe$_{.970}$		−4726.6	5.6	−4721.891	0.012	0.8	o			MA8	1.0	05He.A	
		−4724.8	7.0			0.4	U			MA8	1.0	06He29	
^{130}Cd−^{133}Cs$_{.977}$		26761	24				2			MA8	1.0	15At03	
^{130}In−^{133}Cs$_{.977}$		17437	93	17325.7	1.9	−1.2	U			CP1	1.0	13Va12	*
		17338	30			−0.4	U			TT1	1.0	18Ba08	*
		17325.7	2.0			−0.0	1	92	92 ^{130}In	CP2	1.0	18Or.A	
^{130}In^{m}−^{133}Cs$_{.977}$		17399.0	2.4	17397.1	2.1	−0.8	1	79	79 ^{130}In^{m}	CP2	1.0	18Or.A	
^{130}In^{n}−^{133}Cs$_{.977}$		17757	31	17739.4	2.2	−0.6	U			TT1	1.0	18Ba08	
		17739.3	3.6			0.0	–			CP2	1.0	18Or.A	
		17737.6	4.8			0.4	–			JY2	1.0	20Ne06	
	ave.	17738.7	2.9			0.3	1	60	60 ^{130}In^{n}			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{130}Sn−^{133}Cs$_{.977}$		6346	17	6350.8	1.3	0.3	U				MA8	1.0	05Si34	
		6344	12			0.6	U				MA8	1.0	05Si34	*
		6349.5	3.9			0.3	–				CP1	1.0	13Va12	
		6352.5	1.9			−0.9	–				CP2	1.0	26Ca.A	
	ave.	6351.9	1.7			−0.7	1	58	58	^{130}Sn			average	
^{130}Sn^m−^{133}Cs$_{.977}$		8443.8	3.8	8440.8	1.3	−0.8	1	12	12	^{130}Sn^m	CP2	1.0	26Ca.A	
^{130}Te−^{133}Cs$_{0.977}$		−1405.4	2.3	−1403.820	0.013	0.7	U				CP2	1.0	26Ca.A	
^{130}Xe−^{133}Cs$_{.977}$		−4114	13	−4117.217	0.011	−0.2	U				MA6	1.0	04Di18	
^{130}Cs−^{133}Cs$_{.977}$		−928	17	−917	9	0.6	o				MA1	1.0	90St25	
		−916	13			−0.1	1	48	48	^{130}Cs	MA1	1.0	99Am05	
^{130}Nd ^{19}F−^{133}Cs$_{1.120}$		32902	130	32800	30	−0.8	U				MA5	1.0	00Be42	*
^{130}Te ^{35}Cl−^{128}Te ^{37}Cl		4715	2	4711.6	0.8	−0.4	o				H16	4.0	63Ba47	
		4711.7	1.8			−0.0	U				C3	2.5	70Ke05	
		4711.57	0.72			0.1	1	49	49	^{128}Te	H43	1.5	90Dy04	
^{130}Xe−^{129}Xe$_{1.008}$		−509.78	0.34	−509.755	0.009	0.1	U				CP1	1.0	09Sc19	
		−509.96	0.26			0.8	U				SH2	1.0	13El01	*
		−509.02	0.94			−0.8	U				SH1	1.0	13El01	*
^{130}In−^{130}In^n		−413.8	6.5	−413.7	2.8	0.0	1	18	11	^{130}In^n	JY2	1.0	20Ne06	
^{130}In^m−^{130}In^n		−351.8	6.0	−342.3	2.9	1.6	1	23	12	^{130}In^n	JY2	1.0	20Ne06	
^{130}In^m−^{130}Te		18797.9	6.9	18800.9	2.1	0.4	1	10	10	^{130}In^m	JY1	1.0	20Ne06	
^{130}In^n−^{130}Te		19138.3	5.3	19143.2	2.2	0.9	1	18	18	^{130}In^n	JY1	1.0	20Ne06	
^{130}Sn−^{130}Xe		10463.9	3.6	10468.0	1.3	1.1	–				JY1	1.0	12Ha25	
		10465.4	3.1			0.8	–				JY1	1.0	13Ka08	*
	ave.	10464.8	2.3			1.4	1	31	31	^{130}Sn			average	
^{130}Te−^{130}Xe		2706.2	7.	2713.398	0.012	0.4	U				C3	2.5	70Ke05	
		2712.98	3.02			0.1	U				H43	1.5	90Dy04	
		2713.416	0.034			−0.5	–				FS1	1.0	09Re07	
		2713.402	0.026			−0.2	–				FS1	1.0	09Re07	*
		2713.402	0.014			−0.3	o				FS1	1.0	09Re07	*
		2712.86	0.34			1.6	U				CP1	1.0	09Sc19	
		2712.82	0.25			2.3	U				JY1	1.0	11Ra24	
		2713.44	0.14			−0.3	U				SH1	1.0	12Ne10	
	ave.	2713.407	0.021			−0.4	1	35	22	^{130}Te			average	
^{130}Ba−^{130}Xe		2816.71	0.31	2816.7	0.3	0.0	1	100	100	^{130}Ba	SH1	1.0	12Ne10	
^{130}Te−^{129}Xe		1441.885	0.012	1441.888	0.011	0.3	1	78	78	^{130}Te	FS1	1.0	09Re07	
^{130}Xe−^{129}Xe		−1277	12	−1271.510	0.009	0.2	U				M16	2.5	63Da10	
		−1271.517	0.012			0.6	1	52	50	^{130}Xe	FS1	1.0	09Re07	
^{129}Cs−^{130}Cs$^x_{794}$ ^{125}Cs$_{.206}$		−1270	40	−1200	14	0.7	U				P22	2.5	82Au01	
^{130}Ba(p,t)^{128}Ba		−9482	32	−9548.9	1.5	−2.1	U				Win		74De31	*
^{130}Ba(p,t)^{128}Ba−^{144}Sm()^{142}Sm		1095.9	1.0	1095.9	1.0	0.0	1	100	100	^{128}Ba			09Pa25	
^{130}Te(d,^{3}He)^{129}Sb		−4550	30	−4554.9	1.4	−0.2	R				Oak		68Au04	
^{129}I(n,γ)^{130}I		6500.33	0.04				2				ILn		89Sa11	Z
^{129}Xe(n,γ)^{130}Xe		9255.3	1.0	9255.722	0.008	0.4	U						71Gr28	Z
		9256.1	0.8			−0.5	U						74Ge05	Z
		9255.57	0.30			0.5	U				Bdn		06Fi.A	
^{129}Xe(^{3}He,d)^{130}Cs		5	20	−1	8	−0.3	1	17	17	^{130}Cs	ChR		81Ha08	
^{130}Ba(d,t)^{129}Ba		−4001	15	−4010	11	−0.6	1	49	49	^{129}Ba	Tal		74Gr22	
^{130}Eu(p)^{129}Sm		1028.0	15.0				3				Arp		04Da04	
^{130}Cd(β^-)^{130}In		8350	160	8789	22	2.7	B				Bwg		03Di06	*
^{130}In(β^-)^{130}Sn		10249	38	10223.1	2.2	−0.7	U				Stu		87Sp09	
		9880	90			3.8	B				Gsn		90St13	
^{130}In^m(β^-)^{130}Sn		10300	37	10289.6	2.3	−0.3	U				Stu		87Sp09	
		10170	170			0.7	U				Gsn		90St13	
^{130}In^n(β^-)^{130}Sn		10650	49	10608.5	2.4	−0.8	U				Stu		87Sp09	
		9880	200			3.6	B				Gsn		90St13	
^{130}Sn^m(IT)^{130}Sn		1946.88	0.10	1946.88	0.10	0.0	1	100	88	^{130}Sn^m			Ens015	
^{130}Sn(β^-)^{130}Sb		2195	35	2154	14	−1.2	–				Stu		77Lu06	*
		2080	40			1.8	–						77Nu01	*
		2149	18			0.3	–				Gsn		90St13	*
	ave.	2148	15			0.4	1	90	90	^{130}Sb			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{130}Sb(β^-)^{130}Te	5046	100	5070	14	0.2	U					71Ki15	*
	5015	100			0.5	U			Stu		77Lu06	*
	4990	70			1.1	U			Gsn		90St13	*
	5015	45			1.2	1	10	10 ^{130}Sb	Stu		95Me16	*
^{130}I(β^-)^{130}Xe	2983	10	2944	3	−3.9	B					65Da01	*
	2964	50			−0.4	U					70Qa03	*
^{130}Cs(β^+)^{130}Xe	2992	20	2981	8	−0.6	–					52Sm41	
	2972	20			0.4	–					75We23	
ave.	2982	14			−0.1	1	35	35 ^{130}Cs			average	
^{130}Cs^x(IT)^{130}Cs	27	15				2					AHW	*
^{130}Cs(β^-)^{130}Ba	442	50	357	8	−1.7	U					52Sm41	*
^{130}La(β^+)^{130}Ba	5660	70	5629	26	−0.4	R			JAE		98Ko66	
*^{130}In−u	mixture at gs+m is corrected by the author										21Iz01	**
*^{130}Cs−u	$M-A$=–86716(30) keV for mixture gs+m at 163.25 keV										Nub269	**
*^{130}Pr−u	$M-A$=–71125(29) keV for mixture gs+m at 100#100 keV										Nub269	**
*^{130}In−^{133}Cs$_{.977}$	D_M=17599(21) μu for mixture gs+m+n at 66.5(2.7) and 385.4(2.6) keV;										Nub269	**
*^{130}In−^{133}Cs$_{.977}$	$M-A$=–69652(20) keV										Nub269	**
*^{130}In−^{133}Cs$_{.977}$	D_M=17373(20) μu for mixture gs+m at 66.5(2.7) keV; $M-A$=–69862(20) keV										Nub269	**
*^{130}Sn−^{133}Cs$_{.977}$	D_M=8434(12) μu for ^{130}Sn^m at 1946.88 keV; $M-A$=–78189(11) keV										Nub269	**
*^{130}Nd ^{19}F−^{133}Cs$_{1.120}$	Tentative result, low statistics										00Be42	**
*^{130}Xe−^{129}Xe$_{1.008}$	Respectively for PI−ICR and Ramsey ToF−ICR techniques										13El01	**
*^{130}Sn−^{130}Xe	D_M=12555.5(3.1) μu for ^{130}Sn^m at 1946.88 keV; $M-A$=–78185.1(2.9) keV										Nub269	**
*^{130}Te−^{130}Xe	First item 1 ion; second item 2 ions - considered independent										GAu	**
*^{130}Te−^{130}Xe	Combination of ^{130}Xe−^{129}Xe and ^{130}Te−^{129}Xe										GAu	**
*^{130}Ba(p,t)^{128}Ba	Not resolved peak. Original uncertainty 16 increased to 24 keV and										GAu	**
*	added systematic error 21 keV										GAu	**
*^{130}Eu(p)^{129}Sm	region of large deformation; deviation from TMS justified										FKo249	**
*^{130}Cd(β^-)^{130}In	E_{β^-}=6224(+165–157) to 1^+ level at 2120.2 keV										Ens086	**
*^{130}Sn(β^-)^{130}Sb	E_{β^-}=1490(90), 1150(35) to 1^+ levels at 702.32, 1047.67 keV										Ens017	**
*^{130}Sn(β^-)^{130}Sb	E_{β^-}=1280(80), 1060(40) to 1^+ levels at 702.32, 1047.67 keV										Ens017	**
*^{130}Sn(β^-)^{130}Sb	E_{β^-}=1415(30), 1112(18) to 1^+ levels at 702.32, 1047.67 keV										Ens017	**
*	and a 3σ conflicting 3955(50) from ^{130}Sn^m at 1946.88 keV										Nub269	**
*^{130}Sb(β^-)^{130}Te	E_{β^-}=2900(100) to ^{130}Te^m at 2146.41 keV										Nub269	**
*^{130}Sb(β^-)^{130}Te	Q =5020(100) from ^{130}Sb^m at 4.80 keV										Nub269	**
*^{130}Sb(β^-)^{130}Te	Also 4960(25) from ^{130}Sb^m at 4.80, in disagreement										Nub269	**
*^{130}Sb(β^-)^{130}Te	Derived from given average=5008(38) with 90St13=4990(70) keV										GAu	**
*^{130}I(β^-)^{130}Xe	E_{β^-}=1702(10) 1042(10) 618(10) to 4^+ 1204.614, 6^+ 1944.140, 5^+ 2362.073										Ens017	**
*^{130}I(β^-)^{130}Xe	E_{β^-}=2480(50), 1850(80) from ^{130}I^m at 39.9525 to 2^+ levels at 536.068,										GAu	**
*	and 1122.112 keV										Ens017	**
*^{130}Cs^x(IT)^{130}Cs	Combining isomer ratio of reference										82Au01	**
*	with ^{130}Cs^m(IT)=163.25 keV										Nub269	**
*^{130}Cs(β^-)^{130}Ba	Value given without associated error										AHW	**
^{131}Cd−u	−59671	1023	−59272	21	0.2	U			GT3	2.5	16Kn02	
	−59280	110			0.1	U			MR1	1.0	15At03	
	−59300	26			1.1	1	63	63 ^{131}Cd	MR1	1.0	20Ma09	
^{131}In−u	−73056	43	−73027.2	2.4	0.7	U			TR1	1.0	21Iz01	
^{131}In^m−u	−72652	42	−72623.5	2.6	0.7	U			TR1	1.0	21Iz01	
^{131}In^n−u	−69007	41	−69010	40	−0.1	1	84	84 ^{131}In^n	TR1	1.0	21Iz01	
^{131}Sn−u	−82958	26	−82961.7	2.0	−0.1	U			MA8	1.0	05Si34	*
	−82950	130			−0.0	U			GT2	2.5	08Su19	*
	−82951	24			−0.4	U			TR1	1.0	25Mo18	
^{131}Sb−u	−88170	530	−88011.7	2.2	0.1	U			GT2	2.5	08Kn.A	*
C_{10} H_{11}−^{131}Xe	180991.6	3.0	180991.224	0.008	−0.1	U			M16	2.5	63Da10	
^{131}Xe−u	−94925.5	5.7	−94915.873	0.008	0.7	U			ACC	2.5	90Me08	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{131}Xe−C_2 $^{35}Cl_2$ ^{37}Cl	1472.65	0.80	1476.17	0.09	2.9	B			H47	1.5	94Hy01	
^{131}Ba−u	−92955	66	−93053.6	0.4	−1.5	U			GS2	1.0	05Li24	*
^{131}La−u	−89930	30				2			GS2	1.0	05Li24	
^{131}Ce−u	−85579	36	−85570	40	0.2	1	96	96 ^{131}Ce	GS2	1.0	05Li24	*
^{131}Pr−u	−79741	56	−79770	50	−0.4	1	81	81 ^{131}Pr	GS2	1.0	05Li24	*
^{131}Nd−u	−72753	30	−72752	30	0.0	1	97	97 ^{131}Nd	GS2	1.0	05Li24	
^{131}Cd−$^{133}Cs_{.985}$	33905	34	33858	21	−1.4	1	37	37 ^{131}Cd	MA8	1.0	20Ma09	
^{131}In−$^{133}Cs_{.985}$	20262	37	20102.7	2.4	−4.3	C			CP1	1.0	13Va12	*
	20104.1	4.1			−0.4	1	33	33 ^{131}In	CP2	1.0	18Or.A	
^{131}In^m−$^{133}Cs_{.985}$	20507.9	2.8	20506.3	2.6	−0.6	1	88	88 ^{131}In^m	CP2	1.0	18Or.A	
^{131}Sn ^{34}S−$^{133}Cs_{1.241}$	2253	11	2239.4	2.0	−1.2	U			MA8	1.0	08Dw01	*
^{131}Sn−$^{133}Cs_{.985}$	10153	20	10168.1	2.0	0.8	U			CP1	1.0	13Va12	*
	10168.3	2.1			−0.1	1	95	95 ^{131}Sn	CP2	1.0	25Va.A	
^{131}Sn^m−$^{133}Cs_{.985}$	10238.8	2.0	10238.0	2.1	−0.4	U			CP2	1.0	25Va.A	
^{131}Sb−$^{133}Cs_{.985}$	5114	11	5118.2	2.2	0.4	U			CP1	1.0	13Va12	
^{131}Xe−$^{129}Xe_{1.016}$	1826.7855	0.0096	1826.781	0.006	−0.5	U			FS1	1.0	13Ho22	
	1826.7797	0.0078			0.2	1	55	48 ^{131}Xe	Hep	1.0	20Ri04	
^{131}Cs−$^{133}Cs_{.985}$	−1429	18	−1401.72	0.19	1.5	o			MA1	1.0	90St25	
	−1419	14			1.2	o			MA1	1.0	99Am05	
	−1401.72	0.19				2			IS1	1.0	19Ka48	*
^{131}Ba−$^{133}Cs_{.985}$	72	14	76.2	0.4	0.3	U			MA5	1.0	00Be42	
^{131}In−$^{130}Xe_{1.008}$	24234.7	2.9	24235.4	2.4	0.2	1	67	67 ^{131}In	JY1	1.0	12Ha25	
^{131}In^m−$^{130}Xe_{1.008}$	24626.8	7.7	24639.0	2.6	1.6	1	12	12 ^{131}In^m	JY1	1.0	13Ka08	
^{131}Sn−$^{132}Xe_{.992}$	12134	21	12116.4	2.0	−0.8	U			JY1	1.0	12Ha25	*
^{131}Sb−$^{130}Xe_{1.008}$	9250.7	2.3	9250.9	2.2	0.1	1	94	94 ^{131}Sb	JY1	1.0	12Ha25	
^{131}Xe−$^{132}Xe_{.992}$	162.283	0.030	162.283	0.006	−0.0	U			SH1	1.0	14Ne15	
	162.292	0.014			−0.7	U			FS1	1.0	13Ho22	
	162.2842	0.0078			−0.2	1	55	52 ^{131}Xe	Hep	1.0	20Ri04	
^{131}Sn^m−^{131}Sn	68.7	2.0	69.89	0.29	0.6	U			JY2	1.0	24Be18	
^{131}I−^{131}Xe	1043.76	0.21				2			JY2	1.0	22Er01	
^{131}Xe−^{130}Xe	1574	11	1574.781	0.010	0.0	U			M16	2.5	63Da10	
^{128}Cs−$^{131}Cs_{.391}$ $^{126}Cs_{.610}$	−100	30	−48	8	0.7	U			P22	2.5	82Au01	
^{128}Cs−$^{131}Cs_{.244}$ $^{127}Cs_{.756}$	783	21	751	7	−0.6	F			P33	2.5	86Au02	*
^{129}Cs−$^{131}Cs_{.328}$ $^{128}Cs_{.672}$	−1030	30	−871	6	2.1	U			P22	2.5	82Au01	
^{130}Te(n,γ)^{131}Te	5929.7	0.5	5929.38	0.06	−0.6	U					77Ko.A	
	5929.5	0.4			−0.3	U					80Ho29	Z
	5929.38	0.06				2			Prn		03To08	
	5930.16	0.19			−4.1	C			Bdn		06Fi.A	
^{130}Te(d,p)^{131}Te	3703	6	3704.81	0.06	0.3	U			MIT		67Gr21	
^{130}Ba(n,γ)^{131}Ba	7493.5	0.3				2					82Ka.A	
^{130}Ba(d,p)^{131}Ba	5269	15	5268.90	0.30	−0.0	U			ANL		70Vo04	
^{131}Nd(εp)^{130}Ce	4600	400	4370	40	−0.6	U					78Bo.A	
^{131}Eu(p)^{130}Sm	957.4	8.	947	5	−1.3	3					98Da03	
	939.2	7.			1.1	3					99So17	*
^{131}In(β^-)^{131}Sn	8820	200	9254.0	2.9	2.2	U					80De35	
	8930	150			2.2	o			Stu		84Fo19	
	9184	33			2.1	o			Stu		88Fo05	
	9165	30			3.0	o			Stu		95Me16	
	9174	22			3.6	B			Stu		99Fo01	
	9222	18			1.8	U			Stu		04Fo06	
^{131}In^m(β^-)^{131}Sn	9230	220	9630	3	1.8	o			Stu		84Fo19	
	9547	46			1.8	o			Stu		88Fo05	
	9480	70			2.1	o			Stu		95Me16	
	9524	26			4.1	B			Stu		04Fo06	
^{131}In^n(β^-)^{131}Sn	13000	500	13000	30	−0.0	o			Stu		84Fo19	
	13450	163			−2.8	B			Stu		88Fo05	
	13230	80			−2.9	B			Stu		95Me16	
	12986	86			0.1	1	17	16 ^{131}In^n	Stu		04Fo06	
^{131}Sn^m(IT)^{131}Sn	65.1	0.3	65.10	0.27	0.0	2					04Fo06	
	65.1	0.6			0.0	2					19Du12	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{131}Sn(β^-)^{131}Sb	4582	120	4704.0	2.7	1.0	o			Bwg		79Ke02	*
	4640	20			3.2	B			Stu		84Fo19	*
	4610	110			0.9	U			Bwg		87Gr.A	*
	4689	14			1.1	o			Stu		95Me16	
	4688	14			1.1	o			Stu		99Fo01	
	4701	8			0.4	1	12	6 ^{131}Sb	Stu		04Fo06	*
^{131}Sb(β^-)^{131}Te	3190	70	3228.7	2.1	0.6	o			Stu		77Lu06	
	3217	20			0.6	o			Stu		95Me16	
	3200	26			1.1	U			Stu		99Fo01	
^{131}Te(β^-)^{131}I	2275	10	2230.30	0.20	−4.5	B					61Be20	*
	2278	15			−3.2	B					65De22	*
^{131}I(β^-)^{131}Xe	971.0	0.7	972.26	0.20	1.8	U					51Ve05	*
	970.4	1.2			1.5	U					52Ro16	*
^{131}Cs(ε)^{131}Xe	355	10	358.00	0.18	0.3	U					54Sa22	
	355	10			0.3	U					56Ho66	
	360	15			−0.1	U					57Mi63	
^{131}Ba(β^+)^{131}Cs	1370	16	1376.7	0.5	0.4	U					76Ge14	*
	1371	12			0.5	U					78Va04	*
^{131}La(β^+)^{131}Ba	2960	100	2910	28	−0.5	U					60Cr01	
^{131}Ce(β^+)^{131}La	4020	400	4060	40	0.1	U					66No05	*
^{131}Pr(β^+)^{131}Ce	5250	150	5410	60	1.1	1	14	9 ^{131}Pr	IRS		93Al03	
^{131}Nd(β^+)^{131}Pr	6560	150	6530	50	−0.2	1	13	9 ^{131}Pr	IRS		93Al03	
*^{131}Sn−u	$M-A$=–77242(15) keV for mixture gs+m at 65.1 keV										Nub269	**
*	next ^{131}Sn ^{34}S−^{133}Cs$_{1.241}$ also from 1.4 GeV p on UC target										GAu	**
*^{131}Sn−u	$M-A$=–77238(120) keV for mixture gs+m at 65.1 keV										Nub269	**
*^{131}Sb−u	$M-A$=–81291(96) keV for mixture gs+m at 1676.06 keV										Nub269	**
*^{131}Ba−u	$M-A$=–86494(30) keV for mixture gs+m at 187.995 keV										Nub269	**
*^{131}Ce−u	$M-A$=–79685(28) keV for mixture gs+m at 63.09 keV										Nub269	**
*^{131}Pr−u	$M-A$=–74202(28) keV for mixture gs+m at 152.4 keV										Nub269	**
*^{131}In−^{133}Cs$_{.985}$	Freq. ratio mistyped, derived from mass excess in col.3 Table 1 of paper										GAu	**
*^{131}Sn ^{34}S−^{133}Cs$_{1.241}$	D_M=2300.3(3.6) μu for mixture ^{131}Sn gs+m at 65.1 keV with R=0.65(0.15)										Nub269	**
*^{131}Sn−^{133}Cs$_{.985}$	D_M=10188.2(4.7) μu for mixture gs+m at 65.1((0.3) keV; $M-A$=–77259.6(4.3) keV										Nub264	**
*^{131}Cs−^{133}Cs$_{.985}$	freq. ratio from three methods, PI−ICR (dominant), ToF−ICR, Ramsey										19Ka48	**
*^{131}Sn−^{132}Xe$_{.992}$	D_M=12168.7(3.4) μu for mixture gs+m at 65.1 keV; $M-A$=–77229.6(3.2) keV										Nub269	**
*	identical to result given in reference										13Ka08	**
*^{128}Cs−^{131}Cs$_{.244}$ ^{127}Cs.	F : rejection based on line-shape analysis										86Au02	**
*^{131}Eu(p)^{130}Sm	Trends in Mass Surface TMS suggest ^{131}Eu 240 keV less bound										GAu	**
*^{131}Sn(β^-)^{131}Sb	E_{β^-}=3870(120), 2620(180) from ^{131}Sn^m at 65.1 keV										Nub269	**
*	to (5/2$^+$) level at 798.494, (13/2$^-$) at 1980.39 keV										Ens06c	**
*^{131}Sn(β^-)^{131}Sb	Q_{β^-}=4638(20); and 4796(80) from ^{131}Sn^m at 65.1 keV										Nub269	**
*^{131}Sn(β^-)^{131}Sb	Q_{β^-}=4600(110); and 4680(120) from ^{131}Sn^m at 65.1 keV										Nub269	**
*^{131}Sn(β^-)^{131}Sb	Q_{β^-}=4698(11); and 4767(8)from ^{131}Sn^m at 65.1 keV										Nub269	**
*^{131}Te(β^-)^{131}I	Q_{β^-}=2457(10) 2460(15) respectively, from ^{131}Te^m at 182.258 keV										Nub269	**
*^{131}I(β^-)^{131}Xe	E_{β^-}=606.5(0.7) 605.9(1.2) respectively, to 5/2$^+$ level at 364.490 keV										Ens06c	**
*^{131}Ba(β^+)^{131}Cs	p$^+$ =22(11)×10^{-6} to 3/2$^+$ level at 216.086, recalculated p$^+$ and Q										Ens06c	**
*^{131}Ba(β^+)^{131}Cs	L/K=0.165(0.001) to 3/2$^+$ level at 1047.682 keV										Ens06c	**
*^{131}Ce(β^+)^{131}La	E_{β^+}=2800(400) to 7/2$^+$ level at 195.68 keV										Ens06c	**
^{132}Cd−u	−54213	77	−54180	60	0.5	2			MR1	1.0	20Ma09	
	−54092	118			−0.7	2			MR1	1.0	20Ma09	
^{132}In−u	−66984	41	−66990	30	−0.1	1	71	71 ^{132}In	TR1	1.0	21Iz01	
^{132}Sn−u	−82168.0	11.8	−82176.1	2.1	−0.7	U			TR1	1.0	25Mo18	
^{132}Sb−u	−85820	140	−85490.3	0.9	0.9	U			GT2	2.5	08Su19	*
C_{10} H_{12}−^{132}Xe	189740.8	3.3	189745.299	0.006	0.5	U			M16	2.5	63Da10	
^{132}Xe−u	−95856.2	4.0	−95844.917	0.006	1.1	U			ACC	2.5	90Me08	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{132}Xe−C ^{13}C ^{35}Cl$_2$ ^{37}Cl		−2803.73	1.40	−2807.71	0.09	−1.9	U			H47	1.5	94Hy01	
^{132}Xe−C$_3$ O$_6$		−65332.6117	0.0248	−65332.631	0.006	−0.8	o			FS1	1.0	05Sh38	*
		−65332.6238	0.0140			−0.5	1	16	14 ^{132}Xe	FS1	1.0	09Re03	
C$_{10}$ H$_{12}$−^{132}Ba		188863	70	188839.4	1.1	−0.2	U			R07	1.5	68De17	
		188821	88			0.1	U			R07	1.5	68De17	
^{132}La−u		−89874	67	−89880	40	−0.1	1	34	34 ^{132}La	GS2	1.0	05Li24	*
^{132}Ce−u		−88542	30	−88534	22	0.3	1	53	53 ^{132}Ce	GS2	1.0	05Li24	
^{132}Ce O−^{142}Sm$_{1.042}$		−5258	32	−5267	22	−0.3	1	47	47 ^{132}Ce	MA7	1.0	01Bo59	*
^{132}Pr−u		−80760	31				2			GS2	1.0	05Li24	*
^{132}Nd−u		−76690	30	−76679	26	0.4	2			GS2	1.0	05Li24	
^{132}Xe−^{129}Xe$_{1.023}$		1564.20	0.32	1564.269	0.004	0.2	U			CP1	1.0	09Sc19	
		1565.4	1.0			−1.1	U			CP1	1.0	09Sc19	
^{132}Sb−^{130}Xe$_{1.015}$		12445.7	2.9	12447.7	0.9	0.7	1	9	9 ^{132}Sb	JY1	1.0	12Ha25	
^{132}Te−^{130}Xe$_{1.015}$		6482.9	4.3	6485	4	0.4	1	76	76 ^{132}Te	JY1	1.0	12Ha25	
^{132}Sn−^{133}Cs$_{.992}$		11621	19	11615.6	2.1	−0.3	U			MA8	1.0	05Si34	
		11613.1	2.9			0.9	–			CP1	1.0	13Va12	
^{132}Sn ^{34}S−^{133}Cs$_{1.248}$		3686.3	7.7	3686.9	2.1	0.1	–			MA8	1.0	08Dw01	
^{132}Sn−^{133}Cs$_{.992}$	ave.	11613.3	2.7	11615.6	2.1	0.8	1	61	61 ^{132}Sn			average	
^{132}Sb−^{133}Cs$_{.992}$		8301.3	6.5	8301.3	0.9	0.0	–			CP1	1.0	13Va12	
		8298.1	1.6			2.0	o			CP2	1.0	22Ra.A	
		8305.7	2.8			−1.6	–			JY2	1.0	24Ja10	
		8299.9	1.1			1.3	–			CP2	1.0	25Va.A	
	ave.	8300.7	1.0			0.6	1	77	77 ^{132}Sb			average	
^{132}Sb^m−^{133}Cs$_{.992}$		8453.9	1.6	8454.7	1.1	0.5	o			CP2	1.0	22Ra.A	
		8455.9	1.2			−1.0	1	79	79 ^{132}Sb^m	CP2	1.0	25Va.A	
^{132}Cs−^{133}Cs$_{.992}$		223	18	229.4	1.1	0.4	o			MA1	1.0	90St25	
		232	14			−0.2	U			MA1	1.0	99Am05	
		246.9	5.9			−3.0	C			MA8	1.0	09Bo.A	
		230.8	1.3			−1.1	1	73	73 ^{132}Cs	MA8	1.0	17At01	
^{132}Nd−^{133}Cs$_{.992}$		17147	52	17113	26	−0.7	R			MA5	1.0	00Be42	
^{132}Sn−^{132}Xe		13672.3	3.4	13668.8	2.1	−1.0	1	39	39 ^{132}Sn	JY1	1.0	12Ha25	
^{132}Sb^m−^{132}Sb		149.6	2.1	153.4	1.2	1.8	1	35	21 ^{132}Sb^m	JY2	1.0	24Ja10	
^{132}Xe−^{131}Xe		−930	11	−929.044	0.006	0.0	U			M16	2.5	63Da10	
^{132}Xe−C$_{10}$ H$_{10}$		−174095.2367	0.0095	−174095.236	0.006	0.1	1	35	35 ^{132}Xe	FS1	1.0	09Re03	
^{132}Xe−^{130}Xe		645.724	0.014	645.738	0.009	1.0	1	40	38 ^{130}Xe	FS1	1.0	09Re07	
^{132}Ba−^{130}Ba		−1241	4	−1265.1	1.2	−2.4	U			M17	2.5	66Be10	
		−1253	68			−0.1	U			R07	1.5	68De17	
^{132}Xe−^{129}Xe		−625.7755	0.0156	−625.772	0.004	0.2	o			FS1	1.0	05Sh38	*
		−625.7703	0.0119			−0.1	–			FS1	1.0	09Re03	*
		−625.7732	0.0125			0.1	–			FS1	1.0	09Re03	*
		−625.771	0.013			−0.1	–			FS1	1.0	09Re07	
		−625.7771	0.0083			0.6	–			FS1	1.0	10Mo30	
	ave.	−625.774	0.005			0.4	1	44	26 ^{129}Xe			average	
^{132}Xe$_2$−^{84}Kr$_3$		73816.9775	0.0594	73816.989	0.014	0.2	U			FS1	1.0	05Sh38	*
^{132}Xe$_2$−^{86}Kr$_3$		76478.3099	0.0412	76478.296	0.014	−0.3	1	11	6 ^{86}Kr	FS1	1.0	05Sh38	*
^{14}N$_{10}$−^{132}Xe		126584.9632	0.0168	126584.958	0.006	−0.3	1	12	10 ^{132}Xe	FS1	1.0	09Re03	
^{131}Cs−^{132}Cs$_{.794}$ ^{127}Cs$_{.206}$		−1118	16	−1090.9	1.4	0.7	F			P33	2.5	86Au02	*
^{131}Cs−^{132}Cs$_{.744}$ ^{128}Cs$_{.256}$		−1200	30	−1215.4	1.6	−0.2	U			P22	2.5	82Au01	
^{130}Cs^x−^{132}Cs$_{.492}$ ^{128}Cs$_{.508}$		−210	40	−340	17	−1.3	U			P22	2.5	82Au01	
^{131}Xe(n,γ)^{132}Xe		8936.3	1.0	8936.717	0.005	0.4	U					71Ge05	
		8935	2			0.9	U					71Gr28	
		8936.65	0.22			0.3	U			Bdn		06Fi.A	
^{132}In(β^-)^{132}Sn		13600	400	14150	30	1.4	U					86Bj01	
		14135	60			0.2	1	29	29 ^{132}In	Stu		95Me16	
^{132}Sn(β^-)^{132}Sb		3080	40	3087.2	2.1	0.2	o			Stu		77Al09	
		3103	10			−1.6	o			Stu		95Me16	
		3115	10			−2.8	U			Stu		99Fo01	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{132}Sb(β^-)^{132}Te		5530	70	5554	4	0.3	o			Stu		77Al09	
		5486	24			2.9	U			Stu		95Me16	
		5491	20			3.2	B			Stu		99Fo01	
^{132}Te(β^-)^{132}I		493	4	515	3	5.6	B					65Iv01	*
		517	4			−0.4	1	76	52 ^{132}I	Stu		99Fo01	*
^{132}I(β^-)^{132}Xe		3596	15	3575	4	−1.4	–					61De17	*
		3558	15			1.2	–					65Jo13	*
		3580	7			−0.6	–			Stu		99Fo01	
	ave.	3579	6			−0.6	1	48	48 ^{132}I			average	
^{132}I^m(β^-)^{132}Xe		3685	10				2					74Di03	*
^{132}Cs(β^+)^{132}Xe		2090	25	2126.3	1.0	1.5	U					63Ta05	*
		2127.7	6.			−0.2	U					87De33	*
^{132}La(β^+)^{132}Ba		4820	100	4710	40	−1.1	–					60Wa03	
		4680	50			0.6	–					67Fr02	
	ave.	4710	40			0.1	1	66	66 ^{132}La			average	
*^{132}Sb−u	$M-A$=–79870(124) keV for mixture gs+m at 142.8(1.2) keV											Nub269	**
*^{132}Xe−C$_3$ O$_6$	Corrected in reference of same group											09Re03	**
*^{132}La−u	$M-A$=–83623(30) keV for mixture gs+m at 188.20 keV											Nub269	**
*^{132}Ce O−^{142}Sm$_{1.042}$	Original error (22 keV) increased by 23 for BaF contamination in trap											GAu	**
*^{132}Pr−u	$M-A$=–75213(28) keV for mixture gs+m at 30#30 keV											Nub269	**
*^{132}Xe−^{129}Xe	Corrected in reference of same group											09Re03	**
*^{132}Xe−^{129}Xe	First item 5$^+$ ions; second item 3$^+$ ions - considered to be independent											09Re03	**
*^{132}Xe$_2$−^{84}Kr$_3$	Corrected in reference of same group											09Re03	**
*^{132}Xe$_2$−^{86}Kr$_3$	Corrected in reference of same group											09Re03	**
*^{131}Cs−^{132}Cs$_{.794}$ ^{127}Cs.	F : Rejection based on line-shape analysis											86Au02	**
*^{132}Te(β^-)^{132}I	E_{β^-}=215(4) 239(4) respectively, to 1$^+$ level at 277.86 keV											Ens054	**
*^{132}I(β^-)^{132}Xe	E_{β^-}=2156(15) 2118(15) respectively, to 4$^+$ level at 1440.323 keV											Ens054	**
*^{132}I^m(β^-)^{132}Xe	E_{β^-}=1465(10) to 7$^-$ level at 2214.01 level, and other E_{β^-}											Ens054	**
*^{132}Cs(β^+)^{132}Xe	E_{β^+}=400(25) to 2$^+$ level at 667.715 keV											Ens054	**
*^{132}Cs(β^+)^{132}Xe	p$^+$ =0.0042(0.0001) gives E_{β^+}=438(6) recalculated Q											AHW	**
*	to 2$^+$ level at 667.715 keV											Ens054	**
^{133}In−u		−61920	44				2			TR1	1.0	21Iz01	
^{133}In^m−u		−61231	74				2			TR1	1.0	21Iz01	
^{133}Sn−u		−76082	26	−76086.2	2.0	−0.2	U			TR1	1.0	25Mo18	
^{133}Sb−u		−84766	100	−84725.1	1.1	0.2	o			GT2	2.5	08Kn.A	
		−84795	129			0.2	U			GT2	2.5	08Su19	
		−84702	25			−0.9	U			GS3	1.0	12Ch19	
^{133}Te−u		−89031	43	−89032.3	0.6	−0.0	U			GR1	1.0	19An10	
^{133}I−u		−92166	16	−92183.9	0.9	−1.1	U			GR1	1.0	19An10	
C$_{10}$ H$_{13}$−^{133}Cs		196266	64	196273.457	0.009	0.1	U			R07	1.5	68De17	
		196279	25			−0.1	U			R07	1.5	68De17	
C$_9$ ^{13}C H$_{12}$−^{133}Cs		191796	34	191803.260	0.009	0.1	U			R07	1.5	68De17	
C$_8$ O N H$_7$−^{133}Cs		147321	25	147311.889	0.009	−0.2	U			R07	1.5	68De17	
C$_7$ ^{13}C N O H$_6$−^{133}Cs		142835	31	142841.692	0.009	0.1	U			R07	1.5	68De17	
^{133}Cs−^{85}Rb$_{1.565}$		43500	13	43501.028	0.011	0.1	U			MA5	1.0	00Be42	
		43499.3	1.6			1.1	U			MA8	1.0	07Ke09	
		43500.9	6.7			0.0	U			MA8	1.0	02Ke.A	
		43500.1	6.7			0.1	U			MA8	1.0	09Na.A	
		43470	47			0.7	U			MA9	1.0	09Na.A	
		43501.2	1.7			−0.1	U			MA8	1.0	11He10	
^{133}Cs−u		−94548.41	0.41	−94548.042	0.009	0.9	U			ST2	1.0	99Ca46	*
^{133}La−u		−91810	120	−91780	30	0.2	U			GS1	1.0	00Ra23	
		−91782	30				2			GS2	1.0	05Li24	
^{133}Ce−u		−88471	32	−88480	18	−0.3	2			GS2	1.0	05Li24	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{133}Ce O−^{142}Sm$_{1.049}$	−4618	21	−4618	18	−0.0	R			MA7	1.0	01Bo59	*
^{133}Pr−u	−83663	30	−83669	13	−0.2	R			GS2	1.0	05Li24	
^{133}Nd−u	−77652	50				2			GS2	1.0	05Li24	*
^{133}Pm−u	−70218	54				2			GS2	1.0	05Li24	*
^{133}Sb−^{136}Xe$_{.978}$	6022	10	6019.2	1.1	−0.3	U			CP1	1.0	12Va02	
^{133}Sb−^{130}Xe$_{1.023}$	13984.7	4.0	13984.9	1.1	0.0	1	8	8 ^{133}Sb	JY1	1.0	12Ha25	
^{133}Te−^{130}Xe$_{1.023}$	9672.1	2.3	9677.6	0.6	2.4	U			JY1	1.0	12Ha25	
^{133}Sn−^{134}Xe$_{.993}$	17856.5	2.4	17858.5	2.0	0.8	1	73	73 ^{133}Sn	JY1	1.0	12Ha25	
^{133}Sn ^{34}S−^{133}Cs$_{1.256}$	10562	25	10533.1	2.0	−1.2	U			MA8	1.0	08Dw01	
^{133}Sn−^{133}Cs	18467.0	3.9	18461.8	2.0	−1.3	1	27	27 ^{133}Sn	CP1	1.0	13Va12	
^{133}Sb−^{133}Cs	9818	13	9823.0	1.1	0.4	U			CP1	1.0	13Va12	
	9823.0	1.2			−0.0	1	92	92 ^{133}Sb	CP2	1.0	24Va14	
^{133}Te−^{133}Cs	5551.4	6.9	5515.7	0.6	−5.2	B			CP1	1.0	13Va12	
	5514.75	0.80			1.2	1	50	50 ^{133}Te	CP2	1.0	24Va14	
^{133}Te^m−^{133}Cs	5875.59	0.80	5874.6	0.6	−1.2	1	50	50 ^{133}Te^m	CP2	1.0	24Va14	
^{133}I−^{133}Cs	2375.4	6.9	2364.1	0.9	−1.6	U			CP1	1.0	13Va12	
	2364.11	0.93				2			CP2	1.0	24Va14	
^{133}I^m−^{133}Cs	4120.4	6.1	4118.4	0.9	−0.3	U			CP2	1.0	24Va14	
^{133}Pr−^{133}Cs	10877	15	10879	13	0.1	2			MA5	1.0	00Be42	
^{133}Cs−C_3 O_6	−64035.786	0.026	−64035.756	0.009	1.1	1	11	11 ^{133}Cs	MI2	1.0	99Br47	
^{133}Cs−C_{10} H_{12}	−188448.445	0.057	−188448.425	0.009	0.4	U			MI2	1.0	99Br47	
^{133}Cs−^{132}Xe	1296.8803	0.0103	1296.874	0.007	−0.6	1	49	45 ^{133}Cs	FS1	1.0	10Mo30	
^{133}Cs−^{129}Xe	671.1007	0.0103	671.103	0.007	0.2	1	50	44 ^{133}Cs	FS1	1.0	10Mo30	
^{133}Cs(γ,n)^{132}Cs	−8988	33	−8989.6	1.0	−0.0	U			Phi		60Ge01	
	−8986	2			−1.8	1	27	27 ^{132}Cs	MMn		85Ts02	
^{132}Ba(n,γ)^{133}Ba	7189.91	0.36	7189.9	0.4	−0.0	1	100	99 ^{132}Ba	MMn		90Is07	Z
^{132}Ba(d,p)^{133}Ba	4977	15	4965.3	0.4	−0.8	U			ANL		70Vo04	
^{133}Sn(β^-)^{133}Sb	7830	70	8047.0	2.2	3.1	B			Stu		83Bl16	*
	8013	50			0.7	o			Stu		92Sp.A	*
	7990	25			2.3	U			Stu		95Me16	
^{133}Sb(β^-)^{133}Te	3966	50	4012.2	1.2	0.9	o			Stu		70Ru.A	*
	4003	10			0.9	o			Stu		95Me16	
	4002	7			1.5	U			Stu		99Fo01	
^{133}Te^m(IT)^{133}Te	334.274	0.035	334.28	0.03	0.1	1	100	50 ^{133}Te			79Bo26	
^{133}Te(β^-)^{133}I	2960	100	2935.7	1.0	−0.2	U					68Mc09	
	2876	100			0.6	U					68Pa03	*
	3392	100			−4.6	C			Stu		70Ru.A	*
	2890	15			3.0	B			Stu		95Me16	
	2942	24			−0.3	U			Stu		99Fo01	
^{133}I^m(IT)^{133}I	1634.148	0.010				3					Ens10a	
^{133}I(β^-)^{133}Xe	1800	50	1774.8	2.6	−0.5	U					59Ho97	*
	1760	30			0.5	U					66Ei01	*
	1757	4			4.4	B			Stu		99Fo01	
^{133}Xe(β^-)^{133}Cs	428.0	4.	427.4	2.4	−0.2	2					52Be55	*
	427.0	3.			0.1	2					61Er04	*
	424	11			0.3	U			Stu		99Fo01	
^{133}Ba(ε)^{133}Cs	517.3	1.0	517.2	1.0	−0.1	1	99	99 ^{133}Ba			67Sc10	*
	498	5			3.8	F					68Mc06	*
	486	2			15.6	F					69Bo49	*
	521	5			−0.8	U					69To14	*
^{133}La(β^+)^{133}Ba	2230	200	2059	28	−0.9	U					50Na09	*
^{133}Nd(β^+)^{133}Pr	4800	200	5610	50	4.0	B					95Br24	
*^{133}Cs−u	As revised in reference. Original: –94548.20(0.28) μu										02Bf02	**
*^{133}Ce−u	$M-A$=–82392(28) keV for mixture gs+m at 37.2 keV										Nub269	**
*^{133}Ce O−^{142}Sm$_{1.049}$	D_M=–4599(16) μu for mixture gs+m at 37.2 keV; $M-A$=–87150(16) keV										Nub269	**
*^{133}Nd−u	$M-A$=–72268(28) keV for mixture gs+m at 127.97 keV										Nub269	**
*^{133}Pm−u	$M-A$=–65342(33) keV for mixture gs+m at 129.7 keV										Nub269	**
*^{133}Sn(β^-)^{133}Sb	E_{β^-}=6870(70) to 5/2$^+$ level at 962.30 keV										Ens114	**
*^{133}Sn(β^-)^{133}Sb	Private commmunication to reference										92Ch09	**
*^{133}Sb(β^-)^{133}Te	E_{β^-}=1210(50) to (5/2$^+$) level at 2755.51 keV; re-evaluated										Ens114	**
*^{133}Te(β^-)^{133}I	Q_{β^-}=3210(100) from ^{133}Te^m at 334.28 keV										Nub269	**
*	reported as belonging to ground state, reinterpreted										AHW	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value	Adjusted value	v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{133}$Te(β^-)^{133}I	E_{β^-}=850(100) to (3/2$^+$,5/2$^+$) level at 2541.74 keV								Ens114	**
$*^{133}$I(β^-)^{133}Xe	E_{β^-}=1270(50) 1230(30) respectively, to 5/2$^+$ level at 529.872 keV								Ens114	**
$*^{133}$Xe(β^-)^{133}Cs	E_{β^-}=347(4) 346(3) respectively, to 5/2$^+$ level at 80.9979 keV								Ens114	**
$*^{133}$Ba(ε)^{133}Cs	From L/K=0.371(0.007) to 1/2$^+$ level at 437.0113; recalculated Q								Ens114	**
*	and L/K=0.221(0.005) to 3/2$^+$ level at 383.8491 keV: Q =521(5) keV								Ens114	**
$*^{133}$Ba(ε)^{133}Cs	F : badly resolved L−peak								Ens114	**
$*^{133}$Ba(ε)^{133}Cs	L/K=0.67(0.15) LM/K=1.11(0.05) 0.45(0.04) respectively, to 1/2$^+$ 437.0113 keV								Ens114	**
$*^{133}$La(β^+)^{133}Ba	E_{β^+}=1200(200) to 3/2$^+$ level at 12.327 keV								Ens114	**
^{134}In−u	−55669 47				2			TR1	1.0	21Iz01	
^{134}Sn−u	−71332 28	−71320	3	0.4	U			TR1	1.0	25Mo18	
^{134}Sb−u	−79351 131	−79463	3	−0.9	U			GR1	1.0	19An10	
^{134}Te−u	−88844 130	−88603.6	2.9	0.7	U			GT2	2.5	08Su19	
	−88614 44			0.2	U			GR1	1.0	19An10	
^{134}I−u	−90244 58	−90224	5	0.3	U			GR1	1.0	19An10	
C_{10} H_{14}−^{134}Xe	204155.5 3.2	204157.417	0.010	0.2	U			M16	2.5	63Da10	
^{134}Xe−u	−94634.4 5.4	−94606.970	0.010	2.0	U			ACC	2.5	90Me08	
	−94606.86 0.30			−0.4	U			MS1	1.0	19Sa.A	
^{134}Xe−C ^{13}C ^{35}Cl ^{37}Cl$_2$	1381.76 0.60	1380.36	0.12	−1.6	U			H47	1.5	94Hy01	
C_{10} H_{14}−^{134}Ba	205025 20	205042.14	0.22	0.3	U			M17	2.5	66Be10	
	205010 46			0.5	U			R07	1.5	68De17	
C_{11} H_2−^{134}Ba	111125 48	111141.76	0.22	0.2	U			R07	1.5	68De17	
C_8 N O H_8−^{134}Ba	156063 78	156080.57	0.22	0.2	U			R07	1.5	68De17	
C_{12} H_6−^{134}Ba O	147531 64	147527.27	0.22	−0.0	U			R07	1.5	68De17	
^{134}La−u	−91456 34	−91486	21	−0.9	2			GS2	1.0	05Li24	
^{134}Ce−u	−91190 130	−91070	22	0.9	U			GS1	1.0	00Ra23	
	−91056 30			−0.5	1	53	53 ^{134}Ce	GS2	1.0	05Li24	
^{134}Ce O−^{142}Sm$_{1.056}$	−6631 32	−6615	22	0.5	1	47	47 ^{134}Ce	MA7	1.0	01Bo59	*
^{134}Pr−u	−84285 37	−84303	22	−0.5	2			GS2	1.0	05Li24	*
^{134}Nd−u	−81234 30	−81210	13	0.8	R			GS2	1.0	05Li24	
^{134}Pm−u	−71674 45				2			GS2	1.0	05Li24	*
^{134}Sb−^{130}Xe$_{1.031}$	20016.8 2.2	20019	3	1.1	o			JY1	1.0	12Ha25	
	20019.2 3.3				2			JY1	1.0	13Ka08	*
^{134}Te−^{130}Xe$_{1.031}$	10877.1 3.5	10878.2	2.9	0.3	1	71	71 ^{134}Te	JY1	1.0	12Ha25	
^{134}Sb−^{136}Xe$_{.985}$	11906 36	11931	3	0.7	U			CP1	1.0	12Va02	*
^{134}Te−^{136}Xe$_{.985}$	2791.4 6.5	2790.1	2.9	−0.2	1	21	21 ^{134}Te	CP1	1.0	12Va02	
^{134}Xe−^{132}Xe$_{1.015}$	2675.6193 0.0084	2675.619	0.008	−0.0	U			FS1	1.0	13Ho22	
	2675.6191 0.0081			0.0	1	100	100 ^{134}Xe	Hep	1.0	20Ri04	
^{134}Sn ^{34}S−^{133}Cs$_{1.263}$	16080 160	15962	3	−0.7	U			MA8	1.0	08Dw01	
^{134}Sn−^{133}Cs$_{1.008}$	23974 17	23985	3	0.6	U			CP1	1.0	13Va12	
^{134}Sb−^{133}Cs$_{1.008}$	15850 11	15842	3	−0.7	U			CP1	1.0	13Va12	
^{134}I−^{133}Cs$_{1.008}$	5083.0 6.8	5080	5	−0.4	1	59	59 ^{134}I	CP1	1.0	13Va12	
^{134}Cs−^{133}Cs$_{1.008}$	2025 18	2022.929	0.015	−0.1	o			MA1	1.0	90St25	
	2033 14			−0.7	U			MA1	1.0	99Am05	
^{134}Pr−^{133}Cs$_{1.008}$	10992 27	11001	22	0.3	R			MA5	1.0	00Be42	*
^{134}Nd−^{133}Cs$_{1.008}$	14100 14	14095	13	−0.4	2			MA5	1.0	00Be42	
^{134}Sn−^{134}Xe	23287.4 3.4				2			JY1	1.0	12Ha25	
^{134}Ba−C_{10} H_{13}	−197229 20	−197217.11	0.22	0.2	U			M17	2.5	66Be10	
^{134}Ba−^{132}Ba	−553 4	−552.7	1.2	0.0	U			M17	2.5	66Be10	
	−550 121			−0.0	U			R07	1.5	68De17	
^{131}Cs−^{134}Cs$_{.244}$ ^{130}Cs$^x_{.756}$	−1313 50	−1178	13	1.1	U			P22	2.5	82Au01	*
^{133}Cs(n,γ)^{134}Cs	6891.540 0.017	6891.540	0.014	0.0	–			MMn		84Ke11	Z
	6891.540 0.027			0.0	–			ILn		87Bo24	Z
	6891.39 0.14			1.1	U			Bdn		06Fi.A	
	ave. 6891.540 0.014			0.0	1	100	100 ^{134}Cs			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{134}Sn(β^-)^{134}Sb	7370	90	7585	4	2.4	U			Stu		95Me16	
^{134}Sb(β^-)^{134}Te	8306	210	8515	4	1.0	o			Stu		72Ke21	*
	7960	240			2.3	o			Stu		77Lu06	*
	8310	80			2.6	U			Bwg		79Ke02	*
	8390	45			2.8	U			Stu		95Me16	
^{134}Te(β^-)^{134}I	1560	90	1510	5	−0.6	U			Stu		77Lu06	*
	1550	30			−1.3	U			Stu		95Me16	
	1513	7			−0.5	1	50	41 ^{134}I	Stu		99Fo01	
^{134}I(β^-)^{134}Xe	4170	60	4082	5	−1.5	U					61Jo08	*
	4175	15			−6.2	B			Stu		95Me16	
	4052	8			3.8	B			Stu		99Fo01	
^{134}Cs(β^-)^{134}Ba	2058.6	0.4	2058.78	0.21	0.5	1	26	26 ^{134}Ba			68Hs01	*
^{134}La(β^+)^{134}Ba	3772	50	3731	20	−0.8	R					65Bi12	
	3692	30			1.3	R					73Al20	
^{134}Ce(ε)^{134}La	500	200	387	29	−0.6	U					76Gr09	*
^{134}Pr(β^+)^{134}Ce	6190	90	6303	29	1.3	U			Dbn		95Ve08	*
^{134}Nd(β^+)^{134}Pr	2770	150	2882	24	0.7	U					77Ko.B	
^{134}Pm(β^+)^{134}Nd	9170	200	8880	40	−1.4	U			Dbn		95Ve08	*
*^{134}Ce O−^{142}Sm$_{1.056}$	Original error (22 keV) increased by 23 for BaF contamination in trap										GAu	**
*^{134}Pr−u	$M-A$=–78477(28) keV for mixture gs+m at 67.7(0.4) keV										Nub269	**
*^{134}Pm−u	$M-A$=–66739(30) keV for mixture gs+m at 50#(50#) keV										Nub269	**
*^{134}Sb−^{130}Xe$_{1.031}$	D_M=20318.7(3.1) μu for ^{134}Sb^m at 279(1) keV; $M-A$=–73740.0(2.9) keV										Nub269	**
*^{134}Sb−^{136}Xe$_{.985}$	D_M=12206(36) μu for ^{134}Sb^m at 279(1) keV; $M-A$=–73762(33) keV										Nub269	**
*	assuming high spin is favored, as for ^{136}I^m										GAu	**
*^{134}Pr−^{133}Cs$_{1.008}$	Most certainly gs. Mixture with isomer not completely excluded										00Be42	**
*^{134}Pr−^{133}Cs$_{1.008}$	D_M=11029(16) μu for mixture gs+m at 67.7(0.4) keV; $M-A$=–78503(15) keV										Nub269	**
*^{131}Cs−^{134}Cs$_{.244}$ ^{130}Cs^x	D_M=–1330(50) keV for mixture gs+m at 138.7441 keV										Nub269	**
*^{134}Sb(β^-)^{134}Te	E_{β^-}=8400(300), and 6800(300) from ^{134}Sb^m at 279(1) to ^{134}Te^m at 1691.34										Nub269	**
*^{134}Sb(β^-)^{134}Te	E_{β^-}=5840(240) from ^{134}Sb^m at 279(1) to (6)$^+$ level at 2397.70 keV										Ens04a	**
*^{134}Sb(β^-)^{134}Te	E_{β^-}=8420(120), and 6710(210),6980(210),6070(150) from ^{134}Sb^m at 279(1)										Nub269	**
*	to 6$^+$ levels at 1691.34, 1691.34, 2397.70 keV										Ens04a	**
*^{134}Te(β^-)^{134}I	E_{β^-}=730(110) 610(160) to 1$^+$ levels at 846.688 923.432 keV										Ens04a	**
*^{134}I(β^-)^{134}Xe	E_{β^-}=2410(60) 1680(60) 1250(60) to 4$^+$ levels at 1731.17 2588.46 2867.38 keV										Ens04a	**
*^{134}Cs(β^-)^{134}Ba	E_{β^-}=658.0(0.4) to 4$^+$ level at 1400.590 keV										Ens04a	**
*^{134}Ce(ε)^{134}La	LK=0.798(0.04); also Q_{β^+} >375 keV										76Gr09	**
*^{134}Pr(β^+)^{134}Ce	E_{β^+}=4120(90) to 4$^+$ level at 1048.68 keV										Ens04a	**
*^{134}Pm(β^+)^{134}Nd	E_{β^+}=7360(200) to 4$^+$ level at 788.92 keV										Ens04a	**
^{135}Sn−u	−65107.2	16.1	−65091	3	1.0	U			TR1	1.0	25Mo18	
^{135}Sb−u	−74932	103	−74815.6	2.8	0.5	o			GT2	2.5	08Kn.A	
	−74943	130			0.4	U			GT2	2.5	08Su19	
^{135}Te−u	−83643	106	−83445.3	1.8	0.7	U			GT2	2.5	08Kn.A	
	−83441	132			−0.0	U			GT2	2.5	08Su19	
C$_8$ N O H$_9$−^{135}Ba	162731	48	162725.40	0.21	−0.1	U			R07	1.5	68De17	
C$_{11}$ H$_3$−^{135}Ba	117822	77	117786.59	0.21	−0.3	U			R07	1.5	68De17	
C$_{12}$ H$_7$−^{135}Ba O	154160	46	154172.10	0.21	0.2	U			R07	1.5	68De17	
^{135}Ce−u	−90779	30	−90839	11	−2.0	1	13	13 ^{135}Ce	GS2	1.0	05Li24	*
^{135}Pr−u	−86897	30	−86888	13	0.3	R			GS2	1.0	05Li24	
^{135}Nd−u	−81800	130	−81819	21	−0.1	o			GS1	1.0	00Ra23	
	−81811	36			−0.2	R			GS2	1.0	05Li24	*
^{135}Pm−u	−75225	97				2			GS2	1.0	05Li24	*
^{135}Sm−u	−67480	166				2			GS2	1.0	05Li24	*
^{135}Sn−^{130}Xe$_{1.038}$	35065.9	3.3				2			JY1	1.0	12Ha25	
^{135}Sb−^{130}Xe$_{1.038}$	25342.4	3.1	25341.7	2.8	−0.2	1	84	84 ^{135}Sb	JY1	1.0	12Ha25	
^{135}Te−^{130}Xe$_{1.038}$	16713.0	2.9	16712.0	1.8	−0.3	1	41	41 ^{135}Te	JY1	1.0	12Ha25	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{135}Sn−^{133}Cs$_{1.015}$		30927	37	30875	3	−1.4	U			CP1	1.0	13Va12	
^{135}Sb−^{133}Cs$_{1.015}$		21146.8	7.0	21150.6	2.8	0.5	1	16	16 ^{135}Sb	CP1	1.0	13Va12	
^{135}Te−^{133}Cs$_{1.015}$		12520.3	2.4	12521.0	1.8	0.3	1	59	59 ^{135}Te	CP1	1.0	13Va12	
^{135}I−^{133}Cs$_{1.015}$		6026.6	2.3	6025.6	2.2	−0.4	1	92	92 ^{135}I	CP1	1.0	13Va12	
^{135}Cs−^{133}Cs$_{1.015}$		1958	18	1943.2	0.4	−0.8	o			MA1	1.0	90St25	
		1957	14			−1.0	U			MA1	1.0	99Am05	
^{135}Pr−^{133}Cs$_{1.015}$		9080	14	9078	13	−0.1	2			MA5	1.0	00Be42	
^{135}Nd−^{133}Cs$_{1.015}$		14144	25	14148	21	0.1	2			MA5	1.0	00Be42	*
^{135}Te−^{136}Xe$_{.993}$		8686	10	8690.7	1.8	0.5	U			CP1	1.0	12Va02	
^{135}I−^{136}Xe$_{.993}$		2186.3	8.3	2195.4	2.2	1.1	U			CP1	1.0	12Va02	
^{135}Cs−^{135}Ba		288.43	0.32	288.5	0.3	0.1	1	92	89 ^{135}Cs	JY2	1.0	20De20	
^{135}Ba−C$_{10}$ H$_{14}$		−203860	20	−203861.94	0.21	−0.0	U			M17	2.5	66Be10	
^{135}Ba−^{134}Ba		1177	2	1180.20	0.11	0.6	U			M17	2.5	66Be10	
		1161	70			0.2	U			R07	1.5	68De17	
		1168	78			0.1	U			R07	1.5	68De17	
^{134}Cs(n,γ)^{135}Cs		8762	1	8762.1	0.3	0.1	1	11	11 ^{135}Cs	ILn		92Ul.A	
^{134}Ba(n,γ)^{135}Ba		6973.2	0.4	6971.97	0.10	−3.1	C			BNn		77Ko.A	
		6972.17	0.18			−1.1	–			MMn		90Is07	Z
		6971.84	0.17			0.7	–			Ltn		93Bo01	Z
		6973.24	0.22			−5.8	B			BNn		93Ch21	
		6971.87	0.18			0.5	–			Bdn		06Fi.A	
^{134}Ba(d,p)^{135}Ba		4746	15	4747.40	0.10	0.1	U			ANL		70Vo04	
^{134}Ba(n,γ)^{135}Ba	ave.	6971.96	0.10	6971.97	0.10	0.1	1	95	74 ^{134}Ba			average	
^{135}Tb(p)^{134}Gd		1188	7				3			Arp		04Wo07	
^{135}Sb(β^-)^{135}Te		8120	50	8038	3	−1.6	U			Stu		89Ho08	
^{135}Te(β^-)^{135}I		5950	240	6050.4	2.7	0.4	o			Stu		77Lu06	
		5950	100			1.0	o			Bwg		79Ke02	*
		5970	200			0.4	U					85Sa15	*
		5960	100			0.9	U			Bwg		87Gr.A	
		5888	13			12.5	B			Stu		07Fo02	
^{135}I(β^-)^{135}Xe		2780	80	2634	4	−1.8	U					70Ma19	
		2590	50			0.9	U			Stu		76Lu04	*
		2627	6			1.2	1	41	33 ^{135}Xe	Stu		99Fo01	
^{135}Xe(β^-)^{135}Cs		1155	10	1169	4	1.4	–					52Be55	*
		1167	5			0.3	–			Stu		99Fo01	*
	ave.	1165	4			0.9	1	67	67 ^{135}Xe			average	
^{135}Cs(β^-)^{135}Ba		205	5	268.69	0.29	12.7	B					53Li01	
^{135}La(β^+)^{135}Ba		1200	10	1207	9	0.7	1	89	89 ^{135}La			71Ba18	*
^{135}Ce(β^+)^{135}La		2027	5	2027	5	0.0	–					76Ga.A	*
		2016	13			0.9	–					81Sa09	*
	ave.	2026	5			0.3	1	98	87 ^{135}Ce			average	
^{135}Pr(β^+)^{135}Ce		3720	150	3680	16	−0.3	U					54Ha68	*
^{135}Pm^m(β^+)^{135}Nd		6040	150	6390#	50#	2.3	D			Dbn		95Ve08	*
*^{135}Ce−u	$M-A$=–84114(28) keV for ^{135}Ce^m at 445.81 keV											Nub269	**
*^{135}Nd−u	$M-A$=–76174(28) keV for mixture gs+m at 64.95 keV											Nub269	**
*^{135}Pm−u	$M-A$=–69952(28) keV for mixture gs+m at 240#100 keV											Nub269	**
*^{135}Sm−u	$M-A$=–62857(38) keV for mixture gs+m at 0#300 keV											Nub269	**
*^{135}Nd−^{133}Cs$_{1.015}$	D_M=14179(14) μu for gs+m mixture at 64.95 keV; $M-A$=–76185(13) keV											Nub269	**
*^{135}Tb(p)^{134}Gd	Trends in Mass Surface TMS suggest ^{135}Tb 300 keV less bound											GAu	*G
*^{135}Te(β^-)^{135}I	E_{β^-}=5120(120) to 5/2$^-$ level at 870.52 keV and other E_{β^-}											Ens083	**
*^{135}Te(β^-)^{135}I	E_{β^-}=5370(100) to 5/2$^+$ level at 603.68 keV											Ens083	**
*^{135}I(β^-)^{135}Xe	E_{β^-}=1320(50) to 5/2$^+$ level at 1260.416 keV, and other E_{β^-}											Ens083	**
*^{135}Xe(β^-)^{135}Cs	E_{β^-}=905(10) 917(5) respectively, to 5/2$^+$ level at 249.767 keV											Ens083	**
*^{135}La(β^+)^{135}Ba	p$^+$ =7(1)×10^{-5}, from reanalysis											AHW	**
*^{135}La(β^+)^{135}Ba	But 65Mo05 says p$^+$ <0.002% (2×10^{-5}) or E_{β^+} <125 keV											65Mo05	**
*^{135}Ce(β^+)^{135}La	E_{β^+}=705(5) 694(13) respectively, to 1/2$^+$ level at 300.052 keV											Ens083	**
*^{135}Pr(β^+)^{135}Ce	E_{β^+}=2500(100) to levels (5/2$^+$) 296.11 and 3/2($^+$) 82.67 keV, roughly equal											Ens083	**
*^{135}Pm^m(β^+)^{135}Nd	E_{β^+}=4920(150) to mixture ground state and (11/2$^-$) level at 198.5 keV											Ens083	**
*^{135}Pm^m(β^+)^{135}Nd	Trends in Mass Surface TMS suggest ^{135}Pm^m 350 keV less bound											GAu	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{136}Sn−u		−60197.9	31.3				2				TR1	1.0	25Mo18	
^{136}Te−u		−79945	25	−79898.3	2.5	1.9	U				GS3	1.0	12Ch19	
^{136}Xe−^{120}Sn$_{1.133}$		18019.8	3.1	18019.1	1.1	−0.2	U				JY1	1.0	11Ha48	
C_{10} H_{16}−^{136}Xe		217982.	3.9	217986.038	0.008	0.4	U				M16	2.5	63Da10	
^{136}Xe−u		−92793.6	9.0	−92785.527	0.008	0.4	U				ACC	2.5	90Me08	
^{136}Xe−^{28}Si$_4$ D_{12}		−169712.9828	0.0162	−169712.999	0.007	−1.0	1	20	20	^{136}Xe	FS1	1.0	07Re03	
C_{11} H_4−^{136}Ba		126737	56	126724.27	0.20	−0.2	U				R07	1.5	68De17	
C_8 N O H_{10}−^{136}Ba		171635	56	171663.08	0.20	0.3	U				R07	1.5	68De17	
C_{12} H_8−^{136}Ba O		163094	40	163109.77	0.20	0.3	U				R07	1.5	68De17	
^{136}Ba−u		−95424.05	0.32	−95424.14	0.20	−0.3	1	40	40	^{136}Ba	MS1	1.0	19Sa.A	
^{136}La−u		−92394	88	−92370	60	0.3	2				GS2	1.0	05Li24	*
C_{10} H_{16}−^{136}Ce		218128	50	218071.2	0.3	−0.3	U				R05	4.0	65De13	
C_{12} H_8−^{136}Ce O		160563	36	160556.3	0.3	−0.0	U				R05	4.0	65De13	
^{136}Nd−u		−85044	30	−85024	13	0.7	R				GS2	1.0	05Li24	
^{136}Pm−u		−76373	78	−76400	70	−0.3	2				GS2	1.0	05Li24	*
^{136}Sm−u		−71768	30	−71724	13	1.5	R				GS2	1.0	05Li24	
^{136}Sb−^{130}Xe$_{1.046}$		31675.1	6.8	31678	6	0.5	1	85	85	^{136}Sb	JY1	1.0	12Ha25	
^{136}Te−^{130}Xe$_{1.046}$		21029.9	3.1	21030.9	2.5	0.3	1	63	63	^{136}Te	JY1	1.0	12Ha25	
^{136}Sb−^{133}Cs$_{1.023}$		27489	16	27472	6	−1.1	1	15	15	^{136}Sb	CP1	1.0	13Va12	
^{136}Te−^{133}Cs$_{1.023}$		16827.7	6.8	16824.3	2.5	−0.5	1	13	13	^{136}Te	CP1	1.0	13Va12	
^{136}I^m−^{133}Cs$_{1.023}$		11557.1	3.3	11552.4	0.7	−1.4	U				JY2	1.0	24Be25	
^{136}Xe−^{133}Cs$_{1.023}$		3936.5	1.9	3937.120	0.011	0.3	U				MA8	1.0	09Ne11	
^{136}Cs−^{133}Cs$_{1.023}$		4007	18	4021.9	0.5	0.8	o				MA1	1.0	90St25	
		4021	14			0.1	U				MA1	1.0	99Am05	
^{136}Pr−^{133}Cs$_{1.023}$		9418	15	9400	12	−1.2	1	67	67	^{136}Pr	MA5	1.0	00Be42	
^{136}Nd−^{133}Cs$_{1.023}$		11703	14	11699	13	−0.3	2				MA5	1.0	00Be42	
^{136}Pm^m−^{133}Cs$_{1.023}$		20429	100				2				MA5	1.0	00Be42	*
^{136}Sm−^{133}Cs$_{1.023}$		25009	15	24998	13	−0.7	2				MA5	1.0	00Be42	
^{136}Xe−^{134}Xe$_{1.015}$		3245.8	3.8	3240.546	0.012	−1.4	o				MA8	1.0	05He.A	
		3244.3	4.0			−0.9	U				MA8	1.0	06He29	
^{136}Te−^{136}Xe		12887.9	5.0	12887.2	2.5	−0.1	1	24	24	^{136}Te	CP1	1.0	12Va02	
^{136}I−^{136}I^m		−230.9	4.6				3				JY2	1.0	24Be25	
^{136}I^m−^{136}Xe		7611.2	4.9	7615.3	0.7	0.8	U				CP1	1.0	12Va02	*
		7615.33	0.65				2				JY1	1.0	24Be25	
^{136}Xe−^{136}Ba		2639.6	0.6	2638.61	0.20	−0.7	–				H49	2.5	10Mc04	
		2638.62	0.52			−0.0	–				JY1	1.0	11Ko03	
	ave.	2638.7	0.5			−0.2	1	17	17	^{136}Ba			average	
^{136}Cs−^{136}Ba		2723.40	0.48				2				JY2	1.0	23Ge04	
^{136}Ce−^{136}Ba		2553.46	0.29	2553.46	0.23	−0.0	–				JY1	1.0	11Ko03	
		2553.42	0.37			0.1	–				SH1	1.0	12Ne10	
	ave.	2553.44	0.23			0.0	1	100	100	^{136}Ce			average	
^{136}Xe−$^{13}C_3$ O_6		−72337.7553	0.0109	−72337.747	0.007	0.7	–				FS1	1.0	07Re03	
		−72337.7464	0.0109			−0.1	–				FS1	1.0	07Re03	
	ave.	−72337.751	0.008			0.5	1	82	80	^{136}Xe			average	
^{136}Ba−^{135}Ba		−1115	3	−1112.65	0.04	0.3	U				M17	2.5	66Be10	
		−1119	50			0.1	U				R07	1.5	68De17	
		−1074	50			−0.5	U				R07	1.5	68De17	
^{136}Ba−^{134}Ba		67	5	67.56	0.11	0.0	U				M17	2.5	66Be10	
		69	128			−0.0	U				R07	1.5	68De17	
		72	78			−0.0	U				R07	1.5	68De17	
N_{10}−^{136}Xe		123525.5778	0.0235	123525.569	0.007	−0.4	U				FS1	1.0	07Re03	
^{136}Te(β^-n)^{135}I		1285	50	1283	3	−0.0	U						84Kr.B	
^{136}Xe(d,^{3}He)^{135}I		−4438	30	−4445.5	2.1	−0.2	U				Oak		71Wi04	
^{136}Xe(d,t)^{135}Xe		−1723	40	−1830	4	−2.7	U				Oak		68Mo21	
^{135}Ba(n,γ)^{136}Ba		9106.4	0.8	9107.74	0.04	1.7	U						69Ge07	
		9107.74	0.04			0.0	–				MMn		90Is07	Z
		9107.73	0.19			0.1	–				Bdn		06Fi.A	
^{135}Ba(d,p)^{136}Ba		6886	15	6883.18	0.04	−0.2	U				ANL		70Vo04	
^{135}Ba(n,γ)^{136}Ba	ave.	9107.74	0.04	9107.74	0.04	0.1	1	99	75	^{135}Ba			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{136}Te(β^-)^{136}I	5100	150	5126	5	0.2	U					77Sc21	
	5095	100			0.3	U			Bwg		87Gr.A	
	5086	20			2.0	U			Stu		07Fo02	
^{136}I(β^-)^{136}Xe	6960	100	6879	4	−0.8	U					59Jo37	*
	6690	150			1.3	U			Stu		76Lu04	*
	6925	70			−0.7	U			Bwg		87Gr.A	
	6850	20			1.4	U			Stu		07Fo02	
^{136}I^m(β^-)^{136}Xe	7100	230	7093.6	0.6	−0.0	o			Stu		76Lu04	*
	7705	120			−5.1	C			Bwg		87Gr.A	
	7051	12			3.6	B			Stu		07Fo02	
^{136}Cs(β^-)^{136}Ba	2548.1	2.0	2536.8	0.4	−5.6	B					54Ol05	*
	2549	5			−2.4	U					65Re07	*
^{136}La(β^+)^{136}Ba	2870	70	2850	50	−0.3	R					59Gi50	
^{136}Pr(β^+)^{136}Ce	5084	50	5168	11	1.7	U					68Zh04	*
	5114	75			0.7	U					71Ke07	*
	5134	20			1.7	1	33	33 ^{136}Pr	IRS		83Al.B	*
^{136}Nd(β^+)^{136}Pr	2501	50	2141	16	−7.2	B					68Zh04	*
	2211	25			−2.8	B					75Br16	*
^{136}Pm(β^+)^{136}Nd	7850	200	8030	70	0.9	R			IRS		83Al06	*
*^{136}La−u	$M-A$=–85935(32) keV for mixture gs+m at 259.3 keV										Nub269	**
*^{136}Pm−u	$M-A$=–71091(28) keV for mixture gs+m at 100(120) keV										Nub269	**
*^{136}Pm^m−^{133}Cs$_{1.023}$	Slightly contaminated by ground state, original error (20) increased										00Be42	**
*^{136}I^m−^{136}Xe	High spin isomer is preferred (see also ^{134}Sb)										GAu	**
*^{136}I(β^-)^{136}Xe	E_{β^-}=7000(100), 5610(150), 4280(150) to ground state, 2^+ levels 1313.027, 2634.16 keV										Ens026	**
*^{136}I(β^-)^{136}Xe	E_{β^-}=5370(400), 4700(320), 3920(220) to 2^+ levels 1313.027, 2289.53, 2634.16										Ens026	**
*^{136}I^m(β^-)^{136}Xe	E_{β^-}=5170(400) 4670(270) to ^{136}Xe^m 6^+ at 1891.703 and 2444.39 level										Ens026	**
*^{136}Cs(β^-)^{136}Ba	E_{β^-}=341(2) 342(5) respectively, to 6^+ level at 2207.077 keV										Ens026	**
*^{136}Pr(β^+)^{136}Ce	E_{β^+}=2970(50) 3000(75) 3020(20) respectively, to 2^+ level at 1092.09 keV										Ens026	**
*^{136}Nd(β^+)^{136}Pr	E_{β^+}=1330(50) to 1^+ level at 149.11 keV										Ens026	**
*^{136}Nd(β^+)^{136}Pr	K/β^+=13.2(0.5) to 1^+ level at 149.11 keV										Ens026	**
*^{136}Pm(β^+)^{136}Nd	E_{β^-}=4732(70) probably from high spin isomer going to several										AHW	**
*	high spin levels around 2100 keV										AHW	**
^{137}Sn−u	−53795.3	10.7				2			TR1	1.0	25Mo18	
^{137}Sb−u	−64445	215	−64427	6	0.1	o			GT1	1.5	04Ma.A	
	−65068	186			1.4	B			GT3	2.5	16Kn03	
^{137}Te−u	−74528	101	−74400.6	2.3	0.5	o			GT2	2.5	08Kn.A	
	−74386	129			−0.0	U			GT2	2.5	08Su19	
^{137}I−u	−82145	130	−81978.1	0.5	0.5	U			GT2	2.5	08Su19	
C_{11} H_5−^{137}Ba	133366	24	133297.89	0.21	−1.9	U			R07	1.5	68De17	
C_7 ^{13}C N O H_{10}−^{137}Ba	173792	73	173766.51	0.21	−0.2	U			R07	1.5	68De17	
C_{12} H_9−^{137}Ba O	169692	39	169683.40	0.21	−0.1	U			R07	1.5	68De17	
^{137}La−u	−93556	30	−93549.5	1.8	0.2	U			GS2	1.0	05Li24	
^{137}Ce−u	−92101	85	−92237.5	0.3	−1.6	U			GS2	1.0	05Li24	*
^{137}Nd−u	−85438	30	−85437	13	0.0	1	18	18 ^{137}Nd	GS2	1.0	05Li24	
^{137}Pm−u	−79613	64	−79520	14	1.4	U			GS2	1.0	05Li24	*
^{137}Sm−u	−72982	51	−72990	30	−0.2	1	36	36 ^{137}Sm	GS2	1.0	05Li24	*
^{137}Te−^{130}Xe$_{1.054}$	27300.0	2.7	27300.5	2.3	0.2	1	70	70 ^{137}Te	JY1	1.0	12Ha25	
^{137}Sb−^{133}Cs$_{1.030}$	32907	56	32957	6	0.9	U			CP1	1.0	13Va12	
^{137}Te−^{133}Cs$_{1.030}$	22985.0	4.1	22983.8	2.3	−0.3	1	30	30 ^{137}Te	CP1	1.0	13Va12	
^{137}I−^{133}Cs$_{1.030}$	15406.42	0.52				2			JY2	1.0	24Be25	
^{137}Xe−^{133}Cs$_{1.030}$	8943.6	2.0	8942.25	0.11	−0.7	U			MA8	1.0	09Ne11	
^{137}Cs−^{133}Cs$_{1.030}$	4452	19	4473.84	0.28	1.1	o			MA1	1.0	90St25	
	4470	14			0.3	U			MA1	1.0	99Am05	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{137}Pr−^{133}Cs$_{1.030}$		8095	15	8064	9	−2.1	1	34	34	^{137}Pr	MA5	1.0	00Be42	
^{137}Nd−^{133}Cs$_{1.030}$		11947	14	11948	13	0.0	1	81	81	^{137}Nd	MA5	1.0	00Be42	
^{137}Pm−^{133}Cs$_{1.030}$		17864	14				2				MA5	1.0	00Be42	
^{137}Sm−^{133}Cs$_{1.030}$		24393	42	24390	30	−0.0	1	53	53	^{137}Sm	MA5	1.0	00Be42	*
^{137}Eu−^{133}Cs$_{1.030}$		32815.2	4.7				2				MA8	1.0	13Wo05	
^{137}Ba ^{35}Cl−^{135}Ba ^{37}Cl		3089.1	0.6	3088.89	0.11	−0.1	U				H49	2.5	10Mc04	
^{137}Sb−^{136}Xe$_{1.007}$		29010.3	6.8	29008	6	−0.4	1	91	91	^{137}Sb	JY1	1.0	24Be25	
^{137}Te−^{136}Xe$_{1.007}$		19057	18	19034.4	2.3	−1.3	U				CP1	1.0	12Va02	
^{137}I−^{136}Xe$_{1.007}$		11463.2	9.0	11457.0	0.5	−0.7	U				CP1	1.0	12Va02	
^{137}Xe−^{136}Xe$_{1.007}$		5004	11	4992.79	0.11	−1.0	U				CP1	1.0	12Va02	
^{137}Sb−^{137}Cs		28457	21	28483	6	1.2	1	10	9	^{137}Sb	TR1	1.0	26Si.A	
^{137}Ba−^{136}Ba		1249	3	1251.41	0.07	0.3	U				M17	2.5	66Be10	
		1222	50			0.4	U				R07	1.5	68De17	
		1227	44			0.4	U				R07	1.5	68De17	
^{137}Ba−^{135}Ba		143	3	138.76	0.08	−0.6	U				M17	2.5	66Be10	
		69	63			0.7	U				R07	1.5	68De17	
		106	46			0.5	U				R07	1.5	68De17	
^{137}I(β^-n)^{136}Xe		1850	30	1995.8	0.5	4.9	C						84Kr.B	
^{136}Xe(n,γ)^{137}Xe		4025.5	0.5	4025.56	0.10	0.1	U						77Fo02	Z
		4025.8	0.3			−0.8	2						77Pr07	Z
		4025.53	0.11			0.3	2				Bdn		06Fi.A	
^{136}Xe(d,p)^{137}Xe		1637	20	1801.00	0.10	8.2	F				Oak		68Mo21	*
^{136}Xe(^{3}He,d)^{137}Cs		1918	12	1912.04	0.26	−0.5	U				ChR		81Ha08	
^{136}Ba(n,γ)^{137}Ba		6891	5	6905.64	0.07	2.9	U						69Gr31	
		6905.54	0.10			1.0	–				MMn		90Is07	Z
		6905.70	0.12			−0.5	–				Ltn		95Bo03	Z
		6905.74	0.16			−0.6	–				Bdn		06Fi.A	
^{137}Ba(γ,n)^{136}Ba		−6949	38	−6905.64	0.07	1.1	U				Phi		60Ge01	
^{136}Ba(d,p)^{137}Ba		4680	15	4681.07	0.07	0.1	U				ANL		70Vo04	
^{136}Ba(n,γ)^{137}Ba	ave.	6905.63	0.07	6905.64	0.07	0.1	1	98	79	^{137}Ba			average	
^{136}Ce(n,γ)^{137}Ce		7481.3	0.4	7481.53	0.16	0.6	–						81Ko.A	Z
		7481.58	0.17			−0.3	–				Bdn		06Fi.A	
	ave.	7481.54	0.16			−0.0	1	100	100	^{137}Ce			average	
^{137}Te(β^-)^{137}I		7030	300	7058.3	2.2	0.1	U						85Sa15	
		6925	130			1.0	U				Bwg		87Gr.A	
^{137}I(β^-)^{137}Xe		5880	60	6021.3	0.5	2.4	U				Bwg		87Gr.A	
^{137}Xe(β^-)^{137}Cs		4140	70	4162.30	0.28	0.3	U						64On03	
		4150	100			0.1	U						68Ho22	
^{137}Cs(β^-)^{137}Ba		1173.29	0.84	1175.63	0.17	2.8	U						68Wo02	*
		1175.55	0.26			0.3	–						78Ch22	*
		1175.69	0.23			−0.3	–						83Be18	*
	ave.	1175.63	0.17			0.0	1	100	100	^{137}Cs			average	
^{137}Ce(β^+)^{137}La		1222.1	1.6				2						81Ar.A	*
^{137}Pr(β^+)^{137}Ce		2702	10	2717	8	1.5	1	66	66	^{137}Pr			73Bu17	
^{137}Nd(β^+)^{137}Pr		3497	40	3618	14	3.0	B						73Bu18	*
		3690	54			−1.3	U						85Af.A	*
^{137}Pm^m(β^+)^{137}Nd		5690	130	5670	40	−0.2	–				IRS		83Al06	*
		5650	60			0.3	–				Dbn		95Ve08	*
	ave.	5660	50			0.2	1	67	65	^{137}Pm^m			average	
^{137}Sm(β^+)^{137}Pm^m		5900	70	5920	50	0.3	1	45	35	^{137}Pm^m	Dbn		95Ve08	
*^{137}Ce−u	$M-A$=–85665(29) keV for mixture gs+m at 254.29 keV												Nub269	**
*^{137}Pm−u	$M-A$=–74079(28) keV for mixture gs+m at 160(50) keV												Nub269	**
*^{137}Sm−u	$M-A$=–67932(28) keV for mixture gs+m at 100#50 keV												Nub269	**
*^{137}Sm−^{133}Cs$_{1.030}$	Might be a mixture of ground state and isomer say authors												00Be42	**
*	D_M=24447(14) μu for mixture gs+m at 100#50 keV; $M-A$=–67941(13) keV												Nub269	**
*^{136}Xe(d,p)^{137}Xe	F : error severely underestimated and value low, see excitation energies												AHW	**
*^{137}Cs(β^-)^{137}Ba	E_{β^-}=511.63(0.84) 513.89(0.26) 514.03(0.23) to ^{137}Ba^m at 661.659 keV												Nub269	**
*^{137}Ce(β^+)^{137}La	E_{β^+}=189.5(1.6) to 5/2$^+$ level at 10.59 keV												Ens079	**
*^{137}Nd(β^+)^{137}Pr	E_{β^+}=2400(40) E_{β^+}=2592(54) respectively, to 3/2$^+$ level at 75.5 keV												Ens079	**
*^{137}Pm^m(β^+)^{137}Nd	E_{β^+}=4132(+150–115) 4110(60) respectively, to 11/2$^-$ ^{137}Nd^m at 519.43 keV												Nub269	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference
^{138}Sn−u	−48784.0	22.5				2			TR1	1.0	25Mo18
^{138}Sb−u	−58208	457	−58380	70	−0.2	U			GT3	2.5	16Kn03
^{138}Te−u	−70940	247	−70528	4	1.1	o			GT1	1.5	04Ma.A
	−70583	106			0.2	o			GT2	2.5	08Kn.A
	−70591	131			0.2	U			GT2	2.5	08Su19
C_{10} H_{18}−^{138}Ba	235609	20	235603.46	0.21	−0.1	U			M17	2.5	66Be10
C_{11} H_7−^{138}Ba H	141649	51	141703.07	0.21	0.7	U			R07	1.5	68De17
C_{11} H_6−^{138}Ba	141701	30	141703.07	0.21	0.0	U			R07	1.5	68De17
C_{12} H_{10}−^{138}Ba O	178106	15	178088.58	0.21	−0.8	U			R07	1.5	68De17
	178105	49			−0.2	U			R07	1.5	68De17
C_{11} ^{13}C H_9−^{138}Ba O	173612	37	173618.38	0.21	0.1	U			R07	1.5	68De17
C_{10} H_{18}−^{138}Ce	234799	60	234856.3	0.5	0.2	U			R05	4.0	65De13
C_{12} H_{10}−^{138}Ce O	177382	46	177341.5	0.5	−0.2	U			R05	4.0	65De13
C_9 ^{13}C H_{17}−^{138}Ce	230358	60	230386.1	0.5	0.1	U			R05	4.0	65De13
^{138}Pr^m−u	−88896	30	−88867	17	1.0	1	34	34 ^{138}Pr^m	GS2	1.0	05Li24
^{138}Nd−u	−88060	130	−88050	13	0.1	o			GS1	1.0	00Ra23
	−88060	30			0.3	R			GS2	1.0	05Li24
^{138}Pm−u	−80226	140	−80423	13	−1.4	o			GS1	1.0	00Ra23
	−80437	30			0.5	1	18	18 ^{138}Pm	GS2	1.0	05Li24
^{138}Sm−u	−76766	30	−76756	13	0.3	R			GS2	1.0	05Li24
^{138}Eu−u	−66291	30				2			GS2	1.0	05Li24
^{138}Te−^{130}Xe$_{1.062}$	31945.3	4.7	31946	4	0.0	1	75	75 ^{138}Te	JY1	1.0	12Ha25
^{138}I−^{130}Xe$_{1.062}$	25216.97	0.69	25217.1	0.5	0.2	1	62	62 ^{138}I	JY2	1.0	24Be25
^{138}Te−^{133}Cs$_{1.038}$	27614.0	8.1	27613	4	−0.1	1	25	25 ^{138}Te	CP1	1.0	13Va12
^{138}I−^{133}Cs$_{1.038}$	20885.04	0.88	20884.9	0.5	−0.2	1	38	38 ^{138}I	JY2	1.0	24Be25
^{138}Xe−^{133}Cs$_{1.038}$	12284.1	3.5	12287	3	0.9	1	74	74 ^{138}Xe	MA8	1.0	09Ne11
^{138}Cs−^{133}Cs$_{1.038}$	9158	14	9158	10	0.0	1	49	49 ^{138}Cs	MA1	1.0	99Am05
^{138}Ba−^{133}Cs$_{1.038}$	3388	14	3387.99	0.21	−0.0	U			MA1	1.0	99Am05
^{138}Nd−^{133}Cs$_{1.038}$	10093	14	10091	13	−0.2	2			MA5	1.0	00Be42
^{138}Pm−^{133}Cs$_{1.038}$	17721	14	17718	13	−0.2	1	82	82 ^{138}Pm	MA5	1.0	00Be42
^{138}Sm−^{133}Cs$_{1.038}$	21387	14	21385	13	−0.2	2			MA5	1.0	00Be42
^{138}I−^{136}Xe$_{1.015}$	16903.7	6.4	16921.3	0.5	2.8	U			CP1	1.0	12Va02
^{138}Xe−^{136}Xe$_{1.015}$	8332.2	5.9	8324	3	−1.5	1	26	26 ^{138}Xe	CP1	1.0	12Va02
^{138}Ba−^{136}Xe$_{1.015}$	−575.98	0.48	−575.57	0.21	0.9	1	20	20 ^{138}Ba	MS1	1.0	19Sa36
^{138}Ba ^{35}Cl−^{136}Ba ^{37}Cl	3621.1	0.6	3621.39	0.11	0.2	U			H49	2.5	10Mc04
^{138}Ce−^{136}Xe$_{1.015}$	170.9	1.7	171.5	0.5	0.4	1	9	9 ^{138}Ce	MS1	1.0	19Sa36
^{138}Sb−^{138}Ba	36373	72				2			TR1	1.0	26Si.A
^{138}La−^{138}Ba	1877.29	0.40	1877.0	0.4	−0.8	1	82	82 ^{138}La	MS1	1.0	19Sa36
^{138}Ce−^{138}Ba	746.12	0.77	747.1	0.5	1.3	1	38	38 ^{138}Ce	MS1	1.0	19Sa36
^{138}La−^{138}Ce	1129.36	0.51	1129.9	0.4	1.0	1	72	53 ^{138}Ce	MS1	1.0	19Sa36
^{138}Ba−^{137}Ba	−582	2	−580.15	0.04	0.4	U			M17	2.5	66Be10
	−480	27			−2.5	U			R07	1.5	68De17
	−553	40			−0.5	U			R07	1.5	68De17
^{138}Ba−^{136}Ba	676	3	671.26	0.08	−0.6	U			M17	2.5	66Be10
	658	98			0.1	U			R07	1.5	68De17
	628	43			0.7	U			R07	1.5	68De17
^{138}Ce−^{136}Ce	−1040	47	−1135.1	0.5	−0.5	U			R05	4.0	65De13
	−1158	20			0.5	U			M17	2.5	66Be10
^{138}Ba H−^{137}Ba	7399	88	7244.88	0.04	−1.2	U			R07	1.5	68De17
	7280	43			−0.5	U			R07	1.5	68De17
^{137}Ba(n,γ)^{138}Ba	8611.3	0.8	8611.72	0.04	0.5	U					68Ma35
	8611.72	0.04			0.1	1	99	79 ^{138}Ba	MMn		90Is07 Z
	8611.5	0.15			1.5	U			Ltn		95Bo05
	8611.63	0.18			0.5	U			Bdn		06Fi.A
^{137}Ba(d,p)^{138}Ba	6398	15	6387.16	0.04	−0.7	U			ANL		70Vo04
^{138}I(β^-)^{138}Xe	7820	70	8008.7	2.8	2.7	U			Bwg		87Gr.A

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{138}Xe(β^-)^{138}Cs		2700	50	2915	10	4.3	B					72Mo33	*
		2830	80			1.1	U			Trs		78Wo15	
^{138}Cs^x(IT)^{138}Cs		40	23				2					82Au01	*
^{138}Cs(β^-)^{138}Ba		5350	80	5375	9	0.3	U			Trs		78Wo15	
		5388	25			−0.5	–			Gsn		81De25	
		5370	15			0.3	–			McG		84He.A	
	ave.	5375	13			−0.0	1	51	51 ^{138}Cs			average	
^{138}La(ε)^{138}Ba		1620	15	1748.4	0.3	8.6	B					56Tu17	*
^{138}La(β^-)^{138}Ce		994	10	1052.5	0.4	5.8	B					57Gl20	*
		1159	40			−2.7	U					70El.A	*
		1052.7	4.3			−0.1	U					16Qu01	*
^{138}Pr(β^+)^{138}Ce		4437	10				2					71Af05	
^{138}Pr^m(β^+)^{138}Ce		4801	20	4787	16	−0.7	1	66	66 ^{138}Pr^m			64Fu08	*
^{138}Nd(β^+)^{138}Pr		2020	100	1111	15	−9.1	C					61Bo.B	
^{138}Pm(β^+)^{138}Nd		7090	100	7105	17	0.1	U			IRS		83Al06	
		7080	60			0.4	U			Dbn		95Ve08	
		7000	250			0.4	U					81De38	*
*^{138}Xe(β^-)^{138}Cs	E_{β^-}=2460(50), 2270(50) to (1$^-$ 2$^-$) level at 258.400, 1$^-$ at 412.260 keV											Ens035	**
*^{138}Cs^x(IT)^{138}Cs	Based on ^{138}Cs^m(IT)=79.9 keV											Nub269	**
*^{138}La(ε)^{138}Ba	L/K=1.40(0.25) to 2$^+$ level at 1435.816 keV											Ens035	**
*^{138}La(β^-)^{138}Ce	E_{β^-}=205(10) 370(40) 264.0(4.3) respectively, to 2$^+$ level at 788.744 keV											Ens035	**
*^{138}Pr^m(β^+)^{138}Ce	E_{β^+}=1650(20) to 7$^-$ level at 2129.17 keV											Ens035	**
*^{138}Pm(β^+)^{138}Nd	E_{β^+}=3900(200) to 5$^-$ level at 1990.4, 6$^+$ at 2134.3 and (5$^-$) at 2221.8keV											Ens035	**
^{139}Te−u		−64541	333	−64633	4	−0.2	U			GT1	1.5	04Ma.A	
^{139}Te−^{130}Xe$_{1.069}$		38515.7	3.8				2			JY1	1.0	12Ha25	
^{139}I−u		−73838	102	−73505.5	2.1	1.3	U			GT2	2.5	08Kn.A	
		−73567	130			0.2	U			GT2	2.5	08Su19	
C$_6$ ^{13}C O$_3$ H$_6$−^{139}La		128474	41	128686.0	0.7	1.3	U			R05	4.0	65De13	
C$_7$ O$_3$ H$_7$−^{139}La		133063	32	133156.2	0.7	0.7	U			R05	4.0	65De13	
C$_6$ N O$_3$ H$_5$−^{139}La		120496	21	120580.1	0.7	1.0	U			R05	4.0	65De13	
C$_{12}$ H$_{11}$−^{139}La O		184568	66	184797.9	0.7	0.9	U			R05	4.0	65De13	
C$_{11}$ ^{13}C H$_{10}$−^{139}La O		180100	58	180327.7	0.7	1.0	U			R05	4.0	65De13	
^{139}Nd−u		−87840	79	−88049	30	−2.6	U			GS2	1.0	05Li24	*
^{139}Sm−u		−77704	30	−77703	12	0.0	R			GS2	1.0	05Li24	
		−77711	30			0.3	R			GS2	1.0	05Li24	*
^{139}Eu−u		−70215	30	−70208	14	0.2	R			GS2	1.0	05Li24	
^{139}Te−^{133}Cs$_{1.045}$		34185	17	34170	4	−0.9	U			CP1	1.0	13Va12	
^{139}I−^{133}Cs$_{1.045}$		25296.1	4.3	25297.2	2.1	0.3	2			CP1	1.0	13Va12	
		25297.6	2.4			−0.1	2			JY2	1.0	24Be25	
^{139}Xe−^{133}Cs$_{1.045}$		17594.9	2.3				2			MA8	1.0	09Ne11	*
^{139}Cs−^{133}Cs$_{1.045}$		12163	14	12167	3	0.3	U			MA1	1.0	99Am05	
^{139}Ba−^{133}Cs$_{1.045}$		7649	14	7643.92	0.22	−0.4	U			MA1	1.0	99Am05	
^{139}Pm−^{133}Cs$_{1.045}$		15604	15	15602	15	−0.1	1	95	95 ^{139}Pm	MA5	1.0	00Be42	
^{139}Sm−^{133}Cs$_{1.045}$		21101	14	21099	12	−0.1	2			MA5	1.0	00Be42	
^{139}Eu−^{133}Cs$_{1.045}$		28597	16	28595	14	−0.1	2			MA5	1.0	00Be42	
^{139}I−^{136}Xe$_{1.022}$		21333	31	21321.3	2.1	−0.4	U			CP1	1.0	12Va02	
^{139}Xe−^{136}Xe$_{1.022}$		13618	12	13619.0	2.3	0.1	U			CP1	1.0	12Va02	
^{139}La−^{136}Xe$_{1.022}$		1189.96	0.67	1189.7	0.7	−0.4	1	94	94 ^{139}La	MS1	1.0	19Sa39	
^{139}La−^{138}La		−622	132	−761.2	0.8	−0.3	U			R05	4.0	65De13	
^{139}La−^{138}Ce		485	74	368.6	0.8	−0.4	U			R05	4.0	65De13	
^{133}Cs−^{139}Cs$_{.239}$ ^{131}Cs$_{.761}$		−1774	24	−1773.1	0.8	0.0	F			P33	2.5	86Au02	*
^{138}Cs^x−^{139}Cs$_{.496}$ ^{137}Cs$_{.504}$		770	40	800	25	0.3	U			P23	2.5	82Au01	
^{138}Ba(n,γ)^{139}Ba		4723.4	0.7	4723.43	0.04	0.0	U					69Mo13	
		4723.4	0.3			0.1	U					80Ba.A	
		4723.43	0.04				2			MMn		90Is07	Z
		4723.20	0.14			1.6	U			Bdn		06Fi.A	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{138}Ba(d,p)^{139}Ba	2495	10	2498.86	0.04	0.4	U			MIT		64Sp12	
	2496	15			0.2	U			Hei		67Wi08	
	2493	10			0.6	U			ANL		70Vo04	
^{139}La(γ,n)^{138}La	−8775	25	−8780.4	0.7	−0.2	U			Phi		60Ge01	
^{138}La(d,p)^{139}La	6553	3	6555.8	0.7	0.9	U			Tal		71Du02	
^{139}La(d,t)^{138}La	−2522	5	−2523.2	0.7	−0.2	U			Tal		72La20	
^{139}I(β^-)^{139}Xe	6815	100	7174.7	2.9	3.6	C			Bwg		87Gr.A	
	6806	23			16.0	B			Bwg		92Gr06	
^{139}Xe(β^-)^{139}Cs	5020	60	5056	4	0.6	U			Trs		78Wo15	
	5062	22			−0.3	U			Bwg		92Gr06	
^{139}Cs(β^-)^{139}Ba	4290	70	4213	3	−1.1	U			Trs		78Wo15	
	4190	25			0.9	o			Gsn		80Bl.A	
	4213	5			−0.0	o			Gsn		81De25	
	4214	4			−0.3	3			McG		84He.A	
	4211	5			0.4	3			Gsn		92Pr04	
^{139}Ba(β^-)^{139}La	2307	5	2308.6	0.6	0.3	U					75Fl07	*
	2336	25			−1.1	U			Gsn		81De25	
	2316	4			−1.9	U			McG		84He.A	
^{139}Ce(β^+)^{139}La	264.6	2.0	264.6	2.0	0.0	1	100	100 ^{139}Ce			96Hi14	*
^{139}Ce(ε)^{139}La	278	7			−1.9	U					Averag	*
^{139}Pr(β^+)^{139}Ce	2129	3	2129.1	3.0	0.0	1	100	100 ^{139}Pr			81Ar.A	
^{139}Nd(β^+)^{139}Pr	2787	50	2812	28	0.5	1	31	30 ^{139}Nd			75Vy02	*
^{139}Pm(β^+)^{139}Nd	4450	100	4516	26	0.7	–					77De06	*
	4540	40			−0.6	–			IRS		83Al06	
	4470	50			0.9	–			Dbn		95Ve08	
ave.	4507	30			0.3	1	75	70 ^{139}Nd			average	
^{139}Sm(β^+)^{139}Pm	5430	150	5121	17	−2.1	U					82De06	*
	5510	150			−2.6	U			IRS		83Al06	*
^{139}Eu(β^+)^{139}Sm	6080	50	6982	17	18.0	C			Dbn		95Ve08	*
*^{139}Nd−u	$M-A$=–81707(30) keV for mixture gs+m at 231.16 keV										Nub269	**
*^{139}Sm−u	$M-A$=–71930(28) keV for ^{139}Sm^m at 457.38 keV										Nub269	**
*^{139}Xe−^{133}Cs$_{1.045}$	Typo in original paper, ratio should read 1.045 245 4357(175)										GAu	**
*^{133}Cs−^{139}Cs$_{.239}$ ^{131}Cs.	F : rejection based on line-shape analysis										86Au02	**
*^{139}Ba(β^-)^{139}La	E_{β^-}=2141(5) to 5/2$^+$ level at 165.8576 keV										Ens014	**
*^{139}Ce(β^+)^{139}La	Q_{β^+}=98.7(1.7stat)(1.0syst) to 5/2$^+$ level at 165.8576 keV										Ens014	**
*^{139}Ce(ε)^{139}La	Average pK=0.73(0.01) to 5/2$^+$ level at 165.8576 keV in 10 references:										Ens014	**
*	pK=0.76 (0.04)										54Pr31	**
*	pK=0.73 (0.01)										56Ke23	**
*	pK=0.68 (0.02)										67Ma07	**
*	pK=0.75 (0.01)										68Ad08	**
*	pK=0.69 (0.02)										68Va08	**
*	pK=0.716(0.02)										72Ca07	**
*	pK=0.78 (0.02)										72Sc08	**
*	pK=0.726(0.010)										75Ha43	**
*	pK=0.801(0.034)										75Pl06	**
*	pK=0.705(0.020)										76Ha36	**
*^{139}Nd(β^+)^{139}Pr	E_{β^+}=1770(50); and 1170(50) from ^{139}Nd^m at 231.15 to 1/2$^-$ level at 821.98										Ens014	**
*^{139}Pm(β^+)^{139}Nd	E_{β^+}=3020(120), 2990(100) to 3/2$^+$ levels at 463.10, 402.77 keV										Ens014	**
*^{139}Sm(β^+)^{139}Pm	E_{β^+}=4100(150) to (1/2,3/2)$^+$ level at 306.69 keV										Ens014	**
*^{139}Sm(β^+)^{139}Pm	E_{β^+}=4735(+180–130) from ^{139}Sm^m at 457.38 to ^{139}Pm^m at 188.7 keV										Nub269	**
*^{139}Eu(β^+)^{139}Sm	E_{β^+}=4600(50) to ^{139}Sm^m at 457.38 keV										Nub269	**
^{140}Te−u	−60827	225	−60513	15	0.9	U			GT1	1.5	04Ma.A	
^{140}Te−^{130}Xe$_{1.077}$	43419	30	43407	15	−0.4	1	26	26 ^{140}Te	JY1	1.0	12Ha25	
^{140}I−u	−68181	193	−68310	50	−0.5	o			GT1	1.5	04Ma.A	
	−68463	102			0.6	o			GT2	2.5	08Kn.A	
	−68273	130			−0.1	o			GT2	2.5	08Su19	
	−68202	186			−0.2	U			GT2	2.5	16Kn03	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{140}Xe−u		−78449	103	−78354.2	2.5	0.4	o			GT2	2.5	08Kn.A	
		−78229	130			−0.4	U			GT2	2.5	08Su19	
C_{11} H_8−^{140}Ce		157116	29	157152.5	1.4	0.3	U			R05	4.0	65De13	
C_{10} ^{13}C H_7−^{140}Ce		152553	17	152682.3	1.4	1.9	U			R05	4.0	65De13	
C_{10} N H_6−^{140}Ce		144599	35	144576.4	1.4	−0.2	U			R05	4.0	65De13	
C_{10} N_2 H_8−^{140}Ce O		168207	48	168385.9	1.4	0.9	U			R05	4.0	65De13	
^{140}Nd−u		−90448	30	−90454	4	−0.2	U			GS2	1.0	05Li24	
^{140}Pm^m−u		−83532	30	−83503	14	1.0	1	22	22 ^{140}Pm^m	GS2	1.0	05Li24	
^{140}Sm−u		−81018	30	−81005	13	0.4	R			GS2	1.0	05Li24	
^{140}Gd−u		−66326	30				2			GS2	1.0	05Li24	
^{140}Te−^{133}Cs$_{1.053}$		38822	67	39046	15	3.3	B			CP1	1.0	13Va12	
		39042	18			0.2	1	74	74 ^{140}Te	CP2	1.0	18Or.A	
^{140}I−^{133}Cs$_{1.053}$		31275	13	31240	50	−2.3	U			CP1	1.0	13Va12	
		31245	73			−0.0	2			JY1	1.0	24Be25	*
		31244	68			0.0	2			JY2	1.0	24Be25	*
^{140}Xe−^{133}Cs$_{1.053}$		21204.9	2.5				2			MA8	1.0	09Ne11	
^{140}Cs−^{133}Cs$_{1.053}$		16837	14	16843	9	0.4	–			MA1	1.0	99Am05	
		16857	14			−1.0	–			MA4	1.0	99Am05	
	ave.	16847	10			−0.4	1	80	80 ^{140}Cs			average	
^{140}Ba−^{133}Cs$_{1.053}$		10150	14	10167	9	1.2	1	40	40 ^{140}Ba	MA1	1.0	99Am05	
^{140}Ce O−^{133}Cs$_{1.173}$		11270.3	2.5	11267.3	1.4	−1.2	1	31	31 ^{140}Ce	MA8	1.0	19Hu15	
^{140}Nd O−^{133}Cs$_{1.173}$		15365.6	3.5				2			MA8	1.0	19Hu15	
^{140}Pm^m−^{133}Cs$_{1.053}$		16064	16	16056	14	−0.5	1	78	78 ^{140}Pm^m	MA5	1.0	00Be42	
^{140}Sm−^{133}Cs$_{1.053}$		18557	15	18554	13	−0.2	2			MA5	1.0	00Be42	
^{140}Xe−^{136}Xe$_{1.029}$		17134	11	17122.1	2.5	−1.1	U			CP1	1.0	12Va02	
C_{11} H_9−^{140}Ce		164956	40	164977.5	1.4	0.2	U			M17	2.5	66Be10	
^{140}Ce−^{139}La		−1029	80	−915.1	1.4	0.4	U			R05	4.0	65De13	
C_{11} H_{10}−^{140}Ce		172765	40	172802.5	1.4	0.4	U			M17	2.5	66Be10	
^{140}Ce−^{138}Ce		−497	83	−546.4	1.5	−0.1	U			R05	4.0	65De13	
		−543	8			−0.2	U			M17	2.5	66Be10	
^{139}Cs−^{140}Cs$_{.883}$ ^{131}Cs$_{.118}$		−2280	40	−2275	8	0.0	U			P23	2.5	82Au01	
^{139}Cs−^{140}Cs$_{.869}$ ^{132}Cs$_{.132}$		−2210	40	−2241	8	−0.3	U			P23	2.5	82Au01	
^{138}Ce(t,p)^{140}Ce		8184	15	8169.8	1.4	−0.9	U			LAl		72Mu09	
^{140}Ce(p,t)^{138}Ce		−8167	20	−8169.8	1.4	−0.1	U			Brk		77Sh06	
^{139}La(n,γ)^{140}La		5161.1	1.0	5160.997	0.019	−0.1	U					70Ju04	
		5160	1			1.0	U					72Fu10	
		5160.97	0.05			0.5	–			MMn		90Is09	Z
		5161.00	0.10			−0.0	o			Bdn		06Fi.A	
		5161.001	0.021			−0.2	–			Bdn		19Hu07	
^{139}La(d,p)^{140}La		2938	3	2936.430	0.019	−0.5	U			Tal		67Ke02	
^{139}La(n,γ)^{140}La	ave.	5160.996	0.019	5160.997	0.019	0.0	1	100	94 ^{140}La			average	
^{140}Ho(p)^{139}Dy		1093.9	10.				3			Orp		99Ry04	*
^{140}Xe(β^-)^{140}Cs		4060	60	4063	9	0.1	U			Trs		78Wo15	
^{140}Cs(β^-)^{140}Ba		6100	100	6219	10	1.2	U			Trs		78Wo15	
		6235	25			−0.7	o			Gsn		80Bl.A	
		6220	15			−0.1	o			Gsn		81De25	
		6212	20			0.3	–			Gsn		92Pr04	
		6199	25			0.8	–			Ida		93Gr17	
	ave.	6207	16			0.7	1	41	21 ^{140}Ba			average	
^{140}Ba(β^-)^{140}La		1056	20	1044	8	−0.6	–					49Be36	*
		1055	30			−0.4	–					59Bo61	*
		1060	30			−0.5	–					65Bu07	*
	ave.	1057	15			−0.9	1	32	32 ^{140}Ba			average	
^{140}La(β^-)^{140}Ce		3760.2	2.0	3762.7	1.3	1.3	1	43	38 ^{140}Ce			72Na04	*
^{140}Pr(β^+)^{140}Ce		3388	6				2					68Ab17	
^{140}Nd(ε)^{140}Pr		160	60	430	7	4.5	B					72Ba91	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{140}Pm(β^+)^{140}Nd	6080	100	6045	24	−0.3	U					75Ke09	
	6090	40			−1.1	3			IRS		83Al06	
	6020	30			0.8	3			Dbn		95Ve08	
^{140}Pm^m(β^+)^{140}Nd	6484	70	6474	14	−0.1	U					75Ke09	*
^{140}Sm(ε)^{140}Pm	3400	300	2756	27	−2.1	U					87De04	
^{140}Eu(β^+)^{140}Sm	8400	400	8660#	100#	0.7	U			LBL		91Fi03	*
	8470	50			3.8	D			Dbn		95Ve08	*
^{140}Gd(β^+)^{140}Eu	4800	400	5010#	110#	0.5	U			LBL		91Fi03	
^{140}Tb(β^+)^{140}Gd	11300	800				3			LBL		91Fi03	*
*^{140}I−^{133}Cs$_{1.053}$	D_M=31300.35(0.68) μu for mixture gs+m+n at 50(50)# and										Nub269	**
*^{140}I−^{133}Cs$_{1.053}$	100(100)# keV; $M-A$=–63582.61(0.63) keV										Nub269	**
*^{140}I−^{133}Cs$_{1.053}$	D_M=31298.3(1.6) μu for mixture gs+m+n at 50(50)# and										Nub269	**
*^{140}I−^{133}Cs$_{1.053}$	100(100)# keV; $M-A$=–63584.5(1.5) keV										Nub269	**
*^{140}Ho(p)^{139}Dy	Trends in Mass Surface TMS suggest ^{140}Ho 300 keV less bound										GAu	**
*^{140}Ho(p)^{139}Dy	region of large deformation; deviation from TMS justified										FKo249	**
*^{140}Ba(β^-)^{140}La	E_{β^-}=1022(20), 480(20) to 2$^-$ level at 29.9641, 0$^-$ at 581.07 keV										FKo25a	**
*^{140}Ba(β^-)^{140}La	E_{β^-}=1020(30), 480(30) to 2$^-$ level at 29.9641, 0$^-$ at 581.07 keV										FKo25a	**
*^{140}Ba(β^-)^{140}La	E_{β^-}=1030(30), 480(30) to 2$^-$ level at 29.9641, 0$^-$ at 581.07 keV										FKo25a	**
*^{140}La(β^-)^{140}Ce	E_{β^-}=2164(2) to 2$^+$ level at 1596.237 keV										Ens077	**
*^{140}Pm^m(β^+)^{140}Nd	E_{β^+}=3240(70) to 7$^-$ level at 2221.4 keV										Ens077	**
*^{140}Eu(β^+)^{140}Sm	From p$^+$. May be lower limit										91Fi03	**
*^{140}Eu(β^+)^{140}Sm	Trends in Mass Surface TMS suggest ^{140}Eu 190 keV less bound										HWJ266	**
*^{140}Tb(β^+)^{140}Gd	Lower limit										91Fi03	**
^{141}I−u	−64316	419	−64319	5	−0.0	o			GT1	1.5	04Ma.A	
	−64549	120			0.8	o			GT2	2.5	08Kn.A	
	−64736	137			1.2	B			GT2	2.5	08Su19	
	−64445	186			0.3	U			GT3	2.5	16Kn03	
^{141}Xe−u	−73092	126	−73213	3	−0.4	o			GT2	2.5	08Kn.A	
	−73560	136			1.0	U			GT2	2.5	08Su19	
^{141}Ba−u	−85603.5	7.5	−85596	6	0.9	1	58	58 ^{141}Ba	CP1	1.0	06Sa56	
C$_{11}$ H$_9$−^{141}Pr	162852	41	162766.7	1.5	−0.5	U			R05	4.0	65De13	
C$_{10}$ N H$_7$−^{141}Pr	150229	37	150190.7	1.5	−0.3	U			R05	4.0	65De13	
C$_9$ ^{13}C N H$_6$−^{141}Pr	145722	65	145720.5	1.5	−0.0	U			R05	4.0	65De13	
^{141}Pr−u	−92374	30	−92341.4	1.5	1.1	U			GS2	1.0	05Li24	
^{141}Nd−u	−90401	30	−90384	3	0.6	U			GS2	1.0	05Li24	
	−90365	30			−0.6	U			GS2	1.0	05Li24	*
^{141}Sm−u	−81496	62	−81522	8	−0.4	U			GS2	1.0	05Li24	*
^{141}Eu−u	−75048	42	−75069	14	−0.5	U			GS2	1.0	05Li24	*
^{141}Gd−u	−67881	30	−67874	21	0.2	2			GS2	1.0	05Li24	
	−67867	30			−0.2	2			GS2	1.0	05Li24	*
^{141}Tb−u	−58552	113				2			GS2	1.0	05Li24	*
^{141}I−^{133}Cs$_{1.060}$	35887	17	35902	5	0.9	2			CP1	1.0	13Va12	
	35903.6	4.8			−0.3	2			JY1	1.0	24Be25	
^{141}Xe−^{133}Cs$_{1.060}$	27008.1	3.1				2			MA8	1.0	09Ne11	
^{141}Cs−^{133}Cs$_{1.060}$	20269	16	20266	10	−0.2	1	38	38 ^{141}Cs	MA4	1.0	99Am05	
^{141}Ba−^{133}Cs$_{1.060}$	14625	15	14625	6	−0.0	–			MA1	1.0	99Am05	
	14631	16			−0.4	–			MA4	1.0	99Am05	
ave.	14628	11			−0.3	1	27	27 ^{141}Ba			average	
^{141}Pm−^{133}Cs$_{1.060}$	13776	15				2			MA5	1.0	00Be42	
^{141}Sm−^{133}Cs$_{1.060}$	18694	14	18698	8	0.3	1	30	30 ^{141}Sm	MA5	1.0	00Be42	
^{141}Sm^m−^{133}Cs$_{1.060}$	18878	14	18887	8	0.7	1	30	30 ^{141}Sm^m	MA5	1.0	00Be42	
^{141}Eu−^{133}Cs$_{1.060}$	25164	15	25152	14	−0.8	1	81	81 ^{141}Eu	MA5	1.0	00Be42	*
^{141}Xe−^{136}Xe$_{1.037}$	23003	10	23006	3	0.3	U			CP1	1.0	12Va02	
^{141}Cs−^{136}Xe$_{1.037}$	16277	22	16264	10	−0.6	1	20	20 ^{141}Cs	CP1	1.0	12Va02	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{139}Cs−^{141}Cs$_{.789}$ ^{131}Cs$_{.212}$		−3190	40	−3271	8	−0.8	U			P23	2.5	82Au01	
^{140}Cs−^{141}Cs$_{.894}$ ^{131}Cs$_{.107}$		−970	40	−1045	12	−0.8	U			P23	2.5	82Au01	
^{139}Cs−^{141}Cs$_{.767}$ ^{132}Cs$_{.234}$		−3210	40	−3183	8	0.3	U			P23	2.5	82Au01	
^{141}Cs(β^-n)^{140}Ba		735	30	720	12	−0.5	1	16	9 ^{141}Cs			84Kr.B	
^{140}Ce(n,γ)^{141}Ce		5428.6	0.6	5428.15	0.10	−0.7	U			BNn		70Ge03	Z
		5428.01	0.20			0.7	–			Ptn		80Ba.A	Z
		5428.19	0.12			−0.3	–			Bdn		06Fi.A	
^{140}Ce(d,p)^{141}Ce		3210	10	3203.58	0.10	−0.6	U			MIT		64Sp12	
		3202	15			0.1	U			Hei		67Wi08	
^{140}Ce(n,γ)^{141}Ce	ave.	5428.14	0.10	5428.15	0.10	0.1	1	100	72 ^{141}Ce			average	
^{141}Pr(γ,n)^{140}Pr		−9361	23	−9400	6	−1.7	U			Phi		60Ge01	
^{141}Ho(p)^{140}Dy		1177.4	8.	1177	6	−0.0	3			Arp		98Da03	
		1177	10			0.0	3			ORp		08Ka16	*
^{141}Ho^m(p)^{140}Dy		1239	20	1243	7	0.2	o			ORp		99Ry04	
		1244	9			−0.1	U			Arp		01Se03	
		1240	10			0.3	U			ORp		08Ka16	*
^{141}Xe(β^-)^{141}Cs		6150	90	6280	10	1.4	U			Trs		78Wo15	
^{141}Cs(β^-)^{141}Ba		5200	80	5255	10	0.7	U			Trs		78Wo15	
		5264	15			−0.6	o			Gsn		80Bl.A	*
		5252	15			0.2	o			Gsn		81De25	
		5242	15			0.9	1	41	33 ^{141}Cs	Gsn		92Pr04	
^{141}Ba(β^-)^{141}La		3010	60	3198	7	3.1	B			Trs		78Wo15	
		3208	35			−0.3	U			Gsn		81De25	
		3217	20			−1.0	1	11	7 ^{141}Ba	McG		84He.A	
^{141}La(β^-)^{141}Ce		2430	30	2501	4	2.4	U					51Du19	
		2502	4			−0.2	1	96	96 ^{141}La	McG		84He.A	
^{141}Ce(β^-)^{141}Pr		584	3	583.9	1.2	−0.0	–					50Fr58	*
		585	4			−0.3	–					52Ko27	*
		576.4	2.0			3.7	B					55Jo02	*
		581.4	2.0			1.2	–					68Be06	*
		582.2	2.6			0.6	–					79Ha09	*
	ave.	582.5	1.3			1.0	1	78	51 ^{141}Pr			average	
^{141}Nd(β^+)^{141}Pr		1816	8	1823.0	2.8	0.9	2					73Bu21	
		1824	3			−0.3	2					76Ga.A	*
^{141}Pm(β^+)^{141}Nd		3730	60	3670	14	−1.0	U					70Ch29	*
		3640	70			0.4	U					75Ke09	
^{141}Sm(β^+)^{141}Pm		4580	50	4585	16	0.1	U					77Ke03	*
		4463	60			2.0	U			IRS		83Al06	*
		4524	80			0.8	U			IRS		93Al03	*
^{141}Sm^m(IT)^{141}Sm		175.8	0.3	175.80	0.30	−0.0	1	100	70 ^{141}Sm^m			76Ke06	
^{141}Eu(β^+)^{141}Sm		6030	100	6011	14	−0.2	U					77De25	*
		5950	40			1.5	–			IRS		83Al06	*
		6035	60			−0.4	U					85Af.A	
		5550	100			4.6	B			IRS		93Al03	
		5980	40			0.8	–			Dbn		95Ve08	*
	ave.	5965	28			1.6	1	24	19 ^{141}Eu			average	
^{141}Ho^m(IT)^{141}Ho		66	2				4			Arp		01Se03	
*^{141}Nd−u	$M-A$=–83418(28) keV for ^{141}Nd^m at 756.51 keV											Nub269	**
*^{141}Sm−u	$M-A$=–75825(28) keV for mixture gs+m at 175.8 keV											Nub269	**
*^{141}Eu−u	$M-A$=–69858(28) keV for mixture gs+m at 96.45 keV											Nub269	**
*^{141}Gd−u	$M-A$=–62840(28) keV for ^{141}Gd^m at 377.76 keV											Nub269	**
*^{141}Tb−u	$M-A$=–54541(34) keV for mixture gs+m at 0#200 keV											Nub269	**
*^{141}Eu−^{133}Cs$_{1.060}$	Slight (< 10%) isomeric contamination cannot be excluded											00Be42	**
*^{141}Ho(p)^{140}Dy	E_p=968(10) to the 2^+ level at 202.1 keV;											Nub25a	**
*^{141}Ho(p)^{140}Dy	Ep rebuilt from the energy difference 201(6) keV to the peak at 1169(8)											Nub25a	**
*^{141}Ho^m(p)^{140}Dy	E_p=1030(10) to the 2^+ level at 202.1 keV using E_p=1235(9) as caibrat											Nub25a	**
*^{141}Cs(β^-)^{141}Ba	E_{β^-}=5215(15) to $(5/2)^-$ level at 48.53 keV											Ens141	**
*^{141}Ce(β^-)^{141}Pr	E_{β^-}=442(3) 444(4) 432(2) 436(2) 436.7(2.6) respectively, to $7/2^+$ level at 145.4434											Ens141	**
*^{141}Nd(β^+)^{141}Pr	Was erroneously quoted 77Ga.A in the 1993 tables											GAu	**
*^{141}Pm(β^+)^{141}Nd	Original error 40 increased due to lack of information on calibration											GAu	**
*^{141}Sm(β^+)^{141}Pm	E_{β^+}=3180(50), 3100(50) to $3/2^+$ level at 403.82, $(1/2)^+$ at 438.69 keV											Ens141	**
*	and E_{β^+}=1670(70), 1600(70) from ^{141}Sm^m											77Ke03	**
*	at 175.9 to $11/2^-$ at 2091.71, $(9/2,11/2,13/2)^-$ at 2119.05											Ens141	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{141}$Sm(β^+)^{141}Pm		E_{β^+}=3020(60) 32% to 3/2$^+$ level at 403.82, 31% to (1/2)$^+$ 438.29 keV										Ens141	**
$*^{141}$Sm(β^+)^{141}Pm		Q_{β^+}=4700(80) from ^{141}Sm^m at 175.8 keV										Nub269	**
$*^{141}$Eu(β^+)^{141}Sm		E_{β^+}=4620(110) to (5/2)$^+$ level at 395.56 keV, and other E_{β^+} (not given)										Ens141	**
$*^{141}$Eu(β^+)^{141}Sm		E_{β^+}=4925(40) to 3/2$^+$ level at 1.58 keV										Ens141	**
$*^{141}$Eu(β^+)^{141}Sm		E_{β^+}=4960(40) to 3/2$^+$ level at 1.58 keV										Ens141	**
^{142}I−u		−58798	268	−58833	5	−0.1	U			GT1	1.5	04Ma.A	
^{142}I−^{133}Cs$_{1.068}$		42143.9	5.3				2			CP2	1.0	20Or02	
		42101	64	42144	5	0.7	U			JY1	1.0	24Be25	
^{142}Xe−u		−70247	111	−70026.9	2.9	0.8	U			GT2	2.5	08Kn.A	
^{142}Xe−^{133}Cs$_{1.068}$		30950.4	2.9				2			MA8	1.0	09Ne11	
^{142}Cs−^{133}Cs$_{1.068}$		25270	16	25272	7	0.1	–			MA4	1.0	99Am05	
		25304	23			−1.4	–			CP1	1.0	13Va12	
	ave.	25281	13			−0.7	1	27	27 ^{142}Cs			average	
^{142}Ba−^{133}Cs$_{1.068}$		17410	15	17413	6	0.2	–			MA1	1.0	99Am05	
		17420	16			−0.4	–			MA4	1.0	99Am05	
	ave.	17415	11			−0.1	1	30	30 ^{142}Ba			average	
^{142}Ba−u		−83576.8	9.1	−83564	6	1.4	1	43	43 ^{142}Ba	CP1	1.0	06Sa56	
C_{11} H_{10}−^{142}Ce		169111	15	169001.6	2.6	−1.8	U			R05	4.0	65De13	
		168955	40			0.5	U			M17	2.5	66Be10	
C_{10} ^{13}C H_9−^{142}Ce		164528	82	164531.4	2.6	0.0	U			R05	4.0	65De13	
C_{10} N H_8−^{142}Ce		156558	42	156425.5	2.6	−0.8	U			R05	4.0	65De13	
C_{11} H_{10}−^{142}Nd		170509	36	170523.2	1.1	0.1	U			R05	4.0	65De13	
C_{10} N H_8−^{142}Nd		157870	43	157947.1	1.1	0.4	U			R05	4.0	65De13	
C_{10} O H_6−^{142}Nd		134076	36	134137.7	1.1	0.4	U			R05	4.0	65De13	
C_{10} ^{13}C H_9−^{142}Nd		166021	32	166053.0	1.1	0.2	U			R05	4.0	65De13	*
^{142}Pm−u		−87136	30	−87109	25	0.9	–			GS2	1.0	05Li24	
	ave.	−87124	27			0.6	1	89	89 ^{142}Pm			average	
^{142}Sm−^{133}Cs$_{1.068}$		16173	14	16185.3	1.9	0.9	U			MA5	1.0	00Be42	
^{142}Eu^m−^{133}Cs$_{1.068}$		24909	15	24910	13	0.1	2			MA5	1.0	00Be42	
^{142}Eu^m−u		−76063	30	−76067	13	−0.1	R			GS2	1.0	05Li24	
^{142}Gd−u		−71884	30				2			GS2	1.0	05Li24	
^{142}Cs−^{136}Xe$_{1.044}$		21171	11	21163	7	−0.7	1	39	39 ^{142}Cs	CP1	1.0	12Va02	
^{142}Ce−C_{11} H_9		−161176	40	−161176.5	2.6	−0.0	U			M17	2.5	66Be10	
^{142}Nd−C_{11} H_9		−162665	30	−162698.2	1.1	−0.4	U			M17	2.5	66Be10	
^{142}Ce−^{140}Ce		3818	3	3801.0	2.6	−2.3	U			M17	2.5	66Be10	
^{142}Ce−^{138}Ce		3644	35	3254.5	2.6	−2.8	B			R05	4.0	65De13	
^{139}Cs−^{142}Cs$_{.685}$ ^{132}Cs$_{.316}$		−4840	40	−4855	5	−0.2	U			P23	2.5	82Au01	
^{140}Cs−^{142}Cs$_{.789}$ ^{132}Cs$_{.212}$		−2950	40	−2934	10	0.2	U			P23	2.5	82Au01	
^{141}Cs−^{142}Cs$_{.794}$ ^{137}Cs$_{.206}$		−580	40	−657	10	−0.8	U			P23	2.5	82Au01	
^{138}Cs^x−^{142}Cs$_{.194}$ ^{137}Cs$_{.806}$		550	40	590	25	0.4	U			P23	2.5	82Au01	
^{140}Cs−^{142}Cs$_{.329}$ ^{139}Cs$_{.671}$		260	40	301	9	0.4	U			P23	2.5	82Au01	
^{141}Cs−^{142}Cs$_{.662}$ ^{139}Cs$_{.338}$		−410	40	−517	10	−1.1	U			P23	2.5	82Au01	
^{141}Cs−^{142}Cs$_{.496}$ ^{140}Cs$_{.504}$		−640	40	−667	10	−0.3	U			P23	2.5	82Au01	
		−663	19			−0.1	U			P33	2.5	86Au02	
^{142}Ce(α)^{138}Ba		1545	200	1302.6	2.4	−1.2	U					57Ri43	
^{140}Ce(t,p)^{142}Ce		4112	5	4120.3	2.4	1.7	1	24	20 ^{142}Ce	LAl		72Mu09	
^{142}Ce(p,t)^{140}Ce		−4170	20	−4120.3	2.4	2.5	U			Osa		70Ya05	
^{142}Nd(p,t)^{140}Nd		−9150	20	−9355	3	−10.3	B			Osa		71Ya10	*
^{142}Ce(γ,n)^{141}Ce		−7240	70	−7173.9	2.4	0.9	U			Phi		60Ge01	
^{142}Ce(d,t)^{141}Ce		−909	15	−916.7	2.4	−0.5	U			Mtr		72Le17	
^{141}Pr(n,γ)^{142}Pr		5843.14	0.10	5843.15	0.08	0.1	–			MMn		81Ke11	Z
		5843.16	0.12			−0.1	–			Bdn		06Fi.A	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{141}Pr(d,p)^{142}Pr		3626	10	3618.59	0.08	−0.7	U			MIT		64Sp12	
^{141}Pr(n,γ)^{142}Pr	ave.	5843.15	0.08	5843.15	0.08	0.1	1	100	51 ^{142}Pr			average	
^{142}Xe(β^-)^{142}Cs		5040	100	5289	7	2.5	U			Trs		78Wo15	
^{142}Cs(β^-)^{142}Ba		7230	70	7321	7	1.3	U			Trs		78Wo15	
		7329	20			−0.4	o			Gsn		81De25	
		7280	40			1.0	U			Bwg		87Gr.A	
		7315	15			0.4	–			Gsn		92Pr04	
		7308	11			1.1	–					25Al22	
	ave.	7310	9			1.1	1	56	34 ^{142}Cs			average	
^{142}Ba(β^-)^{142}La		2200	25	2186	8	−0.6	1	11	6 ^{142}La			83Ch39	
		2216	5			−6.1	C			McG		84He.A	
^{142}La(β^-)^{142}Ce		4517	25	4509	6	−0.3	U					65Pr03	
		4510	6			−0.1	1	95	94 ^{142}La	McG		84He.A	
^{142}Pr(β^-)^{142}Nd		2164	2	2164.3	1.3	0.1	–					66Be12	
		2158	3			2.1	–					75Ra09	
	ave.	2162.2	1.7			1.3	1	65	49 ^{142}Pr			average	
^{142}Pm(β^+)^{142}Nd		4800	80	4810	24	0.1	R					60Ma.A	
		4880	80			−0.9	R			IRS		83Al06	
		4880	160			−0.4	U			LBL		91Fi03	
^{142}Sm(β^+)^{142}Pm		2050	70	2158	24	1.5	1	11	11 ^{142}Pm			60Ma.A	
		2100	400			0.1	U			LBL		91Fi03	
^{142}Eu(β^+)^{142}Sm		8000	300	7670	30	−1.1	U					75Ke08	
		7400	100			2.7	U					82Gr.A	
		7000	300			2.2	U			LBL		91Fi03	
		7673	30				2			Dbn		94Po26	
^{142}Eu^m(β^+)^{142}Sm		8150	100	8127	13	−0.2	U					75Ke08	*
		8174	50			−0.9	U			IRS		83Al06	*
		7480	100			6.5	B			IRS		93Al03	*
		8150	60			−0.4	U			Dbn		94Po26	*
^{142}Gd(β^+)^{142}Eu		4200	300	4350	40	0.5	U			LBL		91Fi03	
^{142}Tb(β^+)^{142}Gd		10400	700				3			LBL		91Fi03	
^{142}Dy(β^+)^{142}Tb		7100	200	6580#	200#	−2.6	D			LBL		91Fi03	*
*C_{10} ^{13}C H_9−^{142}Nd	Original 1002055(32) is certainly a typo; rebuilt from M=141.907760(36)u											WgM127	**
*^{142}Nd(p,t)^{140}Nd	Disagrees strongly with ^{140}Nd−u											AHW	**
*^{142}Eu^m(β^+)^{142}Sm	E_{β^+}=4760(100) 4782(50) respectively, to 7$^-$ level at 2372.1 keV											Ens118	**
*^{142}Eu^m(β^+)^{142}Sm	Measured half-life 73.4(0.5) s corresponds to ^{142}Eu^m											GAu	**
*^{142}Eu^m(β^+)^{142}Sm	E_{β^+}=4756(60) to 7$^-$ level at 2372.1 keV											Ens118	**
*^{142}Dy(β^+)^{142}Tb	Trends in Mass Surface TMS suggest ^{142}Dy 520 keV more bound											GAu	**
^{143}I−u		−53849	495	−54530#	220#	−0.5	D			GT3	2.5	16Kn03	*
^{143}Xe−u		−64649	290	−64630	5	0.0	o			GT1	1.5	04Ma.A	
		−64858	108			0.8	o			GT2	2.5	08Kn.A	
		−64684	133			0.2	U			GT2	2.5	08Su19	
^{143}Xe−^{133}$Cs_{1.075}$		37008.7	5.0				2			MA8	1.0	09Ne11	
^{143}Cs−u		−72771	117	−72652	8	0.4	U			GT2	2.5	08Kn.A	
^{143}Cs−^{133}$Cs_{1.075}$		28985.6	8.5	28987	8	0.2	1	91	91 ^{143}Cs	CP1	1.0	13Va12	
^{143}Ba−^{133}$Cs_{1.075}$		22268	16	22272	6	0.3	1	14	14 ^{143}Ba	MA1	1.0	99Am05	
^{143}Ba−u		−79375.0	8.5	−79367	6	0.9	–			CP1	1.0	06Sa56	
		−79352.1	10.1			−1.5	–			GR1	1.0	25Sp03	
	ave.	−79366	7			−0.2	1	82	82 ^{143}Ba			average	
^{143}La−u		−83918.1	8.7	−83921	8	−0.3	1	82	82 ^{143}La	CP1	1.0	06Sa56	
C_{10} N H_9−^{143}Nd		163719	31	163681.2	1.1	−0.3	U			R05	4.0	65De13	
C_{10} O H_7−^{143}Nd		139814	42	139871.7	1.1	0.3	U			R05	4.0	65De13	
^{143}Pm−^{133}$Cs_{1.075}$		12567	15	12575	3	0.6	U			MA5	1.0	00Be42	
^{143}Sm−^{133}$Cs_{1.075}$		16268	15	16272.2	2.8	0.3	U			MA5	1.0	00Be42	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{143}Sm−u		−85347	30	−85367.0	2.8	−0.7	U			GS2	1.0	05Li24	*
^{143}Eu−^{133}Cs$_{1.075}$		21947	14	21938	12	−0.7	2			MA5	1.0	00Be42	
^{143}Eu−u		−79706	30	−79701	12	0.2	R			GS2	1.0	05Li24	
^{143}Gd−u		−73012	56	−73250	220	−4.2	C			GS2	1.0	05Li24	*
^{143}Tb−u		−64906	46	−64890	30	0.4	U			GS2	1.0	05Li24	*
^{143}Tb−^{85}Rb$_{1.682}$		83480	32				2			SH1	1.0	07Ra37	*
^{143}Dy−^{85}Rb$_{1.682}$		92364	14				2			SH1	1.0	07Ra37	*
^{143}Nd ^{35}Cl−^{141}Pr ^{37}Cl		5116	4	5109.7	1.4	−0.6	U			H21	2.5	70Ma05	
^{143}Nd−C$_{11}$ H$_{10}$		−168422	30	−168432.2	1.1	−0.1	U			M17	2.5	66Be10	
^{143}Nd−^{142}Nd		2322	46	2090.99	0.07	−1.3	U			R05	4.0	65De13	
		2084	2			1.4	U			M17	2.5	66Be10	
^{143}Nd−C$_{11}$ H$_9$		−160594	30	−160607.2	1.1	−0.2	U			M17	2.5	66Be10	
^{141}Cs−^{143}Cs$_{.493}$ ^{139}Cs$_{.507}$		−230	40	−198	10	0.3	U			P23	2.5	82Au01	
		−115	22			−1.5	U			P33	2.5	86Au02	
^{142}Cs−^{143}Cs$_{.497}$ ^{141}Cs$_{.504}$		647	15	652	9	0.1	U			P33	2.5	86Au02	
^{143}Nd(n,α)^{140}Ce		9699	15	9717.3	1.4	1.2	U			ILL		75Em.A	
^{143}Nd(p,t)^{141}Nd		−7450	20	−7472	3	−1.1	U			Osa		71Ya10	
^{142}Ce(n,γ)^{143}Ce		5145.9	0.5	5144.80	0.09	−2.2	U					76Ge02	
		5144.78	0.15			0.1	–			Ptn		80Ba.A	Z
		5144.81	0.12			−0.1	–			Bdn		06Fi.A	
^{142}Ce(d,p)^{143}Ce		2945	15	2920.23	0.09	−1.7	U			Mtr		72Le17	
^{142}Ce(n,γ)^{143}Ce	ave.	5144.80	0.09	5144.80	0.09	0.0	1	100	79 ^{142}Ce			average	
^{142}Nd(n,γ)^{143}Nd		6123.62	0.08	6123.57	0.07	−0.6	–			MMn		82Is05	Z
		6123.41	0.14			1.2	–			Bdn		06Fi.A	
^{142}Nd(d,p)^{143}Nd		3916	15	3899.01	0.07	−1.1	U			Kop		67Ch16	
		3902	15			−0.2	U			Tal		67Ne04	
		3902	15			−0.2	U			Hei		67Wi08	
^{142}Nd(n,γ)^{143}Nd	ave.	6123.57	0.07	6123.57	0.07	0.1	1	100	83 ^{142}Nd			average	
^{142}Nd(^{3}He,d)^{143}Pm		−1099	25	−1193.8	2.7	−3.8	B			Oak		71Wi04	
		−1195	5			0.2	1	29	29 ^{143}Pm	McM		80St10	*
^{143}Cs(β^-)^{143}Ba		6250	90	6255	9	0.1	o			Gsn		81De25	*
		6240	70			0.2	U			Bwg		87Gr.A	
		6270	25			−0.6	1	13	9 ^{143}Cs	Gsn		92Pr04	
^{143}Ba(β^-)^{143}La		4240	50	4242	9	0.0	U					79Sc11	
		4259	40			−0.4	U			Gsn		81De25	
		4210	70			0.5	U			Bwg		87Gr.A	
^{143}La(β^-)^{143}Ce		3425	17	3436	8	0.6	1	20	18 ^{143}La			84Is09	*
^{143}Ce(β^-)^{143}Pr		1460.6	2.	1462.0	1.9	0.7	1	87	78 ^{143}Ce			77Ra18	*
^{143}Pr(β^-)^{143}Nd		932	2	934.2	1.4	1.1	–					49Fe18	
		935	2			−0.4	–					76Ra33	
	ave.	933.5	1.4			0.5	1	93	91 ^{143}Pr			average	
^{143}Pm(β^+)^{143}Nd		1000	70	1041.6	2.7	0.6	U					67Va01	*
^{143}Sm(β^+)^{143}Pm		3492	30	3443	4	−1.6	U					66Be21	
		3437	30			0.2	U			IRS		83Al06	
		3500	60			−0.9	U			IRS		93Al03	
		3461	40			−0.4	U			Dbn		94Po26	
^{143}Eu(β^+)^{143}Sm		5100	50	5278	11	3.6	B					74Ch21	
		5160	60			2.0	o			IRS		83Ve.A	
		5240	70			0.5	o			IRS		83Al06	
		5250	80			0.3	U			IRS		93Al03	
		5236	30			1.4	R			Dbn		94Po26	
^{143}Gd(β^+)^{143}Eu		6010	200				3			IRS		93Al03	*
*^{143}I−u	Trends in Mass Surface TMS suggest ^{143}I 630 keV more bound											GAu	**
*^{143}Sm−u	$M-A$=–78746(28) keV for ^{143}Sm^m at 753.99 keV											Nub269	**
*^{143}Gd−u	$M-A$=–67934(28) keV for mixture gs+m at 152.6 keV											Nub269	**
*^{143}Tb−u	$M-A$=–60434(32) keV for mixture gs+m at 50#50 keV											Nub269	**
*	outweighed by next item before correcting for isomeric mixture											GAu	**
*^{143}Tb−^{85}Rb$_{1.682}$	D_M=83506.7(8.3) μu for mixture gs+m at 50#50 keV;											Nub269	**
*^{143}Tb−^{85}Rb$_{1.682}$	$M-A$=–60419.5(7.8) keV											Nub269	**
*^{143}Dy−^{85}Rb$_{1.682}$	D_M=92354(17) and D_M=92705(14) for ^{143}Dy^m at 310.7 keV											Nub269	**
*^{142}Nd(^{3}He,d)^{143}Pm	Based on ^{146}Nd(^{3}He,d)^{147}Pm Q =–87.6(0.9) keV											AHW	**
*^{143}Cs(β^-)^{143}Ba	E_{β^-}=6070(50) and 5847(100) to 3/2$^-$ level at 228.83 keV											Ens123	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{143}$La(β^-)^{143}Ce		E_{β^-}=3419(17) 64% to 3/2$^-$ ground state, 29% to 7/2$^-$ level at 18.9 keV										Ens123	**
$*^{143}$Ce(β^-)^{143}Pr		E_{β^-}=1110(2) to 3/2$^+$ level at 350.622 keV										Ens123	**
$*^{143}$Pm(β^+)^{143}Nd		pK=0.806(0.023) to 3/2$^-$ level at 742.05 keV, and p$^+$ <1×10^{-6}										Ens123	**
$*^{143}$Gd(β^+)^{143}Eu		Q_{β^+}=6160(200) from ^{143}Gd^m at 152.6 keV										Nub269	**
^{144}Xe−^{133}Cs$_{1.083}$		41340.6	5.7				2			MA8	1.0	09Ne11	
^{144}Cs−^{133}Cs$_{1.083}$		34488	33	34469	22	−0.6	1	43	43 ^{144}Cs	CP1	1.0	13Va12	
^{144}Ba−^{133}Cs$_{1.083}$		25347	15	25347	6	0.0	1	17	17 ^{144}Ba	MA1	1.0	99Am05	
^{144}Ba−u		−77045.3	9.1	−77048	6	−0.3	–			CP1	1.0	06Sa56	
		−77053.7	10.3			0.5	–			GR1	1.0	25Sp03	
	ave.	−77049	7			0.1	1	81	81 ^{144}Ba			average	
^{144}La−u		−80337.1	19.3	−80354	14	−0.9	2			CP1	1.0	06Sa56	
		−80373	20			0.9	2			GS3	1.0	12Ch19	
^{144}Ce−u		−86387.0	19.3	−86348.9	3.0	2.0	U			GR1	1.0	25Sp03	
C_{10} O H_8−^{144}Nd		147408	28	147423.8	1.1	0.1	U			R05	4.0	65De13	
		147384	29			0.3	U			R05	4.0	65De13	
C_9 ^{13}C N H_9−^{144}Nd		166777	28	166763.0	1.1	−0.1	U			R05	4.0	65De13	
C_{10} H_8 O−^{144}Sm		145450	50	145510.4	1.4	0.3	U			R04	4.0	64De15	
C_9 ^{13}C H_9 N−^{144}Sm		164955	46	164849.6	1.4	−0.6	U			R04	4.0	64De15	
^{144}Eu−^{133}Cs$_{1.083}$		21223	17	21214	12	−0.5	1	46	46 ^{144}Eu	MA5	1.0	00Be42	
^{144}Eu−u		−81117	30	−81181	12	−2.1	1	15	15 ^{144}Eu	GS2	1.0	05Li24	
^{144}Gd−u		−77037	30				2			GS2	1.0	05Li24	
^{144}Tb^m−u		−66528	30				2			GS2	1.0	05Li24	*
^{144}Dy−u		−60746	33	−60730	8	0.5	U			GS2	1.0	05Li24	
^{144}Dy−^{85}Rb$_{1.694}$		88697.7	7.7				2			SH1	1.0	07Ra37	
^{144}Ho−^{85}Rb$_{1.694}$		101537.9	9.1				2			SH1	1.0	07Ra37	
^{144}Nd ^{35}Cl−^{142}Nd ^{37}Cl		5329	3	5314.10	0.12	−1.2	U			H12	4.0	64Ba15	
		5308	3			0.8	U			H21	2.5	70Ma05	
^{144}Sm−^{144}Nd		1951	3	1913.4	0.8	−3.1	B			H19	4.0	64Mc11	
		1911.9	1.1			0.5	–			H25	2.5	72Ba08	
		1913.68	0.94			−0.3	–			SH1	1.0	11Go23	
	ave.	1913.5	0.9			−0.1	1	91	86 ^{144}Sm			average	
^{144}Nd−^{143}Nd		269	25	272.98	0.06	0.0	U			R05	4.0	65De13	
		273	3			−0.0	U			M17	2.5	66Be10	
^{144}Nd−^{142}Nd		2366	3	2363.97	0.09	−0.3	U			M17	2.5	66Be10	
^{142}Cs−^{144}Cs$_{.592}$ ^{139}Cs$_{.409}$		−60	40	−55	13	0.1	U			P23	2.5	82Au01	
^{143}Cs−^{144}Cs$_{.745}$ ^{140}Cs$_{.255}$		−920	50	−889	17	0.2	U			P23	2.5	82Au01	
^{142}Cs−^{144}Cs$_{.329}$ ^{141}Cs$_{.671}$		290	40	272	11	−0.2	U			P23	2.5	82Au01	
^{143}Cs−^{144}Cs$_{.662}$ ^{141}Cs$_{.338}$		−651	21	−615	16	0.7	U			P33	2.5	86Au02	
^{143}Cs−^{144}Cs$_{.497}$ ^{142}Cs$_{.504}$		−790	50	−687	13	0.8	U			P23	2.5	82Au01	
^{144}Nd(α)^{140}Ce		1882.4	30.	1900.3	1.4	0.6	U					61Ma05	
		1882.4	20.			0.9	U					65Is01	
^{144}Sm(^{3}He,^{6}He)^{141}Sm		−8693	12	−8691	7	0.2	1	36	35 ^{141}Sm	MSU		78Pa11	
^{142}Ce(t,p)^{144}Ce		3582	15	3560	3	−1.5	U			LAl		72Mu09	
^{142}Nd(t,p)^{144}Nd		5450	30	5458.81	0.09	0.3	U			Ald		72Ch11	
^{144}Nd(p,t)^{142}Nd		−5470	20	−5458.81	0.09	0.6	U			Osa		71Ya10	
^{144}Sm(p,t)^{142}Sm		−10649	15	−10644.8	1.2	0.3	U			Ham		73Oe02	
^{143}Nd(n,γ)^{144}Nd		7817.11	0.07	7817.04	0.05	−1.0	–			MMn		82Is05	Z
		7816.93	0.08			1.3	–			ILn		91Ro.A	Z
		7816.94	0.23			0.4	U			Bdn		06Fi.A	
^{144}Nd(d,t)^{143}Nd		−1555	15	−1559.81	0.05	−0.3	U			Ors		73Ga01	
^{143}Nd(n,γ)^{144}Nd	ave.	7817.03	0.05	7817.04	0.05	0.1	1	100	70 ^{143}Nd			average	
^{143}Nd(^{3}He,d)^{144}Pm		−804	5	−790.6	2.6	2.7	B			McM		80St10	*
^{143}Nd(^{3}He,d)^{144}Pm−^{142}Nd()^{143}Pm		402.7	1.6	403.2	1.5	0.3	1	91	49 ^{143}Pm			75Ma04	
^{144}Sm(t,α)^{143}Pm		13542	25	13519.9	2.7	−0.9	U			Ald		68Ha13	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{144}Sm(d,t)^{143}Sm		−4262	10	−4262.6	2.3	−0.1	U					72Ja28	
^{144}Sm(p,d)^{143}Sm−^{148}Gd()^{147}Gd		−1536	2	−1536.0	2.0	0.0	1	100	100 ^{143}Sm			86Ru04	
^{144}Tm(p)^{143}Er		1712.0	16.				3			ORp		05Gr32	
^{144}Cs(β^-)^{144}Ba		8451	30	8497	20	1.5	o			Gsn		81De25	
		8560	80			−0.8	–			Bwg		87Gr.A	
		8462	35			1.0	–			Gsn		92Pr04	
	ave.	8480	30			0.6	1	40	38 ^{144}Cs			average	
^{144}Ba(β^-)^{144}La		3055	70	3080	14	0.4	U			Bwg		87Gr.A	
^{144}La(β^-)^{144}Ce		4300	100	5584	13	12.8	B					79Ik07	
		5435	90			1.7	U			Bwg		87Gr.A	
		5540	100			0.4	o			Kur		02Sh.B	
		5540	100			0.4	U			Kur		02Sh16	
^{144}Ce(β^-)^{144}Pr		315.6	1.5	318.6	0.8	2.0	3					66Da04	
		320	1			−1.4	3					76Ra33	
^{144}Pr(β^-)^{144}Nd		2996	3	2997.4	2.4	0.5	2					59Po77	
		3000	4			−0.6	2					66Da04	
^{144}Eu(β^+)^{144}Sm		6330	30	6347	11	0.6	–			IRS		83Al06	
		6400	80			−0.7	U			IRS		93Al03	
		6287	30			2.0	–			Dbn		94Po26	
^{144}Sm(p,n)^{144}Eu		−7110.0	30.	−7130	11	−0.7	–					65Me12	
^{144}Eu(β^+)^{144}Sm	ave.	6315	17	6347	11	1.9	1	39	39 ^{144}Eu			average	
^{144}Gd(β^+)^{144}Eu		4300	400	3860	30	−1.1	U					70Ar04	
^{144}Tb^m(IT)^{144}Tb		396.9	0.5				3					86Re11	
$*^{144}$Tb^m−u	Re-assigned to the iso. state by the evaluator											HWJ267	**
$*^{143}$Nd(^{3}He,d)^{144}Pm	Based on ^{146}Nd(^{3}He,d)^{147}Pm Q =–87.6(0.9) keV											AHW	**
$*^{144}$Tm(p)^{143}Er	Trends in Mass Surface TMS suggest ^{144}Tm 160 keV less bound											GAu	*G
^{145}Xe−^{133}Cs$_{1.090}$		47777	12				2			MA8	1.0	09Ne11	
^{145}Cs−^{133}Cs$_{1.090}$		38588	12	38586	10	−0.1	–			MA8	1.0	08We02	
		38583	17			0.2	–			CP1	1.0	13Va12	
	ave.	38586	10			−0.0	1	99	99 ^{145}Cs			average	
^{145}Ba−u		−72481.6	9.1	−72482	8	−0.0	2			CP1	1.0	06Sa56	
		−72482.5	14.0			0.0	2			GR1	1.0	25Sp03	
^{145}La−u		−78188.8	13.3	−78196	12	−0.5	2			CP1	1.0	06Sa56	
		−78233.5	30.1			1.2	2			GR1	1.0	25Sp03	
^{145}Ce−u		−82771.8	92.2	−82733	18	0.4	U			CP1	1.0	06Sa56	
		−82728.4	19.3			−0.2	1	89	89 ^{145}Ce	GR1	1.0	25Sp03	
^{145}Pr−u		−85492.8	25.8	−85484	8	0.4	U			GR1	1.0	25Sp03	
C_{10} O H_9−^{145}Nd		152641	55	152762.4	1.2	0.6	U			R05	4.0	65De13	
		152653	30			0.9	U			R05	4.0	65De13	
C_9 ^{13}C O H_8−^{145}Nd		148231	31	148292.2	1.2	0.5	U			R05	4.0	65De13	
^{145}Pm−u		−87255	30	−87246.0	2.9	0.3	U			GS2	1.0	05Li24	
^{145}Sm−u		−86535	30	−86584.6	1.4	−1.7	U			GS2	1.0	05Li24	
^{145}Eu−^{133}Cs$_{1.090}$		19338	17	19328	3	−0.6	U			MA5	1.0	00Be42	
^{145}Gd−u		−78287	30	−78291	21	−0.1	2			GS2	1.0	05Li24	
		−78294	30			0.1	2			GS2	1.0	05Li24	*
^{145}Tb−u		−70726	30				2			GS2	1.0	05Li24	*
^{145}Dy−u		−62575	49	−62526	7	1.0	U			GS2	1.0	05Li24	*
^{145}Dy−^{85}Rb$_{1.706}$		87960.7	7.0				2			SH1	1.0	07Ra37	*
^{145}Ho−^{85}Rb$_{1.706}$		97754.1	8.0				2			SH1	1.0	07Ra37	
^{145}Nd ^{35}Cl$_2$−^{141}Pr ^{37}Cl$_2$		10828	7	10819.2	1.5	−0.5	U			H21	2.5	70Ma05	
^{145}Nd ^{35}Cl−^{143}Nd ^{37}Cl		5744	5	5709.50	0.26	−1.7	U			H12	4.0	64Ba15	
		5703	4			0.7	U			H21	2.5	70Ma05	
^{145}Nd−^{144}Nd		2582	21	2486.39	0.25	−1.1	U			R05	4.0	65De13	
		2480	2			1.3	U			M17	2.5	66Be10	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{145}Nd−^{143}Nd		2862	40	2759.37	0.25	−0.6	U			R05	4.0	65De13	
		2751	3			1.1	U			M17	2.5	66Be10	
^{142}Cs−^{145}Cs$_{.490}$ ^{139}Cs$_{.511}$		240	50	146	8	−0.8	U			P23	2.5	82Au01	
^{144}Cs−^{145}Cs$_{.828}$ ^{139}Cs$_{.173}$		450	50	414	21	−0.3	U			P23	2.5	82Au01	
^{143}Cs−^{145}Cs$_{.592}$ ^{140}Cs$_{.409}$		−700	80	−609	10	0.5	U			P23	2.5	82Au01	
^{143}Cs−^{145}Cs$_{.493}$ ^{141}Cs$_{.507}$		−310	40	−308	10	0.0	U			P23	2.5	82Au01	
^{144}Cs−^{145}Cs$_{.662}$ ^{142}Cs$_{.338}$		320	18	319	21	−0.0	1	21	19 ^{144}Cs	P33	2.5	86Au02	
^{144}Cs−^{145}Cs$_{.497}$ ^{143}Cs$_{.503}$		600	40	615	21	0.1	U			P23	2.5	82Au01	
^{145}Pm(α)^{141}Pr		2303.6	40.	2321.4	2.9	0.4	U					62Nu01	
^{145}Nd(n,α)^{142}Ce		8706	30	8747.1	2.2	1.4	U			ILL		75Em04	
^{145}Nd(p,t)^{143}Nd		−5100	20	−5090.50	0.24	0.5	U			Osa		71Ya10	
^{144}Nd(n,γ)^{145}Nd		5755.3	0.7	5755.26	0.23	−0.1	–					75Na.A	
		5756.9	2.0			−0.8	U					77Mc09	
		5755.26	0.25			0.0	–			Bdn		06Fi.A	
^{144}Nd(d,p)^{145}Nd		3521	15	3530.70	0.23	0.6	U			Hei		67Wi08	
		3538	15			−0.5	U			Ors		73Ga01	
^{144}Nd(n,γ)^{145}Nd	ave.	5755.26	0.24	5755.26	0.23	−0.0	1	96	75 ^{145}Nd			average	
^{144}Nd(^{3}He,d)^{145}Pm		−680	5	−685.0	2.5	−1.0	1	26	25 ^{145}Pm	McM		80St10	*
^{144}Nd(^{3}He,d)^{145}Pm−^{143}Nd()^{144}Pm		105.2	1.6	105.7	1.5	0.3	1	91	57 ^{144}Pm			75Ma04	
^{144}Sm(n,γ)^{145}Sm		6757.1	0.3	6757.10	0.30	−0.0	1	99	92 ^{145}Sm			79Wa22	
^{144}Sm(d,p)^{145}Sm		4533	12	4532.53	0.30	−0.0	U			Tal		65Ke09	
		4547	15			−1.0	U			Kop		67Ch16	
^{144}Sm(^{3}He,d)^{145}Eu		−2184	4	−2178.5	2.7	1.4	–			Mun		82Sc25	
		−2174	4			−1.1	–					84Ru.A	
	ave.	−2179.0	2.8			0.2	1	91	91 ^{145}Eu			average	
^{145}Dy(εp)^{144}Gd		6000	500	6228	29	0.5	U					83La.A	*
^{145}Tm(p)^{144}Er		1740.1	10.	1736	7	−0.4	3			ORp		98Ba13	
		1732.1	10.			0.4	3			Arp		07Se06	
^{145}Cs(β^-)^{145}Ba		7358	70	7462	12	1.5	U			Gsn		81De25	
		7930	75			−6.2	C			Bwg		87Gr.A	
		7865	50			−8.1	B			Gsn		92Pr04	
^{145}Ba(β^-)^{145}La		4925	80	5323	13	5.0	C			Bwg		87Gr.A	
^{145}La(β^-)^{145}Ce		4110	80	4226	20	1.4	U			Bwg		87Gr.A	
^{145}Ce(β^-)^{145}Pr		2490	100	2562	18	0.7	U					67Ho19	*
		2600	100			−0.4	U					80Ya07	*
		2530	50			0.6	1	13	11 ^{145}Ce	Bwg		87Gr.A	
^{145}Pr(β^-)^{145}Nd		1805	10	1806	7	0.1	1	49	49 ^{145}Pr			59Dr.A	
^{145}Pm(ε)^{145}Nd		143	15	164.4	2.5	1.4	U					59Br65	*
		150	5			2.9	B					74To04	*
^{145}Sm(ε)^{145}Pm		607	6	616.1	2.5	1.5	–					71My01	*
		622	5			−1.2	–					83Vo10	*
	ave.	616	4			0.1	1	44	41 ^{145}Pm			average	
^{145}Eu(β^+)^{145}Sm		2710	15	2659.8	2.7	−3.3	B					68Ad04	*
		2647	12			1.1	U					83Sc28	*
^{145}Gd(β^+)^{145}Eu		5070	60	5066	20	−0.1	U					79Fi07	
		5090	90			−0.3	o			IRS		83Ve.A	*
		5070	80			−0.0	U			IRS		85Al13	
^{145}Gd(ε)^{145}Eu		5000	70			0.9	U					77Ho18	
^{145}Tb(β^+)^{145}Gd		6700	200	7050	30	1.7	U					86Ve.A	*
		6400	150			4.3	B			IRS		93Al03	
^{145}Dy(β^+)^{145}Tb^m		7300	200				3			IRS		93Al03	
*^{145}Gd−u	$M-A$=–72181(28) keV for ^{145}Gd^m at 749.1 keV											Nub269	**
*^{145}Tb−u	TMS suggests isomer is not presented											HWJ234	**
*^{145}Dy−u	$M-A$=–58230(30) keV for mixture gs+m at 118.2 keV											Nub269	**
*^{145}Dy−^{85}Rb$_{1.706}$	D_M=88054.7(6.8) μu for mixture gs+m at 118.2 keV with ratio R=0.741(13)											Nub269	**
*^{144}Nd(^{3}He,d)^{145}Pm	Based on ^{146}Nd(^{3}He,d)^{147}Pm Q =–87.6(0.9) keV											AHW	**
*^{145}Dy(εp)^{144}Gd	As read from graph											AHW	**
*^{145}Ce(β^-)^{145}Pr	E_{β^-}=1700(100) 1810(100) respectively, to (3/2)$^-$ level at 786.91; and other E_{β^-}											Ens092	**
*^{145}Pm(ε)^{145}Nd	LM/K=0.85(0.03) to 3/2$^-$ level at 67.167 keV											Ens092	**
*^{145}Pm(ε)^{145}Nd	pK=0.554(0.025) to 5/2$^-$ level at 72.486 keV, and other pK											Ens092	**
*^{145}Sm(ε)^{145}Pm	pK=0.27(0.03) 0.35(0.025) respectively, to 3/2$^+$ level at 492.31 keV											Ens092	**
*^{145}Eu(β^+)^{145}Sm	E_{β^+}=794(15) to 3/2$^+$ level at 893.788 keV											Ens092	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
$*^{145}$Eu(β^+)^{145}Sm		pK=0.72(0.02) to (5/2$^-$,7/2$^-$) level at 2508.31 and 9$^-$ at 2513.37 levels											Ens092	**
$*^{145}$Gd(β^+)^{145}Eu		E_{β^+}=2310(90) to 3/2$^+$ level at 1758.03 keV, and other E_{β^+}											Ens092	**
$*^{145}$Tb(β^+)^{145}Gd		E_{β^+}=3300(200) to (9/2$^-$) level at 2382.3(0.2) keV											Ens092	**
^{146}Xe−^{133}Cs$_{1.098}$		52332	26				2				MA8	1.0	09Ne11	
^{146}Cs−^{133}Cs$_{1.098}$		44437.4	3.3	44436	3	−0.5	2				MA8	1.0	17At01	
		44421.8	9.2			1.5	2				CP1	1.0	13Va12	
^{146}Ba−u		−69618	112	−69636.8	1.9	−0.1	o				GT2	2.5	08Kn.A	
		−69963	141			0.9	U				GT2	2.5	08Su19	
		−69717.5	23.7			3.4	B				CP1	1.0	06Sa56	
		−69636.8	1.9				2				CP2	1.0	18Or.A	*
		−69634.4	19.3			−0.1	U				GR1	1.0	25Sp03	
^{146}La−u		−74259	54	−74311.9	1.8	−1.0	U				CP1	1.0	06Sa56	*
		−74311.9	1.8				2				CP2	1.0	20Or02	*
		−74236.7	54.8			−1.4	U				GR1	1.0	25Sp03	*
^{146}La^m−u		−74160.1	1.8				2				CP2	1.0	20Or02	*
^{146}Ce−u		−81191.8	20.8	−81200	13	−0.4	–				CP1	1.0	06Sa56	
		−81171	40			−0.7	–				GS3	1.0	12Ch19	
		−81199.7	20.4			−0.0	–				GR1	1.0	25Sp03	
	ave.	−81193	14			−0.5	1	95	95	^{146}Ce			average	
^{146}Pr−u		−82404.2	58.0	−82340	30	1.0	1	28	28	^{146}Pr	GR1	1.0	25Sp03	
C_{12} H_2−^{146}Nd		102453	31	102529.3	1.2	0.6	U				R05	4.0	65De13	
C_{10} O H_{10}−^{146}Nd		160017	27	160044.1	1.2	0.3	U				R05	4.0	65De13	
		159971	50			0.4	U				R05	4.0	65De13	
C_9 ^{13}C O H_9−^{146}Nd		155525	35	155573.9	1.2	0.3	U				R05	4.0	65De13	
^{146}Pm−u		−85289	30	−85299	5	−0.3	U				GS2	1.0	05Li24	
^{146}Eu−^{133}Cs$_{1.098}$		21029	15	21023	6	−0.4	1	18	18	^{146}Eu	MA5	1.0	00Be42	
^{146}Tb^m−u		−72464	77				2				GS2	1.0	05Li24	*
^{146}Dy−u		−67150	30	−67155	7	−0.2	U				GS2	1.0	05Li24	
^{146}Dy−^{85}Rb$_{1.718}$		84390.0	7.2	84390	7	−0.0	1	100	100	^{146}Dy	SH1	1.0	07Ra37	
^{146}Ho−^{133}Cs$_{1.098}$		48797	10	48807	7	1.0	1	50	50	^{146}Ho	SH1	1.0	07Ra37	
^{146}Ho−^{85}Rb$_{1.718}$		96549	10	96539	7	−1.0	1	50	50	^{146}Ho	SH1	1.0	07Ra37	
^{146}Er−^{85}Rb$_{1.718}$		103960.4	9.2	103964	7	0.4	1	61	61	^{146}Er	SH1	1.0	07Ra37	
^{146}Nd ^{35}Cl−^{144}Nd ^{37}Cl		6003	3	5979.83	0.27	−1.9	U				H12	4.0	64Ba15	
		5966	4			1.4	U				H21	2.5	70Ma05	
		5982.8	1.1			−1.1	U				H25	2.5	72Ba08	
^{146}Nd−^{145}Nd		526	33	543.31	0.09	0.1	U				R05	4.0	65De13	
		536	2			1.5	U				M17	2.5	66Be10	
^{146}Nd−^{144}Nd		3147	36	3029.70	0.26	−0.8	U				R05	4.0	65De13	
		3026	3			0.5	U				M17	2.5	66Be10	
^{145}Cs−^{146}Cs$_{.828}$ ^{140}Cs$_{.173}$		−580	80	−928	9	−1.7	U				P23	2.5	82Au01	
^{144}Cs−^{146}Cs$_{.329}$ ^{143}Cs$_{.671}$		320	50	334	21	0.1	U				P23	2.5	82Au01	
^{145}Cs−^{146}Cs$_{.662}$ ^{143}Cs$_{.338}$		−440	30	−565	10	−1.7	U				P33	2.5	86Au02	
^{145}Cs−^{146}Cs$_{.497}$ ^{144}Cs$_{.503}$		−730	30	−739	13	−0.1	U				P33	2.5	86Au02	
^{146}Sm(α)^{142}Nd		2529.5	20.	2528.8	2.8	−0.0	U						64Nu02	
		2622.0	30.			−3.1	B						66Fr11	
		2524.2	4.			1.1	1	47	46	^{146}Sm			87Me08	Z
^{144}Nd(t,p)^{146}Nd		4834	30	4838.69	0.24	0.2	U				Ald		72Ch11	
^{144}Sm(t,p)^{146}Sm		6681	25	6691.5	2.8	0.4	U				Ald		66Bj01	
^{144}Sm(^{3}He,p)^{146}Eu		2797	12	2794	6	−0.3	1	25	24	^{146}Eu			84Ru.A	
^{144}Sm(^{3}He,n)^{146}Gd		977	30	980	4	0.1	U				Bld		79Al07	
^{144}Sm(^{12}C,^{10}Be)^{146}Gd		−18476	25	−18487	4	−0.4	U				MSU		80Pa07	
^{146}Nd(d,^{3}He)^{145}Pr		−3095	10	−3095	7	−0.0	1	49	49	^{145}Pr	KVI		79Sa.A	
^{145}Nd(n,γ)^{146}Nd		7565.28	0.10	7565.23	0.09	−0.5	–				MMn		82Is05	Z
		7565.05	0.18			1.0	–				Bdn		06Fi.A	
	ave.	7565.23	0.09			−0.0	1	99	74	^{146}Nd			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{146}Sm(^{3}He,α)^{145}Sm	12161	5	12161.4	2.9	0.1	1	33	30 ^{146}Sm			86Ru04	*
^{146}Tm(p)^{145}Er	895.2	8.	896	6	0.1	o			ORp		03Gi10	
	896.2	8.			−0.1	3			Arp		05Ro40	*
	895.2	8.			0.1	3			ORp		06Ta08	*
^{146}Tm^{m}(p)^{145}Er	1197.3	5.	1199.3	1.0	0.4	U			Dap		93Li18	
	1198.3	10.			0.1	o			ORp		01Ry01	
	1200.3	8.			−0.1	U			Arp		05Ro40	
	1199.3	1.				3			ORp		06Ta08	
^{146}Tm^{m}(p)^{145}Er^{m}	944.5	4.				4			ORp		06Ta08	
^{146}Tm^{n}(p)^{145}Er^{m}	1126.8	5.	1127.8	1.0	0.2	U			Dap		93Li18	
	1127.8	10.			−0.0	o			ORp		01Ry01	
	1129.8	8.			−0.3	U			Arp		05Ro40	
	1127.8	1.				5			ORp		06Ta08	
^{146}Cs(β^-)^{146}Ba	9300	900	9556	3	0.3	o			Gsn		81De25	
	9310	60			4.1	B			Bwg		87Gr.A	
	9375	50			3.6	B			Gsn		92Pr04	
^{146}Ba(β^-)^{146}La	4280	100	4354.8	2.4	0.7	U			Gsn		81De25	*
^{146}La(β^-)^{146}Ce	6175	100	6416	13	2.4	B			Gsn		81De25	*
^{146}La^{m}(β^-)^{146}Ce	6380	30	6558	13	5.9	B			Trs		82Br23	*
	6640	50			−1.6	U			Trs		82Br23	*
	6620	70			−0.9	U			Bwg		86Gr11	
	6580	80			−0.3	U					01Ko07	*
^{146}Ce(β^-)^{146}Pr	1100	80	1066	28	−0.4	–					54Be10	*
	1050	100			0.2	–					67Ho19	*
	951	50			2.3	–					80Ya07	*
	1065	100			0.0	–					81Eb01	*
ave.	1010	40			1.5	1	60	55 ^{146}Pr			average	
^{146}Pr(β^-)^{146}Nd	4150	200	4224	29	0.4	U					54Be10	*
	4250	200			−0.1	U					65Ra02	*
	4080	100			1.4	–					68Da13	*
	4140	100			0.8	–					78Ik03	*
ave.	4110	70			1.6	1	16	16 ^{146}Pr			average	
^{146}Pm(β^-)^{146}Sm	1542	3				2					74Sc06	*
^{146}Eu(β^+)^{146}Sm	3871	10	3879	6	0.8	–					62Fu16	*
	3871	20			0.4	–					64Ta11	*
	3896	20			−0.8	–			Got		88Sa06	*
ave.	3875	8			0.5	1	52	46 ^{146}Eu			average	
^{146}Gd(β^+)^{146}Eu	1757	30	1031	7	−24.2	B					70Ag01	*
	1300	200			−1.3	U					81Ka07	*
^{146}Tb(β^+)^{146}Gd	8240	150	8320	40	0.6	o			IRS		83Al06	
	7790	128			4.2	U			IRS		93Al03	*
	8310	50			0.3	1	80	80 ^{146}Tb	Dbn		94Po26	
^{146}Dy(β^+)^{146}Tb	5160	100	5210	50	0.5	1	20	20 ^{146}Tb	IRS		93Al03	
*^{146}Ba−u	Represents frequency ratio $^{146}Ba^{++}/(C_6H_6)^+$ =0.934887856(12)										HWJ208	**
*^{146}La−u	D_M=–74182.5(30.6) μu for mixture gs+m at 141.4(2.4) keV;										Nub269	**
*^{146}La−u	$M-A$=–69100.6(28.5) keV										Nub269	**
*^{146}La−u	Represents frequency ratio $^{146}La^{++}/(C_6H_6)^+$ =0.934857905(12)										HWJ206	**
*^{146}La−u	Corrected for unresolved isomeric state using exc. energy in 23Ga02										25Sp03	**
*^{146}La^{m}−u	Represents frequency ratio $^{146}Lam^{++}/(C_6H_6)^+$ =0.934858878(12)										HWJ206	**
*^{146}Tb^{m}−u	Re-assigned to the iso. state by the evaluator										HWJ267	**
*^{146}Sm(^{3}He,α)^{145}Sm	$Q-Q(^{148}Gd(^{3}He,\alpha))$=–567(5) keV										AHW	**
*^{146}Tm(p)^{145}Er	Trends in Mass Surface TMS suggest ^{146}Tm 200 keV less bound										HWJ266	**
*^{146}Ba(β^-)^{146}La	E_{β^-}=3910(100) to 1^+ level at 372.4 keV, and other E_{β^-}										Ens97c	**
*^{146}La(β^-)^{146}Ce	E_{β^-}=5919(100) 6120(30) respectively, to 2^+ level at 258.46 keV, and other E_{β^-}										Ens97c	**
*^{146}La^{m}(β^-)^{146}Ce	corrected 82Br23 data as E_{β^-}=6380 keV to 2^+ level at 258.46 keV										86Gr11	**
*^{146}La^{m}(β^-)^{146}Ce	E_{β^-}=6580(100) and 6320(80) to ground state and 2^+ level at 258.46 keV										01Ko07	**
*^{146}Ce(β^-)^{146}Pr	E_{β^-}=750(80) 700(100) 600(50) 715(100) respectively, to 1^+ at 351.78 keV										Ens97c	**
*^{146}Pr(β^-)^{146}Nd	E_{β^-}=3700(200) 3800(200) respectively, to 2^+ level at 453.77 keV										Ens97c	**
*^{146}Pr(β^-)^{146}Nd	E_{β^-}=4100(200), 3600(100), 2100(100) to ground state, 2^+ 453.77, 2^+ 1978.45 levels										Ens97c	**
*^{146}Pr(β^-)^{146}Nd	E_{β^-}=4150(150), 3700(100), 2160(100) to ground state, 2^+ 453.77, 2^+ 1978.45 levels										Ens97c	**
*^{146}Pm(β^-)^{146}Sm	E_{β^-}=795(3) to 2^+ level at 747.2 keV										Ens97c	**
*^{146}Eu(β^+)^{146}Sm	E_{β^+}=2107(11) 2100(20) respectively, to 2^+ level at 747.2 keV, and other E_{β^+}										Ens97c	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{146}$Eu(β^+)^{146}Sm		e/β^+ to 2045.8 level										Ens97c	**
$*^{146}$Gd(β^+)^{146}Eu		E_{β^+}=350(30) to 1^- level at 384.79 keV										Ens97c	**
$*^{146}$Gd(β^+)^{146}Eu		pK to 690.7 level, $p^+ < 1\times10^{-4}$ to 384.8 level, see ^{150}Dy(β^+)										Ens97c	**
$*^{146}$Tb(β^+)^{146}Gd		Reported half-life 24.1(0.5)s corresponds to ^{146}Tb^m										GAu	**
*		Q_{β^-}=8060(100) keV from ^{146}Tb^m at 270(80) keV										Nub269	**
^{147}Cs−^{133}Cs$_{1.105}$		48640	64	48737	9	1.5	U			MA8	1.0	08We02	
		48737.1	9.0				2			MA8	1.0	17At01	
^{147}Ba−u		−64696.1	21.2	−64736.8	2.2	−1.9	U			CP1	1.0	06Sa56	
		−64715.4	12.9			−1.7	U			GR1	1.0	25Sp03	
^{147}Ba−^{12}C$_6$ ^{1}H$_6$		−111687.0	2.2				2			CP2	1.0	24Ra.A	
^{147}La−u		−71582.2	11.5	−71571.9	2.8	0.9	U			CP1	1.0	06Sa56	
		−71516.3	39.7			−1.4	U			GR1	1.0	25Sp03	
^{147}La−^{12}C$_6$ ^{1}H$_6$		−118522.1	2.8				2			CP2	1.0	24Ra.A	
^{147}Ce−u		−77309.2	9.6	−77334	3	−2.5	U			CP1	1.0	06Sa56	
		−77328.5	14.0			−0.4	U			GR1	1.0	25Sp03	
^{147}Ce−^{12}C$_6$ ^{1}H$_6$		−124283.8	3.3				2			CP2	1.0	24Ra.A	
^{147}Pr−^{12}C$_6$ ^{1}H$_6$		−127980.6	3.1				2			CP2	1.0	24Ra.A	
^{147}Pr−u		−81020.4	22.5	−81030	3	−0.4	U			GR1	1.0	25Sp03	
^{147}Nd−^{12}C$_6$ ^{1}H$_6$		−130846.7	3.6	−130845.9	1.2	0.2	1	10	10 ^{147}Nd	CP2	1.0	24Ra.A	
^{147}Nd−u		−83784.8	75.1	−83895.7	1.2	−1.5	U			GR1	1.0	25Sp03	
C$_8$ H$_5$ N O$_2$−^{147}Sm		117197	40	117126.0	1.2	−0.4	U			R04	4.0	64De15	
C$_9$ H$_7$ O$_2$−^{147}Sm		129703	17	129702.0	1.2	−0.0	U			R04	4.0	64De15	
^{147}Eu−^{133}Cs$_{1.105}$		21215	16	21226.1	2.7	0.7	U			MA5	1.0	00Be42	
^{147}Tb−u		−75934	34	−75946	8	−0.3	U			GS2	1.0	05Li24	*
^{147}Tb−^{133}Cs$_{1.105}$		28533	12	28530	8	−0.3	1	48	48 ^{147}Tb	SH1	1.0	07Ra37	*
^{147}Dy−u		−68909	30	−68917	10	−0.3	U			GS2	1.0	05Li24	
		−68908	30			−0.3	U			GS2	1.0	05Li24	*
^{147}Dy−^{133}Cs$_{1.105}$		35558.3	9.5				2			SH1	1.0	07Ra37	*
^{147}Ho−u		−59944	30	−59858	5	2.9	B			GS2	1.0	05Li24	
^{147}Ho−^{133}Cs$_{1.105}$		44613.7	7.8	44618	5	0.5	1	47	47 ^{147}Ho	SH1	1.0	07Ra37	
^{147}Ho−^{85}Rb$_{1.729}$		92661.6	7.4	92658	5	−0.5	1	53	53 ^{147}Ho	SH1	1.0	07Ra37	
^{147}Er−^{133}Cs$_{1.105}$		54467	22	54496	21	1.3	o			SH1	1.0	07Ra37	*
^{147}Er−^{85}Rb$_{1.729}$		102536	21				2			SH1	1.0	07Ra37	*
^{147}Tm−^{85}Rb$_{1.729}$		113900	11	113895	7	−0.4	1	44	44 ^{147}Tm	SH1	1.0	07Ra37	
^{147}Eu−^{142}Sm$_{1.035}$		4516	17	4510.3	2.9	−0.3	U			MA7	1.0	01Bo59	
^{147}Sm ^{35}Cl−^{145}Nd ^{37}Cl		5305	4	5275.1	0.6	−1.9	U			H12	4.0	64Ba15	
		5264	4			1.1	U			H21	2.5	70Ma05	
^{145}Cs−^{147}Cs$_{.705}$ ^{140}Cs$_{.296}$		−170	170	−644	11	−1.1	U			P23	2.5	82Au01	
^{144}Cs−^{147}Cs$_{.490}$ ^{141}Cs$_{.511}$		80	250	226	21	0.2	U			P23	2.5	82Au01	
^{145}Cs−^{147}Cs$_{.493}$ ^{143}Cs$_{.507}$		−87	22	−147	11	−1.1	U			P33	2.5	86Au02	
^{147}Sm(α)^{143}Nd		2292.5	10.	2311.1	0.5	1.9	U					62Si14	Z
		2296.7	5.			2.9	U					66Ma05	Z
		2300.8	5.			2.1	U					70Gu14	Z
		2310.5	0.5			1.2	o					16Ca43	*
^{147}Eu(α)^{143}Pm		2990.6	10.	2991	3	0.0	U					62Si14	Z
		2981.5	20.			0.5	U					64To04	Z
		2987.2	5.			0.7	1	37	22 ^{143}Pm	Dba		67Go32	Z
^{147}Sm(n,α)^{144}Nd		10114	8	10128.1	0.5	1.8	U			ILL		74Em01	
^{144}Sm(^{12}C,^{9}Be)^{147}Gd		−17832	30	−17957.1	1.2	−4.2	B			MSU		80Pa07	
		−17921	25			−1.4	U			Ors		85Be24	
^{144}Sm(^{14}N,^{11}Be)^{147}Tb		−28280	50	−28539	8	−5.2	B			Hei		85Gy01	
^{147}Sm(p,t)^{145}Sm		−6287	8	−6275.7	1.0	1.4	U			Min		72De47	
^{146}Nd(n,γ)^{147}Nd		5292.19	0.15	5292.19	0.09	0.0	–			ILn		75Ro16	Z
		5292.19	0.11			0.0	–			Bdn		06Fi.A	
^{146}Nd(d,p)^{147}Nd		3070	15	3067.63	0.09	−0.2	U			Hei		67Wi08	
^{146}Nd(n,γ)^{147}Nd	ave.	5292.19	0.09	5292.19	0.09	0.0	1	99	75 ^{147}Nd			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{147}Sm(d,t)^{146}Sm		−98	10	−84.0	2.8	1.4	U			McM		75Si03	
^{147}Tb(p)^{146}Gd		−1945	18	−1944	8	0.0	1	21	17 ^{147}Tb			87Sc.A	
^{147}Tm(p)^{146}Er		1062.2	6.	1058.2	2.7	−0.7	o					82Kl03	
		1058.2	3.3			0.0	–			Dap		93Se04	*
		1052.3	10.			0.6	U					93To02	
		1067.3	15.			−0.6	U			ORp		03Gi10	
		1057.2	5.0			0.2	–			Jyp		23Au03	
	ave.	1057.9	2.8			0.1	1	96	56 ^{147}Tm			average	
^{147}Tm^{m}(p)^{146}Er		1124.7	6.	1123.0	2.7	−0.3	o					84Ho.A	
		1118.5	3.9			1.2	2			Dap		93Se04	
		1126.1	5.3			−0.6	2					95Ho24	
		1127.7	5.0			−0.9	2			Jyp		23Au03	
^{147}Ba(β^-)^{147}La		5750	50	6367	3	12.3	C			Bwg		87Gr.A	
^{147}La(β^-)^{147}Ce		4945	55	5367	4	7.7	C			Bwg		87Gr.A	
		5150	40			5.4	B			Kur		95Ik03	
		5370	100			−0.0	o			Kur		02Sh.B	
		5366	40			0.0	U			Kur		09Ha.B	
^{147}Ce(β^-)^{147}Pr		3290	40	3444	4	3.8	C			Bwg		87Gr.A	
		3426	20			0.9	U			Kur		95Ik03	
		3380	100			0.6	U			Kur		02Sh.B	
^{147}Pr(β^-)^{147}Nd		2700	200	2669	3	−0.2	U					64Ho03	
		2790	100			−1.2	U					81Ya06	*
		2669	40			−0.0	U					148Nd-1	
		2711	28			−1.5	U			Kur		95Ik03	
^{147}Nd(β^-)^{147}Pm		894.6	1.0	894.6	0.6	−0.0	1	32	18 ^{147}Pm			67Ca18	*
^{147}Pm(β^-)^{147}Sm		223.2	0.5	224.99	0.29	3.6	o					50La04	
		224.3	1.3			0.5	–					58Ha32	
		224.5	0.4			1.2	–					66Hs01	
		225.84	0.49			−1.7	–					25Cr03	
	ave.	225.0	0.3			−0.0	1	94	82 ^{147}Pm			average	
^{147}Eu(β^+)^{147}Sm		1767	10	1721.5	2.3	−4.5	B					67Ad03	
		1723	3			−0.5	1	58	57 ^{147}Eu			80Bu04	
		1702	13			1.5	U					84Sc18	*
		1692	18			1.6	U					84Sc18	
^{147}Gd(β^+)^{147}Eu		2185	5	2187.8	2.5	0.6	1	25	19 ^{147}Eu			80Vy01	*
		2199	17			−0.7	U					84Sc18	*
^{147}Tb(β^+)^{147}Gd		4700	90	4616	8	−0.9	U					83Ve06	*
		4490	60			2.1	U			Got		85Ti01	
		4560	50			1.1	U					Averag	*
		4609	15			0.4	1	27	26 ^{147}Tb	GSI		91Ke11	*
		4509	60			1.8	U			IRS		93Al03	*
		4560	80			0.7	U					97Wa04	
^{147}Dy(β^+)^{147}Tb		6334	60	6547	12	3.5	B			IRS		83Al06	*
		6480	100			0.7	U			IRS		83Al18	*
		6334	60			3.5	C					85Af.A	*
		6480	100			0.7	U			IRS		85Al08	*

*^{147}Tb−u	$M-A$=–70707(28) keV for mixture gs+m at 50.6 keV	Nub269 **
*^{147}Tb−^{133}Cs$_{1.105}$	D_M=28574(12) μu for mixture gs+m at 50.6 keV with ratio R=0.741(13)	Nub269 **
*^{147}Dy−u	$M-A$=–63437(28) keV for ^{147}Dy^{m} at 750.5 keV	Nub269 **
*^{147}Dy−^{133}Cs$_{1.105}$	D_M=35567(14) and D_M=36358.4(9.5) for ^{147}Dy^{m} at 750.5 keV	Nub269 **
*^{147}Er−^{133}Cs$_{1.105}$	D_M=54531(11) μu for mixture gs+m at 60#35 keV with ratio R=0.741(13)	Nub269 **
*	error due to excitation energy, use only next item	GAu **
*^{147}Er−^{85}Rb$_{1.729}$	D_M=102559.5(8.3) μu for mixture gs+m at 60#35 keV with R=0.741(13)	Nub269 **
*^{147}Sm(α)^{143}Nd	Used as calibrant, without the origin	16Ca43 **
*^{147}Tm(p)^{146}Er	Q_p from E_p=1051.0(3.3), no screening correction should be applied	GAu **
*^{147}Pr(β^-)^{147}Nd	E_{β^-}=2760(100) to 49.93, 1450(100) to 1310.7 and 1350.5 keV	Ens092 **
*^{147}Nd(β^-)^{147}Pm	E_{β^-}=803.5(1.0) to 5/2$^+$ level at 91.1047 keV	Ens092 **
*^{147}Eu(β^+)^{147}Sm	p$^+$ =2.9(0.3)×10^{-3} to 3/2$^-$ level at 197.284 keV	Ens092 **
*^{147}Eu(β^+)^{147}Sm	pK=0.724(0.026) to (3/2$^+$,5/2$^+$) level at 1548.634 keV	Ens092 **
*^{147}Gd(β^+)^{147}Eu	E_{β^+}=933(5) to 7/2$^+$ level at 229.323 keV	Ens092 **

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{147}$Gd(β^+)^{147}Eu		pK=0.694(0.016) to 2073 level, recalculated by AHW										84Sc18	**
$*^{147}$Tb(β^+)^{147}Gd		E_{β^+}=2460(80) to 3/2$^+$ level at 1152.56 and 1/2$^+$ at 1292.3 keV, reinterpreted										Ens092	**
$*^{147}$Tb(β^+)^{147}Gd		Average KLM/β^+=2.03(0.15) → E_{β^+}=2190(50) from ^{147}Tb^m at 50.6(0.9) to 9/2$^-$										Ens092	**
*		level at 1397.00 from 3 references (no side-feeding correction applied):										AHW	**
*		p$^+$ =0.32(0.07) gives KLM/β^+=2.2(0.8)										84Ha.B	**
*		KLM/β^+=2.17(0.30)										85Al08	**
*		β^+/K=0.59(0.05) gives KLM/β^+=1.99(0.17)										85Sc09	**
$*^{147}$Tb(β^+)^{147}Gd		Q_{β^+}=4660(15) 4560(60) respectively, from ^{147}Tb^m at 50.6(0.9) keV										Nub269	**
$*^{147}$Dy(β^+)^{147}Tb		E_{β^+}=6012(60) from ^{147}Dy^m at 750.5 to ^{147}Tb^m at 50.6(0.9) keV										Nub269	**
$*^{147}$Dy(β^+)^{147}Tb		Q_{β^+}=7180(100) from ^{147}Dy^m at 750.5 to ^{147}Tb^m at 50.6(0.9) keV										Nub269	**
$*^{147}$Dy(β^+)^{147}Tb		E_{β^+}=6012(60) from ^{147}Dy^m at 750.5 to ^{147}Tb^m at 50.6(0.9) keV										Nub269	**
$*^{147}$Dy(β^+)^{147}Tb		Q_{β^+}=7180(100) from ^{147}Dy^m at 750.5 to ^{147}Tb^m at 50.6(0.9) keV										Nub269	**
^{148}Cs−^{133}Cs$_{1.113}$		54871	14				2			MA8	1.0	17At01	
^{148}Ba−u		−61777.0	1.6				2			CP2	1.0	18Or.A	*
		−61765.3	19.3	−61777.0	1.6	−0.6	U			GR1	1.0	25Sp03	
^{148}La−^{133}Cs$_{1.113}$		37922.4	2.9				2			JY2	1.0	25Ja05	
^{148}La−u		−67320.6	20.9	−67309.6	2.9	0.5	U			CP1	1.0	06Sa56	
		−67319.8	56.9			0.2	U			GR1	1.0	25Sp03	
^{148}Ce−u		−75578.2	13.0	−75567	9	0.9	–			CP1	1.0	06Sa56	
		−75561.4	14.0			−0.4	–			GR1	1.0	25Sp03	
	ave.	−75570	10			0.4	1	89	89 ^{148}Ce			average	
^{148}Ce ^{40}Ar$_2$−^{152}Pr$_{1.500}$		−48076.9	27.3	−48106	10	−1.1	1	13	11 ^{148}Ce	RI1	1.0	24Ki16	*
^{148}Pr−u		−77781	41	−77812	18	−0.8	–			CP1	1.0	06Sa56	*
		−77770.8	30.1			−1.4	–			GR1	1.0	25Sp03	*
	ave.	−77774	24			−1.6	1	57	57 ^{148}Pr			average	
C_{12} H_4−^{148}Nd		114261	34	114402.6	2.1	1.0	U			R05	4.0	65De13	
C_9 N O H_{10}−^{148}Nd		159186	59	159341.4	2.1	0.7	U			R05	4.0	65De13	
^{148}Nd−u		−83050.5	51.5	−83102.4	2.1	−1.0	U			GR1	1.0	25Sp03	
C_9 H_8 O_2−^{148}Sm		137540	26	137602.2	1.1	0.6	U			R04	4.0	64De15	
C_9 H_{10} N O−^{148}Sm		161275	31	161411.6	1.1	1.1	U			R04	4.0	64De15	
C_8 ^{13}C H_7 O_2−^{148}Sm		133030	60	133132.0	1.1	0.4	U			R04	4.0	64De15	
^{148}Eu−^{133}Cs$_{1.113}$		23315	15	23322	11	0.5	1	51	51 ^{148}Eu	MA5	1.0	00Be42	
^{148}Tb−u		−75692	41	−75720	13	−0.7	U			GS2	1.0	05Li24	*
^{148}Dy−^{133}Cs$_{1.113}$		32394	16	32384	9	−0.6	–			MA5	1.0	00Be42	
		32380	14			0.3	–			SH1	1.0	07Ra37	
	ave.	32386	11			−0.2	1	78	78 ^{148}Dy			average	
^{148}Ho−u		−62201	100	−62260	90	−0.6	U			GS2	1.0	05Li24	*
^{148}Ho−^{85}Rb$_{1.741}$		91318	90				2			SH1	1.0	07Ra37	*
^{148}Er−^{133}Cs$_{1.113}$		49967	11				2			SH1	1.0	07Ra37	
^{148}Tm−^{133}Cs$_{1.113}$		63616	11				2			SH1	1.0	07Ra37	
^{148}Eu−^{142}Sm$_{1.042}$		6451	17	6444	11	−0.4	1	40	39 ^{148}Eu	MA7	1.0	01Bo59	
^{148}Nd ^{35}Cl$_2$−^{144}Nd ^{37}Cl$_2$		12690	9	12706.7	1.8	0.7	U			H21	2.5	70Ma05	
		12703.6	2.1			0.6	1	12	11 ^{148}Nd	H25	2.5	72Ba08	
^{148}Sm ^{35}Cl$_2$−^{144}Sm ^{37}Cl$_2$		8710	10	8723.1	1.0	0.3	U			H12	4.0	64Ba15	
		8721.4	2.6			0.3	U			H25	2.5	72Ba08	
^{148}Nd ^{35}Cl−^{146}Nd ^{37}Cl		6740	5	6726.9	1.8	−0.7	U			H12	4.0	64Ba15	
		6721	4			0.6	U			H21	2.5	70Ma05	
		6723.8	2.7			0.5	U			H25	2.5	72Ba08	
		6725.7	0.9			0.5	1	62	61 ^{148}Nd	H26	2.5	73Me28	
^{148}Sm ^{35}Cl−^{146}Nd ^{37}Cl		4656	3	4656.6	0.5	0.1	U			H12	4.0	64Ba15	
^{148}Sm−^{147}Sm		110	44	−75.11	0.29	−1.1	U			R04	4.0	64De15	
^{148}Nd−^{146}Nd		3866	50	3776.8	1.8	−0.4	U			R05	4.0	65De13	
		3773	3			0.5	U			M17	2.5	66Be10	
^{145}Cs−^{148}Cs$_{.392}$ ^{143}Cs$_{.608}$		−370	90	−519	11	−0.7	U			P33	2.5	86Au02	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{148}Sm(α)^{144}Nd		2014.6	20.	1986.8	0.4	−1.4	U					70Gu14	
		1987.3	0.5			−0.9	1	78	44 ^{144}Nd			16Ca43	*
^{148}Eu(α)^{144}Pm		2703.2	30.	2695	10	−0.3	1	11	10 ^{148}Eu			64To04	
^{148}Gd(α)^{144}Sm		3271.29	0.03	3271.29	0.03	−0.0	1	100	96 ^{148}Gd			73Go29	
^{146}Nd(t,p)^{148}Nd		4139	30	4142.8	1.7	0.1	U			Ald		72Ch11	
^{148}Sm(p,t)^{146}Sm		−6011	8	−6000.7	2.8	1.3	1	13	12 ^{146}Sm	Min		72De47	
		−6018	15			1.2	U			Ham		74Oe03	
^{148}Gd(p,t)^{146}Gd		−7844	14	−7844	4	−0.0	U			LAl		83Fl05	
^{148}Gd(p,t)^{146}Gd−^{65}Cu()^{63}Cu		1500	4	1500	4	0.1	1	89	88 ^{146}Gd	Liv		86Ma40	
^{148}Nd(d,^{3}He)^{147}Pr		−3726	40	−3726	3	0.0	R			KVI		79Sa.A	
^{148}Nd(d,t)^{147}Nd		−1072	4	−1075.2	1.7	−0.8	1	17	17 ^{148}Nd	McM		77St22	
^{147}Sm(n,γ)^{148}Sm		8139.8	1.2	8141.29	0.27	1.2	U					69Re04	Z
		8141.1	1.5			0.1	U					70Bu19	Z
		8141.8	0.8			−0.6	–					71Gr37	Z
		8141.3	0.3			−0.0	–			Bdn		06Fi.A	
^{147}Sm(d,p)^{148}Sm		5920	10	5916.72	0.27	−0.3	U			Tal		64Ke03	
^{148}Sm(d,t)^{147}Sm		−1890	15	−1884.06	0.27	0.4	U			Kop		67Ve04	
^{147}Sm(n,γ)^{148}Sm	ave.	8141.36	0.28	8141.29	0.27	−0.3	1	93	71 ^{147}Sm			average	
^{148}Gd(p,d)^{147}Gd		−6755	5	−6759.2	1.2	−0.8	U					86Ru04	
^{148}Gd(p,d)^{147}Gd−^{148}Sm()^{147}Sm		−842	2	−842.5	1.2	−0.2	–					86Ru04	
^{148}Gd(d,t)^{147}Gd−^{148}Sm()^{147}Sm		−843	2			0.3	–					86Ru04	
^{148}Gd(^{3}He,α)^{147}Gd−^{148}Sm()^{147}Sm		−842	3			−0.2	–					86Ru04	
^{148}Gd(p,d)^{147}Gd−^{148}Sm()^{147}S	ave.	−842.4	1.3			−0.1	1	83	79 ^{147}Gd			average	
^{148}Ba(β^-)^{148}La		5115	60	5154	3	0.6	U			Bwg		90Gr10	
^{148}La(β^-)^{148}Ce		7310	140	7692	9	2.7	U			Trs		82Br23	*
		7255	55			7.9	B			Bwg		90Gr10	
		7650	100			0.4	U			Kur		02Sh.B	
		7732	70			−0.6	U			Kur		09Ha.B	
^{148}Ce(β^-)^{148}Pr		2060	75	2091	19	0.4	U			Bwg		87Gr.A	
		2140	14			−3.5	B			Kur		95Ik03	
^{148}Pr(β^-)^{148}Nd		4800	200	4928	17	0.6	U					79Ik06	
		4965	100			−0.4	U			Bwg		87Gr.A	
		4890	50			0.8	–					88Ka14	
		4880	30			1.6	–			Kur		95Ik03	
		4930	100			−0.0	U			Kur		02Sh.B	
	ave.	4883	26			1.8	1	44	43 ^{148}Pr			average	
^{148}Pm(β^-)^{148}Sm		2480	15	2470	6	−0.7	R					62Sc04	*
		2475	30			−0.2	U					63Ba31	*
^{148}Eu(β^+)^{148}Sm		3122	30	3040	10	−2.7	U					63Ba32	*
		3150	30			−3.7	B					70Ag01	*
^{148}Tb(β^+)^{148}Gd		5630	80	5738	12	1.4	F					76Cr.B	*
		5835	70			−1.4	U					83Ve06	*
		5710	100			0.3	U			Got		85Sc09	*
		5390	100			3.5	B			Got		85Ti01	*
		5760	80			−0.3	U			IRS		93Al03	*
		5752	40			−0.3	1	9	9 ^{148}Tb	GSI		95Ke05	*
^{148}Dy(β^+)^{148}Tb		2660	60	2675	9	0.3	U					81Sc21	
		2805	60			−2.2	U					81Sp03	*
		2700	60			−0.4	U			IRS		82Al.A	
		2700	60			−0.4	U					82Ve.A	
		2722	60			−0.8	U					83Ve06	*
		2835	95			−1.7	U					84Ha.B	*
		2740	60			−1.1	U			Got		85Sc09	*
		2682	10			−0.7	1	79	73 ^{148}Tb	GSI		95Ke05	*
^{148}Ho^{m}(β^+)^{148}Dy		9400	250	10120#	130#	2.9	B			IRS		93Al03	
*^{148}Ba−u	Represents frequency ratio $^{148}Ba^{++}/(C_6H_6)^+$ =0.947751100(10)											HWJ208	**
*^{148}Ce ^{40}Ar$_2$−^{152}Pr$_{1.500}$	Reported $M-A$=–140456(39) keV, author confirmed mistake in the unc.											24Ki16	**
*^{148}Pr−u	D_M=–77739.3(30.6) μu for mixture gs+m at 76.80; $M-A$=–72413.7(28.5) keV											Nub266	**
*^{148}Pr−u	Corrected for unresolved isomeric state using exc. energy in 23Ga02											25Sp03	**
*^{148}Tb−u	$M-A$=–70462(28) keV for mixture gs+m at 90.1 keV											Nub269	**
*^{148}Ho−u	$M-A$=–57815(30) keV for mixture gs+m at 250#100 keV											Nub269	**
*	outweighed by next item before correcting for isomeric mixture											GAu	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value	Adjusted value	v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{148}$Ho−^{85}Rb$_{1.741}$	D_M=91517.5(9.5) μu for mixture gs+m at 250#100 keV with R=0.74(15)								Nub269	**
$*^{148}$Sm(α)^{144}Nd	Using Q_α(^{147}Sm)=2310.5(0.5) as calibrant								16Ca43	**
$*^{148}$La(β^-)^{148}Ce	E_{β^-}=5862(100) supposed to go to levels around E=1450(100) keV								90Gr10	**
$*^{148}$Pm(β^-)^{148}Sm	E_{β^-}=2460(20) 1020(15) to ground state, 1^- level at 1465.137 keV								Ens144	**
$*^{148}$Pm(β^-)^{148}Sm	E_{β^-}=2480(30) 1930(30) 1020(30) to ground state, 2^+ at 550.255, 1^- at 1465.137 keV								Ens144	**
*	and E_{β^-}=400(30) from ^{148}Pm^m at 137.9 to 6^+ level at 2194.061 keV								Nub269	**
$*^{148}$Eu(β^+)^{148}Sm	E_{β^+}=920(30) to 1180.261 keV 4^+ level								Ens144	**
$*^{148}$Eu(β^+)^{148}Sm	E_{β^+}=540(30) to 1594.247 keV 5^- level, and other E_{β^+}								Ens144	**
$*^{148}$Tb(β^+)^{148}Gd	E_{β^+}=4610(80) assumed to ground state								76Cr.B	**
*	F : since ^{148}Tb ground state 2^-, transition to ^{148}Gd ground state weak								AHW	**
$*^{148}$Tb(β^+)^{148}Gd	E_{β^+}=2210(70) from ^{148}Tb^m at 90.1 to 8^+ level at 2693.35 keV and								Nub269	**
*	E_{β^+}=4560(80) mainly to 2^+ at 784.433 keV. Conflicting, not used								Ens144	**
$*^{148}$Tb(β^+)^{148}Gd	p^+ =0.271(0.10) → E_{β^+}=1920(30) from ^{148}Tb^m at 90.1 to 8^+ at 2693.35 keV								Ens144	**
*	but assuming 5(5)% side-feeding; see reference								90Sa32	**
$*^{148}$Tb(β^+)^{148}Gd	KL/β^+=1.54(0.09) to 1863.42 level; yields Q_{β^+}=5295(45) keV								85Ti01	**
*	but assuming 7(7)% side-feeding; see 1990Sa32								AHW	**
$*^{148}$Tb(β^+)^{148}Gd	Q_{β^+}=5700(80); and 5910(80) from ^{148}Tb^m at 90.1 keV								Nub269	**
$*^{148}$Tb(β^+)^{148}Gd	Q_{β^+}=5750(40); and 5846(50) from ^{148}Tb^m at 90.1 keV								Nub269	**
$*^{148}$Dy(β^+)^{148}Tb	p^+ =0.069(0.014) to 1^+ level at 620.24 keV, recalculated Q								Ens144	**
$*^{148}$Dy(β^+)^{148}Tb	E_{β^+}=1040(60), 1120(60) to 1^+ level at 620.24 keV								Ens144	**
$*^{148}$Dy(β^+)^{148}Tb	p^+ =0.055(0.015) to 1^+ level at 620.24 keV								Ens144	**
$*^{148}$Dy(β^+)^{148}Tb	β^+/K=0.045(0.005) to 1^+ level at 620.24 keV, gives Q_{β^+}=2680(30) keV								Ens144	**
$*^{148}$Dy(β^+)^{148}Tb	GSI average of E_{β^+}=1043(10) and 1036(10) of reference								91Ke11	**
*	to 1^+ level at 620.24 keV								Ens144	**
^{149}Cs−^{149}Sm	36211	23				2			TR1 1.0	26Ye.A *
^{149}Ba−u	−57027	188	−56715.2	2.6	0.7	U			GT3 2.5	16Kn03
	−56716.0	2.7			0.3	2			CP2 1.0	18Or.A *
	−56706.2	9.0			−1.0	2			RKM 1.0	26Ki04
^{149}Ba−^{149}Sm	26116	45	26095.7	2.8	−0.5	U			TR1 1.0	26Ye.A *
^{149}La−^{133}Cs$_{1.120}$	41493.7	3.1	41255.6	1.6	−76.8	B			JY2 1.0	25Ja05
^{149}La−u	−64637.8	5.4	−64638.2	1.6	−0.1	2			RKM 1.0	26Ki04
	−64638.2	1.7			0.0	2			CP2 1.0	26Li.A *
^{149}Ce−u	−71573.1	11.0	−71577.5	1.3	−0.4	2			CP1 1.0	06Sa56
	−71570.0	15.0			−0.5	U			GR1 1.0	25Sp03
	−71577.6	1.3			0.0	2			CP2 1.0	26Li.A *
^{149}Pr−u	−76263.9	10.6	−76282.6	1.4	−1.8	U			CP1 1.0	06Sa56
	−76252.8	14.0			−2.1	B			GR1 1.0	25Sp03
	−76285.5	5.6			0.5	2			RKM 1.0	26Ki04
	−76282.4	1.4			−0.1	2			CP2 1.0	26Li.A *
^{149}Nd−u	−79806.2	25.8	−79846.9	2.1	−1.6	U			GR1 1.0	25Sp03
	−79843	14			−0.3	U			RKM 1.0	26Ki04
C_8 ^{13}C H_8 O_2−^{149}Sm	138597	29	138595.3	1.0	−0.0	U			R04 4.0	64De15
C_9 H_{11} N O−^{149}Sm	166820	33	166874.9	1.0	0.4	U			R04 4.0	64De15
C_8 ^{13}C H_{10} N O−^{149}Sm	162408	46	162404.7	1.0	−0.0	U			R04 4.0	64De15
^{149}Eu−^{133}Cs$_{1.120}$	23849	17	23829	4	−1.2	U			MA5 1.0	00Be42
^{149}Tb−u	−76730	32	−76748	4	−0.6	U			GS2 1.0	05Li24 *
^{149}Dy−^{133}Cs$_{1.120}$	33278	109	33216	9	−0.6	U			MA5 1.0	00Be42
^{149}Dy−u	−72698	30	−72678	9	0.7	U			GS2 1.0	05Li24 *
^{149}Ho−u	−66179	34	−66172	12	0.2	1	13	13 ^{149}Ho	GS2 1.0	05Li24 *
^{149}Er^m−u	−56898	30	−56750	26	4.9	B			GS2 1.0	05Li24
^{149}Er^m−^{149}Dy	15928	24				2			TR1 1.0	25Ko14
^{149}Eu−^{142}Sm$_{1.049}$	6909	18	6882	4	−1.5	U			MA7 1.0	01Bo59
^{149}Dy−^{142}Sm$_{1.049}$	16249	16	16269	9	1.3	1	34	34 ^{149}Dy	MA7 1.0	01Bo59
^{149}Sm ^{35}Cl−^{147}Sm ^{37}Cl	5257	4	5236.8	0.9	−1.3	U			H12 4.0	64Ba15
	5231	3			0.8	U			H21 2.5	70Ma05
	5239.8	0.8			−1.5	1	22	16 ^{147}Sm	M21 2.5	75Ka25

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{149}Er−^{149}Dy		15147	49	15132	24	−0.3	U			TR1	1.0	25Ko14	
^{149}Tm−^{149}Dy		25638	54				2			TR1	1.0	25Ko14	*
^{149}Sm−^{148}Sm		2282	31	2361.8	0.9	0.6	U			R04	4.0	64De15	
^{149}Sm−^{147}Sm		2320	60	2286.6	0.9	−0.1	U			R04	4.0	64De15	
^{149}Gd(α)^{145}Sm		3102.3	10.	3099	3	−0.3	–					65Ma51	Z
		3096.1	10.3			0.3	–			ORa		66Wi12	Z
		3099.1	5.			0.1	–			Dba		67Go32	Z
	ave.	3099	4			0.1	1	56	53 ^{149}Gd			average	
^{149}Tb(α)^{145}Eu		4074.4	3.	4077.9	2.2	1.2	–			Dba		67Go32	Z
		4073.8	7.			0.6	U			ORa		74To07	*
		4074.6	10.			0.3	U					81Ho.A	Z
		4081.8	5.			−0.8	–			Bka		82Bo04	Z
		4082.8	4.			−1.2	–			Daa		96Pa01	
	ave.	4078.2	2.2			−0.1	1	95	86 ^{149}Tb			average	
^{149}Sm(n,α)^{146}Nd		9429	4	9436.0	0.9	1.7	U			McM		67Oa01	
		9421	15			1.0	U			ILL		75Em.A	
^{149}Sm(p,t)^{147}Sm		−5532	8	−5530.8	0.9	0.1	U			Min		72De47	
		−5532	7			0.2	U			McM		73Ga04	
^{148}Nd(n,γ)^{149}Nd		5038.76	0.10	5038.79	0.07	0.3	2			ILn		76Pi04	Z
		5038.82	0.11			−0.3	2			Bdn		06Fi.A	
^{148}Nd(^{3}He,d)^{149}Pm		455	5	452.4	2.5	−0.5	1	24	13 ^{149}Pm	McM		80St10	*
^{149}Sm(d,^{3}He)^{148}Pm		−2064	6	−2066	6	−0.3	2					88No02	
^{148}Sm(n,γ)^{149}Sm		5872.5	1.8	5871.4	0.9	−0.6	1	23	16 ^{148}Sm			70Sm.A	
		5850.8	0.6			34.3	C					82Ba15	
^{149}Sm(γ,n)^{148}Sm		−5890	160	−5871.4	0.9	0.1	U			Phi		60Ge01	
^{148}Sm(d,p)^{149}Sm		3656	15	3646.8	0.9	−0.6	U			Kop		67Ve04	
^{149}Er(εp)^{148}Dy		5758	900	6965	25	1.3	U					83La.A	*
		7080	470			−0.2	U			LBL		89Fi01	
^{149}Lu(p)^{148}Yb		1922.9	20.1				3					22Au01	
^{149}La(β^-)^{149}Ce		6450	200	6464.0	1.9	0.1	U			Kur		02Sh.B	
^{149}Ce(β^-)^{149}Pr		4190	75	4382.7	1.7	2.6	U			Bwg		87Gr.A	
		4380	60			0.0	U			Kur		95Ik03	
		4310	100			0.7	U			Kur		02Sh.B	
^{149}Pr(β^-)^{149}Nd		3000	200	3320.1	2.3	1.6	U					67Va14	
		3390	90			−0.8	U			Kur		95Ik03	
^{149}Nd(β^-)^{149}Pm		1669	10	1689.4	2.5	2.0	U					64Go08	*
^{149}Pm(β^-)^{149}Sm		1072	2	1071.6	1.9	−0.2	1	88	87 ^{149}Pm			60Ar05	
		1062	2			4.8	B					78Re01	
^{149}Eu(ε)^{149}Sm		680	10	695	4	1.5	1	14	14 ^{149}Eu			85Ad.A	
^{149}Gd(ε)^{149}Eu		1308	6	1314	4	1.1	1	47	30 ^{149}Eu	Got		84Sc.B	
^{149}Tb(β^+)^{149}Gd		3575	50	3638	4	1.3	U			Got		85Sc09	*
		3635	10			0.3	1	19	10 ^{149}Tb	GSI		91Ke06	*
^{149}Dy(β^+)^{149}Tb		3930	150	3792	9	−0.9	U					84Al36	*
		3925	65			−2.0	U			Got		90Sa32	*
		3797	13			−0.4	1	46	42 ^{149}Dy	GSI		91Ke11	*
		3950	100			−1.6	U			IRS		93Al03	
^{149}Ho(β^+)^{149}Dy		6043	50	6060	11	0.3	–			IRS		83Al06	*
		5330	100			7.3	C					84Ha.B	
		6000	90			0.7	U			IRS		93Al03	
		6009	20			2.6	–			GSI		91Ke11	*
	ave.	6014	19			2.5	1	37	26 ^{149}Ho			average	
^{149}Er(ε)^{149}Ho		8610	650	8035	25	−0.9	U			LBL		89Fi01	*
^{149}Er^m(IT)^{149}Er		741.69	0.23				3					Ens228	
*^{149}Cs−^{149}Sm	from frequency ratio ^{149}Cs$^+$/(^{149}Sm)$^+$ =1.000243163(153)											26Ye.A	**
*^{149}Ba−u	Represents frequency ratio ^{149}Ba^{++}/(C$_6$H$_6$)$^+$ =0.954189968(17)											HWJ208	**
*^{149}Ba−^{149}Sm	from frequency ratio ^{149}Ba$^+$/(^{149}Sm)$^+$ =1.000175372(304)											26Ye.A	**
*^{149}La−u	derived from frequency ratio (C$_6$H$_6$)$^+$/(^{149}La)$^{++}$ =0.954139215(11)											26Li.A	**
*^{149}Ce−u	derived from frequency ratio (C$_6$H$_6$)$^+$/(^{149}Ce)$^{++}$ =0.9540947582(84)											26Li.A	**
*^{149}Pr−u	derived from frequency ratio (C$_6$H$_6$)$^+$/(^{149}Pr)$^{++}$ =0.9540646172(89)											26Li.A	**
*^{149}Tb−u	$M-A$=–71456(28) keV for mixture gs+m at 35.75 keV											Nub269	**
*^{149}Dy−u	$M-A$=–65057(28) keV for ^{149}Dy^m at 2660.94(0.34) keV											Nub269	**
*^{149}Ho−u	$M-A$=–61621(28) keV for mixture gs+m at 48.80 keV											Nub269	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value	Adjusted value	v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{149}$Tm−^{149}Dy	frequency ratio was corrected for the isomer by the author								25Ko14	**
$*^{149}$Tb(α)^{145}Eu	E_α=3999(7) from ^{149}Tb^m at 35.75 keV								Nub269	**
$*^{148}$Nd(^{3}He,d)^{149}Pm	Based on ^{146}Nd(^{3}He,d)^{147}Pm Q =–87.6(0.9) keV								AHW	**
$*^{149}$Er(εp)^{148}Dy	As read from graph; Q =6500 from ^{149}Er^m at 741.69 keV								Nub269	**
$*^{149}$Nd(β^-)^{149}Pm	E_{β^-}=1555(10) to 5/2$^+$ level at 114.312 level								Ens046	**
$*^{149}$Tb(β^+)^{149}Gd	β^+/K=0.31(0.03) from ^{149}Tb^m at 35.78 to 9/2$^-$ level at 795.82 keV								Ens046	**
$*^{149}$Tb(β^+)^{149}Gd	E_{β^+}=1853(10) from ^{149}Tb^m at 35.78 to 9/2$^-$ level at 795.82 keV								Ens046	**
$*^{149}$Dy(β^+)^{149}Tb	E_{β^+}=1030(150) to 1728.31−1876.96 levels								Ens046	**
$*^{149}$Dy(β^+)^{149}Tb	KL/β^+=29.4(10.6), 14.5(3.7), 17.6(4.9) to 1883, 1735, 1728 levels								90Sa32	**
$*^{149}$Dy(β^+)^{149}Tb	Original Q =3812(10) from E_{β^+}=1965(10) to 825.16 level corrected								GAu	**
*	to E_{β^+}=1950(13) for background subtraction								GAu	**
$*^{149}$Ho(β^+)^{149}Dy	E_{β^+}=3930(50) to 9/2$^-$ level at 1090.76 keV								Ens046	**
$*^{149}$Ho(β^+)^{149}Dy	E_{β^+}=3896(20) to 9/2$^-$ level at 1090.76 keV								Ens046	**
$*^{149}$Er(ε)^{149}Ho	KLM/β^+=0.68(0.34) from ^{149}Er^m at 741.8 to 4699.7 level								Ens046	**
^{150}Cs−^{150}Sm	41650	25			2			TR1	1.0	26Ye.A *
^{150}Ba−u	−55309	371	−53559	6	1.9	U		GT3	2.5	16Kn03
	−53558.9	6.1			2			CP2	1.0	18Or.A *
^{150}Ba−^{150}Sm	29156	32	29161	6	0.2	U		TR1	1.0	26Ye.A *
^{150}La−u	−60258	187	−60453.8	2.0	−0.4	U		GT3	2.5	16Kn03
	−60452.5	2.7			−0.5	1	56	56 ^{150}La	CP2	1.0	18Or.A *
^{150}La−^{133}Cs$_{1.128}$	46199.6	3.7	46196.4	2.0	−0.9	1	30	30 ^{150}La	JY2	1.0	25Ja05
^{150}La ^{1}H−^{151}Pr	19047.9	5.3	19059.5	2.2	2.2	1	18	14 ^{150}La	RI1	1.0	24Ki16 *
^{150}Ce−u	−69618.6	13.1	−69637.3	2.2	−1.4	U			CP1	1.0	06Sa56
	−69614.0	16.1			−1.4	U			GR1	1.0	25Sp03
^{150}Ce−^{12}C$_6$ ^{1}H$_6$	−116585.3	3.1	−116587.4	2.2	−0.7	1	52	52 ^{150}Ce	CP2	1.0	24Ra.A
^{150}Ce ^{1}H−^{151}Pr	9874.0	3.0	9876.0	2.2	0.7	1	55	48 ^{150}Ce	RI1	1.0	24Ki16 *
^{150}Pr−u	−73322.9	10.6	−73332.5	1.3	−0.9	U			CP1	1.0	06Sa56
	−73312.3	15.0			−1.3	U			GR1	1.0	25Sp03
C$_{12}$ H$_6$−^{150}Nd	126194	43	126051.1	0.9	−0.8	U			R05	4.0	65De13
C$_8$ ^{13}C N O H$_{11}$−^{150}Nd	166439	34	166519.7	0.9	0.6	U			R05	4.0	65De13
C$_9$ N O H$_{12}$−^{150}Nd	170931	46	170989.9	0.9	0.3	U			R05	4.0	65De13
^{150}Nd−u	−79073.0	19.3	−79100.9	0.9	−1.4	U			GR1	1.0	25Sp03
C$_{12}$ H$_6$−^{150}Sm	129810	140	129670.4	0.9	−0.2	U			R04	4.0	64De15
C$_8$ ^{13}C H$_{11}$ N O−^{150}Sm	170029	25	170139.0	0.9	1.1	U			R04	4.0	64De15
C$_9$ H$_{12}$ N O−^{150}Sm	174612	47	174609.2	0.9	−0.0	U			R04	4.0	64De15
^{150}Tb^m−u	−75850	30	−75841	28	0.3	1	89	89 ^{150}Tb^m	GS2	1.0	05Li24
^{150}Ho−^{133}Cs$_{1.128}$	40150	29	40152	14	0.1	–			MA5	1.0	00Be42
ave.	40132	21			1.0	1	43	43 ^{150}Ho			average
^{150}Ho−u	−66504	40	−66498	14	0.1	U			GS2	1.0	05Li24 *
^{150}Er−u	−62060	30	−62077	13	−0.6	1	20	20 ^{150}Er	GS2	1.0	05Li24
^{150}Nd ^{35}Cl$_2$−^{146}Nd ^{37}Cl$_2$	13654	9	13678.6	1.0	1.1	U			H21	2.5	70Ma05
	13672.5	1.8			1.3	U			H25	2.5	72Ba08
^{150}Nd ^{35}Cl−^{148}Nd ^{37}Cl	7006	4	6951.7	2.0	−3.4	B			H12	4.0	64Ba15
	6939	4			1.3	U			H21	2.5	70Ma05
^{150}Sm ^{35}Cl−^{148}Sm ^{37}Cl	5452	8	5402.6	0.9	−1.5	U			H12	4.0	64Ba15
	5400	4			0.3	U			H21	2.5	70Ma05
	5404.8	0.6			−1.4	1	37	28 ^{148}Sm	M21	2.5	75Ka25
^{150}Pr−^{150}Nd	5768.4	0.9				2			CP2	1.0	24Ra.A
^{150}Nd−^{150}Sm	3633	4	3619.28	0.21	−0.9	U			H19	4.0	64Mc11
	3617.0	1.2			0.8	U			H25	2.5	72Ba08
	3619.33	0.21			−0.3	1	99	68 ^{150}Nd	JY1	1.0	10Ko28
^{150}Er−^{150}Dy	12349	24	12332	13	−0.7	1	31	29 ^{150}Er	TR1	1.0	25Ko14
^{150}Tm−^{150}Dy	24490	93				2			TR1	1.0	25Ko14 *
	24520	110	24490	90	−0.3	U			TR1	1.0	25Ko14 *

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{150}Yb−^{150}Dy		32931	48				2			TR1	1.0	21Be23	*
^{150}Sm−^{149}Sm		149	30	90.8	0.4	−0.5	U			R04	4.0	64De15	
^{150}Nd−^{148}Nd		3860	46	4001.5	2.0	0.8	U			R05	4.0	65De13	
		3988	3			1.8	U			M17	2.5	66Be10	
^{150}Sm−^{148}Sm		2430	50	2452.5	0.9	0.1	U			R04	4.0	64De15	
^{150}Nd−^{146}Nd		7719	67	7778.3	1.0	0.2	U			R05	4.0	65De13	
^{150}Gd(α)^{146}Sm		2804.9	10.	2808	6	0.2	–					62Si14	
		2792.6	18.			0.8	–					65Og01	
	ave.	2802	9			0.6	1	45	39 ^{150}Gd			average	
^{150}Tb(α)^{146}Eu		3585.5	5.	3587	5	0.3	1	92	80 ^{150}Tb	Dba		67Go32	Z
^{150}Dy(α)^{146}Gd		4345.8	5.	4351.2	1.5	1.1	–			Dba		67Go32	Z
		4349.5	5.			0.3	–			GSa		79Ho10	Z
		4351.3	3.			−0.0	–			Bka		82Bo04	*
		4352.5	2.1			−0.6	–			Ora		82De11	Z
	ave.	4351.3	1.5			−0.1	1	99	90 ^{150}Dy			average	
^{148}Nd(t,p)^{150}Nd		3935	30	3933.4	1.9	−0.1	U			Ald		72Ch11	
^{148}Sm(t,p)^{150}Sm		5372	25	5376.3	0.9	0.2	U			Ald		66Bj01	
^{150}Sm(p,t)^{148}Sm		−5379	8	−5376.3	0.9	0.3	U			Min		72De47	
		−5378	15			0.1	U			Ham		74Oe03	
^{150}Nd(d,^{3}He)^{149}Pr		−4501	10	−4420.7	1.5	8.0	C			KVI		79Sa.A	
^{150}Nd(d,t)^{149}Nd		−1122	10	−1119.2	1.9	0.3	U			McM		73Bu02	
^{149}Sm(n,γ)^{150}Sm		7984.9	0.6	7986.8	0.4	3.1	B					69Re04	Z
		7986.7	1.5			0.1	–					70Bu19	Z
		7986.7	0.4			0.2	–			Bdn		06Fi.A	
^{149}Sm(d,p)^{150}Sm		5764	4	5762.2	0.4	−0.4	U			Tal		64Ke03	
^{150}Sm(d,t)^{149}Sm		−1738	15	−1729.5	0.4	0.6	U			Kop		67Ve04	
^{149}Sm(n,γ)^{150}Sm	ave.	7986.7	0.4	7986.8	0.4	0.2	1	95	85 ^{149}Sm			average	
^{150}Lu(p)^{149}Yb		1269.6	4.	1269.6	2.3	−0.0	3					84Ho.A	
		1269.6	4.			−0.0	3			Dap		93Se04	
		1269.6	4.			−0.0	3			ORp		03Gi10	
^{150}Lu^m(p)^{149}Yb		1303.8	15.	1291	5	−0.8	o			ORp		00Gi01	
		1285.6	8.			0.7	3			ORp		03Gi10	
		1294.7	6.			−0.5	3			Arp		03Ro21	
^{150}Ce(β^-)^{150}Pr		3010	90	3442.1	2.4	4.8	C			Bwg		87Gr.A	
		3480	40			−0.9	U			Kur		95Ik03	
^{150}Pr(β^-)^{150}Nd		5690	80	5373.2	0.8	−4.0	C			Bwg		87Gr.A	
		5386	26			−0.5	U			Kur		95Ik03	
		5290	100			0.8	U			Kur		02Sh.B	
^{150}Pm(β^-)^{150}Sm		3454	20				2					77Ho09	
^{150}Eu(β^+)^{150}Sm		2222	25	2259	6	1.5	U					65Gu03	*
^{150}Eu(β^-)^{150}Gd		978	10	972	4	−0.6	–					63Yo07	*
		968	4			0.9	–					65Gu03	*
	ave.	969	4			0.6	1	91	53 ^{150}Eu			average	
^{150}Tb(β^+)^{150}Gd		4720	40	4658	8	−1.5	U					68Wi21	
		4670	15			−0.8	1	31	20 ^{150}Tb			76Cr.B	
		4760	50			−2.0	U					77Ha31	*
		4620	60			0.6	U					83Ve06	
^{150}Tb^m(β^+)^{150}Gd		5040	100	5120	27	0.8	U			IRS		93Al03	
^{150}Dy(β^+)^{150}Tb		1760	40	1796	8	0.9	U					81Ka07	*
^{150}Ho(β^+)^{150}Dy		6980	150	7369	13	2.6	U					84Al36	*
		6560	100			8.1	B			IRS		93Al03	
^{150}Ho(ε)^{150}Dy		7400	200			−0.2	U					98Ag.A	
		7372	27			−0.1	1	23	21 ^{150}Ho			00Ca.A	
		7444	126			−0.6	U					01Ro35	
^{150}Ho^m(β^+)^{150}Dy		7360	50				2			IRS		83Al06	*
		6575	75	7360	50	10.5	C					84Ha.B	*
		6625	120			6.1	B			Got		85Sc09	*
		6900	130			3.5	B			Got		90Sa32	*
		7060	80			3.7	C			IRS		93Al03	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{150}Er(β^+)^{150}Ho	4010	80	4118	12	1.4	o					82No08	*
	4105	75			0.2	U					84Ha.B	*
	4108	15			0.7	1	68	36 ^{150}Ho	GSI		91Ke11	*
*^{150}Cs−^{150}Sm	from frequency ratio $^{150}Cs^+/(^{150}Sm)^+$ =1.000277818(167)										26Ye.A	**
*^{150}Ba−u	Represents frequency ratio $^{150}Ba^{++}/(C_6H_6)^+$ =0.960616639(39)										HWJ208	**
*^{150}Ba−^{150}Sm	from frequency ratio $^{150}Ba^+/(^{150}Sm)^+$ =1.000194481(211)										26Ye.A	**
*^{150}La−u	Represents frequency ratio $^{150}La^{++}/(C_6H_6)^+$ =0.960572476(17)										HWJ208	**
*^{150}La ^{1}H−^{151}Pr	Reported $M-A$=–49037(24) keV, author confirmed mistake in the unc.										24Ki16	**
*^{150}Ce ^{1}H−^{151}Pr	Reported $M-A$=–57582(24) keV, author confirmed mistake in the unc.										24Ki16	**
*^{150}Ho−u	$M-A$=–61948(28) keV for mixture gs+m at –0(50) keV										Nub211	**
*^{150}Tm−^{150}Dy	frequency ratio was corrected for the isomer by the author										25Ko14	**
*^{150}Tm−^{150}Dy	frequency ratio was corrected for the isomer by the author										25Ko14	**
*^{150}Dy(α)^{146}Gd	Recalibrated as in reference										91Ry01	**
*^{150}Eu(β^+)^{150}Sm	E_{β^+}=1242(25) from ^{150}Eu^m at 41.7 keV										Nub269	**
*^{150}Eu(β^-)^{150}Gd	Q_{β^-}=1020(10) 1010(4) respectively, from ^{150}Eu^m at 41.7 keV										Nub269	**
*^{150}Tb(β^+)^{150}Gd	E_{β^+}=1655(80) to 2^+ at 2091.623, and other E_{β^+}. Orig. error 35 increased										Ens136	**
*^{150}Dy(β^+)^{150}Tb	p^+ =2(10)×10^{-3} to 1^+ 397.2 level combined with lower limit through ^{146}Gd(ε)										Ens136	**
*^{150}Ho(β^+)^{150}Dy	E_{β^+}=4550(150) to 1395.0 and 1456.8 levels										82No08	**
*^{150}Ho^m(β^+)^{150}Dy	E_{β^+}=3940(50) to 8^+ level at 2401.6 keV										Ens136	**
*^{150}Ho^m(β^+)^{150}Dy	p^+ =0.56(0.02) 0.58(0.07) respectively to 8^+ level at 2401.6 keV										Ens136	**
*^{150}Ho^m(β^+)^{150}Dy	Q_{β^+}=6819(+117–100) from p^+ to 2401.6 level; could be raised 140										90Sa32	**
*	if 4% feeding of higher Dy levels										90Sa32	**
*^{150}Er(β^+)^{150}Ho	p^+ =0.36(0.04) 0.39(0.04) to 1^+ level at 476.16 keV										Ens136	**
*^{150}Er(β^+)^{150}Ho	E_{β^+}=2610(15) to 1^+ level at 476.16 keV										Ens136	**
^{151}Cs−^{151}Pm	41789	28				2			TR1	1.0	26Ye.A	*
^{151}Ba−^{151}Pm	30493	23				2			TR1	1.0	26Ye.A	*
^{151}La−u	−58734	397	−57465.0	1.6	2.1	C			GT1	1.5	04Ma.A	
	−57231	187			−0.5	U			GT3	2.5	16Kn03	
^{151}La−^{133}Cs$_{1.135}$	49850.5	3.1	49847.0	1.6	−1.1	1	27	27 ^{151}La	JY2	1.0	25Ja05	
^{151}La−^{12}C$_6$ ^{1}H$_6$	−104416.2	1.9	−104415.2	1.6	0.5	1	71	71 ^{151}La	CP2	1.0	24Ra.A	
^{151}La ^{1}H−^{152}Pr	18814.8	9.4	18823	3	0.9	1	12	10 ^{152}Pr	RI1	1.0	24Ki16	*
^{151}Ce−u	−65727.8	19.0	−65721.7	1.2	0.3	U			CP1	1.0	06Sa56	
	−65654.7	34.4			−1.9	U			GR1	1.0	25Sp03	
^{151}Ce−^{12}C$_6$ ^{1}H$_6$	−112671.5	1.4	−112671.9	1.2	−0.3	1	70	70 ^{151}Ce	CP2	1.0	24Ra.A	
^{151}Ce−^{133}Cs$_{1.135}$	41589.9	2.9	41590.4	1.2	0.2	1	16	16 ^{151}Ce	JY2	1.0	25Ja05	
^{151}Ce ^{1}H−^{152}Pr	10560.1	3.0	10567	3	2.2	B			RI1	1.0	24Ki16	*
^{151}Ce ^{40}Ar$_2$−^{154}Nd$_{1.500}$	−35333.1	12.0	−35350.3	2.0	−1.4	U			RI1	1.0	24Ki16	*
^{151}Pr−u	−71697.5	14.3	−71688.2	1.1	0.6	U			CP1	1.0	06Sa56	
	−71654.8	22.5			−1.5	U			GR1	1.0	25Sp03	
^{111}Ru ^{40}Ar−^{151}Pr	−48347.0	6.9	−48354	5	−1.0	1	45	44 ^{111}Ru	RI1	1.0	24Ki16	*
^{111}Pd ^{40}Ar−^{151}Pr	−58199	59	−58238.2	1.4	−0.7	U			RI1	1.0	24Ki16	*
^{151}Pr−^{12}C$_6$ ^{1}H$_6$	−118639.2	1.3	−118638.4	1.1	0.6	1	74	74 ^{151}Pr	CP2	1.0	24Ra.A	
^{151}Pr ^{40}Ar$_2$−^{154}Nd$_{1.500}$	−41303.0	10.4	−41316.9	1.9	−1.3	U			RI1	1.0	24Ki16	*
^{151}Nd−^{12}C$_6$ ^{1}H$_6$	−123115.6	1.7	−123113.1	0.9	1.5	1	30	30 ^{151}Nd	CP2	1.0	24Ra.A	
^{151}Nd ^{40}Ar$_2$−^{154}Nd$_{1.500}$	−45792.5	12.7	−45791.5	1.8	0.1	U			RI1	1.0	24Ki16	*
^{151}Nd−u	−76146.5	24.7	−76162.9	0.9	−0.7	U			GR1	1.0	25Sp03	
^{151}Pm−^{12}C$_6$ ^{1}H$_6$	−125747.9	1.5	−125747.1	1.4	0.5	1	93	93 ^{151}Pm	CP2	1.0	24Ra.A	
^{151}Pm−u	−78765.9	56.9	−78796.9	1.4	−0.5	U			GR1	1.0	25Sp03	
C$_{12}$ H$_7$−^{151}Eu	134920	37	134920.6	0.9	0.0	U			R04	4.0	64De15	
C$_{10}$ H$_{15}$ O−^{151}Eu	192490	70	192435.4	0.9	−0.2	U			R04	4.0	64De15	
^{151}Eu−^{85}Rb$_{1.776}$	76520	15	76516.1	0.9	−0.3	U			MA5	1.0	00Be42	
^{151}Tb−u	−76866	43	−76893	4	−0.6	U			GS2	1.0	05Li24	*
^{151}Dy−u	−73809	30	−73811	3	−0.1	U			GS2	1.0	05Li24	
^{151}Ho−u	−68323	33	−68302	8	0.6	U			GS2	1.0	05Li24	*
^{151}Er−u	−62528	30	−62532	12	−0.1	–			GS2	1.0	05Li24	
	−62540	30			0.3	–			GS2	1.0	05Li24	*
ave.	−62551	18			1.1	1	45	45 ^{151}Er			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{151}Eu ^{35}Cl−^{149}Sm ^{37}Cl		5620.3	2.6	5615.7	0.4	−0.7	U			H25	2.5	72Ba08	
^{151}La−^{151}Pr		14217.2	13.4	14223.2	2.0	0.4	U			RI1	1.0	24Ki16	*
^{151}Ce−^{151}Pr		5965.5	2.9	5966.6	1.5	0.4	1	26	14 ^{151}Ce	RI1	1.0	24Ki16	*
^{151}Nd−^{151}Pr		−4472.4	2.7	−4474.6	1.5	−0.8	U			RI1	1.0	24Ki16	*
^{151}Pm−^{151}Pr		−7091.8	7.1	−7108.7	1.8	−2.4	U			RI1	1.0	24Ki16	*
^{151}Er−^{151}Dy		11277	23	11280	12	0.1	1	26	25 ^{151}Er	TR1	1.0	25Ko14	
^{151}Tm−^{151}Dy		19300	39	19309	16	0.2	1	17	16 ^{151}Tm	TR1	1.0	25Ko14	*
^{151}Yb−^{151}Er		18220	120	18190	110	−0.3	1	88	88 ^{151}Yb	TR1	1.0	21Be23	
^{151}Yb^{m}−^{151}Er		18947	50				2			TR1	1.0	21Be23	
^{151}Eu−^{150}Sm		2800	60	2574.84	0.14	−0.9	U			R04	4.0	64De15	
^{151}Eu(α)^{147}Pm		1960	30	1963.1	0.9	0.1	U					07Be48	
		1948.9	8.6			1.6	U					14Ca13	
^{151}Gd(α)^{147}Sm		2670.8	30.	2652.3	2.9	−0.6	U					65Si06	
^{151}Tb(α)^{147}Eu		3499.6	5.	3496	4	−0.7	1	58	48 ^{151}Tb	Dba		67Go32	
^{151}Dy(α)^{147}Gd		4175.5	5.	4179.0	2.6	0.7	–			Dba		67Go32	Z
		4181.1	3.			−0.7	–			Bka		82Bo04	Z
	ave.	4179.6	2.6			−0.2	1	97	96 ^{151}Dy			average	
^{151}Ho(α)^{147}Tb		4696.3	5.	4695.0	1.8	−0.3	–			GSa		79Ho10	*
		4695.8	3.			−0.3	–			Bka		82Bo04	*
		4693.8	3.			0.4	–			Ora		82De11	*
		4694.9	5.			0.0	–			Daa		96Pa01	*
	ave.	4695.0	1.8			0.0	1	100	91 ^{151}Ho			average	
^{151}Eu(n,α)^{148}Pm		7870	20	7859	6	−0.5	U			ILL		74Em01	
^{151}Sm(p,t)^{149}Sm		−5100	4	−5101.4	0.4	−0.4	U			McM		73Ga04	
^{151}Eu(p,t)^{149}Eu		−5872	5	−5873	4	−0.1	1	57	56 ^{149}Eu	Min		75Ta12	
^{150}Nd(n,γ)^{151}Nd		5334.55	0.2	5334.56	0.10	0.0	–			ILn		76Pi13	Z
		5334.55	0.11			0.1	–			Bdn		06Fi.A	
^{150}Nd(d,p)^{151}Nd		3084	15	3109.99	0.10	1.7	U			Tal		67Ne08	
^{150}Nd(n,γ)^{151}Nd	ave.	5334.55	0.10	5334.56	0.10	0.1	1	100	70 ^{151}Nd			average	
^{150}Nd(^{3}He,d)^{151}Pm		1503	5	1512.4	1.6	1.9	1	10	7 ^{151}Pm	McM		80St10	*
^{150}Sm(n,γ)^{151}Sm		5596.5	1.8	5596.44	0.11	−0.0	U					70Sm.A	*
		5596.	1.5			0.3	U					71Gr22	
		5596.42	0.20			0.1	–			ILn		86Va08	Z
		5596.44	0.13			−0.0	–			Bdn		06Fi.A	
^{150}Sm(d,p)^{151}Sm		3369	16	3371.87	0.11	0.2	U			Tal		65Ke09	
^{150}Sm(n,γ)^{151}Sm	ave.	5596.43	0.11	5596.44	0.11	0.0	1	100	50 ^{150}Sm			average	
^{151}Eu(γ,n)^{150}Eu		−8040	110	−7932	6	1.0	U			Phi		60Ge01	
^{151}Eu(p,d)^{150}Eu		−5721	9	−5708	6	1.5	1	47	47 ^{150}Eu			82So.B	
^{151}Yb(εp)^{150}Er		9000	300	9230	110	0.8	1	12	12 ^{151}Yb	ORa		86To12	*
^{151}Lu(p)^{150}Yb		1239.2	3.	1241.0	1.8	0.6	o			Dap		82Ho04	
		1241.0	2.8			−0.0	3			Dap		93Se04	
		1241.3	3.0			−0.1	3			ORp		03Gi10	
		1240.3	4.0			0.2	3			Man		15Ta12	
^{151}Lu^{m}(p)^{150}Yb		1318.8	10.	1298	3	−2.1	B			Dap		98Ba.B	
		1293.6	4.			1.0	3			Man		15Ta12	
		1303.7	5.			−1.2	3			Jyp		17Wa18	
^{151}Ce(β^-)^{151}Pr		5270	100	5557.8	1.4	2.9	U			Kur		02Sh.B	
^{151}Pr(β^-)^{151}Nd		4170	75	4168.1	1.4	−0.0	U			Bwg		90Gr10	
		4136	40			0.8	U			Ida		93Gr17	*
		4210	30			−1.4	U			Kur		95Ik03	
^{151}Nd(β^-)^{151}Pm		2510	50	2453.6	1.6	−1.1	U					73Se12	*
		2480	50			−0.5	U			Kur		95Ik03	
^{151}Pm(β^-)^{151}Sm		1195	10	1179.6	1.6	−1.5	U					64Be10	*
^{151}Sm(β^-)^{151}Eu		75.9	0.6	76.44	0.07	0.9	U					59Ac28	
		76.430	0.068			0.1	1	100	79 ^{151}Eu			22Ko10	
^{151}Gd(ε)^{151}Eu		463	3	464.2	2.8	0.4	1	86	85 ^{151}Gd			83Vo10	*
^{151}Tb(β^+)^{151}Gd		2562	5	2565	4	0.7	–					77Cr05	*
		2566	12			−0.1	–					84Sc18	*
	ave.	2563	5			0.6	1	66	52 ^{151}Tb			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{151}Ho(β^+)^{151}Dy	5080	50	5132	8	1.0	U			IRS		83Al06	*
	5100	80			0.4	U			IRS		93Al03	
^{151}Er(β^+)^{151}Ho	5130	110	5375	13	2.2	U					98Fo06	
^{151}Tm(β^+)^{151}Er	6025	145	7480	18	10.0	C					84Ha.B	*
	7074	50			8.1	F			GSI		91Ke11	*
^{151}Tm^m(IT)^{151}Tm	96.4	7.0	95	6	−0.2	o					97Da07	*
^{151}Lu^m(IT)^{151}Lu	77	5	57	4	−4.1	B			Dap		99Bi14	
*^{151}Cs−^{151}Pm	from frequency ratio ^{151}Cs$^+$/(^{151}Pm)$^+$ =1.000276896(186)										26Ye.A	**
*^{151}Ba−^{151}Pm	from frequency ratio ^{151}Ba$^+$/(^{151}Pm)$^+$ =1.000202049(154)										26Ye.A	**
*^{151}La ^{1}H−^{152}Pr	Reported $M-A$=–46256(22) keV, author confirmed mistake in the unc.										24Ki16	**
*^{151}Ce ^{1}H−^{152}Pr	Reported $M-A$=–53945(20) keV, author confirmed mistake in the unc.										24Ki16	**
*^{151}Ce ^{40}Ar$_2$−^{154}Nd$_{1.500}$	Reported $M-A$=–131282(41) keV, author confirmed mistake in the unc.										24Ki16	**
*^{111}Ru ^{40}Ar−^{151}Pr	Reported $M-A$=–111815(24) keV, author confirmed mistake in the unc.										24Ki16	**
*^{111}Pd ^{40}Ar−^{151}Pr	D_M=–58107(20) μu, $M-A$=–120906(22) for mixture of ground state and m at										Nub269	**
*^{111}Pd ^{40}Ar−^{151}Pr	172.17(0.01) keV; reported $M-A$=–120906(30), author confirmed mistake										Nub269	**
*^{111}Pd ^{40}Ar−^{151}Pr	in the uncertainty										Nub269	**
*^{151}Pr ^{40}Ar$_2$−^{154}Nd$_{1.500}$	Reported $M-A$=–136843(41) keV, author confirmed mistake in the unc.										24Ki16	**
*^{151}Nd ^{40}Ar$_2$−^{154}Nd$_{1.500}$	Reported $M-A$=–141025(42) keV, author confirmed mistake in the unc.										24Ki16	**
*^{151}Tb−u	$M-A$=–71551(28) keV for mixture gs+m at 99.53 keV										Nub269	**
*^{151}Ho−u	$M-A$=–63622(28) keV for mixture gs+m at 41.0 keV										Nub269	**
*^{151}Er−u	$M-A$=–55670(28) keV for ^{151}Er^m at 2586.0 keV										Nub269	**
*^{151}La−^{151}Pr	Reported $M-A$=–53537(26) keV, author confirmed mistake in the unc.										24Ki16	**
*^{151}Ce−^{151}Pr	Reported $M-A$=–61223(24) keV, author confirmed mistake in the unc.										24Ki16	**
*^{151}Nd−^{151}Pr	Reported $M-A$=–70946(23) keV, author confirmed mistake in the unc.										24Ki16	**
*^{151}Pm−^{151}Pr	Reported $M-A$=–73386(24) keV, author confirmed mistake in the unc.										24Ki16	**
*^{151}Tm−^{151}Dy	frequency ratio corrected for the unresolved isomer at 93(6) keV										25Ko14	**
*^{151}Ho(α)^{147}Tb	E=4523.8(5,Z) to ^{147}Tb^m at 50.6(0.9); 4610.8(5,Z) from ^{151}Ho^m 41.0(0.2)										Nub269	**
*^{151}Ho(α)^{147}Tb	E=4521.5(3,Z) to ^{147}Tb^m at 50.6(0.9); 4611.5(3,Z) from ^{151}Ho^m 41.0(0.2)										Nub269	**
*^{151}Ho(α)^{147}Tb	E=4521.2(3,Z) to ^{147}Tb^m at 50.6(0.9); 4607.2(4,Z) from ^{151}Ho^m 41.0(0.2)										Nub269	**
*^{151}Ho(α)^{147}Tb	E_α=4521(5,Z) to ^{147}Tb^m at 50.6(0.9)										Nub269	**
*^{150}Nd(^{3}He,d)^{151}Pm	Based on ^{146}Nd(^{3}He,d)^{147}Pm Q =–87.6(0.9) keV										AHW	**
*^{150}Sm(n,γ)^{151}Sm	E(γ)=5591.7(1.8) to 3/2$^-$ level at 4.821 keV										Ens091	**
*^{151}Yb(εp)^{150}Er	E_p estimated 7300(300) to levels around 1700 keV										GAu	**
*	"Statistical p's originate from 11/2$^-$ isomer."										86To12	**
*^{151}Pr(β^-)^{151}Nd	Two highest Q_{β^-}=4135(50),4137(40) keV										AHW	**
*^{151}Nd(β^-)^{151}Pm	E_{β^-}=2260(90) 2100(90) 1210(50) to 3/2$^+$ level at 255.692, 1/2$^+$ at 426.451,										Ens091	**
*	and 5/2$^+$ at 1297.682 keV										Ens091	**
*^{151}Pm(β^-)^{151}Sm	E_{β^-}=1190(10) to ground state and 3/2$^-$ level at 4.821 keV, and other E_{β^-}										Ens091	**
*^{151}Gd(ε)^{151}Eu	pK=0.652(0.007) to (5/2$^-$,7/2$^-$) level at 353.64 keV										Ens091	**
*^{151}Tb(β^+)^{151}Gd	E_{β^+}=700(5) p$^+$ =104(5)×10^{-4} respectively, to 1/2$^-$ level at 839.320 keV, and other E_{β^+}										Ens091	**
*^{151}Ho(β^+)^{151}Dy	E_{β^+}=3530(50) to 9/2$^-$ level at 527.40 keV										Ens091	**
*^{151}Tm(β^+)^{151}Er	p$^+$ =0.71(0.02) to 9/2$^-$ level at 801.52 keV										Ens091	**
*^{151}Tm(β^+)^{151}Er	F : lower limit: positrons escape from detector; E_{β^+}=5250(50) to 801.6 level										91Ke11	**
*^{151}Tm^m(IT)^{151}Tm	Only α-decay energies are used										GAu	**
^{152}Ba−^{152}Sm	35476	41				2			TR1	1.0	26Ye.A	*
^{152}La−^{133}Cs$_{1.143}$	55274	11	55260	5	−1.3	1	24	24 ^{152}La	JY2	1.0	25Ja05	
^{152}La−^{12}C$_6$ ^{1}H$_6$	−99762.9	6.2	−99758	5	0.7	1	76	76 ^{152}La	CP2	1.0	24Ra.A	
^{152}Ce−u	−63173	66	−63211.1	2.1	−0.6	U			RI1	1.0	19Ki.B	
	−63208.6	2.5			−1.0	1	74	74 ^{152}Ce	CP2	1.0	22Or02	*
	−63224.2	44.0			0.3	U			GR1	1.0	25Sp03	
^{152}Ce−^{12}C$_6$ ^{1}H$_6$	−110158.7	2.5	−110161.3	2.1	−1.1	o			CP2	1.0	22Ra.A	
^{152}Ce ^{1}H−^{153}Nd	16884.6	5.5	16890.7	2.6	1.1	1	23	13 ^{152}Ce	RI1	1.0	24Ki16	*
^{152}Pr−u	−68447.1	19.9	−68463.3	2.9	−0.8	–			CP1	1.0	06Sa56	
	−68472.7	10.7			0.9	–			CP2	1.0	22Or02	*
	−68461.0	22.5			−0.1	U			GR1	1.0	25Sp03	
^{152}Pr−^{12}C$_6$ ^{1}H$_6$	−115423	11	−115413.5	2.9	0.9	o			CP2	1.0	22Ra.A	
^{152}Pr−u ave.	−68467	9	−68463.3	2.9	0.4	1	10	10 ^{152}Pr			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{152}Pr ^{1}H$-^{153}$Nd	11599.5	22.9	11639	3	1.7	U			RI1	1.0	24Ki16	*
^{152}Nd$-$u	−75289.8	16.1	−75298	4	−0.5	U			GR1	1.0	25Sp03	
C_{12} H_8-^{152}Sm	142764	32	142862.8	0.9	0.8	U			R04	4.0	64De15	
	142867.0	5.0			−0.3	U			M22	2.5	75Ka25	
^{152}Eu$-$u	−78347	50	−78250.9	0.9	1.9	U			GS2	1.0	05Li24	*
C_{12} H_8-^{152}Gd	142870	50	142803.0	0.9	−0.3	U			R04	4.0	64De15	
^{152}Gd O$-C_{14}$	−85290.7	3.5	−85288.1	0.9	0.5	U			TG1	1.5	11Ke03	
^{152}Tb$-$u	−76212	159	−75920	40	1.8	U			GS2	1.0	05Li24	*
^{152}Dy$-$u	−75278	30	−75277	5	0.0	U			GS2	1.0	05Li24	
^{152}Ho$-$u	−68247	58	−68278	13	−0.5	U			GS2	1.0	05Li24	*
^{152}Er$-$u	−64962	30	−64948	9	0.5	U			GS2	1.0	05Li24	
^{152}Tm$-$u	−55599	77	−55520	60	1.0	1	66	66 ^{152}Tm	GS2	1.0	05Li24	*
^{152}Yb$-^{136}$Ce O	48508	49	48500	50	−0.1	1	93	93 ^{152}Yb	TR1	1.0	21Be23	
^{152}Sm $^{35}Cl_2-^{148}$Sm $^{37}Cl_2$	10802	10	10810.4	1.1	0.3	U			H21	2.5	70Ma05	
	10810.8	2.0			−0.1	U			H25	2.5	72Ba08	
	10807.9	1.4			0.7	U			M21	2.5	75Ka25	
^{152}Sm ^{35}Cl$-^{150}$Sm ^{37}Cl	5429	4	5407.8	0.6	−1.3	U			H12	4.0	64Ba15	
	5396	4			1.2	o			H21	2.5	70Ma05	
	5402.7	0.8			2.5	U			M21	2.5	75Ka25	
^{152}Ce$-^{152}$Pr	5250.7	2.7	5252.2	2.4	0.6	1	82	69 ^{152}Pr	RI1	1.0	24Ki16	*
^{152}Nd$-^{152}$Pr	−6834.4	2.6	−6834.5	2.6	−0.0	1	100	99 ^{152}Nd	RI1	1.0	24Ki16	*
^{152}Pm$-^{152}$Pr	−8057	67	−8025	19	0.5	1	8	8 ^{152}Pm	RI1	1.0	24Ki16	*
^{152}Gd$-^{152}$Sm	59.80	0.19	59.79	0.19	−0.1	1	98	55 ^{152}Sm	SH1	1.0	11El02	
^{152}Er$-^{152}$Dy	10314	27	10329	11	0.6	U			TR1	1.0	25Ko14	
^{152}Tm$-^{152}$Dy	19950	140	19750	60	−1.4	1	20	20 ^{152}Tm	TR1	1.0	25Ko14	*
^{152}Sm$-^{151}$Eu	95	42	−117.2	0.6	−1.3	U			R04	4.0	64De15	
^{152}Sm$-^{150}$Sm	2563	31	2457.6	0.6	−0.8	U			R04	4.0	64De15	
^{152}Gd(α)^{148}Sm	2197.9	30.	2204.5	1.0	0.2	U					61Ma05	
^{152}Dy(α)^{148}Gd	3728.0	8.	3726	4	−0.2	–					65Ma51	Z
	3726.0	5.			0.1	–			Dba		67Go32	Z
ave.	3727	4			−0.0	1	100	100 ^{152}Dy			average	
^{152}Ho(α)^{148}Tb	4506.9	3.	4507.4	1.3	0.2	–			Bka		82Bo04	*
	4508.0	2.			−0.3	–			Ora		82De11	Z
	4505.8	3.			0.5	–					82To14	
	4507.9	3.			−0.1	–					87St.A	Z
ave.	4507.3	1.3			0.1	1	100	82 ^{152}Ho			average	
^{152}Er(α)^{148}Dy	4935.2	5.	4934.3	1.6	−0.2	–			GSa		79Ho10	
	4934.6	3.			−0.1	–			Bka		82Bo04	Z
	4934.3	2.			−0.0	–			Ora		82De11	Z
ave.	4934.5	1.6			−0.1	1	99	83 ^{152}Er			average	
^{150}Nd(t,p)^{152}Nd	4125	30	4118	4	−0.2	U			Ald		72Ch11	
^{150}Sm(t,p)^{152}Sm	5376	25	5371.6	0.6	−0.2	U			Ald		66Bj01	
^{152}Sm(p,t)^{150}Sm	−5378	8	−5371.6	0.6	0.8	U			Min		72De47	
	−5376	4			1.1	U			McM		73Ga04	
	−5379	15			0.5	U			Ham		74Oe03	
^{151}Sm(n,γ)^{152}Sm	8257.6	0.8	8256.9	0.6	−0.8	1	54	30 ^{151}Sm			71Gr22	Z
^{151}Sm(p,γ)^{152}Eu	5604	4	5600.80	0.12	−0.8	U					75Jo.A	
^{151}Eu(n,γ)^{152}Eu	6306.70	0.10	6306.71	0.10	0.1	1	99	79 ^{152}Eu	ILn		85Vo15	Z
	6307.11	0.14			−2.8	C			Bdn		06Fi.A	
^{152}Gd(d,t)^{151}Gd	−2338	10	−2331.8	2.8	0.6	U			Kop		67Tj01	
^{152}Pr(β^-)^{152}Nd	6350	120	6366.3	2.4	0.1	U			Kur		95Ik03	
^{152}Nd(β^-)^{152}Pm	1088	27	1109	17	0.8	–					93Sh23	
	1120	30			−0.4	–			Kur		95Ik03	
ave.	1102	20			0.3	1	74	73 ^{152}Pm			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{152}Pm(β^-)^{152}Sm	3600	200	3515	17	−0.4	U					71Da19	
	3520	150			−0.0	U					72Wa04	
	3400	200			0.6	U					75Wi08	
	3500	100			0.2	U					77Ya07	
	3500	40			0.4	1	19	19 ^{152}Pm	Kur		95Ik03	
^{152}Pm^m(β^-)^{152}Sm	3603	100	3650	80	0.5	2					71Da19	*
	3753	150			−0.7	2					72Wa04	*
^{152}Eu(β^+)^{152}Sm	1871	5	1873.8	0.6	0.6	U					58Al99	*
	1866	5			1.6	U					62Lo10	*
	1870.8	2.			1.5	–					72Sv02	*
	1872.8	1.5			0.7	–					77Mi.A	*
ave.	1872.1	1.2			1.4	1	24	14 ^{152}Eu			average	
^{152}Eu(β^-)^{152}Gd	1809	10	1818.1	0.6	0.9	U					58Al99	*
	1827	7			−1.3	U					60La04	*
	1836	20			−0.9	U					60Sc14	*
	1806	4			3.0	B					69An18	*
^{152}Tb(β^+)^{152}Gd	3990	40				2					76Cr.B	*
^{152}Ho(β^+)^{152}Dy	6690	100	6519	13	−1.7	U			IRS		83Al06	*
	6270	140			1.8	U					Averag	*
	6225	90			3.3	B			IRS		93Al03	*
^{152}Tm(β^+)^{152}Er	8680	240	8780	60	0.4	U					16Na02	
	6850	110			17.5	C					84Ha.B	*
^{152}Tm^m(β^+)^{152}Er	8820	240	8930	70	0.4	U					16Na02	
^{152}Tm^m(IT)^{152}Tm	140	50	150	50	0.1	1	91	77 ^{152}Tm^m			FKo256	*
^{152}Yb(β^+)^{152}Tm^m	5065	165	5510	80	2.7	C					84Ha.B	*
	5465	195			0.2	–			Got		90Sa.A	
	5434	200			0.4	–			GSI		04Na.A	*
ave.	5450	140			0.4	1	30	23 ^{152}Tm^m			average	
*^{152}Ba−^{152}Sm	from frequency ratio ^{152}Ba$^+$/(^{152}Sm)$^+$ =1.000233521(268)										26Ye.A	**
*^{152}Ce−u	Represents frequency ratio (C_6H_6)$^+$/^{152}Ce^{++} =0.973367710(16)										HWJ234	**
*^{152}Ce ^{1}H−^{153}Nd	Reported $M-A$=–51602(33) keV, author confirmed mistake in the unc.										24Ki16	**
*^{152}Pr−u	Represents frequency ratio (C_6H_6)$^+$/^{152}Pr^{++} =0.973333983(69)										HWJ234	**
*^{152}Pr ^{1}H−^{153}Nd	Reported $M-A$=–56526(39) keV, author confirmed mistake in the unc.										24Ki16	**
*^{152}Eu−u	$M-A$=–72915(35) keV for mixture gs+m+r at 45.5998 and 147.86 keV										Nub269	**
*^{152}Tb−u	$M-A$=–70740(29) keV for mixture gs+n at 501.74 keV										Nub269	**
*^{152}Ho−u	$M-A$=–63492(28) keV for mixture gs+m at 160(3) keV										Nub269	**
*^{152}Ce−^{152}Pr	Reported $M-A$=–58891(20) keV, author confirmed mistake in the unc.										24Ki16	**
*^{152}Nd−^{152}Pr	Reported $M-A$=–70148(20) keV, author confirmed mistake in the unc.										24Ki16	**
*^{152}Pm−^{152}Pr	D_M=–7981.8(3.8) μu, $M-A$=–71217.2(10.4) keV for a mixture of ground state and m										24Ki16	**
*^{152}Pm−^{152}Pr	at 140(90) keV; reported $M-A$=–71217(20) keV, author confirmed										24Ki16	**
*^{152}Pm−^{152}Pr	mistake in the uncertainty										24Ki16	**
*^{152}Tm−^{152}Dy	frequency ratio corrected for the unresolved isomer at 43.2(0.2) keV										25Ko14	**
*^{152}Ho(α)^{148}Tb	E_α=4389.1(3,Z); and 4455.1(3,Z) from ^{152}Ho^m to ^{148}Tb^m										82Bo04	**
*	combined with ^{152}Ho^m(IT)-^{148}Tb^m(IT)=160(1)-90.1(0.3) keV										87St.A	**
*^{152}Pm^m(β^-)^{152}Sm	E_{β^-}=1800(100) 1950(150) respectively, to 5$^-$ level at 1803.94 keV										Ens13b	**
*^{152}Eu(β^+)^{152}Sm	E_{β^+}=895(5) 890(5) respectively, from ^{152}Eu^m at 45.5998 keV										Nub269	**
*^{152}Eu(β^+)^{152}Sm	E_{β^+}=727(2) 729(1.5) respectively, to 2$^+$ level at 121.7818 keV										Ens13b	**
*^{152}Eu(β^-)^{152}Gd	Q_{β^-}=1855(10) from ^{152}Eu^m at 45.5998 keV										Nub269	**
*^{152}Eu(β^-)^{152}Gd	E_{β^-}=1483(7) to 2$^+$ level at 344.2790 keV										Ens13b	**
*^{152}Eu(β^-)^{152}Gd	E_{β^-}=1840(30) 1490(20) 1072(20) to ground state, 2$^+$ at 344.2790, 4$^+$ at 755.3961 keV										Ens13b	**
*^{152}Eu(β^-)^{152}Gd	Q_{β^-}=1852(4) from ^{152}Eu^m at 45.5998 keV										Nub269	**
*^{152}Tb(β^+)^{152}Gd	E_{β^+}=2830(15) 8(4)% to ground state, 5.2(1)% to 2$^+$ level at 344.2790 keV										Ens13b	**
*^{152}Ho(β^+)^{152}Dy	E_{β^+}=3390(100) from ^{152}Ho^m at 160(1) to 8$^+$ level at 2437.42 keV										Ens13b	**
*^{152}Ho(β^+)^{152}Dy	From adopted KLM/β^+=0.97(0.13)										AHW	**
*	from ^{152}Ho^m at 160(1) to 8$^+$ level at 2437.42 keV										Ens13b	**
*	after extra 3(2)% side-feeding correction; see reference										90Sa32	**
*	p$^+$ =0.52(0.04)/.967 gives KLM/β^+=0.86(0.14)										85Sc09	**
*	KLM/β^+=1.12(0.10) after 0.967(0.008) side-feeding correction										90Sa32	**
*^{152}Ho(β^+)^{152}Dy	Q_{β^+}=6270(90); and 6330(100) from ^{152}Ho^m at 160(3) keV										Nub269	**
*^{152}Tm(β^+)^{152}Er	p$^+$ =0.64(0.02) to 8$^+$ level at 2183.3 keV										Ens13b	**
*^{152}Tm^m(IT)^{152}Tm	from QEC(iso)−QEC(gs)=8820−8680 keV in 16Na02; unc by evaluator										FKo256	**
*^{152}Yb(β^+)^{152}Tm^m	p$^+$ =0.57(0.03) (1)$^+$ level at 482.32 keV										Ens13b	**
*^{152}Yb(β^+)^{152}Tm^m	As reported in reference										11Es03	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{153}La−^{133}Cs$_{1.150}$		59314	20				2			JY2	1.0	25Ja05	
		59306	40	59314	20	0.2	U			JY2	1.0	25Ja05	
^{153}Ce−^{12}C$_6$ ^{1}H$_6$		−105685.5	2.7	−105684.9	2.4	0.2	o			CP2	1.0	22Ra.A	
^{153}Ce−u		−58735.3	2.8	−58734.7	2.4	0.2	1	70	70 ^{153}Ce	CP2	1.0	22Or02	*
		−58873.2	204.0			0.7	U			GR1	1.0	25Sp03	
^{153}Ce ^{1}H−^{154}Nd		19500.3	6.8	19493.7	2.5	−1.0	1	14	12 ^{153}Ce	RI1	1.0	24Ki16	*
^{153}Pr−u		−66110.5	15.3	−66076.1	2.0	2.3	U			CP1	1.0	06Sa56	
		−66065	40			−0.3	U			CP1	1.0	12Va02	*
^{153}Pr−^{12}C$_6$ ^{1}H$_6$		−113024.3	2.7	−113026.3	2.0	−0.7	o			CP2	1.0	22Ra.A	
^{153}Pr−u		−66074.1	2.6	−66076.1	2.0	−0.8	1	60	60 ^{153}Pr	CP2	1.0	22Or02	*
		−66050.9	29.0			−0.9	U			GR1	1.0	25Sp03	
^{153}Pr−^{80}Kr$_{1.913}$		93906	40	93892.2	2.4	−0.3	U			CP1	1.0	12Va02	
^{153}Pr−^{86}Kr$_{1.779}$		92958	40	92947.6	2.0	−0.3	U			CP1	1.0	12Va02	
^{153}Pr ^{1}H−^{154}Nd		12128.6	11.5	12152.4	2.3	2.1	U			RI1	1.0	24Ki16	*
^{153}Nd−u		−72283.3	5.2	−72276.8	1.9	1.2	1	13	13 ^{153}Nd	CP1	1.0	12Va02	*
^{153}Nd−^{80}Kr$_{1.913}$		87687.9	4.7	87691.5	2.2	0.8	1	22	14 ^{153}Nd	CP1	1.0	12Va02	
^{153}Nd−^{86}Kr$_{1.779}$		86740.7	5.3	86746.9	1.9	1.2	1	12	12 ^{153}Nd	CP1	1.0	12Va02	
^{153}Nd−u		−72259.2	22.5	−72276.8	1.9	−0.8	U			GR1	1.0	25Sp03	
^{153}Pm−u		−75833	23	−75875	3	−1.8	U			CP1	1.0	12Va02	*
^{153}Pm−^{80}Kr$_{1.913}$		84139	23	84094	4	−2.0	U			CP1	1.0	12Va02	
^{153}Pm−^{86}Kr$_{1.779}$		83192	23	83149	3	−1.9	U			CP1	1.0	12Va02	
^{153}Pm−u		−75841.6	30.1	−75875	3	−1.1	U			GR1	1.0	25Sp03	
^{153}Sm−u		−77887.8	60.1	−77897.6	0.9	−0.2	U			GR1	1.0	25Sp03	
C$_{12}$ H$_9$−^{153}Eu		149103	18	149190.4	0.9	1.2	U			R04	4.0	64De15	
C$_{11}$ ^{13}C H$_8$−^{153}Eu		144606	30	144720.2	0.9	1.0	U			R04	4.0	64De15	
C$_9$ ^{13}C H$_{16}$ O−^{153}Eu		201934	38	202235.1	0.9	2.0	U			R04	4.0	64De15	
^{153}Eu−^{85}Rb$_{1.800}$		80021	16	80013.3	0.9	−0.5	U			MA5	1.0	00Be42	
^{153}Ho−u		−69814	37	−69795	5	0.5	U			GS2	1.0	05Li24	*
^{153}Er−u		−64942	30	−64919	9	0.8	U			GS2	1.0	05Li24	
^{153}Eu ^{35}Cl−^{151}Eu ^{37}Cl		4334	4	4330.32	0.18	−0.4	U			H21	2.5	70Ma05	*
^{153}Ce−^{153}Nd		13540.3	5.2	13542.1	2.8	0.3	1	28	18 ^{153}Ce	RI1	1.0	24Ki16	*
^{153}Pr−^{153}Nd		6199.2	2.3	6200.7	1.9	0.7	1	69	40 ^{153}Pr	RI1	1.0	24Ki16	*
^{153}Pm−^{153}Nd		−3597.9	2.8	−3597.9	2.8	−0.0	1	98	98 ^{153}Pm	RI1	1.0	24Ki16	*
^{153}Sm−^{153}Nd		−5622.8	5.5	−5620.8	2.0	0.4	1	13	11 ^{153}Nd	RI1	1.0	24Ki16	*
^{153}Er−^{153}Dy		9273	52	9311	10	0.7	U			TR1	1.0	25Ko14	
^{153}Tm−^{153}Dy		16303	57	16296	13	−0.1	U			TR1	1.0	25Ko14	*
^{153}Yb−^{153}Dy		23662	49				2			TR1	1.0	21Be23	
^{138}La O−^{153}Eu		−19266	123	−19196.1	1.0	0.1	U			R05	4.0	65De13	
^{153}Eu O−C$_{14}$		−83849.6	5.8	−83850.5	0.9	−0.1	U			TG1	1.5	11Ke03	
^{153}Eu−^{152}Sm		1544	42	1497.4	0.6	−0.3	U			R04	4.0	64De15	
^{153}Eu−^{151}Eu		1567	33	1380.19	0.17	−1.4	U			R04	4.0	64De15	
^{153}Dy(α)^{149}Gd		3560.0	8.	3559	4	−0.1	–					65Ma51	Z
		3554.9	5.			0.9	–			Dba		67Go32	Z
	ave.	3556	4			0.7	1	69	48 ^{153}Dy			average	
^{153}Ho(α)^{149}Tb		4052.3	5.	4052	4	−0.1	2					68Go.C	*
		4051.0	5.			0.1	2			ORa		71To01	*
^{153}Er(α)^{149}Dy		4799.8	10.	4802.3	1.4	0.2	U					81Ho.A	
		4804.5	3.			−0.7	–			Bka		82Bo04	Z
		4802.0	2.			0.1	–			Ora		82De11	Z
		4802.8	3.			−0.2	–					87Sc.A	Z
		4799.7	4.			0.6	–			Daa		96Pa01	
	ave.	4802.4	1.4			−0.1	1	100	86 ^{153}Er			average	
^{153}Tm(α)^{149}Ho		5252.3	5.	5248.4	1.5	−0.8	U			GSa		79Ho10	*
		5246.1	3.			0.8	–			Bka		82Bo04	*
		5249.2	2.			−0.4	–			Ora		82De11	*
		5247.7	3.			0.2	o					87Sc.A	*
		5247.7	3.			0.2	–					88Sc.A	
		5249.5	5.			−0.2	U			Daa		96Pa01	
	ave.	5248.1	1.5			0.2	1	100	62 ^{149}Ho			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{153}Gd(n,α)^{150}Sm		9790	30	9815.7	0.6	0.9	U			ILL		81Wa31	
^{153}Eu(p,t)^{151}Eu		−6374	5	−6375.20	0.16	−0.2	U			Min		75Ta12	
^{152}Sm(n,γ)^{153}Sm		5867.1	0.4	5868.40	0.13	3.3	B					69Re04	Z
		5868.4	0.3			0.0	–					71Be41	Z
		5868.4	0.7			0.0	U					82Ba15	Z
		5868.40	0.15			0.0	–			Bdn		06Fi.A	
^{152}Sm(d,p)^{153}Sm		3645	12	3643.84	0.13	−0.1	U			Tal		65Ke09	
^{152}Sm(n,γ)^{153}Sm	ave.	5868.40	0.13	5868.40	0.13	0.0	1	100	98 ^{153}Sm			average	
^{152}Eu(n,γ)^{153}Eu		8550.28	0.12	8550.28	0.12	0.0	1	100	93 ^{153}Eu	ILn		85Vo15	Z
^{153}Eu(γ,n)^{152}Eu		−8650	130	−8550.28	0.12	0.8	U			Phi		60Ge01	
^{152}Gd(n,γ)^{153}Gd		6247.27	0.35	6246.95	0.13	−0.9	–			ILn		85Vo15	Z
		6246.89	0.14			0.4	–			ILn		93Sp.A	
		6247.48	0.21			−2.5	B			Bdn		06Fi.A	
^{152}Gd(d,p)^{153}Gd		4015	10	4022.38	0.13	0.7	U			Kop		67Tj01	
^{152}Gd(n,γ)^{153}Gd	ave.	6246.94	0.13	6246.95	0.13	0.1	1	99	57 ^{152}Gd			average	
^{152}Gd(^{3}He,d)^{153}Tb		−1634	30	−1598	4	1.2	U			McM		76St10	
^{153}Pr(β^-)^{153}Nd		5720	100	5776.0	1.8	0.6	U			Kur		02Sh.B	
^{153}Nd(β^-)^{153}Pm		3336	25	3351.5	2.6	0.6	U			Ida		93Gr17	
		3260	100			0.9	U			Kur		02Sh.B	
^{153}Pm(β^-)^{153}Sm		1777	50	1884	3	2.1	U					62Ko10	*
		1863	15			1.4	U			Ida		93Gr17	
^{153}Sm(β^-)^{153}Eu		810	10	808.1	0.6	−0.2	U					54Gr19	
		795	10			1.3	U					54Le08	
		820	10			−1.2	U					55Ma62	
		825	10			−1.7	U					56Du31	
		792	10			1.6	U					57Jo24	
^{153}Tb(β^+)^{153}Gd		1573	5	1569	4	−0.8	1	59	59 ^{153}Tb			78Cr02	*
^{153}Dy(β^+)^{153}Tb		2171	2	2170.4	1.9	−0.3	1	93	52 ^{153}Dy			78Gr13	*
^{153}Ho(β^+)^{153}Dy		4153	50	4131	6	−0.4	o			IRS		83Al06	*
		4160	60			−0.5	U			IRS		93Al03	
^{153}Lu^m(IT)^{153}Lu		80	5				4					97Ir01	*
*^{153}Ce−u	Represents frequency ratio $(C_6H_6)^+/^{153}Ce^{++}$ =0.979802813(17)											HWJ234	**
*^{153}Ce ^{1}H−^{154}Nd	Reported $M-A$=–47415(27) keV, author confirmed mistake in the unc.											24Ki16	**
*^{153}Pr−u	Represents frequency ratio $^{153}Pr^{++}/(C_{12}H_4)^+$ =0.99430250(26)											WgM124	**
*^{153}Pr−u	Represents frequency ratio $(C_6H_6)^+/^{153}Pr^{++}$ =0.979755797(17)											HWJ234	**
*^{153}Pr ^{1}H−^{154}Nd	Reported $M-A$=–54282(29) keV, author confirmed mistake in the unc.											24Ki16	**
*^{153}Nd−u	Represents frequency ratio $^{153}Pr^{++}/(C_{12}H_4)^+$ =0.994342.931(34)											WgM124	**
*^{153}Pm−u	Represents frequency ratio $^{153}Pm^{++}/(C_{12}H_4)^+$ =0.99436601(15)											WgM124	**
*^{153}Ho−u	$M-A$=–64997(28) keV for mixture gs+m at 68.7 keV											Nub269	**
*^{153}Eu ^{35}Cl−^{151}Eu ^{37}Cl	Increased by 5 for systematic difference of H21 with later data											AHW	**
*^{153}Ce−^{153}Nd	Reported $M-A$=–54718(33) keV, author confirmed mistake in the unc.											24Ki16	**
*^{153}Pr−^{153}Nd	Reported $M-A$=–61556(33) keV, author confirmed mistake in the unc.											24Ki16	**
*^{153}Pm−^{153}Nd	Reported $M-A$=–70682(33) keV, author confirmed mistake in the unc.											24Ki16	**
*^{153}Sm−^{153}Nd	Reported $M-A$=–72568(33) keV, author confirmed mistake in the unc.											24Ki16	**
*^{153}Tm−^{153}Dy	frequency ratio corrected for the unresolved isomer at 150(50) keV											25Ko14	**
*^{153}Ho(α)^{149}Tb	E_α=4013.1(5,Z) from ^{153}Ho^m at 68.7 keV											Nub269	**
*^{153}Ho(α)^{149}Tb	E_α=3910(5) to ^{149}Tb^m at 35.75 keV											Nub269	**
*^{153}Tm(α)^{149}Ho	E_α=5114.2(5,Z) 5108.2(3,Z) 5111.2(2,Z) respectively, contain a 8%											AHW	**
*	^{153}Tm(α)^{149}Ho−^{153}Tm^m(α)^{149}Ho^m=48.80(0.20)-43.2(0.2)=											Nub269	**
*	=5.6(0.3) keV lower ^{153}Tm^m(α)^{149}Ho^m branch, corrected thus +0.5keV											GAu	**
*^{153}Tm(α)^{149}Ho	E_α=5110.6(3,Z); and 5103.6(4,Z) for lower ^{153}Tm^m(α) branch											87Sc.A	**
*^{153}Pm(β^-)^{153}Sm	E_{β^-}=1650(50) to 3/2$^-$ level at 127.298 keV, and other E_{β^-}											Ens062	**
*^{153}Tb(β^+)^{153}Gd	E_{β^+}=339(5) to 3/2$^+$ level at 212.0078 keV											Ens062	**
*^{153}Dy(β^+)^{153}Tb	E_{β^+}=886(2) to 9/2$^-$ level at 262.831 keV											Ens062	**
*^{153}Ho(β^+)^{153}Dy	E_{β^+}=2835(50) to 9/2$^-$ level at 295.84 keV											Ens062	**
*^{153}Lu^m(IT)^{153}Lu	Trends in Mass Surface TMS suggest ^{153}Lu 130 keV less bound											GAu	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{154}Ce−u		−56404	619	−55898.4	2.8	0.3	B			GT3	2.5	16Kn03	
^{154}Ce−^{12}C$_6$ ^{1}H$_6$		−102848.5	2.8	−102848.6	2.8	−0.0	o			CP2	1.0	22Ra.A	
^{154}Ce−u		−55898.4	2.8				2			CP2	1.0	22Or02	*
^{154}Ce ^{1}H−^{155}Pm		23782.4	12.4	23787	3	0.4	U			RI1	1.0	24Ki16	*
^{154}Pr−^{12}C$_6$ ^{1}H$_6$		−109216.6	2.8	−109217.1	2.7	−0.2	o			CP2	1.0	22Ra.A	
^{154}Pr−u		−62266.4	2.8	−62266.9	2.7	−0.2	1	94	94 ^{154}Pr	CP2	1.0	22Or02	*
		−62277.4	26.8			0.4	U			GR1	1.0	25Sp03	
^{154}Pr ^{1}H−^{155}Pm		17411.2	11.0	17418	3	0.7	1	9	6 ^{154}Pr	RI1	1.0	24Ki16	*
^{154}Nd−^{133}Cs$_{1.158}$		39061	27	39083.2	1.0	0.8	U			JY1	1.0	20Vi04	
^{154}Nd−^{82}Kr$_{1.878}$		92079.8	1.1	92079.0	1.0	−0.7	1	91	91 ^{154}Nd	CP2	1.0	18Or02	
^{154}Nd−u		−70405.2	18.3	−70403.4	1.0	0.1	U			GR1	1.0	25Sp03	
^{154}Pm−u		−73322.0	24.7	−73327.3	2.5	−0.2	U			GR1	1.0	25Sp03	*
C_{12} H_{10}−^{154}Sm		155830	29	156035.5	1.2	1.8	U			R04	4.0	64De15	
		156035.7	4.0			−0.0	U			M22	2.5	75Ka25	
^{154}Sm−u		−77769.7	31.1	−77785.1	1.2	−0.5	U			GR1	1.0	25Sp03	
C_{12} H_{10}−^{154}Gd		157149	40	157378.5	0.9	1.4	U			R04	4.0	64De15	
C_{11} ^{13}C H_9−^{154}Gd		152550	110	152908.3	0.9	0.8	U			R04	4.0	64De15	
C_{10} $^{13}C_2$ H_8−^{154}Gd		148030	90	148438.1	0.9	1.1	U			R04	4.0	64De15	
C_{10} H_6 N_2−^{154}Gd		131980	240	132226.4	0.9	0.3	U			R04	4.0	64De15	
^{154}Gd−^{138}La O		19005	80	18833.1	1.0	−0.5	U			R05	4.0	65De13	
^{154}Tb−u		−75380	110	−75320	50	0.6	R			GS2	1.0	05Li24	*
^{154}Dy−^{133}Cs$_{1.158}$		33903	19	33914	8	0.6	1	18	18 ^{154}Dy	MA5	1.0	00Be42	*
^{154}Ho−u		−69350	83	−69395	9	−0.5	U			GS2	1.0	05Li24	*
^{154}Tm−u		−58479	49	−58426	14	1.1	U			GS2	1.0	05Li24	*
^{154}Sm $^{35}Cl_2$−^{150}Sm $^{37}Cl_2$		10832.9	5.2	10835.3	1.1	0.2	U			M21	2.5	75Ka25	
^{154}Sm ^{35}Cl−^{152}Sm ^{37}Cl		5480	4	5427.5	0.9	−3.3	B			H12	4.0	64Ba15	
		5417	4			1.1	U			H21	2.5	70Ma05	
		5427.2	0.4			0.3	1	73	64 ^{154}Sm	M21	2.5	75Ka25	
^{154}Gd ^{35}Cl−^{152}Gd ^{37}Cl		4019.5	2.	4024.71	0.23	1.0	U			H25	2.5	72Ba08	
		4016	30			0.1	U			H12	4.0	64Ba15	
^{154}Ce−^{154}Nd		14474.4	8.9	14505.0	3.0	3.4	B			RI1	1.0	24Ki16	*
^{154}Pr−^{154}Nd		8131.3	2.5	8136.5	2.9	2.1	U			RI1	1.0	24Ki16	*
^{154}Pm−^{154}Nd		−2930	11	−2923.9	2.3	0.6	2			CP2	1.0	24Ra.A	*
		−2923.6	2.3			−0.1	2			RI1	1.0	24Ki16	*
^{154}Sm−^{154}Nd		−7375.0	3.8	−7381.8	1.5	−1.8	1	16	9 ^{154}Sm	RI1	1.0	24Ki16	*
^{154}Sm−^{154}Gd		1338.2	3.8	1343.0	0.9	0.5	U			H25	2.5	72Ba08	
		1342.8	0.8			0.1	1	19	17 ^{154}Sm	M21	2.5	75Ka25	
^{154}Er−^{154}Dy		8401	25	8361	9	−1.6	U			TR1	1.0	25Ko14	
^{154}Tm−^{154}Dy		17169	45	17146	16	−0.5	U			TR1	1.0	25Ko14	*
^{154}Yb−^{138}Ce O		45485	48	45494	13	0.2	1	8	8 ^{154}Yb	TR1	1.0	21Be23	
^{154}Sm−C_{12} H_9		−148211.0	8.0	−148210.4	1.2	0.0	U			M21	2.5	75Ka25	
^{139}La O−^{154}Gd		−19616	55	−19594.3	1.1	0.1	U			R05	4.0	65De13	
^{154}Sm−^{153}Eu		1082	42	980.0	1.1	−0.6	U			R04	4.0	64De15	
^{154}Sm−^{152}Sm		2664	43	2477.4	0.9	−1.1	U			R04	4.0	64De15	
^{154}Gd−^{152}Gd		1400	50	1074.58	0.22	−1.6	U			R04	4.0	64De15	
^{154}Gd O−C_{15}		−84207.4	5.9	−84213.6	0.9	−0.7	U			TG1	1.5	09Ke.A	
		−84206.6	4.3			−1.1	U			TG1	1.5	11Ke03	
^{154}Dy(α)^{150}Gd		2946.4	5.	2945	5	−0.2	1	93	82 ^{154}Dy	Dba		67Go32	Z
^{154}Ho(α)^{150}Tb		4041.3	5.	4041	4	0.0	2					68Go.C	Z
		4041.7	5.			−0.0	2			ORa		74Sc19	Z
^{154}Ho^m(α)^{150}Tb^m		3819.2	10.	3823	5	0.4	–			ORa		71To01	Z
		3824.0	5.			−0.1	–			ORa		74Sc19	Z
	ave.	3823	5			0.1	1	100	89 ^{154}Ho^m			average	
^{154}Er(α)^{150}Dy		4280.5	5.	4279.8	2.6	−0.1	–					68Go.C	Z
		4279.5	3.			0.1	–			Bka		82Bo04	Z
	ave.	4279.8	2.6			−0.0	1	98	92 ^{154}Er			average	
^{154}Tm(α)^{150}Ho		5096.9	5.1	5093.8	2.6	−0.6	2			GSa		79Ho10	Z
		5092.7	3.			0.4	2			Bka		82Bo04	
^{154}Tm^m(α)^{150}Ho^m		5174.8	5.	5171.8	1.6	−0.6	3			GSa		79Ho10	Z
		5170.8	3.			0.3	3			Bka		82Bo04	Z
		5171.8	2.1			0.0	3			Ora		82De11	Z
^{154}Yb(α)^{150}Er		5473.4	5.	5474.3	1.7	0.2	–			GSa		79Ho10	Z

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{154}Yb(α)^{150}Er		5474.7	2.	5474.3	1.7	−0.2	–			Ora		82De11	Z
		5473.4	4.			0.2	–			Daa		96Pa01	
	ave.	5474.3	1.7			−0.0	1	100	81 ^{154}Yb			average	
^{152}Sm(t,p)^{154}Sm		5361	25	5353.2	0.8	−0.3	U			Ald		66Bj01	
^{154}Sm(p,t)^{152}Sm		−5357	8	−5353.2	0.8	0.5	U			Min		72De47	
		−5353	15			−0.0	U			Ham		74Oe03	
^{154}Gd(p,t)^{152}Gd		−6660	5	−6659.87	0.21	0.0	U			Min		73Oo01	
^{154}Sm(d,^{3}He)^{153}Pm		−3623	25	−3575	3	1.9	U					76Su.B	
^{154}Sm(t,α)^{153}Pm		10748	20	10745	3	−0.1	1	3	2 ^{153}Pm	LAl		78Bu18	
^{153}Eu(n,γ)^{154}Eu		6442.2	0.3	6442.20	0.24	0.0	–			ILn		87Ba52	Z
		6442.2	0.4			0.0	–			Bdn		06Fi.A	
	ave.	6442.20	0.24			0.0	1	99	91 ^{154}Eu			average	
^{153}Gd(n,γ)^{154}Gd		8895.25	0.30	8894.72	0.16	−1.8	–			ILn		85Vo15	Z
		8894.47	0.20			1.2	–			ILn		93Sp.A	Z
^{154}Gd(d,t)^{153}Gd		−2642	10	−2637.49	0.16	0.5	U			Kop		67Tj01	
^{153}Gd(n,γ)^{154}Gd	ave.	8894.71	0.17	8894.72	0.16	0.0	1	98	58 ^{153}Gd			average	
^{154}Pr(β^-)^{154}Nd		7720	100	7579.1	2.7	−1.4	U			Kur		02Sh.B	*
^{154}Nd(β^-)^{154}Pm^m		2687	25	2709	4	0.9	U			Ida		93Gr17	*
^{154}Pm^m(IT)^{154}Pm		15	3				3					24Ra.A	*
^{154}Pm(β^-)^{154}Sm		3900	200	4152.5	2.5	1.3	U					71Da28	
		4396	180			−1.4	U					72Ta13	
		3880	200			1.4	U					74Ya07	*
^{154}Pm^m(β^-)^{154}Sm		3900	200	4167	4	1.3	U					71Da28	*
		4190	170			−0.1	U					72Ta13	*
		3940	50			4.5	B					73Pr05	*
		3940	200			1.1	U					74Ya07	*
		4056	100			1.1	U			Ida		93Gr17	
^{154}Eu(β^-)^{154}Gd		1978	5	1967.3	0.7	−2.1	U					60La04	*
		1967	2			0.1	1	11	7 ^{154}Eu			77Ra08	*
		1975	3			−2.6	U					81Bu.A	*
^{154}Tb(β^+)^{154}Gd		3562	50	3550	50	−0.2	2					70Ag02	*
^{154}Ho(β^+)^{154}Dy		5700	80	5755	10	0.7	U			IRS		83Al06	*
		5750	80			0.1	U			IRS		93Al03	
^{154}Ho^m(β^+)^{154}Dy		5994	100	5998	27	0.0	o			IRS		83Al.A	*
		6070	80			−0.9	1	12	11 ^{154}Ho^m	IRS		93Al03	
^{154}Tm^m(β^+)^{154}Er		8234	150	8250	50	0.1	U			Dbn		94Po26	*
^{154}Lu(β^+)^{154}Yb		7556	450	10310	50	6.1	C					84Ha.B	*
^{154}Lu^m(IT)^{154}Lu		58.7	9.3	62	12	0.4	o			Ara		97Da07	*
*^{154}Ce−u	Represents frequency ratio $(C_6H_6)^+/^{154}Ce^{++}$ =0.986227433(18)											HWJ234	**
*^{154}Ce ^{1}H−^{155}Pm	Reported $M-A$=–44787(18) keV, author confirmed mistake in the unc.											24Ki16	**
*^{154}Pr−u	Represents frequency ratio $(C_6H_6)^+/^{154}Pr^{++}$ =0.986186636(18)											HWJ234	**
*^{154}Pr ^{1}H−^{155}Pm	Reported $M-A$=–50722(18) keV, author confirmed mistake in the unc.											24Ki16	**
*^{154}Pm−u	Corrected for unresolved isomeric state using exc. energy in 12So10											25Sp03	**
*^{154}Tb−u	$M-A$=–70142(43) keV for mixture gs+m+n at 13(7) and 200#150 keV											Nub269	**
*^{154}Dy−^{133}Cs$_{1.158}$	No contamination observed, but contamination by ^{154}Tb											00Be42	**
*	cannot be excluded											00Be42	**
*^{154}Ho−u	$M-A$=–64478(28) keV for mixture gs+m at 244(28) keV											Nub269	**
*^{154}Tm−u	$M-A$=–54438(32) keV for mixture gs+m at 70(50) keV											Nub269	**
*^{154}Ce−^{154}Nd	Reported $M-A$=–52097(28) keV, author confirmed mistake in the unc.											24Ki16	**
*^{154}Pr−^{154}Nd	Reported $M-A$=–58005(27) keV, author confirmed mistake in the uncertainty											24Ki16	**
*^{154}Pm−^{154}Nd	possible mixture of gs. and m at 10−20 keV											24Ra.A	**
*^{154}Pm−^{154}Nd	Reported $M-A$=–68303(27) keV, author confirmed mistake in the uncertainty											24Ki16	**
*^{154}Pm−^{154}Nd	possible mixture of gs. and m at 10−20 keV											Nub269	**
*^{154}Sm−^{154}Nd	Reported $M-A$=–72449(27) keV, author confirmed mistake in the unc.											24Ki16	**
*^{154}Tm−^{154}Dy	frequency ratio corrected for the unresolved isomer at 70(50) keV											25Ko14	**
*^{154}Pr(β^-)^{154}Nd	E_{β^-}=7490(100) to 4$^+$ level at 233.2 keV											96To05	**
*^{154}Nd(β^-)^{154}Pm^m	Decay correspond to the excited (low-spin, 1-) isomer in ^{154}Pm											FGK	*K
*^{154}Pm^m(IT)^{154}Pm	From E=10−20 keV											24Ra.A	**
*^{154}Pm(β^-)^{154}Sm	E_{β^-}=2410(180) to 3$^-$ level at 1986.59 keV											Ens09a	*W
*^{154}Pm(β^-)^{154}Sm	E_{β^-}=2400(200), 1850(200) to 2$^+$ levels at 1440.04, 2069.07 keV											Ens09a	**
*^{154}Pm^m(β^-)^{154}Sm	E_{β^-}=3270, 3090, 2810 (all 170) to 921.345 1$^-$, 1099.26 0$^+$, 1475.81 1$^-$											Ens09a	**
*^{154}Pm^m(β^-)^{154}Sm	E_{β^-}=2810(170) to 1$^-$ level at 1475.81 keV, and other E_{β^-}											Ens09a	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value	Adjusted value	v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{154}Pm^m(\beta^-)^{154}Sm$	E_{β^-}=3950(50) 3010(80) to ground state, 1^- level at 921.345 keV								Ens09a	**
$*^{154}Pm^m(\beta^-)^{154}Sm$	E_{β^-}=3000(200), 1900(200), 1800(200) to 1^- level at 921.345, 2^+ at 2069.07,								Ens09a	**
*	and $(1,2^+)$ at 2139.82 keV								Ens09a	**
$*^{154}Eu(\beta^-)^{154}Gd$	E_{β^-}=1855(5) 1844(2) respectively, to 2^+ level at 123.0709 keV								Ens09a	**
$*^{154}Eu(\beta^-)^{154}Gd$	E_{β^-}=257(3) to 2^- level at 1719.5593 keV, and other E_{β^-}								Ens09a	**
$*^{154}Tb(\beta^+)^{154}Gd$	E_{β^+}=2540(50) 1860(50) to ground state and 0^+ level at 680.6673 keV								70Ag02	**
$*^{154}Ho(\beta^+)^{154}Dy$	E_{β^+}=4340(80) to 2^+ level at 334.34 keV								Ens09a	**
$*^{154}Ho^m(\beta^+)^{154}Dy$	E_{β^+}=2500(100) to 7^+ level at 2472.40 keV								Ens09a	**
$*^{154}Tm^m(\beta^+)^{154}Er$	E_{β^+}=4882(150) to 8^+ level 2329.5 keV								Ens09a	**
$*^{154}Lu(\beta^+)^{154}Yb$	p^+ =0.75(0.05) Q =5710(450) from $^{154}Lu^m$ at 200#150 to 8^+ level at 2046.2 keV								Ens09a	**
$*^{154}Lu^m(IT)^{154}Lu$	Use only their Q_α's								GAu	**
$^{155}Ce-{}^{12}C_6\ {}^1H_6$	−98067.9	6.4			2			CP2	1.0	24Ra.A	
$^{155}Pr-u$	−59492	31	−59504.2	1.9	−0.4	U		CP1	1.0	12Va02	*
$^{155}Pr-{}^{80}Kr_{1.938}$	102571	33	102554.6	2.4	−0.5	U		CP1	1.0	12Va02	
$^{155}Pr-{}^{86}Kr_{1.802}$	101588	32	101575.4	1.9	−0.4	U		CP1	1.0	12Va02	
$^{155}Pr-{}^{12}C_6\ {}^1H_6$	−106453.8	2.3	−106454.4	1.9	−0.3	1	71	71 ^{155}Pr	CP2	1.0	24Ra.A	
$^{155}Nd-u$	−66866	17	−66850	3	0.9	U			CP1	1.0	12Va02	*
$^{155}Nd-{}^{80}Kr_{1.938}$	95197	17	95209	3	0.7	U			CP1	1.0	12Va02	
$^{155}Nd-{}^{86}Kr_{1.802}$	94215	17	94230	3	0.9	U			CP1	1.0	12Va02	
$^{155}Nd-u$	−66832.4	22.5	−66850	3	−0.8	U			GR1	1.0	25Sp03	
$^{155}Pm-u$	−71863.8	8.8	−71860.2	2.0	0.4	–			CP1	1.0	12Va02	*
	−71849.1	20.4			−0.5	–			GR1	1.0	25Sp03	
ave.	−71861	8			0.2	1	6	6 ^{155}Pm			average	
$^{155}Pm-{}^{80}Kr_{1.938}$	90197.8	8.6	90198.6	2.4	0.1	1	8	5 ^{155}Pm	CP1	1.0	12Va02	
$^{155}Pm-{}^{86}Kr_{1.802}$	89216.0	8.8	89219.4	2.0	0.4	1	5	5 ^{155}Pm	CP1	1.0	12Va02	
$^{155}Sm-u$	−75357	24	−75354.3	1.2	0.1	U			CP1	1.0	12Va02	*
$^{155}Sm-{}^{80}Kr_{1.938}$	86704	24	86704.6	1.8	0.0	U			CP1	1.0	12Va02	
$^{155}Sm-{}^{86}Kr_{1.802}$	85722	24	85725.4	1.2	0.1	U			CP1	1.0	12Va02	
$^{155}Sm-u$	−75328.5	24.7	−75354.3	1.2	−1.0	U			GR1	1.0	25Sp03	
$C_{12}\ H_{11}-{}^{155}Gd$	163530	70	163447.1	0.9	−0.3	U			R04	4.0	64De15	
$C_{11}\ {}^{13}C\ H_{10}-{}^{155}Gd$	158921	42	158976.9	0.9	0.3	U			R04	4.0	64De15	
$C_{10}\ {}^{13}C_2\ H_9-{}^{155}Gd$	154450	140	154506.7	0.9	0.1	U			R04	4.0	64De15	
$C_{10}\ H_7\ N_2-{}^{155}Gd$	138213	38	138295.0	0.9	0.5	U			R04	4.0	64De15	
$^{155}Gd-{}^{139}La\ O$	21252	32	21350.7	1.1	0.8	U			R05	4.0	65De13	
$^{155}Tb-u$	−76431	30	−76496.9	0.9	−2.2	U			GS2	1.0	05Li24	
$^{155}Dy-u$	−74227	30	−74248.2	2.2	−0.7	U			GS2	1.0	05Li24	
$^{155}Ho-u$	−70867	30	−70901	18	−1.1	1	34	34 ^{155}Ho	GS2	1.0	05Li24	
$^{155}Er-u$	−66785	30	−66790	6	−0.2	U			GS2	1.0	05Li24	
$^{155}Tm-u$	−60814	33	−60790	10	0.7	U			GS2	1.0	05Li24	*
$^{155}Gd\ {}^{35}Cl_3-{}^{149}Sm\ {}^{37}Cl_3$	14282.4	6.3	14289.6	0.8	0.5	U			M21	2.5	75Ka25	
$^{155}Gd\ {}^{35}Cl-{}^{153}Eu\ {}^{37}Cl$	4345.4	2.4	4343.5	0.7	−0.3	U			H25	2.5	72Ba08	
$^{155}Ce-{}^{155}Pm$	20788.1	29.4	20743	7	−1.6	U			RI1	1.0	24Ki16	*
$^{155}Pr-{}^{155}Pm$	12355.1	2.8	12356.0	2.1	0.3	1	58	29 ^{155}Pm	RI1	1.0	24Ki16	*
$^{155}Nd-{}^{155}Pm$	5010.3	2.3				2			RI1	1.0	24Ki16	*
$^{155}Sm-{}^{155}Pm$	−3494.5	2.5	−3494.0	1.9	0.2	1	59	49 ^{155}Pm	RI1	1.0	24Ki16	*
$^{155}Tb-{}^{155}Gd$	874.88	0.19	874.88	0.19	0.0	1	100	100 ^{155}Tb	JY2	1.0	22Ge07	
$^{155}Tm-{}^{155}Er$	6004	26	5999	11	−0.2	1	18	14 ^{155}Tm	TR1	1.0	25Ko14	*
$^{155}Yb-{}^{155}Er$	12638	23	12593	12	−2.0	1	29	24 ^{155}Yb	TR1	1.0	25Ko14	
$^{155}Yb-{}^{155}Eu$	22871	48	22905	12	0.7	1	6	6 ^{155}Yb	TR1	1.0	21Be23	
$^{155}Gd-{}^{138}La\ O$	20558	49	20589.5	1.0	0.2	U			R05	4.0	65De13	
$^{155}Gd-{}^{154}Gd$	1480	60	1756.41	0.19	1.2	U			R04	4.0	64De15	
$^{155}Gd\ O-C_{15}$	−82452.8	5.0	−82457.1	0.9	−0.6	o			TG1	1.5	09Ke.A	
	−82452.2	2.6			−1.3	U			TG1	1.5	11Ke03	
$^{155}Er(\alpha)^{151}Dy$	4118.3	5.	4116	5	−0.5	1	93	91 ^{155}Er	ORa		74To07	Z
$^{155}Tm(\alpha)^{151}Ho$	4578.3	10.3	4572	5	−0.6	–			ORa		71To01	*
	4568.1	10.			0.4	–			ORa		71To01	*
	4570.1	8.			0.2	–					92Ha10	*
ave.	4572	5			0.0	1	96	86 ^{155}Tm			average	
$^{155}Yb(\alpha)^{151}Er$	5344.1	5.	5339.1	2.1	−1.0	–			GSa		79Ho10	
	5336.6	5.			0.5	–			Bka		82Bo04	Z

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{155}Yb(α)^{151}Er		5344.2	5.	5339.1	2.1	−1.0	–						87Ka.A	
		5331.8	4.			1.8	–				ORa		91To08	
		5340.1	4.			−0.2	–				Daa		96Pa01	
	ave.	5338.8	2.1			0.2	1	99	69	^{155}Yb			average	
^{155}Lu(α)^{151}Tm		5796.9	5.	5804.3	2.3	1.4	–						89Ho12	*
		5797.9	5.			1.2	–				ORa		91To08	
		5805.1	5.			−0.2	–				Daa		96Pa01	
		5811.2	5.			−1.4	–				Ara		97Da07	
		5810.2	5.1			−1.2	–				Jya		26Br.B	
	ave.	5804.3	2.3			0.0	1	100	84	^{151}Tm			average	
^{155}Lu^{n}(α)^{151}Tm		7586.1	5.1	7582.6	2.5	−0.7	o				Jya		26Br.B	*
^{155}Lu^{m}(α)^{151}Tm^{m}		5723.0	10.	5730.6	2.8	0.7	3						89Ho12	
		5727.1	5.			0.7	3				ORa		91To08	
		5732.2	5.			−0.3	3				Daa		96Pa01	
		5734.2	5.			−0.7	3				Ara		97Da07	
^{155}Lu^{n}(α)^{151}Tm		7574.9	15.	7582.6	2.5	0.1	U						89Ho12	*
		7586.1	5.			−0.7	o				Daa		96Pa01	*
		7579.0	4.1			0.9	U						18Pa37	
^{155}Gd(n,α)^{152}Sm		8331	6	8339.1	0.3	1.4	U				McM		69Be17	
^{155}Gd(p,t)^{153}Gd		−6850	7	−6848.16	0.24	0.3	U				McM		73Lo08	
		−6853	5			1.0	U				Min		73Oo01	
^{154}Sm(n,γ)^{155}Sm		5806.8	0.6	5806.97	0.27	0.3	–						82Ba15	Z
		5807.0	0.3			−0.1	–				ILn		82Sc03	Z
^{154}Sm(d,p)^{155}Sm		3584	12	3582.40	0.27	−0.1	U				Tal		65Ke09	
^{154}Sm(n,γ)^{155}Sm	ave.	5806.96	0.27	5806.97	0.27	0.0	1	99	90	^{155}Sm			average	
^{154}Eu(n,γ)^{155}Eu		8151.3	0.4	8151.3	0.4	−0.0	1	100	99	^{155}Eu	ILn		86Pr03	
^{154}Gd(n,γ)^{155}Gd		6435.11	0.30	6435.24	0.18	0.4	–				ILn		86Sc25	Z
		6435.29	0.23			−0.2	–				Bdn		06Fi.A	
^{154}Gd(d,p)^{155}Gd		4217	10	4210.67	0.18	−0.6	U				Kop		67Tj01	
^{155}Gd(d,t)^{154}Gd		−190	10	−178.01	0.18	1.2	U				Kop		67Tj01	
^{154}Gd(n,γ)^{155}Gd	ave.	6435.22	0.18	6435.24	0.18	0.1	1	99	53	^{154}Gd			average	
^{155}Ta(p)^{154}Hf		1776	10	1453	15	−32.3	B				Arp		99Uu01	*
		1453	15				3				Jya		07Pa27	
^{155}Nd(β^-)^{155}Pm		4222	150	4667.1	2.1	3.0	B				Ida		93Gr17	
^{155}Pm(β^-)^{155}Sm		3224	30	3254.7	1.8	1.0	U				Ida		93Gr17	
^{155}Sm(β^-)^{155}Eu		1634	15	1628.1	1.1	−0.4	U						63Kr04	*
		1624	15			0.3	U						65Fu13	*
		1607	25			0.8	U				Ida		93Gr17	
^{155}Eu(β^-)^{155}Gd		252	5	251.2	0.8	−0.2	U						54Le08	
		245	5			1.2	U						58Gl56	
		245	5			1.2	U						59Am16	
^{155}Dy(β^+)^{155}Tb		2099	6	2094.6	1.9	−0.7	–						63Pe13	*
		2094	2			0.3	–						80Bu04	*
	ave.	2094.5	1.9			0.1	1	100	100	^{155}Dy			average	
^{155}Ho(β^+)^{155}Dy		3102	20	3118	16	0.8	1	66	66	^{155}Ho			72To07	*
^{155}Lu^{m}(IT)^{155}Lu		23.0	6.2	21	4	−0.2	2						96Pa01	
		19.9	6.2			0.3	2						97Da07	
^{155}Lu^{n}(IT)^{155}Lu		1781	2	1778.3	1.0	−1.3	2						96Pa01	
		1787.0	1.7			−5.1	o						21Br.A	*
		1777.5	1.1			0.7	2				Jya		26Br.B	*
*^{155}Pr−u	Represents frequency ratio $^{155}Pr^{++}/(C_{12}H_4)^{+}$ =0.98142559(20)												WgM124	**
*^{155}Nd−u	Represents frequency ratio $^{155}Nd^{++}/(C_{12}H_4)^{+}$ =0.98147230(11)												WgM124	**
*^{155}Pm−u	Represents frequency ratio $^{155}Pm^{++}/(C_{12}H_4)^{+}$ =0.981503965(56)												WgM124	**
*^{155}Sm−u	Represents frequency ratio $^{155}Sm^{++}/(C_{12}H_4)^{+}$ =0.98152610(15)												WgM124	**
*^{155}Tm−u	$M-A$=–56627(28) keV for mixture gs+m at 41(6) keV												Nub269	**
*^{155}Ce−^{155}Pm	Reported $M-A$=–47576(31) keV, author confirmed mistake in the unc.												24Ki16	**
*^{155}Pr−^{155}Pm	Reported $M-A$=–55431(14) keV, author confirmed mistake in the unc.												24Ki16	**
*^{155}Nd−^{155}Pm	Reported $M-A$=–62273(14) keV, author confirmed mistake in the unc.												24Ki16	**
*^{155}Sm−^{155}Pm	Reported $M-A$=–70195(14) keV, author confirmed mistake in the unc.												24Ki16	**
*^{155}Tm−^{155}Er	frequency ratio corrected for the unresolved isomer at 41(6) keV												25Ko14	**
*^{155}Tm(α)^{151}Ho	First assigned to ^{156}Tm^{m} but belonging to ^{155}Tm ground state												94To10	**
*^{155}Tm(α)^{151}Ho	Doublet from ground state and isomer, less than 5 keV apart												90Po13	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value	Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
$*^{155}$Lu(α)^{151}Tm	Original value E=5656(6) (Q =5806.1) recalibrated											79Ho10	**
$*^{155}$Lu^n(α)^{151}Tm	Replaced by authors' value for ^{155}Lu^n(IT)											RCh266	**
$*^{155}$Lu^n(α)^{151}Tm	Original value E=7408(10) recalibrated											81Ho.A	**
$*^{155}$Lu^n(α)^{151}Tm	Replaced by authors' value for ^{155}Lu^n(IT)											AHW	**
$*^{155}$Ta(p)^{154}Hf	E_p=1765(10) for (11/2$^-$) state; ground state may be 1/2$^+$, slightly lower											99Uu01	**
*	1776 keV proton not observed in coincidence with feeding α											07Pa27	**
$*^{155}$Sm(β^-)^{155}Eu	E_{β^-}=1530(15) E_{β^-}=1520(15) respectively, to 5/2$^-$ level at 104.334 keV											Ens051	**
$*^{155}$Dy(β^+)^{155}Tb	E_{β^+}=850(6) 845(2) respectively, to 5/2$^-$ level at 226.918 keV, and other E_{β^+}											Ens051	**
$*^{155}$Ho(β^+)^{155}Dy	E_{β^+}=1840(20) to 3/2$^+$ level at 240.196 keV											Ens051	**
$*^{155}$Lu^n(IT)^{155}Lu	E(γ)=295(1)+685(1)+807(1) in a cascade											21Br.A	**
$*^{155}$Lu^n(IT)^{155}Lu	E(γ)=286(1),684.8(0.3),806.7(0.3) in a cascade											26Br.B	**
^{156}Pr−u		−55233.1	1.1				2				CP2	1.0	22Or02	*
^{156}Nd−^{136}Xe$_{1.147}$		41786.8	2.6	41795.4	1.4	3.3	B				JY1	1.0	18Vi02	
^{156}Nd−^{82}Kr$_{1.902}$		99929.2	1.4				2				CP2	1.0	18Or02	
^{156}Nd−u		−64596.2	23.6	−64629.6	1.4	−1.4	U				GR1	1.0	25Sp03	
^{156}Pm−u		−68883.4	6.9	−68886.0	1.3	−0.4	–				CP1	1.0	12Va02	*
		−68887.8	2.5			0.7	–				CP2	1.0	20Or03	*
	ave.	−68887.3	2.4			0.5	1	31	31	^{156}Pm			average	
^{156}Pm−^{80}Kr$_{1.950}$		94181.7	6.4	94176.3	1.9	−0.8	1	9	5	^{80}Kr	CP1	1.0	12Va02	
^{156}Pm−^{86}Kr$_{1.814}$		93269.1	6.8	93266.3	1.3	−0.4	U				CP1	1.0	12Va02	
^{156}Pm−u		−68919.4	53.7	−68886.0	1.3	0.6	U				GR1	1.0	25Sp03	*
^{156}Pm^m−u		−68724.4	1.6	−68724.7	1.3	−0.2	1	66	66	^{156}Pm^m	CP2	1.0	20Or03	*
^{156}Sm−u		−74444.9	20.4	−74460	8	−0.8	1	17	17	^{156}Sm	GR1	1.0	25Sp03	
C_{12} H_{12}−^{156}Gd		171923	44	171771.4	0.9	−0.9	U				R04	4.0	64De15	
C_{11} ^{13}C H_{11}−^{156}Gd		167384	43	167301.2	0.9	−0.5	U				R04	4.0	64De15	
C_{10} $^{13}C_2$ H_{10}−^{156}Gd		162810	60	162831.0	0.9	0.1	U				R04	4.0	64De15	
C_{10} H_8 N_2−^{156}Gd		146661	38	146619.3	0.9	−0.3	U				R04	4.0	64De15	
^{156}Tb−u		−75181	52	−75247	4	−1.3	U				GS2	1.0	05Li24	*
C_{10} H_8 N_2−^{156}Dy		145130	100	144465.8	0.9	−1.7	U				R04	4.0	64De15	
^{156}Dy−^{133}Cs$_{1.173}$		35195.1	4.5	35187.3	0.9	−1.7	U				MA8	1.0	19Hu15	
^{156}Ho−u		−70130	120	−70360	40	−1.9	o				GS1	1.0	00Ra23	*
^{156}Ho^n−u		−70107	30				2				GS2	1.0	05Li24	*
^{156}Er−u		−68907	30	−68933	26	−0.9	1	77	77	^{156}Er	GS2	1.0	05Li24	
^{156}Tm−u		−61044	30	−61008	14	1.2	U				GS2	1.0	05Li24	
^{156}Yb−u		−57202	30	−57181	10	0.7	U				GS2	1.0	05Li24	
^{156}Gd ^{35}Cl−^{154}Gd ^{37}Cl		4199	5	4207.30	0.22	0.4	U				H12	4.0	64Ba15	
		4206	10			0.1	U				H21	2.5	70Ma05	
		4204.8	1.4			0.7	U				H25	2.5	72Ba08	
		4203.0	1.0			1.7	U				M21	2.5	75Ka25	
^{156}Dy−^{156}Gd		2153.47	0.11	2153.47	0.11	0.0	1	100	100	^{156}Dy	SH1	1.0	11El05	
^{156}Tm−^{156}Yb		−3814	23	−3827	12	−0.6	1	26	22	^{156}Tm	TR1	1.0	25Ko14	
^{156}Yb−^{156}Tm		3761	58	3827	12	1.1	U				TR1	1.0	21Be23	
^{156}Gd−^{139}La O		20618	71	20851.5	1.1	0.8	U				R05	4.0	65De13	
^{156}Gd−^{155}Gd		−584	33	−499.23	0.07	0.6	U				R04	4.0	64De15	
^{156}Gd O−C_{15}		−82946.5	5.8	−82956.4	0.9	−1.1	o				TG1	1.5	09Ke.A	
		−82945.6	3.6			−2.0	U				TG1	1.5	11Ke03	
^{156}Er(α)^{152}Dy		3109.9	70.	3485	25	5.4	C						95Ka.A	
^{156}Tm(α)^{152}Ho		4341.6	10.	4347	7	0.5	–				ORa		71To10	
		4345.6	10.			0.1	–						81Ga36	
	ave.	4344	7			0.4	1	91	72	^{156}Tm			average	
^{156}Tm^m(α)^{152}Ho		4737.5	10.	*			F				ORa		71To01	*
^{156}Yb(α)^{152}Er		4813.6	10.	4810	3	−0.4	–						77Ha48	
		4809.6	10.			0.0	–				GSa		79Ho10	
		4810.6	4.			−0.2	–				Daa		96Pa01	
	ave.	4811	4			−0.3	1	96	78	^{156}Yb			average	
^{156}Lu(α)^{152}Tm		5713.7	5.	5710.4	2.2	−0.6	2				GSa		79Ho10	Z
		5709.7	5.			0.1	2				Dba		92Po14	
		5709.7	8.			0.1	2						92Ha10	
		5711.7	4.			−0.3	2				Daa		96Pa01	
		5707.6	4.1			0.7	2				Jya		19Pa27	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{156}Lu$^{m}(\alpha)^{152}$Tm^{m}		5593.7	10.	5596	3	0.2	U				GSa		79Ho10	
		5592.7	5.			0.6	2				Dba		92Po14	
		5597.9	4.			−0.5	2				Daa		96Pa01	
^{156}Hf$(\alpha)^{152}$Yb		6033.0	10.	6026	3	−0.7	–				GSa		79Ho10	
		6027.9	4.			−0.5	–				Daa		96Pa01	
	ave.	6029	4			−0.7	1	65	65	^{156}Hf			average	
^{156}Hf$^{m}(\alpha)^{152}$Yb		8009.8	15.	7985	3	−1.7	U				GSa		81Ho.A	
		7987.2	4.			−0.6	o				Daa		96Pa01	*
		7980.0	5.1			0.9	1	37	37	^{156}Hf^{m}			18Pa37	
^{154}Sm(t,p)^{156}Sm		4556	25	4564	8	0.3	1	10	10	^{156}Sm	Ald		66Bj01	
^{154}Eu(t,p)^{156}Eu		6003	4	6005	3	0.4	1	69	69	^{156}Eu	LAl		84La06	*
^{154}Gd(t,p)^{156}Gd		6495.1	3.6	6489.79	0.19	−1.5	U				McM		89Lo07	
^{156}Gd(p,t)^{154}Gd		−6490	7	−6489.79	0.19	0.0	U				McM		73Lo08	
		−6490	5			0.0	U				Min		73Oo01	
^{155}Gd(n,γ)^{156}Gd		8536.8	0.5	8536.35	0.07	−0.9	U				ILn		82Ba28	
		8536.39	0.07			−0.6	–				MMn		82Is05	Z
		8536.04	0.19			1.6	–				Bdn		06Fi.A	
^{155}Gd(d,p)^{156}Gd		6319	10	6311.78	0.07	−0.7	U				Kop		67Tj01	
^{156}Gd(d,t)^{155}Gd		−2287	10	−2279.12	0.07	0.8	U				Kop		67Tj01	
^{155}Gd(n,γ)^{156}Gd	ave.	8536.35	0.07	8536.35	0.07	0.0	1	100	52	^{155}Gd			average	
^{155}Gd(α,t)^{156}Tb−^{158}Gd()^{159}Tb		−821.9	3.6	−822	4	0.0	1	100	100	^{156}Tb	McM		75Bu02	
^{156}Dy(d,t)^{155}Dy		−3184	10	−3182.8	1.9	0.1	U				Kop		70Gr46	
^{156}Ta(p)^{155}Hf		1028.6	13.	1020	4	−0.7	o				Dap		92Pa05	
		1013.6	5.			1.2	o				Dap		96Pa01	
		1017.9	5.			0.4	3				Dap		11Da12	
^{156}Ta^{m}(p)^{155}Hf		1110.2	12.	1114	7	0.3	3				Dap		93Li34	
		1115.2	8.			−0.2	3				Dap		96Pa01	
^{156}Nd$(\beta^-)^{156}$Pm		3690	200	3964.8	1.8	1.4	U				Kur		02Sh.B	
^{156}Pm^{m}(IT)^{156}Pm		150.3	0.1	150.30	0.10	0.0	1	100	66	^{156}Pm			07Sh05	
^{156}Pm$(\beta^-)^{156}$Sm		5155	35	5192	8	1.1	U				Stu		90He11	
		5110	100			0.8	U				Kur		02Sh.B	
^{156}Sm$(\beta^-)^{156}$Eu		721	10	725	7	0.4	–						63Gu04	*
		721	15			0.2	–						65Wi08	*
	ave.	721	8			0.4	1	78	74	^{156}Sm			average	
^{156}Eu$(\beta^-)^{156}$Gd		2430	10	2452	3	2.2	–						62Ew01	
		2460	10			−0.8	–						63Th02	
		2450	15			0.2	–						64Pe17	
		2478	20			−1.3	U						67Va23	
	ave.	2446	6			1.0	1	28	27	^{156}Eu			average	
^{156}Tb$(\beta^+)^{156}$Gd		3570	50	2445	4	−22.5	B						70Ag02	*
^{156}Ho$(\beta^+)^{156}$Dy		4400	400	4990	40	1.5	U						76Gr20	*
		5050	90			−0.7	o						02Iz01	
		5050	60			−1.0	2						04Iz02	*
		4950	50			0.8	2						96By.A	
^{156}Er$(\beta^+)^{156}$Ho		1670	70	1330	50	−4.9	B						82Vy06	*
^{156}Tm$(\beta^+)^{156}$Er		7458	50	7381	26	−1.5	1	28	23	^{156}Er	Dbn		94Po26	*
		7390	100			−0.1	U						95Ga.A	
^{156}Hf^{m}(IT)^{156}Hf		1959	1	1958.8	1.0	−0.2	1	98	63	^{156}Hf^{m}			96Pa01	
*^{156}Pr−u	Represents frequency ratio $(C_6H_6)^+/^{156}Pr^{++}$ =0.9990445852(69)												HWJ208	**
*^{156}Pm−u	Represents frequency ratio $^{156}Pm^{++}/(C_{12}H_4)^+$ =0.975190689(43)												WgM124	**
*^{156}Pm−u	Represents frequency ratio $^{156}Pm^{++}/(C_6H_6)^+$ =0.998957107(16)												HWJ208	**
*^{156}Pm−u	Corrected for unresolved isomeric state using exc. energy in 23Ga02												25Sp03	**
*^{156}Pm^{m}−u	Represents frequency ratio $^{156}Pmm^{++}/(C_6H_6)^+$ =0.998958154(10)												HWJ208	**
*^{156}Tb−u	$M-A$=–69968(32) keV for mixture gs+m+n at 88.4(0.2) and 100#50 keV												Nub269	**
*^{156}Ho−u	$M-A$=–65230(100) keV for mixture gs+m+n at 52.37 and 240(50) keV												Nub269	**
*^{156}Ho^{n}−u	Assuming high spin isomer is favored												GAu	**
*^{156}Tm$^{m}(\alpha)^{152}$Ho	F : originally E_α=4460(10) to ^{152}Ho^{m} at 160(1), reassigned to ^{155}Tm												94To10	**
*^{156}Hf$^{m}(\alpha)^{152}$Yb	Replaced by authors' value for ^{156}Hf^{m}(IT)												AHW	**
*^{154}Eu(t,p)^{156}Eu	Q =5569(4) to 3^- level at 434.23 keV												91Ba06	**
*^{156}Sm$(\beta^-)^{156}$Eu	E_{β^-}=430(10) 430(15) respectively, to 1^+ level at 291.3037 keV												Ens12a	**
*^{156}Tb$(\beta^+)^{156}$Gd	E_{β^+}=2640(50) from ^{156}Tb^{m} at 88.4 to ground state												Nub269	**
*^{156}Ho$(\beta^+)^{156}$Dy	E_{β^+}=1800(50) to levels around 1600												Ens12a	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{156}$Ho(β^+)^{156}Dy		Original error 20 is for statistics only, increased by evaluator										GAu	**
$*^{156}$Er(β^+)^{156}Ho		p^+ =0.0036(0.0017) to 2^- level at 82.23 keV, reanalyzed										Ens12a	**
$*^{156}$Tm(β^+)^{156}Er		E_{β^+}=6091(50) to 2^+ level at 344.53 keV										Ens12a	**
^{157}Pr−u		−51996.9	3.4				2			CP2	1.0	22Or02	*
^{157}Nd−u		−60614	47	−60649.1	2.3	−0.7	U			CP1	1.0	12Va02	*
		−60649.1	2.3				2			CP2	1.0	22Or02	*
^{157}Nd−^{80}Kr$_{1.963}$		103537	46	103500.3	2.7	−0.8	U			CP1	1.0	12Va02	
^{157}Nd−^{86}Kr$_{1.826}$		102610	46	102575.9	2.3	−0.7	U			CP1	1.0	12Va02	
^{157}Pm−u		−66880	13	−66878	8	0.1	1	33	33 ^{157}Pm	CP1	1.0	12Va02	*
^{157}Pm−^{80}Kr$_{1.963}$		97273	13	97271	8	−0.2	1	34	33 ^{157}Pm	CP1	1.0	12Va02	
^{157}Pm−^{86}Kr$_{1.826}$		96346	13	96347	8	0.0	1	33	33 ^{157}Pm	CP1	1.0	12Va02	
^{157}Pm−u		−66870.0	33.3	−66878	8	−0.3	U			GR1	1.0	25Sp03	
^{157}Sm−u		−71582.2	8.3	−71581	5	0.1	1	33	33 ^{157}Sm	CP1	1.0	12Va02	*
^{157}Sm−^{80}Kr$_{1.963}$		92570.0	8.0	92568	5	−0.2	1	36	34 ^{157}Sm	CP1	1.0	12Va02	
^{157}Sm−^{86}Kr$_{1.826}$		91643.0	8.3	91644	5	0.1	1	33	33 ^{157}Sm	CP1	1.0	12Va02	
^{157}Sm−u		−71575.3	21.5	−71581	5	−0.3	U			GR1	1.0	25Sp03	
^{157}Eu−u		−74575.9	54.8	−74569	5	0.1	U			GR1	1.0	25Sp03	
C_{10} H_9 N_2−^{157}Gd		152720	60	152607.0	0.9	−0.5	U			R04	4.0	64De15	
C_9 ^{13}C H_8 N_2−^{157}Gd		148170	70	148136.8	0.9	−0.1	U			R04	4.0	64De15	
C_{10} H_5 O_2−^{157}Gd		105080	60	104988.1	0.9	−0.4	U			R04	4.0	64De15	
^{157}Ho−u		−71724	30	−71748	25	−0.8	1	71	71 ^{157}Ho	GS2	1.0	05Li24	
^{157}Er−u		−68084	30	−68077	28	0.2	1	90	90 ^{157}Er	GS2	1.0	05Li24	
^{157}Tm−u		−63027	30	−63034	21	−0.2	1	50	50 ^{157}Tm	GS2	1.0	05Li24	
^{157}Yb−u		−57389	30	−57356	11	1.1	U			GS2	1.0	05Li24	
^{157}Lu−u		−49842	31	−49847	12	−0.2	1	15	15 ^{157}Lu	GS2	1.0	05Li24	*
^{157}Gd ^{35}Cl−^{155}Gd ^{37}Cl		4318	4	4288.22	0.19	−1.9	U			H12	4.0	64Ba15	
		4287	3			0.2	U			H21	2.5	70Ma05	
		4289.0	0.7			−0.4	U			M21	2.5	75Ka25	
		4288.83	0.66			−0.4	U			H41	2.5	85Dy04	
^{157}Yb−^{157}Tm		5705	50	5677	21	−0.6	o			TR1	1.0	21Be23	
^{157}Tm−^{157}Yb		−5683	28	−5677	21	0.2	1	57	50 ^{157}Tm	TR1	1.0	25Ko14	*
^{157}Lu−^{157}Yb		7554	25	7509	13	−1.8	1	27	16 ^{157}Lu	TR1	1.0	25Ko14	*
^{157}Gd−^{156}Gd		1860	60	1837.33	0.16	−0.1	U			R04	4.0	64De15	
^{157}Gd O−C_{15}		−81114.2	5.4	−81119.1	0.9	−0.6	o			TG1	1.5	09Ke.A	
		−81113.6	3.3			−1.1	U			TG1	1.5	11Ke03	
^{157}Yb(α)^{153}Er		4622.0	7.	4620	6	−0.3	–					77Ha48	
		4623.0	10.			−0.3	–			GSa		79Ho10	
	ave.	4622	6			−0.5	1	93	79 ^{157}Yb			average	
^{157}Lu(α)^{153}Tm		5097.2	5.	5108.3	2.8	2.2	o			Dba		91Le15	*
		5096.2	20.			0.6	U			Bka		91To09	*
		5111.5	5.			−0.6	o			Dba		92Po14	*
^{157}Lu^m(α)^{153}Tm		5128.9	10.	5129.0	2.0	0.0	U			IRa		79Al16	Z
		5131.8	5.			−0.5	–			GSa		79Ho10	Z
		5133.7	5.			−0.9	–			ORa		83To01	Z
		5128.9	5.			0.0	o			Dba		91Le15	
		5118.7	5.			2.0	–			Bka		91To09	
		5125.8	6.			0.5	–					92Ha10	
		5132.0	5.			−0.6	–			Dba		92Po14	
		5127.9	4.			0.3	–			Daa		96Pa01	
	ave.	5128.4	2.1			0.3	1	99	62 ^{153}Tm			average	
^{157}Hf(α)^{153}Yb		5869.4	10.	5880	3	1.0	3					73Ea01	Z
		5884.1	5.			−0.8	3			GSa		79Ho10	Z
		5879.1	4.			0.2	3			Daa		96Pa01	
^{157}Ta(α)^{153}Lu^m		6277.2	4.	6275	8	−0.6	o			Ara		97Ir01	*
^{157}Ta^m(α)^{153}Lu		6381.9	10.	6377	4	−0.5	3			GSa		79Ho10	
		6375.8	4.			0.2	3			Daa		96Pa01	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{157}Ta$^{n}(\alpha)^{153}$Lu	7946.9	8.	7948	8	0.0	o			Daa		96Pa01	*
^{155}Gd(t,p)^{157}Gd	6417.8	2.9	6414.41	0.16	−1.2	U			McM		89Lo07	
^{157}Gd(p,t)^{155}Gd	−6414	7	−6414.41	0.16	−0.1	U			McM		73Lo08	
	−6417	5			0.5	U			Min		73Oo01	
^{156}Gd(n,γ)^{157}Gd	6359.6	0.8	6359.86	0.15	0.3	U					70Bo29	
	6360	1			−0.1	U					71Gr42	
	6359.80	0.15			0.4	o			ILn		87Sp.A	Z
	6359.86	0.15			0.0	1	98	55 ^{156}Gd	ILn		03Bo25	
^{157}Gd(γ,n)^{156}Gd	−6350	80	−6359.86	0.15	−0.1	U			Phi		60Ge01	
^{156}Gd(d,p)^{157}Gd	4136	10	4135.29	0.15	−0.1	U			Kop		67Tj01	
^{157}Gd(d,t)^{156}Gd	−112	10	−102.63	0.15	0.9	U			Kop		67Tj01	
^{156}Gd(α,t)^{157}Tb−^{158}Gd()^{159}Tb	−616.2	2.0	−614.0	0.8	1.1	1	14	10 ^{159}Tb	McM		75Bu02	
^{156}Dy(d,p)^{157}Dy	4748	10	4742	5	−0.6	–			Tal		68Be.A	
	4753	10			−1.1	–			Kop		70Gr46	
ave.	4751	7			−1.2	1	52	52 ^{157}Dy			average	
^{157}Ta(p)^{156}Hf	925.0	17.	935	10	0.6	o			Dap		96Pa01	
	933.0	7.			0.2	o			Ara		97Ir01	*
^{157}Pm(β^-)^{157}Sm	4360	100	4381	8	0.2	U			Kur		02Sh.B	
^{157}Sm(β^-)^{157}Eu	2700	200	2783	6	0.4	U					73Ka23	*
	2734	50			1.0	U			Ida		93Gr17	
^{157}Eu(β^-)^{157}Gd	1350	20	1365	4	0.7	U					64Sh21	*
	1370	20			−0.3	U					66Fu05	*
^{157}Tb(ε)^{157}Gd	62.4	0.6	60.19	0.12	−3.7	B					67Na08	*
	62.2	0.6			−3.4	B					83Be42	*
	60.0	0.3			0.6	–					92Ra18	
	60.2	0.2			−0.1	–					98Pa.1	
	60.23	0.18			−0.2	–					24Ri04	
ave.	60.18	0.12			0.1	1	100	96 ^{157}Tb			average	
^{157}Ho(β^+)^{157}Dy	2540	50	2592	24	1.0	1	23	22 ^{157}Ho			72To05	*
^{157}Er(β^+)^{157}Ho	3470	80	3420	30	−0.6	1	18	10 ^{157}Er			75Al.A	
	3805	100			−3.9	B			Dbn		94Po26	*
^{157}Tm(β^+)^{157}Er	4480	100	4700	30	2.2	B			IRS		93Al03	
	4482	100			2.2	B			Dbn		94Po26	
^{157}Yb(β^+)^{157}Tm	5074	100	5288	20	2.1	U			Dbn		94Po26	
^{157}Lu^{m}(IT)^{157}Lu	32	2	20.7	2.0	−5.7	B			Dba		91Le15	
	21	2			−0.2	1	99	69 ^{157}Lu	Dba		92Po14	*
^{157}Ta^{m}(IT)^{157}Ta	22	5				3					97Ir01	
^{157}Ta^{n}(IT)^{157}Ta^{m}	1571	7				3			Daa		96Pa01	
$*^{157}$Pr−u	Represents frequency ratio $(C_6H_6)^+/^{157}Pr^{++}$ =1.005471763(22)										HWJ208	**
$*^{157}$Nd−u	Represents frequency ratio $^{157}Nd^{++}/(C_{12}H_4)^+$ =0.96892546(29)										WgM124	**
$*^{157}$Nd−u	Represents frequency ratio $(C_6H_6)^+/^{157}Nd^{++}$ =1.005416333(17)										HWJ208	**
$*^{157}$Pm−u	Represents frequency ratio $^{157}Pm^{++}/(C_{12}H_4)^+$ =0.968964141(81)										WgM124	**
$*^{157}$Sm−u	Represents frequency ratio $^{157}Sm^{++}/(C_{12}H_4)^+$ =0.968993178(51)										WgM124	**
$*^{157}$Lu−u	$M-A$=–46417(28) keV for mixture gs+m at 20.7(2.0) keV										Nub269	**
$*^{157}$Tm−^{157}Yb	Could be a mixture of gs. and iso. at 100#(50#) keV										HWJ25b	**
$*^{157}$Lu−^{157}Yb	frequency ratio corrected for the unresolved isomer at 21(2) keV										25Ko14	**
$*^{157}$Lu(α)^{153}Tm	E_α=4925(5) to ^{153}Tm^{m} at 43.2(0.2) keV										Nub269	**
$*^{157}$Lu(α)^{153}Tm	E_α=4924(20) to ^{153}Tm^{m} at 43.2(0.2) keV										Nub269	**
$*^{157}$Lu(α)^{153}Tm	E_α=4939(5) to ^{153}Tm^{m} at 43.2(0.2) keV; replaced by ^{157}Lu^{m}(IT)										Nub269	**
$*^{157}$Ta(α)^{153}Lu^{m}	Replaced by ^{153}Lu^{m}(IT)										AHW	**
$*^{157}$Ta$^{m}(\alpha)^{153}$Lu	Reassigned										97Ir01	**
$*^{157}$Ta$^{n}(\alpha)^{153}$Lu	Replaced by authors' value for ^{157}Ta^{n}(IT)										AHW	**
$*^{157}$Ta(p)^{156}Hf	Use instead ^{157}Ta^{m}(IT)										AHW	**
$*^{157}$Sm(β^-)^{157}Eu	E_{β^-}=2400(200) to 5/2$^-$ level at 197.863 and 3/2$^+$ at 394.334 keV										Ens162	**
$*^{157}$Eu(β^-)^{157}Gd	E_{β^-}=870(30) 910(20) respectively, to 3/2$^+$ level at 474.630 keV, and other E_{β^-}										Ens162	**
$*^{157}$Tb(ε)^{157}Gd	LK=2.65(0.20); original value 66(6) recalculated										92Ha03	**
$*^{157}$Tb(ε)^{157}Gd	LK=2.69(0.20); original value 62.9(0.7) recalculated										85Vo09	**
$*^{157}$Ho(β^+)^{157}Dy	E_{β^+}=1180(50) to 5/2$^-$ level at 341.118 keV										Ens162	**
$*^{157}$Er(β^+)^{157}Ho	E_{β^+}=2525(100) to ground state yielding 3547(100), rather 24% to (3/2$^+$)										94Po26	**
*	level at 174.55 keV, 15% to 5/2$^-$ at 391.32 keV → +258 keV										Ens162	**
$*^{157}$Lu^{m}(IT)^{157}Lu	Derived from ^{157}Lu$^{m}(\alpha)$-^{157}Lu(α) difference										92Po14	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{158}Nd−^{136}Xe$_{1.162}$		49956	39	50022.4	1.4	1.7	U			JY1	1.0	18Vi02	
^{158}Nd−^{84}Kr$_{1.881}$		108678.4	1.4				2			CP2	1.0	18Or02	
^{158}Pm−u		−63436	25	−63453.5	0.9	−0.7	U			CP1	1.0	12Va02	*
		−63453.7	1.1			0.2	1	71	71 ^{158}Pm	CP2	1.0	20Or03	*
^{158}Pm−^{80}Kr$_{1.975}$		101720	25	101699.4	1.7	−0.8	U			CP1	1.0	12Va02	
^{158}Pm−^{86}Kr$_{1.837}$		100773	25	100754.8	0.9	−0.7	U			CP1	1.0	12Va02	
^{158}Pm−u		−63451.8	32.2	−63453.5	0.9	−0.1	U			GR1	1.0	25Sp03	
^{158}Sm−u		−70049.2	9.5	−70051	5	−0.1	1	29	29 ^{158}Sm	CP1	1.0	12Va02	*
^{158}Sm−^{80}Kr$_{1.975}$		95106.5	9.1	95102	5	−0.5	1	33	31 ^{158}Sm	CP1	1.0	12Va02	
^{158}Sm−^{86}Kr$_{1.837}$		94159.3	9.4	94158	5	−0.2	1	30	30 ^{158}Sm	CP1	1.0	12Va02	
^{158}Sm−u		−70060.6	22.5	−70051	5	0.4	U			GR1	1.0	25Sp03	
^{158}Eu−u		−72208	25	−72217.8	2.2	−0.4	U			CP1	1.0	12Va02	*
		−72218.2	2.2			0.2	1	98	98 ^{158}Eu	CP2	1.0	20Or03	*
		−72219.5	34.4			0.0	U			GR1	1.0	25Sp03	
^{158}Eu−^{80}Kr$_{1.975}$		92949	25	92935.1	2.6	−0.6	U			CP1	1.0	12Va02	
^{158}Eu−^{86}Kr$_{1.837}$		92001	25	91990.5	2.2	−0.4	U			CP1	1.0	12Va02	
C_{10} H_6 O_2−^{158}Gd		112444	33	112669.3	0.9	1.7	U			R04	4.0	64De15	
C_{10} H_6 O_2−^{158}Dy		112870	100	112365.0	2.5	−1.3	U			R04	4.0	64De15	
^{158}Ho−u		−71101	67	−71056	29	0.7	R			GS2	1.0	05Li24	*
^{158}Er−u		−70220	110	−70107	27	1.0	U			GS1	1.0	00Ra23	
		−70107	30			0.0	1	81	81 ^{158}Er	GS2	1.0	05Li24	
^{158}Tm−u		−63080	110	−63020	27	0.5	U			GS1	1.0	00Ra23	
		−63020	30			−0.0	1	81	81 ^{158}Tm	GS2	1.0	05Li24	
^{158}Yb−^{142}Sm$_{1.113}$		34252	22	34243	9	−0.4	1	15	15 ^{158}Yb	MA7	1.0	01Bo59	
^{158}Lu−u		−50720	30	−50681	15	1.3	R			GS2	1.0	05Li24	
^{158}Gd ^{35}Cl−^{156}Gd ^{37}Cl		4956	4	4931.23	0.19	−1.5	U			H12	4.0	64Ba15	
		4929	3			0.3	U			H21	2.5	70Ma05	
		4926.2	1.4			1.4	U			H25	2.5	72Ba08	
		4930.8	0.7			0.2	U			M21	2.5	75Ka25	
		4930.13	1.36			0.3	U			H41	2.5	85Dy04	
^{158}Dy ^{35}Cl−^{156}Dy ^{37}Cl		3081.4	3.3	3082.0	2.5	0.1	U			H25	2.5	72Ba08	
^{158}Pm−^{158}Gd		12436.8	1.4	12436.4	1.0	−0.3	1	53	29 ^{158}Pm	JY1	1.0	18Vi02	
^{158}Gd−^{157}Gd		392	48	143.78	0.07	−1.3	U			R04	4.0	64De15	
^{158}Gd O−C_{15}		−80968.3	5.4	−80975.3	0.9	−0.9	o			TG1	1.5	09Ke.A	
		−80967.8	3.2			−1.6	U			TG1	1.5	11Ke03	
^{158}Gd O−C_{14}		−80964.7	8.2			−0.9	U			TG1	1.5	11Ke03	
^{158}Yb(α)^{154}Er		4174.9	10.	4171	7	−0.4	–					77Ha48	
		4164.6	12.			0.5	–					92Ha10	
	ave.	4171	8			−0.0	1	80	71 ^{158}Yb			average	
^{158}Lu(α)^{154}Tm		4792.2	10.	4790	5	−0.2	3			IRa		79Al16	Z
		4789.5	5.			0.1	3			ORa		83To01	Z
^{158}Hf(α)^{154}Yb		5406.0	5.	5404.7	2.7	−0.3	–			GSa		79Ho10	Z
		5401.4	5.			0.6	–			ORa		83To01	Z
		5406.1	4.			−0.3	–			Daa		96Pa01	
	ave.	5404.8	2.7			−0.0	1	100	88 ^{158}Hf			average	
^{158}Ta(α)^{154}Lu		6124.4	8.	6124	4	−0.1	9			Daa		96Pa01	
		6123.3	5.			0.1	9			Ara		97Da07	
^{158}Ta^m(α)^{154}Lu^m		6208.5	6.	6202.8	2.3	−0.9	10			GSa		79Ho10	
		6203.4	4.			−0.1	10			Daa		96Pa01	
		6205.4	5.			−0.5	10			Ara		97Da07	
		6198.2	4.1			1.1	10			Jya		19Pa27	
^{158}Ta^n(α)^{154}Lu^m		8869.0	11.3				11			Jya		14Ca03	
^{158}W(α)^{154}Hf		6600.4	30.	6612.5	2.6	0.4	U			GSa		81Ho10	*
		6609.7	30.			0.1	U			Daa		96Pa01	
		6612.7	3.			−0.1	3			Ara		00Ma95	
		6611.7	5.1			0.1	3			Jya		19Hi06	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$^{158}W^{m}(\alpha)^{154}Hf$		8495.5	30.	8502	7	0.2	U			GSa		89Ho12	
		8506.8	24.			−0.2	U			Daa		96Pa01	
		8501.6	7.				3			Ara		00Ma95	
$^{158}Gd(p,t)^{156}Gd$		−5818	5	−5815.45	0.16	0.5	U			Min		73Oo01	
$^{158}Dy(p,t)^{156}Dy$		−7535	15	−7538.0	2.4	−0.2	U			Pri		77Ko04	
$^{158}Gd(t,\alpha)^{157}Eu-^{156}Gd()^{155}Eu$		−512	5	−514	4	−0.5	1	69	68 ^{157}Eu	LAl		79Bu05	
$^{157}Gd(n,\gamma)^{158}Gd$		7937.39	0.07	7937.39	0.06	−0.0	–			MMn		82Is05	Z
		7937.39	0.17			−0.0	–			Bdn		06Fi.A	
$^{157}Gd(d,p)^{158}Gd$		5724	10	5712.82	0.06	−1.1	U			Kop		67Tj01	
		5706	5			1.4	U			Tal		71Sh04	
$^{158}Gd(d,t)^{157}Gd$		−1688	10	−1680.16	0.06	0.8	U			Kop		67Tj01	
$^{157}Gd(n,\gamma)^{158}Gd$	ave.	7937.39	0.06	7937.39	0.06	−0.0	1	99	53 ^{158}Gd			average	
$^{158}Gd(d,t)^{157}Gd-^{159}Tb()^{158}Tb$		195.0	1.5	195.6	0.6	0.4	1	18	18 ^{158}Tb	McM		84Bu14	
$^{157}Gd(\alpha,t)^{158}Tb-^{158}Gd()^{159}Tb$		−198.3	1.0	−195.6	0.6	2.7	o			McM		75Bu02	
		−196.6	1.0			1.0	1	41	40 ^{158}Tb	McM		84Bu14	
$^{158}Tb(p,d)^{157}Tb$		−4560.3	4.2	−4553.7	1.0	1.6	U			Pri		85Al02	*
$^{158}Dy(d,t)^{157}Dy$		−2804	10	−2796	5	0.8	–			Tal		68Be.A	
		−2804	10			0.8	–			Kop		70Gr46	
	ave.	−2804	7			1.1	1	52	47 ^{157}Dy			average	
$^{158}Pm(\beta^-)^{158}Sm$		6120	100	6145	5	0.3	o			Kur		02Sh.A	
		6085	80			0.8	o			Kur		07Ha57	
		6060	80			1.1	U			Kur		10Ha38	
$^{158}Sm(\beta^-)^{158}Eu$		1999	15	2019	5	1.3	1	12	10 ^{158}Sm	Ida		93Gr17	
$^{158}Eu(\beta^-)^{158}Gd$		3550	120	3420.5	2.2	−1.1	U					65Sc19	*
		3440	100			−0.2	U					66Da06	*
$^{158}Tb(\varepsilon)^{158}Gd$		1237.542	0.018	1219.3	1.0	******	F					83Ra25	*
		1220	13			−0.1	U					87Br33	
		1222.1	3.			−0.9	U					85Vo13	*
$^{158}Tb(\beta^-)^{158}Dy$		952	10	935.9	2.4	−1.6	U					68Sc04	*
		933	6			0.5	1	17	14 ^{158}Dy			85Vo03	*
$^{158}Ho(\beta^+)^{158}Dy$		4350	100	4220	27	−1.3	U					61Bo24	*
		4230	30			−0.3	2					68Ab14	*
$^{158}Er(\beta^+)^{158}Ho$		1710	40	880	40	−20.7	F					82Vy06	*
$^{158}Tm(\beta^+)^{158}Er$		6530	100	6600	30	0.7	–			IRS		93Al03	
		6624	60			−0.4	–			Dbn		94Po26	*
	ave.	6600	50			0.0	1	37	19 ^{158}Er			average	
$^{158}Lu(\varepsilon)^{158}Yb$		8960	200	8802	16	−0.8	U					95Ga.A	
$*^{158}Pm-u$	Represents frequency ratio $^{158}Pm^{++}/(C_{12}H_4)^+$ =0.96280782(15)											WgM124	**
$*^{158}Pm-u$	Represents frequency ratio $^{158}Pm^{++}/(C_6H_6)^+$ =1.0118048112(69)											HWJ208	**
$*^{158}Sm-u$	Represents frequency ratio $^{158}Sm^{++}/(C_{12}H_4)^+$ =0.962848141(58)											WgM124	**
$*^{158}Eu-u$	Represents frequency ratio $^{158}Eu^{++}/(C_{12}H_4)^+$ =0.96286130(15)											WgM124	**
$*^{158}Eu-u$	Represents frequency ratio $^{158}Eu^{++}/(C_6H_6)^+$ =1.011748662(14)											HWJ208	**
$*^{158}Ho-u$	$M-A$=–66148(29) keV for mixture gs+m+n at 67.2 and 180#70 keV											Nub269	**
$*^{158}W(\alpha)^{154}Hf$	Original value E=6450(30) (Q =6617.8) recalibrated to E=6433(30) keV											89Ho12	**
$*^{158}Tb(p,d)^{157}Tb$	$Q-Q(^{158}Gd(p,d))$=1152.5(4.2) keV											AHW	**
$*^{158}Eu(\beta^-)^{158}Gd$	E_{β^-}=2520(120) 2430(100) respectively, to 2^- level at 1023.6974 keV											Ens043	**
*	and 3^- level at 1041.6376 keV, and other E_{β^-}											Ens043	**
$*^{158}Tb(\varepsilon)^{158}Gd$	pK=0.00009(2) to 2^+ level at 1187.143, recalculated Q											Ens043	**
*	F : pK<0.00002											87Br33	**
$*^{158}Tb(\varepsilon)^{158}Gd$	pL=0.689(0.01) to 2^+ level at 1187.143 keV, recalculated Q											Ens043	**
$*^{158}Tb(\beta^-)^{158}Dy$	E_{β^-}=853(10) 834(6) respectively, to 2^+ level at 98.9180 keV											Ens043	**
$*^{158}Ho(\beta^+)^{158}Dy$	E_{β^+}=780(80) to 2436−2605 levels; originally assigned to $^{158}Er(\beta^+)$;											Ens043	**
*	reinterpreted by evaluator											AHW	**
$*^{158}Ho(\beta^+)^{158}Dy$	E_{β^+}=2890(20), 700(60) to 317.139−637.712 and 2436.52−2605.96 levels, and											Ens043	**
*	E_{β^+}=1300(30), 1850(25) keV from $^{158}Ho^{m}$ at 67.2 to 1920.43−1940.75											Nub269	**
*	and 1441.75 levels; E_{β^+}=700(60) was originally assigned to $^{158}Er(\beta^+)$;											68Ab14	**
*	reinterpreted by evaluator											AHW	**
$*^{158}Er(\beta^+)^{158}Ho$	p^+ =0.3(0.1) from annih. γ coinc. to 146.90 level											96Go06	**
*	F : Q <1550 from upper limit on p^+											75Bu.A	**
$*^{158}Tm(\beta^+)^{158}Er$	E_{β^+}=5410(60) to 2^+ level at 192.15 keV											Ens07a	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{159}Nd−^{133}Cs$_{1.195}$	59604	32				2			CP2	1.0	18Or02	
^{159}Pm−u	−60715	18	−60713	11	0.1	1	36	36 ^{159}Pm	CP1	1.0	12Va02	*
^{159}Pm−^{80}Kr$_{1.988}$	105529	19	105527	11	−0.1	1	32	32 ^{159}Pm	CP1	1.0	12Va02	
^{159}Pm−^{86}Kr$_{1.849}$	104567	19	104568	11	0.0	1	32	32 ^{159}Pm	CP1	1.0	12Va02	
^{159}Pm−u	−60638.1	66.6	−60713	11	−1.1	U			GR1	1.0	25Sp03	
^{159}Sm−u	−66784	11	−66783	6	0.1	1	34	34 ^{159}Sm	CP1	1.0	12Va02	*
^{159}Sm−^{80}Kr$_{1.988}$	99459	11	99457	6	−0.2	1	34	33 ^{159}Sm	CP1	1.0	12Va02	
^{159}Sm−^{86}Kr$_{1.849}$	98498	11	98498	6	0.0	1	34	34 ^{159}Sm	CP1	1.0	12Va02	
^{159}Sm−u	−66800.2	24.7	−66783	6	0.7	U			GR1	1.0	25Sp03	
^{159}Eu−u	−70899	10	−70901	5	−0.2	1	21	21 ^{159}Eu	CP1	1.0	12Va02	*
	−70897.9	26.8			−0.1	U			GR1	1.0	25Sp03	
^{159}Eu−^{80}Kr$_{1.988}$	95344	10	95339	5	−0.5	1	23	21 ^{159}Eu	CP1	1.0	12Va02	
^{159}Eu−^{86}Kr$_{1.849}$	94382	10	94380	5	−0.2	1	21	21 ^{159}Eu	CP1	1.0	12Va02	
C_9 ^{13}C H_6 O_2−^{159}Tb	114840	50	114781.4	1.0	−0.3	U			R04	4.0	64De15	
C_{10} H_7 O_2−^{159}Tb	119238	25	119251.6	1.0	0.1	U			R04	4.0	64De15	
^{159}Dy−u	−74285	30	−74255.6	1.0	1.0	U			GS2	1.0	05Li24	
^{159}Ho−u	−72365	71	−72283	3	1.2	U			GS2	1.0	05Li24	*
^{159}Er−u	−69290	30	−69311	4	−0.7	U			GS2	1.0	05Li24	
^{159}Tm−u	−65025	30				2			GS2	1.0	05Li24	
^{159}Yb−^{142}Sm$_{1.120}$	35035	24	35026	19	−0.4	2			MA7	1.0	01Bo59	
^{159}Yb−u	−59960	30	−59941	19	0.6	R			GS2	1.0	05Li24	
^{159}Lu−u	−53420	61	−53360	40	0.9	2			GS2	1.0	05Li24	*
^{159}Hf−u	−46044	32	−45984	12	1.9	R			GS2	1.0	05Li24	
^{159}Tb $^{35}Cl_2$−^{155}Gd $^{37}Cl_2$	8625.64	1.03	8624.9	0.8	−0.3	1	10	7 ^{159}Tb	H41	2.5	85Dy04	
^{159}Tb ^{35}Cl−^{157}Gd ^{37}Cl	4333.3	1.2	4336.7	0.8	1.1	U			H25	2.5	72Ba08	
	4337.01	0.61			−0.2	1	26	20 ^{159}Tb	H41	2.5	85Dy04	
^{159}Dy−^{159}Tb	391.559	0.023	391.559	0.023	−0.0	1	100	86 ^{159}Dy	JY2	1.0	21Ge11	
^{159}Lu(α)^{155}Tm	4534.3	10.	4490	40	−0.8	R			IRa		80Al14	
	4531.3	10.			−0.8	R					92Ha10	
^{159}Hf(α)^{155}Yb	5221.2	10.	5225.1	2.7	0.4	U					73Ea01	Z
	5226.2	5.			−0.2	2			GSa		79Ho10	Z
	5223.0	5.			0.4	2			ORa		83To01	Z
	5219.6	6.			0.9	2					92Ha10	
	5229.8	5.			−0.9	2			Daa		96Pa01	
^{159}Ta(α)^{155}Lu^m	5658.6	5.	5660	7	0.2	o			Daa		96Pa01	*
	5661.7	5.			−0.4	o			Ara		97Da07	*
^{159}Ta^m(α)^{155}Lu	5745.8	6.	5745	3	−0.2	–			GSa		79Ho10	
	5743.8	5.			0.2	–			Daa		96Pa01	
	5744.8	5.			−0.0	–			Ara		97Da07	
ave.	5745	3			0.0	1	99	84 ^{155}Lu			average	
^{159}W(α)^{155}Hf	6444.5	6.	6451	4	1.0	3			GSa		81Ho10	*
	6440.3	5.1			2.1	o			Daa		92Pa05	
	6454.7	5.			−0.7	3			Daa		96Pa01	
	6452.6	8.2			−0.2	3			Jya		19Hi06	
^{159}Re^m(α)^{155}Ta	6951.2	26.7	6968	23	0.6	R			Daa		07Pa27	
^{157}Gd(t,p)^{159}Gd	5398.9	2.3	5398.80	0.11	−0.0	U			McM		89Lo07	
^{158}Gd(n,γ)^{159}Gd	5942	1	5943.21	0.08	1.2	U					71Gr42	
	5943.07	0.15			0.9	–			ILn		87Sp.A	Z
	5943.1	0.2			0.5	–			Dbn		03Gr13	
	5943.32	0.12			−1.0	–			BNn		03Gr27	
^{158}Gd(d,p)^{159}Gd	3717	10	3718.64	0.08	0.2	U			Kop		67Tj01	
^{158}Gd(n,γ)^{159}Gd ave.	5943.20	0.08	5943.21	0.08	0.1	1	100	94 ^{159}Gd			average	
^{158}Gd(α,t)^{159}Tb	−13686.6	10.	−13682.5	0.7	0.4	U			McM		75Bu02	
^{158}Gd(α,t)^{159}Tb−^{164}Dy()^{165}Ho	−85.7	2.2	−88.0	1.0	−1.0	1	21	9 ^{159}Tb	McM		84Bu14	
^{159}Tb(γ,n)^{158}Tb	−8141	39	−8133.0	0.6	0.2	U			Phi		60Ge01	
^{159}Tb(d,t)^{158}Tb	−1870	15	−1875.8	0.6	−0.4	U			Tal		70Jo22	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{159}Tb(d,t)^{158}Tb−^{164}Dy()^{163}Dy	−474.3	1.0	−474.9	0.6	−0.6	1	41	40 ^{158}Tb	McM		84Bu14	
^{158}Dy(d,p)^{159}Dy	4608	10	4607.8	2.4	−0.0	U			Tal		68Be.A	
	4600	10			0.8	U			Kop		70Gr46	
^{159}Re^{m}(p)^{158}W	1816.4	20.	1809	17	−0.4	4			Dap		06Jo10	
^{159}Pm(β^-)^{159}Sm	5460	140	5653	12	1.4	o			Kur		07Ha57	
	5430	140			1.6	U			Kur		10Ha38	
^{159}Sm(β^-)^{159}Eu	3840	100	3836	7	−0.0	o			Kur		02Sh.A	
	3805	65			0.5	o			Kur		07Ha57	
	3800	65			0.6	U			Kur		10Ha38	
^{159}Eu(β^-)^{159}Gd	2600	50	2519	4	−1.6	U					65Iw01	*
^{159}Gd(β^-)^{159}Tb	969.0	1.5	970.5	0.7	1.0	1	24	18 ^{159}Tb			77Bo.A	
^{159}Dy(ε)^{159}Tb	365.9	1.3	364.735	0.021	−0.9	U					68My.A	*
^{159}Ho(β^+)^{159}Dy	1837.6	6.	1837.6	2.7	0.0	2					79Ad08	*
	1837.6	3.			0.0	2					82Vy02	*
^{159}Er(β^+)^{159}Ho	2768.5	2.0				3					84Ka.A	*
	2810	100	2768.5	2.0	−0.4	U			IRS		93Al03	
^{159}Tm(β^+)^{159}Er	3400	300	3992	28	2.0	U					75St07	
	3850	100			1.4	U			IRS		93Al03	
	3670	100			3.2	B			Dbn		94Po26	
^{159}Yb(β^+)^{159}Tm	5050	200	4740	30	−1.6	U			IRS		93Al03	
	4554	150			1.2	U			Dbn		94Po26	*
^{159}Lu(β^+)^{159}Yb	5850	150	6130	40	1.8	U			IRS		93Al03	
	5803	150			2.2	U			Dbn		94Po26	
^{159}Ta^{m}(IT)^{159}Ta	63.7	5.2				2			Ara		97Da07	
*^{159}Pm−u	Represents frequency ratio $^{159}Pm^{++}/(C_{12}H_4)^+$ =0.95673359(11)										WgM124	**
*^{159}Sm−u	Represents frequency ratio $^{159}Sm^{++}/(C_{12}H_4)^+$ =0.956770122(65)										WgM124	**
*^{159}Eu−u	Represents frequency ratio $^{159}Eu^{++}/(C_{12}H_4)^+$ =0.956794898(63)										WgM124	**
*^{159}Ho−u	$M-A$=–67304(28) keV for mixture gs+m at 205.91 keV										Nub269	**
*^{159}Lu−u	$M-A$=–49710(28) keV for mixture gs+m at 100#80 keV										Nub269	**
*^{159}Ta(α)^{155}Lu^{m}	Replaced by ^{155}Lu^{m}(IT)										AHW	**
*^{159}W(α)^{155}Hf	Original value E_α=6299(6) recalibrated to E_α=6282(6) keV										89Ho12	**
*^{159}Eu(β^-)^{159}Gd	E_{β^-}=2350(50) to 7/2$^-$ level at 227.412 level, and other E_{β^-}										Ens121	**
*^{159}Dy(ε)^{159}Tb	From intensity of feeding 5/2$^-$ level at 363.5449 keV										Ens121	**
*^{159}Ho(β^+)^{159}Dy	E_{β^+}=506(6) 506(3) respectively, to 5/2$^-$ level at 309.593 keV										Ens121	**
*^{159}Er(β^+)^{159}Ho	E_{β^+}=1122(3) to 13/2$^+$ level at 624.5 keV, and other E_{β^+}										Ens121	**
*^{159}Yb(β^+)^{159}Tm	E_{β^+}=3366(150) to 7/2$^-$ level at 166.17 keV										Ens121	**
^{160}Nd−^{84}Kr$_{1.905}$	118436	50				2			CP2	1.0	18Or02	
^{160}Pm−^{136}Xe$_{1.176}$	52377	18	52331.1	2.2	−2.6	B			JY1	1.0	18Vi02	*
^{160}Pm−^{84}Kr$_{1.905}$	111812.1	2.2				2			CP2	1.0	20Or03	
^{160}Pm^{m}−^{84}Kr$_{1.905}$	112017	12				2			CP2	1.0	20Or03	
^{160}Sm−u	−64666	11	−64663.0	2.1	0.3	U			CP1	1.0	12Va02	*
^{160}Sm−^{80}Kr$_{2.000}$	102581	11	102580.4	2.6	−0.1	U			CP1	1.0	12Va02	
^{160}Sm−^{86}Kr$_{1.860}$	101599	11	101601.3	2.1	0.2	U			CP1	1.0	12Va02	
^{160}Sm−^{82}Kr$_{1.951}$	104135.3	2.1				2			CP2	1.0	18Or.A	
^{160}Sm−u	−64689.6	29.0	−64663.0	2.1	0.9	U			GR1	1.0	25Sp03	
^{160}Eu−u	−68150	17	−68163.0	1.0	−0.8	U			CP1	1.0	12Va02	*
	−68179.7	41.9			0.4	U			GR1	1.0	25Sp03	*
^{160}Eu−^{80}Kr$_{2.000}$	99096	18	99080.4	1.8	−0.9	U			CP1	1.0	12Va02	
^{160}Eu−^{86}Kr$_{1.860}$	98115	18	98101.2	1.0	−0.8	U			CP1	1.0	12Va02	
^{160}Eu−^{84}Kr$_{1.905}$	100433.81	0.97				2			CP2	1.0	18Ha19	
^{160}Eu^{m}−^{84}Kr$_{1.905}$	100533.70	0.86				2			CP2	1.0	18Ha19	
C_{12} H_{16}−^{160}Gd	198150	50	198140.3	1.1	−0.0	U			R04	4.0	64De15	
C_{12} H_{16}−^{160}Dy	200050	70	199997.3	0.7	−0.2	U			R04	4.0	64De15	
^{160}Er−u	−70916	30	−70923	26	−0.2	–			GS2	1.0	05Li24	
ave.	−70914	27			−0.3	1	95	95 ^{160}Er			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{160}Tm−u		−64770	120	−64740	40	0.3	U				GS1	1.0	00Ra23	*
		−64755	37			0.4	1	90	90	^{160}Tm	GS2	1.0	05Li24	*
^{160}Yb−^{142}Sm$_{1.127}$		33120	20	33120	6	−0.0	U				MA7	1.0	01Bo59	
^{160}Yb−u		−62440	120	−62441	6	−0.0	U				GS1	1.0	00Ra23	
		−62438	30			−0.1	U				GS2	1.0	05Li24	
^{160}Yb−^{133}Cs$_{1.203}$		51300.5	5.9				2				MA8	1.0	19Hu15	
^{160}Lu−u		−53994	43				2				GS2	1.0	05Li24	*
^{160}Hf−u		−49334	30	−49315	10	0.6	U				GS2	1.0	05Li24	
^{160}Gd ^{35}Cl$_2$−^{156}Gd ^{37}Cl$_2$		10831.70	1.27	10831.5	0.8	−0.1	U				H41	2.5	85Dy04	
^{160}Gd ^{35}Cl−^{158}Gd ^{37}Cl		5890	5	5900.3	0.8	0.5	U				H12	4.0	64Ba15	
		5899	3			0.2	U				H21	2.5	70Ma05	
		5900.0	0.5			0.2	–				M21	2.5	75Ka25	
		5899.88	0.96			0.2	–				H41	2.5	85Dy04	
	ave.	5900.0	1.1			0.3	1	49	42	^{160}Gd			average	
^{160}Dy ^{35}Cl−^{158}Dy ^{37}Cl		3731.8	2.3	3739.0	2.4	1.3	1	18	17	^{158}Dy	H25	2.5	72Ba08	
^{160}Gd−^{160}Dy		1854.5	0.8	1857.0	1.1	1.2	1	31	24	^{160}Gd	H25	2.5	72Ba08	
^{160}Gd O−C$_{15}$		−78020.1	5.8	−78025.2	1.1	−0.6	o				TG1	1.5	09Ke.A	
		−78019.9	3.6			−1.0	U				TG1	1.5	11Ke03	
^{160}Hf(α)^{156}Yb		4892.2	10.	4902.0	2.6	1.0	–						73Ea01	Z
		4905.0	5.			−0.6	–				GSa		79Ho10	Z
		4904.0	5.			−0.4	–				ORa		83To01	Z
		4901.8	6.			0.0	–						92Ha10	
		4902.8	10.			−0.1	–						95Hi12	
		4900.8	6.			0.2	–				Daa		96Pa01	
	ave.	4902.4	2.6			−0.2	1	99	80	^{160}Hf			average	
^{160}Ta(α)^{156}Lu^m		5449.5	5.	5451	5	0.3	3				Daa		96Pa01	
		5456.6	10.			−0.6	3				Jya		09Ha42	
^{160}Ta^m(α)^{156}Lu		5550.9	5.	5548.5	3.0	−0.5	3				GSa		79Ho10	Z
		5538.7	6.			1.6	3						92Ha10	
		5552.1	5.			−0.7	3				Daa		96Pa01	
		5551.0	10.			−0.3	3				Jya		09Ha42	
^{160}W(α)^{156}Hf		6072.1	10.	6066	5	−0.6	–				GSa		79Ho10	
		6063.9	5.			0.3	–				Daa		96Pa01	
	ave.	6066	5			−0.0	1	100	100	^{160}W			average	
^{160}Re(α)^{156}Ta		6704.9	16.	6698	4	−0.4	o				Daa		92Pa05	
		6711.1	16.			−0.8	o				Daa		96Pa01	
		6697.7	4.				4				Daa		11Da12	
^{160}Os(α)^{156}W		7274.2	15.4	7271	13	−0.2	3				Jya		23Br10	
		7261.9	26.7			0.3	3				Lza		24Ya04	
^{160}Os^m(α)^{156}W		9118.4	10.3				3				Jya		23Br10	
^{158}Gd(t,p)^{160}Gd		4912.0	2.2	4912.8	0.7	0.4	U				McM		89Lo07	
^{160}Gd(p,t)^{158}Gd		−4919	5	−4912.8	0.7	1.2	U				Min		73Oo01	
^{160}Dy(p,t)^{158}Dy		−6924	5	−6926.0	2.2	−0.4	–				Min		73Oo01	
		−6925.1	3.4			−0.3	–				McM		88Bu08	*
	ave.	−6924.8	2.8			−0.4	1	64	63	^{158}Dy			average	
^{160}Gd(t,α)^{159}Eu−^{158}Gd()^{157}Eu		−666	5	−668	4	−0.5	1	69	36	^{159}Eu	LAl		79Bu05	
^{160}Gd(d,t)^{159}Gd		−1200	10	−1194.2	0.7	0.6	U				Kop		67Tj01	
^{159}Tb(n,γ)^{160}Tb		6375.45	0.3	6375.21	0.13	−0.8	–						74Ke01	Z
		6375.13	0.15			0.5	–				Bdn		06Fi.A	
^{159}Tb(d,p)^{160}Tb		4165	20	4150.64	0.13	−0.7	U				MIT		64Sp12	
		4153	5			−0.5	U				Tal		67St14	
^{159}Tb(n,γ)^{160}Tb	ave.	6375.19	0.13	6375.21	0.13	0.1	1	99	88	^{160}Tb			average	
^{160}Dy(d,t)^{159}Dy		−2339	10	−2318.2	1.0	2.1	U				Tal		68Be.A	
		−2323	10			0.5	U				Kop		70Gr46	
^{160}Re(p)^{159}W		1269.1	6.	1267	7	−0.4	o				Dap		92Pa05	
		1271	9			−0.5	o				Dap		96Pa01	*
		1272.2	6.			−0.9	R				Dap		11Da12	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{160}Eu(β^-)^{160}Gd	3900	300	4449.5	1.4	1.8	U					73Da05	
	4200	200			1.2	U					73Mo18	
	4705	60			−4.3	B			Kur		07Ha57	
	4695	60			−4.1	C			Kur		10Ha38	
	4480	35			−0.9	U			Kur		14Ha38	
^{160}Tb(β^-)^{160}Dy	1838	10	1835.5	1.0	−0.3	U					57Na03	*
	1827	10			0.8	U					59Gr93	*
	1825	10			1.0	U					63Wu01	*
^{160}Ho(β^+)^{160}Dy	3290	15				2					66Av03	*
^{160}Er(ε)^{160}Ho	420	150	318	29	−0.7	U					82Vy06	*
^{160}Tm(β^+)^{160}Er	5600	300	5760	40	0.5	U					75St12	*
	5890	100			−1.3	1	15	10 ^{160}Tm	IRS		93Al03	
^{160}Lu(β^+)^{160}Yb	7210	240	7870	40	2.7	U					83Ge08	
	7340	100			5.3	C			IRS		83Vi.A	
	7300	100			5.7	B			IRS		93Al03	
*^{160}Pm−^{136}Xe$_{1.176}$	Could be a mixture of ground state and isomer										HWJ208	**
*^{160}Sm−u	Represents frequency ratio $^{160}Sm^{++}/(C_{12}H_4)^+$ =0.950775181(67)										WgM124	**
*^{160}Eu−u	Represents frequency ratio $^{160}Eu^{++}/(C_{12}H_4)^+$ =0.95079589(10)										WgM124	**
*^{160}Eu−u	Corrected for unresolved isomeric state using exc. energy in 23Ga02										25Sp03	**
*^{160}Tm−u	$M-A$=–60300(110) keV for mixture gs+m at 71(12) keV										Nub269	**
*^{160}Tm−u	$M-A$=–60283(28) keV for mixture gs+m at 71(12) keV										Nub269	**
*^{160}Lu−u	$M-A$=–50270(28) keV for mixture gs+m at 50#50 keV										Nub269	**
*^{160}Dy(p,t)^{158}Dy	$Q-Q(^{164}$Dy(p,t))=–1477.9(3.4), see ^{164}Dy(p,t)										AHW	**
*^{160}Re(p)^{159}W	Corrected : Ame2003 assumed E_p=1271(9) thus Q_p=1279.1(9.) keV										WgM105	**
*^{160}Tb(β^-)^{160}Dy	E_{β^-}=870(10) 858(10) 868(10) respectively, to 8^+ level at 966.85 keV, and other E_{β^-}										Ens059	**
*^{160}Ho(β^+)^{160}Dy	E_{β^+}=570(15) to 4^+ level at 1694.37 keV; and 1045(15) from										Ens059	**
*	^{160}Ho^m at 59.98 to 1^- level at 1285.602 and 3^- at 1286.711 keV										Nub269	**
*^{160}Er(ε)^{160}Ho	pK=0.795(0.2) to 1^+ level at 67.11 keV										Ens059	**
*^{160}Tm(β^+)^{160}Er	E_{β^+}=3700(300) to 854.4−1007.95 levels, reassigned by evaluator										Ens059	**
^{161}Pm−^{136}Xe$_{1.184}$	56065	64	56088	10	0.4	U			JY1	1.0	20Vi04	
^{161}Pm−^{84}Kr$_{1.917}$	115888.7	9.9				2			CP2	1.0	22Or02	
^{161}Sm−u	−60841	13	−60840	7	0.1	1	32	32 ^{161}Sm	CP1	1.0	12Va02	*
^{161}Sm−^{80}Kr$_{2.013}$	107493	12	107491	7	−0.2	1	38	37 ^{161}Sm	CP1	1.0	12Va02	
^{161}Sm−^{86}Kr$_{1.872}$	106496	13	106497	7	0.1	1	32	32 ^{161}Sm	CP1	1.0	12Va02	
^{161}Sm−u	−60875.3	51.5	−60840	7	0.7	U			GR1	1.0	25Sp03	
^{161}Eu−u	−66336	19	−66336	11	0.0	1	35	35 ^{161}Eu	CP1	1.0	12Va02	*
	−66302.1	33.3			−1.0	U			GR1	1.0	25Sp03	
^{161}Eu−^{80}Kr$_{2.013}$	101996	19	101995	11	−0.1	1	35	34 ^{161}Eu	CP1	1.0	12Va02	
^{161}Eu−^{86}Kr$_{1.872}$	101000	20	101001	11	0.1	1	31	31 ^{161}Eu	CP1	1.0	12Va02	
C_{13} H_5−^{161}Dy	112246	25	112186.1	0.7	−0.6	U			R04	4.0	64De15	
^{161}Tm−u	−66451	30				2			GS2	1.0	05Li24	*
^{161}Yb−^{142}Sm$_{1.134}$	34071	19	34065	16	−0.3	2			MA7	1.0	01Bo59	
^{161}Yb−u	−62120	110	−62089	16	0.3	U			GS1	1.0	00Ra23	
	−62107	30			0.6	R			GS2	1.0	05Li24	
^{161}Lu−u	−56428	30				2			GS2	1.0	05Li24	
^{161}Hf−u	−49733	30	−49723	25	0.3	1	70	70 ^{161}Hf	GS2	1.0	05Li24	
^{161}Dy ^{35}Cl−^{159}Tb ^{37}Cl	4535.0	1.0	4536.3	1.0	0.5	1	17	13 ^{159}Tb	H25	2.5	72Ba08	
^{161}Hf(α)^{157}Yb	4717.0	10.	4685	25	−0.6	–					73Ea01	Z
	4725.2	10.			−0.8	–					82Sc15	Z
	4724.2	5.			−0.8	–			ORa		83To01	Z
	4716.4	7.			−0.6	–					92Ha10	
	4721.5	10.			−0.7	–					95Hi12	
ave.	4721	23			−0.7	1	20	18 ^{161}Hf			average	
^{161}Ta^m(α)^{157}Lu^m	5278.9	5.	5276	6	−0.7	F			GSa		79Ho10	*
	5280.4	5.			−1.0	F					92Ha10	
	5271.2	7.			0.6	F			Daa		96Pa01	*
	5282.5	7.			−1.0	F			Jya		05Sc22	*
	5273.2	6.			0.4	1	93	60 ^{161}Ta^m	Jya		12Th13	
^{161}W(α)^{157}Hf	5923.4	5.	5923	4	−0.1	4			GSa		79Ho10	Z
	5922.4	5.			0.1	4			Daa		96Pa01	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{161}Re$^{m}(\alpha)^{157}$Ta^{m}		6439.3	10.	6430	4	−0.9	2				GSa		79Ho10	
		6425.0	6.			0.8	2				Daa		96Pa01	
		6432.1	7.			−0.3	2				Ara		97Ir01	
^{161}Os$(\alpha)^{157}$W		7065.9	12.	7069	11	0.2	3						10Bi03	
		7077.2	20.5			−0.4	3				Jya		19Hi06	
^{161}Os$(\alpha)^{157}$W^{p}		6748.0	30.				4						10Bi03	
^{161}Dy(p,t)^{159}Dy		−6546	5	−6548.0	1.0	−0.4	–				Min		73Oo01	
		−6547.9	2.5			−0.0	–				McM		88Bu08	*
	ave.	−6547.5	2.2			−0.2	1	19	14	^{159}Dy			average	
^{160}Gd(n,γ)^{161}Gd		5635.4	1.0				2						71Gr42	
^{160}Gd(d,p)^{161}Gd		3411	10	3410.8	1.0	−0.0	U				Kop		67Tj01	
^{160}Gd$(\alpha$,t)^{161}Tb−^{158}Gd()^{159}Tb		678.0	1.0	677.2	0.7	−0.8	1	55	33	^{160}Gd	McM		75Bu02	
^{160}Tb(n,γ)^{161}Tb		7696.3	0.6	7696.6	0.5	0.5	1	84	80	^{161}Tb			75He.C	
^{161}Dy(t,α)^{160}Tb		12306	3	12306.3	1.0	0.1	1	10	8	^{160}Tb	McM		07Bu29	
^{160}Dy(n,γ)^{161}Dy		6451.5	2.	6454.39	0.08	1.4	U						77Be03	
		6454.40	0.09			−0.2	–				ILn		86Sc16	Z
		6454.34	0.14			0.3	–				Bdn		06Fi.A	
^{160}Dy(d,p)^{161}Dy		4231	10	4229.82	0.08	−0.1	U				Tal		68Be.A	
		4237	10			−0.7	U				Kop		70Gr46	
^{161}Dy(d,t)^{160}Dy		−205	10	−197.16	0.08	0.8	U				Kop		70Gr46	
^{160}Dy(n,γ)^{161}Dy	ave.	6454.38	0.08	6454.39	0.08	0.0	1	100	93	^{160}Dy			average	
^{160}Dy(^{3}He,d)^{161}Ho−^{164}Dy()^{165}Ho		−1406.5	2.0	−1406.5	2.0	0.0	1	100	100	^{161}Ho	McM		75Bu02	
^{161}Re(p)^{160}W		1199.5	6.	1197	5	−0.4	1	79	79	^{161}Re	Ara		97Ir01	
^{161}Re^{m}(p)^{160}W		1323.3	7.	1321	5	−0.3	o				Ara		97Ir01	*
^{161}Sm$(\beta^{-})^{161}$Eu		5065	130	5120	12	0.4	o				Kur		07Ha57	
		5050	130			0.5	U				Kur		10Ha38	
^{161}Eu$(\beta^{-})^{161}$Gd		3705	60	3716	10	0.2	o				Kur		07Ha57	
		3705	60			0.2	U				Kur		10Ha38	
		3722	35			−0.2	U				Kur		14Ha38	
^{161}Gd$(\beta^{-})^{161}$Tb		1977	30	1955.5	1.4	−0.7	U						66Zy02	*
^{161}Tb$(\beta^{-})^{161}$Dy		584	6	593.3	1.1	1.5	U						63Ko08	
		590	10			0.3	U						64Fu11	
^{161}Er$(\beta^{+})^{161}$Ho		2050	40	1995	9	−1.4	U						65Gr35	*
		1980	18			0.8	R						84Ka.A	*
^{161}Tm$(\beta^{+})^{161}$Er		3100	200	3303	29	1.0	U						75Ad08	*
		3180	100			1.2	U				IRS		93Al03	
^{161}Yb$(\beta^{+})^{161}$Tm		3850	250	4060	30	0.9	U						81Ad02	
		3585	200			2.4	U				Dbn		94Po26	
^{161}Lu$(\beta^{+})^{161}$Yb		5300	100	5270	30	−0.3	o				IRS		83Vi.A	
		5300	100			−0.3	U				IRS		93Al03	
		5255	150			0.1	U				Dbn		94Po26	*
^{161}Re^{m}(IT)^{161}Re		123.8	1.3	123.7	1.3	−0.1	1	99	78	^{161}Re^{m}			97Ir01	
$*^{161}$Sm−u	Represents frequency ratio $^{161}Sm^{++}/(C_{12}H_4)^{+}$ =0.944844874(74)												WgM124	**
$*^{161}$Eu−u	Represents frequency ratio $^{161}Eu^{++}/(C_{12}H_4)^{+}$ =0.94487714(11)												WgM124	**
$*^{161}$Tm−u	$M-A$=–61895(28) keV for mixture gs+m at 7.51 keV												Nub269	**
$*^{161}$Ta$^{m}(\alpha)^{157}$Lu^{m}	F : above 4 items are not unambiguously assigned												WgM151	**
$*^{161}$Dy(p,t)^{159}Dy	$Q-Q(^{164}$Dy(p,t))=–1100.7(2.5) keV												AHW	**
$*^{161}$Re^{m}(p)^{160}W	Replaced by author's result for ^{161}Re^{m}(IT)^{161}Re												AHW	**
$*^{161}$Gd$(\beta^{-})^{161}$Tb	E_{β^-}=1560(30) mainly to 7/2$^-$ level at 417.228 keV												Ens11b	**
$*^{161}$Er$(\beta^{+})^{161}$Ho	E_{β^+}=820(40) 748(18) respectively, to 1/2$^+$ level at 211.15 keV												Ens11b	**
$*^{161}$Tm$(\beta^{+})^{161}$Er	E_{β^+}=1800(100) to several levels around 7/2$^-$ one at 266.44 keV												Ens11b	**
$*^{161}$Lu$(\beta^{+})^{161}$Yb	E_{β^+}=3866(150) to 367.28 level												Ens11b	**
^{162}Sm−^{136}Xe$_{1.191}$		52127.2	5.3	52129	4	0.4	1	51	51	^{162}Sm	JY1	1.0	18Vi02	
^{162}Sm−^{84}Kr$_{1.929}$		112344.7	5.4	112343	4	−0.4	1	49	49	^{162}Sm	CP2	1.0	18Or02	
^{162}Eu−^{136}Xe$_{1.191}$		47449	50	47465.5	1.5	0.3	U				JY1	1.0	18Vi02	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{162}Eu−^{84}Kr$_{1.929}$		107678.2	1.6	107678.9	1.5	0.4	1	82	82	^{162}Eu	CP2	1.0	18Ha19	
^{162}Eu−^{133}Cs$_{1.218}$		52124.0	6.0	52117.5	1.5	−1.1	U				JY1	1.0	20Vi04	
^{162}Eu−u		−63099.7	104.1	−63042.0	1.5	0.6	U				GR1	1.0	25Sp03	*
^{162}Eu^m−^{84}Kr$_{1.929}$		107850.1	2.0	107848.7	1.4	−0.7	1	48	48	^{162}Eu^m	CP2	1.0	18Ha19	
^{162}Eu^m−^{133}Cs$_{1.218}$		52286.4	2.4	52287.4	1.4	0.4	–				JY2	1.0	20Vi04	
		52292.4	8.1			−0.6	–				JY1	1.0	20Vi04	
	ave.	52286.9	2.3			0.2	1	36	36	^{162}Eu^m			average	
^{162}Tb−u		−70724.6	2.2	−70419.5	2.1	138.7	B				CP2	1.0	20Or03	*
		−70418.1	2.7			−0.5	1	60	60	^{162}Tb	CP2	1.0	20Or03	*
C_{13} H_6−^{162}Dy		120115	19	120146.0	0.7	0.4	U				R04	4.0	64De15	
C_{12} H_4 N−^{162}Er		105590	70	105587.2	0.8	−0.0	U				R04	4.0	64De15	
C_{13} H_6−^{162}Er		118430	170	118163.2	0.8	−0.4	U				R04	4.0	64De15	
^{162}Tm−u		−65942	55	−65999	28	−1.0	R				GS2	1.0	05Li24	*
^{162}Yb−^{142}Sm$_{1.141}$		32524	19	32525	16	0.1	2				MA7	1.0	01Bo59	
^{162}Yb−u		−64210	110	−64222	16	−0.1	U				GS1	1.0	00Ra23	
		−64223	30			0.0	R				GS2	1.0	05Li24	
^{162}Lu−u		−56640	140	−56710	50	−0.5	o				GS1	1.0	00Ra23	*
		−56706	60			0.0	2				GS2	1.0	05Li24	*
^{162}Hf−u		−52756	30	−52786	10	−1.0	U				GS2	1.0	05Li24	
^{162}Er ^{35}Cl$_2$−^{158}Gd ^{37}Cl$_2$		10577.5	2.7	10577.1	1.0	−0.1	U				H25	2.5	72Ba08	
^{162}Dy ^{35}Cl−^{160}Dy ^{37}Cl		4555	6	4551.06	0.12	−0.2	U				H12	4.0	64Ba15	
		4552.1	1.1			−0.4	U				H25	2.5	72Ba08	
^{162}Er ^{35}Cl−^{160}Gd ^{37}Cl		4674.6	1.9	4676.9	1.2	0.5	U				H25	2.5	72Ba08	
^{162}Eu^m−^{162}Eu		167.4	3.1	169.9	1.8	0.8	1	33	18	^{162}Eu	JY2	1.0	20Vi04	
^{162}Er−^{162}Dy		1982.79	0.32	1982.8	0.3	0.0	1	100	100	^{162}Er	SH1	1.0	11El04	
^{161}Dy ^{37}Cl−^{162}Dy ^{35}Cl		−3080	70	−2815.21	0.09	0.9	U				R04	4.0	64De15	
^{162}Dy−^{161}Dy		150	70	−134.92	0.06	−1.0	U				R04	4.0	64De15	
		78	23			−2.3	U				R04	4.0	64De15	
		22	40			−1.0	U				R04	4.0	64De15	
^{162}Hf(α)^{158}Yb		4417.2	10.	4416	5	−0.1	–						82Sc15	
		4420.4	10.3			−0.4	–				ORa		83To01	
		4414.2	9.			0.2	–						92Ha10	
		4416.3	10.3			0.0	–						95Hi12	
	ave.	4417	5			−0.1	1	95	81	^{162}Hf			average	
^{162}Ta(α)^{158}Lu		5003.8	10.	5010	60	0.0	4						86Ru05	
		5007.9	5.			−0.0	4						92Ha10	
^{162}W(α)^{158}Hf		5669.9	10.	5678.2	2.4	0.8	U						73Ea01	Z
		5668.0	10.			1.0	U				ORa		75To05	Z
		5677.5	5.			0.2	–				GSa		81Ho10	Z
		5674.5	4.			0.9	–				Ora		82De11	Z
		5681.6	5.			−0.6	–				Daa		96Pa01	
		5681.5	5.1			−0.6	–				Jya		15Li24	
	ave.	5678.3	2.4			−0.0	1	100	88	^{162}W			average	
^{162}Re(α)^{158}Ta		6240.3	5.				8				Ara		97Da07	
^{162}Re^m(α)^{158}Ta^m		6274.2	6.	6274	3	−0.0	9				GSa		79Ho10	
		6278.3	6.			−0.7	9				Daa		96Pa01	
		6271.1	5.			0.6	9				Ara		97Da07	
		6256	16			1.1	U				Jya		16Ca15	*
^{162}Os(α)^{158}W		6778.8	30.	6767.8	2.9	−0.4	U				GSa		89Ho12	*
		6785.8	10.			−1.8	U				ORa		96Bi07	
		6767.4	3.			0.1	4				Ara		00Ma95	
		6781.7	13.			−1.1	U				Jya		04Jo12	
		6770.5	8.2			−0.3	4				Jya		19Hi06	
^{160}Gd(t,p)^{162}Gd		3999.5	3.8				2				McM		89Lo07	
^{160}Dy(t,p)^{162}Dy		6169.5	1.9	6169.58	0.10	0.0	U				McM		88Bu08	*
^{162}Dy(p,t)^{160}Dy		−6168	5	−6169.58	0.10	−0.3	U				Min		73Oo01	
		−6169.7	2.1			0.1	U				McM		88Bu08	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{162}Er(p,t)^{160}Er	−7944	55	−7931	24	0.2	R			Win		74De31	*
^{161}Dy(n,γ)^{162}Dy	8196.99	0.06	8196.99	0.06	0.0	1	100	82 ^{161}Dy	MMn		82Is05	Z
	8193	3			1.3	U			Bdn		06Fi.A	
^{161}Dy(d,p)^{162}Dy	5969	10	5972.43	0.06	0.3	U			Tal		67Ba34	
	5981	10			−0.9	U			Kop		70Gr46	
^{162}Dy(d,t)^{161}Dy	−1944	10	−1939.76	0.06	0.4	U			Kop		70Gr46	
	−1943	10			0.3	U			Tal		77Be03	
^{161}Dy(^{3}He,d)^{162}Ho−^{164}Dy()^{165}Ho	−945.3	3.0	−945	3	0.0	1	100	100 ^{162}Ho	McM		75Bu02	
^{162}Er(d,t)^{161}Er	−2952	10	−2947	9	0.5	2			Kop		69Tj01	
^{162}Eu(β^-)^{162}Gd	5575	60	5558	4	−0.3	o			Kur		07Ha57	
	5585	60			−0.4	o			Kur		10Ha38	
	5577	35			−0.5	U			Kur		14Ha38	
^{162}Gd(β^-)^{162}Tb	1442	100	1314	4	−1.3	U					70Ch02	*
^{162}Tb^m(IT)^{162}Tb	216.0	1.0				2					07Bu29	
^{162}Tb(β^-)^{162}Dy	2448	100	2586.1	2.0	1.4	U					66Fu08	*
	2523	50			1.3	U					66Sc24	*
	2528	80			0.7	U					77Ka08	*
^{162}Ho(β^+)^{162}Dy	2220	50	2141	3	−1.6	U					69Ak01	
^{162}Tm(β^+)^{162}Er	4840	50	4857	26	0.3	2					63Ab02	
	4705	70			2.2	2					74De47	*
	4900	100			−0.4	2			IRS		93Al03	
	4892	50			−0.7	2			Dbn		94Po26	*
^{162}Lu(β^+)^{162}Yb	6740	270	7000	50	1.0	U					83Ge08	
	6990	120			0.1	o			IRS		83Vi.A	
	6960	100			0.4	R			IRS		93Al03	
	7111	150			−0.7	R			Dbn		94Po26	*
*^{162}Eu−^{136}Xe$_{1.191}$	D_M=47536.0(3.8) μu for mixture gs+m at 158.2(1.7) keV; $M-A$=–58657.6(3.5) keV										Nub269	**
*^{162}Eu−u	orrected for unresolved isomeric state using exc. energy in 23Ga02										25Sp03	**
*^{162}Tb−u	Represents frequency ratio ^{162}Tb^{++}/(C$_6$H$_6$)$^+$ =1.037384012(14)										HWJ208	**
*^{162}Tb−u	Trends in Mass Surface TMS suggest ^{162}Tb 400 keV less bound										HWJ23a	**
*^{162}Tb−u	Represents frequency ratio ^{162}Tb^{++}/(C$_6$H$_6$)$^+$ =1.037385975(17)										HWJ208	**
*^{162}Tb−u	isomer re-assigned to the gs.										HWJ258	**
*^{162}Tm−u	$M-A$=–61359(28) keV for mixture gs+m at 130(40) keV										Nub269	**
*^{162}Lu−u	$M-A$=–52730(130) keV for mixture gs+m+n at 60#40 and 150#50 keV										Nub269	**
*^{162}Lu−u	$M-A$=–52751(28) keV for mixture gs+m+n at 60#40 and 150#50 keV										Nub269	**
*^{162}Re^m(α)^{158}Ta^m	E_α=6037(16) keV to 10$^+$ level 66 keV above the 9$^+$ ^{158}Ta^m										16Ca15	**
*^{162}Os(α)^{158}W	Original value E=6640(20) (Q =6808.4) recalibrated										88Ho.B	**
*^{160}Dy(t,p)^{162}Dy	$Q-Q$(^{162}Dy(t,p))=722.3(1.9) keV										AHW	**
*^{162}Dy(p,t)^{160}Dy	$Q-Q$(^{164}Dy(p,t))=–722.5(2.1) keV										AHW	**
*^{162}Er(p,t)^{160}Er	Not resolved peak. Original uncertainty 28 increased to 51 keV and										GAu	**
*	added systematic error 21 keV										GAu	**
*^{162}Gd(β^-)^{162}Tb	E_{β^-}=1000(100) to 1$^+$ level at 442.11 keV										Ens162	**
*^{162}Tb(β^-)^{162}Dy	E_{β^-}=1300(100) 1375(50) 1380(80) respectively, to 2$^-$ level at 1148.232 keV										Ens078	**
*^{162}Tm(β^+)^{162}Er	E_{β^+}=2110(70) to 2$^-$ level at 1572.84 keV										Ens078	**
*^{162}Tm(β^+)^{162}Er	E_{β^+}=3768(50) to 2$^+$ level at 102.04 keV										Ens078	**
*^{162}Lu(β^+)^{162}Yb	E_{β^+}=6006(150) to ground state and 2$^+$ level at 166.8, unknown intensity ratio										Ens078	**
^{163}Sm−^{84}Kr$_{1.940}$	117373.5	7.9				2			CP2	1.0	18Or02	
^{163}Sm−^{136}Xe$_{1.199}$	56980	39	56929	8	−1.3	U			JY1	1.0	20Vi04	
^{163}Eu−^{133}Cs$_{1.226}$	55179.4	4.0	55181.4	1.0	0.5	U			JY1	1.0	20Vi04	
^{163}Eu−^{82}Kr$_{1.988}$	111264.98	0.97				2			CP2	1.0	22Or02	
^{163}Eu−u	−60775.5	52.6	−60734.5	1.0	0.8	U			GR1	1.0	25Sp03	
^{163}Gd−u	−65824	16	−65903.4	0.9	−5.0	B			CP1	1.0	12Va02	*
	−65849.5	3.7			−14.6	F			JY1	1.0	18Vi02	*
^{163}Gd−^{80}Kr$_{2.038}$	104600	16	104517.7	1.7	−5.1	F			CP1	1.0	12Va02	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{163}Gd−^{86}Kr$_{1.895}$		103569	15	103489.5	0.9	−5.3	B				CP1	1.0	12Va02	
^{163}Gd−^{136}Xe$_{1.199}$		45353	11	45346.5	0.9	−0.6	U				JY1	1.0	20Vi04	
^{163}Gd−^{82}Kr$_{1.988}$		106095.6	1.2	106096.1	0.9	0.4	1	51	51	^{163}Gd	CP2	1.0	20Or02	
^{163}Gd^{m}−^{136}Xe$_{1.199}$		45526	14	45494.9	0.9	−2.2	U				JY1	1.0	20Vi04	
^{163}Gd^{m}−^{82}Kr$_{1.988}$		106245.0	1.2	106244.5	0.9	−0.4	1	51	51	^{163}Gd^{m}	CP2	1.0	20Or02	
C_{13} H_7−^{163}Dy		125906	36	126038.3	0.7	0.9	U				R04	4.0	64De15	
^{163}Tm−u		−67327	30	−67342	6	−0.5	U				GS2	1.0	05Li24	
^{163}Yb−^{142}Sm$_{1.148}$		33686	19	33685	16	−0.1	2				MA7	1.0	01Bo59	
^{163}Yb−u		−63663	30	−63656	16	0.2	R				GS2	1.0	05Li24	
^{163}Lu−u		−58730	110	−58820	30	−0.8	U				GS1	1.0	00Ra23	
		−58821	30				2				GS2	1.0	05Li24	
^{163}Hf−u		−52911	30	−52916	28	−0.2	1	84	84	^{163}Hf	GS2	1.0	05Li24	
^{163}Ta−u		−45874	63	−45660	40	3.4	C				GS2	1.0	05Li24	*
^{163}Dy ^{35}Cl−^{161}Dy ^{37}Cl		5200	60	4747.93	0.11	−1.9	U				R04	4.0	64De15	
		4746	3			0.3	U				H23	2.5	70Wh01	
		4744.7	1.2			1.1	U				H25	2.5	72Ba08	
^{163}Eu−^{163}Dy		10693.5	3.7	10528.6	1.2	−44.6	F				JY1	1.0	18Vi02	*
^{163}Ho O−^{163}Dy O		2.72	0.77	3.0738	0.0006	0.3	U				TG1	1.5	15Sc13	
^{163}Ho−^{163}Dy		3.042	0.037	3.0738	0.0006	0.9	U				SH2	1.0	15El03	
		3.07377	0.00064			0.0	1	100	58	^{163}Ho	Hep	1.0	24Sc12	
^{162}Dy ^{37}Cl−^{163}Dy ^{35}Cl		−5069	42	−4882.84	0.08	1.1	U				R04	4.0	64De15	
^{163}Dy−^{162}Dy		2164	35	1932.72	0.05	−1.7	U				R04	4.0	64De15	
		1985	38			−0.3	U				R04	4.0	64De15	
		2174	40			−1.5	U				R04	4.0	64De15	
^{163}Dy O−C_{15}		−76349.06	0.86	−76348.5	0.7	0.4	1	31	31	^{163}Dy	TG1	1.5	15Sc13	
^{163}Ho O−C_{15}		−76346.61	0.97	−76345.4	0.7	0.8	1	24	24	^{163}Ho	TG1	1.5	15Sc13	
^{163}Ta(α)^{159}Lu		4741.5	15.	4749	5	0.5	3						83Sc18	*
		4746.7	10.			0.2	3						86Ru05	
		4751.8	7.			−0.4	3						92Ha10	
^{163}W(α)^{159}Hf		5520.3	5.	5520	60	−0.0	3						73Ea01	Z
		5518.1	5.			0.0	3				GSa		79Ho10	Z
		5519.9	3.			−0.0	3				Ora		82De11	Z
		5525.9	10.3			−0.1	U						84Sc06	*
		5518.7	6.			0.0	3				Daa		96Pa01	
^{163}Re(α)^{159}Ta		6017.9	5.	6012	8	−1.1	o				Ara		97Da07	*
^{163}Re^{m}(α)^{159}Ta^{m}		6067.2	6.	6068	3	0.2	–				GSa		79Ho10	
		6067.2	7.			0.1	–				Daa		96Pa01	
		6069.2	5.			−0.2	–				Ara		97Da07	
	ave.	6068	3			0.0	1	99	85	^{159}Ta^{m}			average	
^{163}Os(α)^{159}W		6674.1	30.	6673	7	−0.0	4				GSa		81Ho10	
		6678.2	10.			−0.5	4				ORa		96Bi07	
		6676.2	19.			−0.2	4				Daa		96Pa01	
		6674.1	30.			−0.0	o				Jya		13Dr06	*
		6662.8	12.3			0.8	4				Jya		19Hi06	
^{161}Dy(t,p)^{163}Dy		5986.3	1.5	5986.20	0.08	−0.1	U				McM		88Bu08	*
^{163}Dy(p,t)^{161}Dy		−5985	5	−5986.20	0.08	−0.2	U				Min		73Oo01	
		−5987.1	2.2			0.4	U				McM		88Bu08	*
^{163}Dy(t,α)^{162}Tb		11741	3	11739.1	2.0	−0.6	1	43	40	^{162}Tb	McM		07Bu29	
^{162}Dy(n,γ)^{163}Dy		6270.98	0.06	6271.01	0.05	0.4	–				MMn		82Is05	Z
		6271.00	0.09			0.1	–				ILn		89Sc31	Z
		6271.14	0.13			−1.0	–				Bdn		06Fi.A	
^{163}Dy(γ,n)^{162}Dy		−6320	110	−6271.01	0.05	0.4	U				Phi		60Ge01	
^{162}Dy(d,p)^{163}Dy		4049	5	4046.44	0.05	−0.5	U				Tal		67Sc05	
		4045	10			0.1	U				Kop		70Gr46	
^{163}Dy(d,t)^{162}Dy		−14	5	−13.78	0.05	0.0	U						67Ba34	
		−27	10			1.3	U				Kop		70Gr46	
^{162}Dy(n,γ)^{163}Dy	ave.	6271.01	0.05	6271.01	0.05	−0.0	1	100	100	^{162}Dy			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{162}Dy(^{3}He,d)^{163}Ho−^{164}Dy()^{165}Ho	−734.3	1.0	−733.5	0.7	0.8	1	55	35 ^{165}Ho	McM		75Bu02	
^{162}Er(d,p)^{163}Er	4682	10	4680	5	−0.2	1	21	21 ^{163}Er	Kop		69Tj01	
^{163}Eu(β^-)^{163}Gd	4828	70	4814.8	1.2	−0.2	o			Kur		07Ha57	*
	4813	70			0.0	o			Kur		10Ha38	*
	4829	65			−0.2	U			Kur		14Ha38	
^{163}Gd^{m}(IT)^{163}Gd	137.8	1.0	138.22	0.20	0.4	U					14Ha38	*
	138.2	0.2			0.1	1	98	49 ^{163}Gd			20Za04	
^{163}Gd(β^-)^{163}Tb	3170	70	3207	4	0.5	o			Kur		07Ha57	
	3150	70			0.8	o			Kur		10Ha38	
	3187	40			0.5	U			Kur		14Ha38	
^{163}Tb(β^-)^{163}Dy	1684	50	1785	4	2.0	U					66Fu08	*
	1721	100			0.6	U					71Ka22	*
^{163}Ho(ε)^{163}Dy	2.83	0.05	2.8632	0.0006	0.7	U					82An19	*
	2.65	0.20			1.1	U					83Ba32	
	2.84	0.10			0.2	U					84La.A	*
	2.56	0.05			6.1	B					85Ha12	*
	2.60	0.03			8.8	B					86Ya17	
	2.561	0.020			15.1	B					92Ha15	
	2.54	0.03			10.8	C					93Bo.A	*
	2.71	0.10			1.5	U					94Ya07	
	2.800	0.050			1.3	U					97Ga12	
	2.858	0.051			0.1	U					17Ra15	
^{163}Er(β^+)^{163}Ho	1210	6	1211	5	0.1	1	58	58 ^{163}Er			63Pe16	
^{163}Tm(β^+)^{163}Er	2439	3				2					82Vy07	*
	2360	100	2439.0	3.0	0.8	U			IRS		93Al03	
^{163}Yb(β^+)^{163}Tm	3370	100	3433	16	0.6	U					75Ad09	*
^{163}Lu(β^+)^{163}Yb	4860	170	4500	30	−2.1	U					83Ge08	
	4600	200			−0.5	o			IRS		83Vi.A	
	4600	200			−0.5	U			IRS		93Al03	
^{163}Re^{m}(IT)^{163}Re	115.1	4.0	120	5	1.2	o			Ara		97Da07	*
*^{163}Gd−u	Represents frequency ratio $^{163}Gd^{++}/(C_{12}H_4)^+$ =0.933275822(91)										WgM124	**
*^{163}Gd−u	$M-A$=–61200.6(3.4) for ^{163}Gd^{m} at 137.8(1.0)										HWJ188	**
*^{163}Gd−u	reference ion ^{163}Dy wrongly identified, most likely ^{146}LaOH or mixture										20Vi04	**
*^{163}Gd−^{80}Kr$_{2.038}$	Probably mixture state										HWJ205	**
*^{163}Gd−^{80}Kr$_{2.038}$	Probably mixture state										HWJ205	**
*^{163}Ta−u	$M-A$=–42644(28) keV for mixture gs+m at 174#25 keV										Nub269	**
*^{163}Eu−^{163}Dy	reference ion ^{163}Dy wrongly identified, most likely ^{146}LaOH or mixture										20Vi04	**
*^{163}Ta(α)^{159}Lu	Original assignment to 13 s ^{164}Ta changed to ^{163}Ta										86Ru05	**
*^{163}W(α)^{159}Hf	Originally assigned to ^{166}Re, re-assigned in reference										92Me10	**
*	original E_α=5372 recalibrated using their ^{168}Os−^{170}Os results										GAu	**
*^{163}Re(α)^{159}Ta	Replaced by author's value for ^{159}Ta^{m}(IT)										AHW	**
*^{163}Os(α)^{159}W	Error not given, estimated by evaluator										GAu	**
*^{161}Dy(t,p)^{163}Dy	$Q-Q$(^{162}Dy(t,p))=539.1(1.5) keV										AHW	**
*^{163}Dy(p,t)^{161}Dy	$Q-Q$(^{164}Dy(p,t))=–539.9(2.2) keV										AHW	**
*^{163}Eu(β^-)^{163}Gd	E_{β^-}=4690(70) to ^{163}Gd^{m} at 137.8 keV										14Ha38	**
*^{163}Eu(β^-)^{163}Gd	E_{β^-}=4675(70) to ^{163}Gd^{m} at 137.8 keV										14Ha38	**
*^{163}Tb(β^-)^{163}Dy	E_{β^-}=800(50) to 1/2$^+$ level at 884.2943 keV										Ens105	**
*^{163}Tb(β^-)^{163}Dy	E_{β^-}=1300(100) to 3/2$^-$ level at 421.8439 keV										Ens105	**
*^{163}Ho(ε)^{163}Dy	Original 2.58(0.10) from partial T=40(12)×10^{+3} y, re-evaluated by authors										84La.A	**
*^{163}Ho(ε)^{163}Dy	Original value 2.82(+0.11–0.08)										84La.A	**
*^{163}Ho(ε)^{163}Dy	Original value 2.60(0.03) corrected to 2.561(0.020) for dynamic effects										87Sp02	**
*	error 0.020 is statistical only										87Sp02	**
*^{163}Ho(ε)^{163}Dy	Original 2616 < Q < 2694 eV 68% CL for charge 66$^+$ Q_{β^+},										92Ju01	**
*	corrected to 2511< Q_{β^+} <2572 eV 68% CL										93Bo.A	**
*^{163}Tm(β^+)^{163}Er	E_{β^+}=884(3) to 1/2$^+$ level at 540.56 keV, and other E_{β^+}										Ens105	**
*^{163}Yb(β^+)^{163}Tm	E_{β^+}=1400(100) to 5/2$^-$ level at 947.29 keV										Ens105	**
*^{163}Re^{m}(IT)^{163}Re	Redundant with ^{167}Ir(α)^{163}Re in same paper										GAu	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{164}Sm−^{84}Kr$_{1.952}$		121306.5	4.4				2				CP2	1.0	18Or02	
^{164}Eu−^{136}Xe$_{1.206}$		54753.5	3.9	54752.3	2.2	−0.3	1	32	32	^{164}Eu	JY1	1.0	20Vi04	
^{164}Eu−^{84}Kr$_{1.952}$		115608.8	2.7	115609.4	2.2	0.2	1	68	68	^{164}Eu	CP2	1.0	20Or03	
^{164}Gd−^{84}Kr$_{1.952}$		108672.5	1.2	108672.6	1.1	0.1	1	80	80	^{164}Gd	CP2	1.0	18Or.A	
^{164}Tb−^{84}Kr$_{1.952}$		106084.0	2.0				2				CP2	1.0	20Or03	
^{164}Tb^{m}−^{84}Kr$_{1.952}$		106240	13				2				CP2	1.0	20Or03	
C_{13} H_8−^{164}Dy		133320	38	133419.8	0.7	0.7	U				R04	4.0	64De15	
C_{12} ^{13}C H_7−^{164}Dy		128920	34	128949.6	0.7	0.2	U				R04	4.0	64De15	
C_{12} H_6 N−^{164}Er		120876	39	120816.8	0.7	−0.4	U				R04	4.0	64De15	
^{164}Tm−u		−66446	31	−66462	27	−0.5	1	75	75	^{164}Tm	GS2	1.0	05Li24	*
^{164}Yb−^{142}Sm$_{1.155}$		32429	19	32434	16	0.3	2				MA7	1.0	01Bo59	
^{164}Yb−u		−65690	104	−65501	16	1.8	U				GS1	1.0	00Ra23	
		−65493	30			−0.3	R				GS2	1.0	05Li24	
^{164}Lu−u		−58750	110	−58660	30	0.8	U				GS1	1.0	00Ra23	
		−58661	30				2				GS2	1.0	05Li24	
^{164}Hf−u		−55620	110	−55630	17	−0.1	U				GS1	1.0	00Ra23	
		−55596	30			−1.1	1	31	31	^{164}Hf	GS2	1.0	05Li24	
^{164}Ta−u		−46466	30				2				GS2	1.0	05Li24	
^{164}Gd−^{171}Yb$_{.959}$		−3025.2	2.4	−3025.7	1.1	−0.2	1	20	20	^{164}Gd	JY1	1.0	18Vi02	
^{164}Tb−^{171}Yb$_{.959}$		−5597.6	3.6	−5614.4	2.0	−4.7	B				JY1	1.0	18Vi02	*
^{164}Dy ^{35}Cl−^{162}Dy ^{37}Cl		5347	5	5326.44	0.11	−1.0	U				H12	4.0	64Ba15	
		5589	19			−3.5	B				R04	4.0	64De15	
		5321	3			0.7	U				H23	2.5	70Wh01	
		5326.5	0.9			−0.0	U				H25	2.5	72Ba08	
^{164}Er ^{35}Cl−^{162}Er ^{37}Cl		3373.3	1.3	3370.6	0.4	−0.8	U				H25	2.5	72Ba08	
^{164}Er−^{164}Dy		26.92	0.12	26.92	0.12	−0.0	1	100	100	^{164}Er	SH1	1.0	11El08	
^{164}Dy ^{35}Cl−^{161}Dy ^{37}Cl		5610	48	5191.53	0.13	−2.2	U				R04	4.0	64De15	
^{163}Dy ^{37}Cl−^{164}Dy ^{35}Cl		−3360	50	−3393.73	0.10	−0.2	U				R04	4.0	64De15	
^{164}Dy−^{163}Dy		392	48	443.60	0.07	0.3	U				R04	4.0	64De15	
		540	25			−1.0	U				R04	4.0	64De15	
		446	28			−0.0	U				R04	4.0	64De15	
^{164}Er−^{162}Er		556	48	420.4	0.4	−0.7	U				R04	4.0	64De15	
^{164}W(α)^{160}Hf		5281.7	5.	5278.4	2.0	−0.6	–						73Ea01	Z
		5274.7	5.			0.7	–				ORa		75To05	Z
		5268.7	10.			1.0	U						78Sc26	*
		5279.0	5.			−0.1	–				GSa		79Ho10	
		5279.2	3.			−0.3	–				Ora		82De11	Z
		5283.0	8.			−0.6	U						84Sc06	*
		5277.0	6.			0.2	–				Daa		96Pa01	
	ave.	5278.6	2.0			−0.1	1	99	80	^{164}W			average	
^{164}Re(α)^{160}Ta^{m}		5763.8	10.				4				Jya		09Ha42	
^{164}Re^{m}(α)^{160}Ta		5922.7	10.	5926	5	0.4	4				GSa		79Ho10	
		5928.9	7.			−0.4	4				Daa		96Pa01	
		5924.7	10.			0.2	4				Jya		09Ha42	
^{164}Os(α)^{160}W		6478.3	20.	6479	5	0.1	U				GSa		81Ho10	
		6473.2	10.			0.6	–				ORa		96Bi07	
		6479.4	7.			0.0	–				Daa		96Pa01	
	ave.	6477	6			0.4	1	80	80	^{164}Os			average	
^{164}Ir^{m}(α)^{160}Re^{m}		7052.3	10.2				6				Jya		14Dr02	
^{162}Dy(t,p)^{164}Dy		5447.3	1.9	5447.32	0.08	0.0	U				McM		88Bu08	*
^{164}Dy(p,t)^{162}Dy		−5450	5	−5447.32	0.08	0.5	U				Min		73Oo01	
^{164}Er(p,t)^{162}Er		−7262	10	−7269.2	0.3	−0.7	U				Min		73Oo01	
^{164}Dy(t,α)^{163}Tb		11153	4				2				McM		92Ga15	*
^{163}Dy(n,γ)^{164}Dy		7658.11	0.07	7658.11	0.07	−0.0	1	99	75	^{164}Dy	MMn		82Is05	Z
		7658.90	0.06			−13.2	C						99Fo.A	
		7655.0	0.9			3.5	C				Bdn		06Fi.A	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{163}Dy(d,p)^{164}Dy	5434	5	5433.54	0.07	−0.1	U			Tal		64Sh06	
	5441	10			−0.7	U			Kop		70Gr46	
^{164}Dy(d,t)^{163}Dy	−1407	10	−1400.88	0.07	0.6	U			Kop		70Gr46	
^{163}Dy(^{3}He,d)^{164}Ho−^{164}Dy()^{165}Ho	−331.6	1.4	−330.7	1.1	0.6	1	67	67 ^{164}Ho	McM		75Bu02	*
^{164}Er(d,t)^{163}Er	−2593	10	−2589	5	0.4	1	21	21 ^{163}Er	Kop		69Tj01	
^{164}Ir^{m}(p)^{163}Os	1828	8	1824	6	−0.5	o			Jyp		01Ke05	
	1818	14			0.4	5			Arp		02Ma61	
	1825	6			−0.2	5			Jyp		14Dr02	
^{164}Eu(β^-)^{164}Gd	6430	70	6461.5	2.3	0.5	o			Kur		07Ha57	
	6440	70			0.3	o			Kur		10Ha38	
	6393	50			1.4	U			Kur		14Ha38	
^{164}Tb(β^-)^{164}Dy	3890	100	3863.0	2.0	−0.3	U					71Gu18	*
^{164}Ho(β^-)^{164}Er	990	30	962.1	1.4	−0.9	U					54Br96	
	965	20			−0.1	U					66Se07	
^{164}Tm(β^+)^{164}Er	3985	20	4034	25	2.4	B					67Vr04	*
	3989	50			0.9	1	25	25 ^{164}Tm	IRS		94Po26	*
^{164}Lu(β^+)^{164}Yb	6390	140	6370	30	−0.1	U					83Ge08	
	6250	90			1.3	o			IRS		83Vi.A	
	6290	90			0.9	U			IRS		93Al03	*
	6255	120			1.0	U			Dbn		94Po26	*
*^{164}Tm−u	$M-A$=–61884(28) keV for mixture gs+m at 20(12) keV										Nub269	**
*^{164}Tb−^{171}Yb$_{.959}$	Could be a mixture of ground state and iso.										HWJ208	**
*^{164}W(α)^{160}Hf	Originally assigned to ^{168}Re										AHW	**
*^{164}W(α)^{160}Hf	Originally assigned to ^{167}Re, re-assigned in reference										92Me10	**
*	original E_α=5136 recalibrated using their ^{168}Os−^{170}Os results										GAu	**
*^{162}Dy(t,p)^{164}Dy	$Q-Q(^{160}$Dy(t,p))=–722.3(1.9) keV, see ^{162}Dy(p,t)										88Bu08	**
*^{164}Dy(t,α)^{163}Tb	$Q-Q(^{162}$Dy(t,α)=–123(4)+54−584=–653(4) keV										AHW	**
*^{163}Dy(^{3}He,d)^{164}Ho−^{164}Dy()^{165}H	See erratum										75Bu02	**
*^{164}Tb(β^-)^{164}Dy	E_{β^-}=1700(100) to 4^+ level at 2194.44 and 4^+ at 2205.63 keV, and other E_{β^-}										Ens017	**
*^{164}Tm(β^+)^{164}Er	E_{β^+}=2940(20) 29 to ground state 10 to 2^+ level at 91.38 keV										Ens017	**
*^{164}Tm(β^+)^{164}Er	E_{β^+}=2944(50) 29 to ground state 10 to 2^+ level at 91.38 keV										Ens017	**
*^{164}Lu(β^+)^{164}Yb	Q_{β^+}=6250(90) partly to 2^+ level at 123.31 keV										Ens017	**
*^{164}Lu(β^+)^{164}Yb	E_{β^+}=5191(120) partly to 2^+ level at 123.31 keV										Ens017	**
^{165}Eu−^{136}Xe$_{1.213}$	58091.2	6.5	58089	6	−0.4	1	74	74 ^{165}Eu	JY1	1.0	20Vi04	
^{165}Eu−^{84}Kr$_{1.964}$	119352	11	119359	6	0.6	1	26	26 ^{165}Eu	CP2	1.0	22Or02	
^{165}Gd−^{84}Kr$_{1.964}$	113134.9	1.5	113135.5	1.4	0.4	1	87	87 ^{165}Gd	CP2	1.0	18Or.A	
^{165}Tb−^{136}Xe$_{1.213}$	47497.3	4.2	47504.0	1.7	1.6	1	16	16 ^{165}Tb	JY1	1.0	20Vi04	
^{165}Tb−^{84}Kr$_{1.964}$	108774.9	1.8	108773.7	1.7	−0.7	1	84	84 ^{165}Tb	CP2	1.0	18Or.A	
C_{13} H_9−^{165}Ho	140043	29	140096.4	0.8	0.5	U			R04	4.0	64De15	
C_{12} H_7 N−^{165}Ho	127537	28	127520.4	0.8	−0.1	U			R04	4.0	64De15	
C_{11} ^{13}C H_6 N−^{165}Ho	122970	50	123050.2	0.8	0.4	U			R04	4.0	64De15	
^{165}Tm−^{142}Sm$_{1.162}$	30970	20	30970.0	2.8	0.0	U			MA7	1.0	01Bo59	
^{165}Yb−u	−64721	30	−64730	28	−0.3	1	90	90 ^{165}Yb	GS2	1.0	05Li24	
^{165}Lu−u	−60602	30	−60593	28	0.3	1	90	90 ^{165}Lu	GS2	1.0	05Li24	
^{165}Hf−u	−55360	140	−55430	30	−0.5	U			GS1	1.0	00Ra23	
	−55433	30				2			GS2	1.0	05Li24	
^{165}Ta−u	−49191	30	−49217	14	−0.9	1	22	22 ^{165}Ta	GS2	1.0	05Li24	
^{165}W−u	−41720	30	−41720	28	0.0	1	85	85 ^{165}W	GS2	1.0	05Li24	
^{165}Gd O−^{171}Yb$_{1.058}$	1597.3	3.9	1593.0	1.4	−1.1	1	13	13 ^{165}Gd	JY1	1.0	18Vi02	
^{165}Ho ^{35}Cl−^{163}Dy ^{37}Cl	4539	4	4542.1	0.8	0.3	U			H23	2.5	70Wh01	
^{165}W(α)^{161}Hf	5031.0	5.	5030	30	−0.0	–			ORa		75To05	Z
	5034.2	10.			−0.1	–					84Sc06	*
ave.	5030	40			−0.0	1	27	15 ^{165}W			average	
^{165}Re(α)^{161}Ta	5629.6	6.	5694	6	10.8	B			Jya		05Sc22	
	5694.3	6.1				6			Jya		12Th13	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{165}Re$^{m}(\alpha)^{161}$Ta^{m}	5631.7	10.	5660.7	2.8	2.9	U					78Sc26	*
	5643.0	10.			1.8	U			GSa		81Ho10	
	5664.5	4.			−0.9	–			Ora		82De11	*
	5655.4	5.			1.0	–			Daa		96Pa01	*
	5657.4	5.			0.6	o			Jya		05Sc22	
	5657.4	6.			0.5	–			Jya		12Th13	
ave.	5660.2	2.8			0.2	1	99	59 ^{165}Re^{m}			average	
^{165}Os$(\alpha)^{161}$W	6354.3	20.	6335	6	−0.9	5			Ora		78Ca11	
	6317.4	10.			1.8	5			GSa		81Ho10	
	6342.1	7.			−0.9	5			Daa		96Pa01	
	6342.1	30.			−0.2	U			Jya		13Dr06	*
^{165}Ir$^{m}(\alpha)^{161}$Re^{m}	6882.1	7.	6879	6	−0.4	1	70	48 ^{165}Ir^{m}	Ara		97Da07	
^{165}Pt$(\alpha)^{161}$Os	7453.0	14.3				4			Jya		19Hi06	
^{163}Dy(t,p)^{165}Dy	4890.6	2.9	4892.27	0.09	0.6	U			McM		88Bu08	*
^{164}Dy(n,γ)^{165}Dy	5716.36	0.20	5715.96	0.05	−2.0	U			ILn		79Br25	Z
	5715.96	0.06			−0.0	2			MMn		82Is05	Z
	5715.70	0.30			0.9	U			ILn		90Ka21	Z
	5715.95	0.12			0.1	2			Bdn		06Fi.A	
^{164}Dy(d,p)^{165}Dy	3488	5	3491.39	0.05	0.7	U			Tal		64Sh13	
	3496	10			−0.5	U			Kop		70Gr46	
^{164}Dy(^{3}He,d)^{165}Ho	717.3	10.	725.8	0.7	0.9	U			McM		75Bu02	
^{165}Ho(γ,n)^{164}Ho	−8160	80	−7988.8	1.1	2.1	U			Phi		60Ge01	
	−7987	2			−0.9	1	33	33 ^{164}Ho	MMn		85Ts01	
^{165}Ho(d,t)^{164}Ho	−1730	15	−1731.6	1.1	−0.1	U			Tal		70Jo11	
^{164}Er(n,γ)^{165}Er	6650.1	0.6	6650.1	0.6	−0.0	1	96	96 ^{165}Er			70Bo29	Z
^{164}Er(d,p)^{165}Er	4431	10	4425.5	0.6	−0.5	U			Kop		69Tj01	
^{164}Er(α,t)^{165}Tm−^{168}Er()^{169}Tm	−1298.0	2.0	−1297.9	1.5	0.1	1	55	48 ^{165}Tm	McM		75Bu02	
^{165}Ir(p)^{164}Os	1463	38				2			Jyp		20Hi.B	
^{165}Ir^{m}(p)^{164}Os	1717.5	7.	1721	6	0.4	1	72	52 ^{165}Ir^{m}	Ara		97Da07	
^{165}Eu$(\beta^{-})^{165}$Gd	5800	120	5797	5	−0.0	o			Kur		07Ha57	
	5800	120			−0.0	o			Kur		10Ha38	
	5729	65			1.0	U			Kur		14Ha38	
^{165}Gd$(\beta^{-})^{165}$Tb	4113	65	4063.1	2.0	−0.8	U			Kur		14Ha38	
^{165}Dy$(\beta^{-})^{165}$Ho	1305	20	1285.7	0.7	−1.0	U					59Bo52	
	1285	10			0.1	U					63Pe11	
^{165}Er$(\varepsilon)^{165}$Ho	370	10	376.6	1.0	0.7	U					63Ry01	
	371	6			0.9	U					63Zy01	
^{165}Tm$(\beta^{+})^{165}$Er	1591.3	2.0	1591.4	1.5	0.1	1	55	52 ^{165}Tm			82Vy03	*
^{165}Yb$(\beta^{+})^{165}$Tm	2762	20	2635	27	−6.4	B					67Pa04	*
^{165}Lu$(\beta^{+})^{165}$Yb	4250	140	3850	40	−2.8	B					83Ge08	
	3920	80			−0.8	o			IRS		83Vi.A	
	3920	80			−0.8	1	20	10 ^{165}Yb	IRS		93Al03	
*^{165}W$(\alpha)^{161}$Hf	Originally assigned to ^{168}Re, re-assigned in reference										92Me10	**
*	original E_{α}=4894 recalibrated using their ^{168}Os−^{170}Os results										GAu	**
*^{165}Re$^{m}(\alpha)^{161}$Ta^{m}	Originally assigned to ^{166}Re										AHW	**
*^{165}Re$^{m}(\alpha)^{161}$Ta^{m}	Originally assigned to ^{166}Re										AHW	**
*^{165}Re$^{m}(\alpha)^{161}$Ta^{m}	Due to a high spin isomer										99Po09	**
*^{165}Os$(\alpha)^{161}$W	Error not given, estimated by evaluator										GAu	**
*^{163}Dy(t,p)^{165}Dy	$Q-Q(^{162}$Dy(t,p))=–556.6(2.9) keV										AHW	**
*^{165}Tm$(\beta^{+})^{165}$Er	$E_{\beta^{+}}$=272(2) to 1/2^{-} level at 297.371 keV										Ens066	**
*^{165}Yb$(\beta^{+})^{165}$Tm	$E_{\beta^{+}}$=1580(20) to 7/2^{-} level at 160.47 keV										Ens066	**
^{166}Gd−^{136}Xe$_{1.221}$	54904.1	3.9	54921.5	1.7	4.5	B			JY1	1.0	18Vi02	
^{166}Gd−^{84}Kr$_{1.976}$	116510.9	1.7				2			CP2	1.0	18Or.A	
^{166}Tb−^{136}Xe$_{1.221}$	51232.1	4.1	51230.9	1.6	−0.3	1	15	15 ^{166}Tb	JY1	1.0	20Vi04	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{166}Tb−^{84}Kr$_{1.976}$		112820.0	1.7	112820.2	1.6	0.1	1	85	85 ^{166}Tb	CP2	1.0	20Or03	
C_{12} H_8 N−^{166}Er		135376	29	135373.5	0.3	−0.0	U			R04	4.0	64De15	
		135420	60			−0.2	U			R04	4.0	64De15	
C_{13} H_{10}−^{166}Er		147740	60	147949.6	0.3	0.9	U			R04	4.0	64De15	
^{166}Lu−u		−60157	108	−60140	30	0.1	U			GS1	1.0	00Ra23	*
		−60141	32				2			GS2	1.0	05Li24	*
^{166}Hf−u		−57860	110	−57820	30	0.4	U			GS1	1.0	00Ra23	
		−57820	30				2			GS2	1.0	05Li24	
^{166}Ta−u		−49488	30				2			GS2	1.0	05Li24	
^{166}W−u		−44957	30	−44970	10	−0.4	1	11	11 ^{166}W	GS2	1.0	05Li24	
^{166}Er ^{35}Cl−^{164}Er ^{37}Cl		4040.9	1.4	4043.4	0.8	0.7	U			H25	2.5	72Ba08	
^{166}Er−^{164}Er		1214	46	1093.3	0.8	−0.7	U			R04	4.0	64De15	
		1110	80			−0.1	U			R04	4.0	64De15	
^{166}W(α)^{162}Hf		4856.0	5.	4856	4	0.0	–			ORa		75To05	Z
		4855.0	10.			0.1	–			GSa		79Ho10	Z
		4858.3	8.2			−0.3	–					89Hi04	
	ave.	4856	4			−0.1	1	97	78 ^{166}W			average	
^{166}Re(α)^{162}Ta		5461.8	10.	5570	50	2.2	B					78Sc26	*
		5574.5	3.				5			Ora		82De11	*
		5504.9	10.			1.4	B					84Sc06	
		5637.0	13.			−1.2	B			Bea		92Me10	*
^{166}Re(α)^{162}Ta^m		5669.9	10.	5665	4	−0.5	6			Daa		96Pa01	*
		5663.7	4.1			0.2	6			Ora		82De11	
^{166}Os(α)^{162}W		6148.5	20.	6143	3	−0.3	U					77Ca23	
		6129.0	6.			2.2	–			GSa		81Ho10	
		6148.5	6.			−0.9	–			Daa		96Pa01	
		6148.4	5.1			−1.1	–			Jya		15Li24	
	ave.	6143	3			−0.0	1	99	87 ^{166}Os			average	
^{166}Ir(α)^{162}Re		6702.8	20.	6722	6	1.0	U			GSa		81Ho10	
		6724.3	6.			−0.3	7			Ara		97Da07	
		6713.1	13.			0.7	7			Jya		04Ke06	*
^{166}Ir^m(α)^{162}Re^m		6718.2	11.	6719	4	0.0	8			Daa		96Pa01	*
		6723.3	5.			−0.9	8			Ara		97Da07	
		6706.9	8.2			1.4	8			Jya		04Ke06	
^{166}Pt(α)^{162}Os		7285.9	15.	7292	7	0.4	5			ORa		96Bi07	
		7294.1	8.2			−0.2	5			Jya		19Hi06	
^{164}Dy(t,p)^{166}Dy		4276.4	4.4	4277.7	0.4	0.3	U			McM		88Bu08	*
^{166}Er(p,t)^{164}Er		−6641	5	−6642.4	0.7	−0.3	U			Min		73Oo01	
^{165}Dy(n,γ)^{166}Dy		7043.5	0.4				3					83Ke.A	
^{165}Ho(n,γ)^{166}Ho		6243.69	0.06	6243.640	0.020	−0.8	U			MMn		82Is05	Z
		6243.64	0.02			−0.0	1	100	50 ^{165}Ho	MMn		84Ke15	Z
		6243.68	0.13			−0.3	U			Bdn		06Fi.A	
^{165}Ho(d,p)^{166}Ho		4025	7	4019.073	0.020	−0.8	U			Tal		65St06	
^{166}Er(d,t)^{165}Er		−2218	10	−2216.9	0.9	0.1	U			Kop		69Tj01	
^{166}Ir(p)^{165}Os		1152.0	8.0				6			Ara		97Da07	
^{166}Ir^m(p)^{165}Os		1324.1	8.	1323	10	−0.1	o			Ara		97Da07	*
^{166}Eu(β^-)^{166}Gd		7322	300	7620#	100#	1.0	D			Kur		14Ha38	*
^{166}Tb(β^-)^{166}Dy		4830	100	4776.0	1.7	−0.5	o			Kur		02Sh.A	
		4695	70			1.2	o			Kur		07Ha57	
		4700	70			1.1	U			Kur		10Ha38	
^{166}Dy(β^-)^{166}Ho		483	5	485.8	0.8	0.6	U					60He09	*
^{166}Ho(β^-)^{166}Er		1859	3	1853.9	0.8	−1.7	–					63Fu17	
		1857	3			−1.0	–					66Da04	
		1854.7	1.5			−0.6	–					74Gr41	
		1851.6	2.0			1.1	–					83Ra.A	
	ave.	1854.7	1.0			−0.8	1	55	50 ^{166}Ho			average	
^{166}Tm(β^+)^{166}Er		3043	20	3038	12	−0.3	2					61Gr33	*
		3031	20			0.3	2					61Zy02	*
		3039	20			−0.1	2					63Pr13	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{166}Yb(ε)^{166}Tm		280	40	293	14	0.3	U					Averag	*
^{166}Lu(β^+)^{166}Yb		5480	160	5570	30	0.6	U					74De09	*
^{166}Ir^m(IT)^{166}Ir		171.5	6.1				7			Ara		97Da07	
*^{166}Lu−u	$M-A$=–56010(100) keV for mixture gs+m+n at 34.37 and 43.0 keV											Nub269	**
*^{166}Lu−u	$M-A$=–55995(28) keV for mixture gs+m+n at 34.37 and 43.0 keV											Nub269	**
*^{166}Re(α)^{162}Ta	Originally assigned to ^{167}Re											AHW	**
*^{166}Re(α)^{162}Ta	Assignment uncertain, no other obvious attribution											AHW	**
*^{166}Re(α)^{162}Ta	Originally assigned to ^{167}Re											AHW	**
*^{166}Re(α)^{162}Ta	Assignment tentative, may be ^{165}Re											92Me10	**
*^{166}Re(α)^{162}Ta^m	Correlated with E_α=6003 keV from the high-spin state in ^{170}Ir											FGK258	**
*^{166}Ir(α)^{162}Re	All Q_α of reference increased by 7 keV for calibration error											04Ke06	**
*^{166}Ir^m(α)^{162}Re^m	Correlated with E_α=6123 of ^{162}Re^m											96Pa01	**
*^{164}Dy(t,p)^{166}Dy	$Q-Q(^{162}$Dy(t,p))=–1170.8(4.4) keV											AHW	**
*^{166}Ir^m(p)^{165}Os	Replaced by author's value for ^{166}Ir^m(IT)^{166}Ir											97Da07	**
*^{166}Eu(β^-)^{166}Gd	Trends in Mass Surface TMS suggest ^{166}Eu 300 keV less bound											GAu	**
*^{166}Dy(β^-)^{166}Ho	E_{β^-}=402(5) to 1^- level at 82.47 keV, and other E_{β^-}											Ens084	**
*^{166}Tm(β^+)^{166}Er	E_{β^+}=1940(20) 1928(20) 1936(20) respectively, to 2^+ level at 80.5776 keV											Ens084	**
*^{166}Yb(ε)^{166}Tm	Average pK=0.712(0.038) to 1^+ level at 82.298 keV from 2 references:											Ens084	**
*	pK=0.74(0.05) to 82.298 level											63Ja06	**
*	pK=0.675(0.059) to 82.298 level											73De22	**
*^{166}Lu(β^+)^{166}Yb	E_{β^+}=2225(160) to (6^-,7^-) level at 2233.36 keV											Ens084	**
^{167}Gd−^{136}Xe$_{1.228}$		59422	13	59431	6	0.7	1	19	19 ^{167}Gd	JY1	1.0	20Vi04	
^{167}Gd−^{84}Kr$_{1.988}$		121434.5	6.2	121433	6	−0.3	1	81	81 ^{167}Gd	CP2	1.0	18Or.A	
^{167}Tb−^{136}Xe$_{1.228}$		53947.0	4.1	53947.4	1.9	0.1	1	22	22 ^{167}Tb	JY1	1.0	20Vi04	
^{167}Tb−^{84}Kr$_{1.988}$		115949.8	2.4	115949.2	1.9	−0.2	1	64	64 ^{167}Tb	CP2	1.0	18Or.A	
^{167}Tb−^{133}Cs$_{1.256}$		58757.1	5.1	58759.1	1.9	0.4	1	14	14 ^{167}Tb	JY2	1.0	24Ja12	
^{167}Dy−^{82}Kr$_{2.037}$		111921.3	4.3	111918	3	−0.8	1	54	54 ^{167}Dy	CP2	1.0	18Or.A	
^{167}Dy−^{133}Cs$_{1.256}$		54427.3	4.7	54431	3	0.9	1	46	46 ^{167}Dy	JY2	1.0	24Ja12	
C$_{13}$ H$_{11}$−^{167}Er		153840	130	154019.50	0.29	0.3	U			R04	4.0	64De15	
		154040.4	6.2			−0.8	U			M23	4.0	79Ha32	
C$_{12}$ H$_9$ N−^{167}Er		141480	27	141443.44	0.29	−0.3	U			R04	4.0	64De15	
		141520	50			−0.4	U			R04	4.0	64De15	
^{167}Lu−u		−61748	32				2			GS2	1.0	05Li24	*
^{167}Hf−u		−57490	110	−57400	30	0.8	U			GS1	1.0	00Ra23	
		−57400	30				2			GS2	1.0	05Li24	
^{167}Ta−u		−51870	120	−51910	30	−0.3	U			GS1	1.0	00Ra23	
		−51907	30				2			GS2	1.0	05Li24	
^{167}W^m−u		−45175	30	−45194	18	−0.6	1	34	34 ^{167}W^m	GS2	1.0	05Li24	*
^{167}Er ^{35}Cl−^{165}Ho ^{37}Cl		4666	3	4677.1	0.8	1.5	U			H23	2.5	70Wh01	
		4679.5	1.2			−0.8	U			H25	2.5	72Ba08	
^{167}Er−^{166}Er		1722	31	1755.12	0.19	0.3	U			R04	4.0	64De15	
^{167}W(α)^{163}Hf		4661.9	20.	4641	29	−0.4	–					89Me02	
		4671.1	13.			−0.6	–					91Me05	
	ave.	4670	40			−0.4	1	22	16 ^{163}Hf			average	
^{167}Re(α)^{163}Ta^m		5138.3	12.				12			Bea		92Me10	
^{167}Re^m(α)^{163}Ta		5408.8	3.	5407.0	2.9	−0.6	4			Ora		82De11	*
		5397.5	10.			0.9	4			ChR		84Sc06	*
		5392.4	12.			1.2	4			Bea		92Me10	
^{167}Os(α)^{163}W		5983.6	5.	5980	60	0.0	4			GSa		81Ho10	Z
		5978.7	2.			0.1	4			Ora		82De11	Z
		5996.9	5.			−0.2	4			Daa		96Pa01	
		5979.5	5.			0.1	4			Bka		02Ro17	
^{167}Ir(α)^{163}Re		6507.1	5.	6504.9	2.6	−0.4	2			Ara		97Da07	
		6504.0	3.			0.3	2			Jya		05Sc22	
^{167}Ir^m(α)^{163}Re^m		6543.0	10.	6561	3	1.7	–			GSa		81Ho10	
		6567.6	11.			−0.6	–			Daa		96Pa01	
		6567.6	5.			−1.4	–			Ara		97Da07	
		6551.2	7.2			1.3	–			Jya		04Ke06	
		6561.5	6.			−0.1	–			Jya		05Sc22	
	ave.	6561	3			0.0	1	99	86 ^{163}Re^m			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{167}Pt(α)^{163}Os		7159.8	10.	7160	60	−0.0	5			ORa		96Bi07	
		7150.6	10.			0.1	o			Jya		04Ke06	
		7156.4	8.2			0.0	5			Jya		19Hi06	
^{167}Er(p,t)^{165}Er		−6427	6	−6428.8	0.9	−0.3	U			Min		73Oo01	
		−6430	5			0.2	U					75St08	
^{166}Er(n,γ)^{167}Er		6436.35	0.50	6436.43	0.18	0.2	–					70Bo29	Z
		6436.51	0.40			−0.2	–					70Mi01	Z
		6436.46	0.22			−0.1	–			Bdn		06Fi.A	
^{167}Er(γ,n)^{166}Er		−6560	80	−6436.43	0.18	1.5	U			Phi		60Ge01	
^{166}Er(d,p)^{167}Er		4209	10	4211.86	0.18	0.3	U			Tal		68Ha10	
		4214	10			−0.2	U			Kop		69Tj01	
^{167}Er(d,t)^{166}Er		−189	12	−179.20	0.18	0.8	U			Kop		69Bu01	
^{166}Er(n,γ)^{167}Er	ave.	6436.46	0.18	6436.43	0.18	−0.1	1	99	96 ^{166}Er			average	
^{166}Er(α,t)^{167}Tm−^{168}Er()^{169}Tm		−666.5	1.0	−666.4	1.0	0.1	1	99	99 ^{167}Tm	McM		75Bu02	
^{167}Ir(p)^{166}Os		1070.5	6.	1070	4	−0.1	–					97Da07	
		1068.5	6.			0.3	–			Jyp		05Sc22	
	ave.	1070	4			0.1	1	76	67 ^{167}Ir			average	
^{167}Ir^{m}(p)^{166}Os		1245.5	7.	1245	4	−0.0	o					97Da07	*
		1256	7			−1.5	o			Jyp		05Sc22	*
^{167}Dy(β^-)^{167}Ho		2350	60	2364	6	0.2	U					77Tu01	*
^{167}Ho(β^-)^{167}Er		970	20	1011	5	2.0	U					68Fu07	
^{167}Yb(β^+)^{167}Tm		1954	4	1953	4	−0.2	1	90	89 ^{167}Yb			77Kr.A	*
^{167}Lu(β^+)^{167}Yb		3130	100	3070	30	−0.6	U					64Ag.A	*
^{167}W(β^+)^{167}Ta		5620	270	6130	30	1.9	U			Got		89Me02	
^{167}W^{m}(IT)^{167}W		127.1	1.7	127.2	1.7	0.0	1	100	66 ^{167}W^{m}			92Th06	
^{167}Ir^{m}(IT)^{167}Ir		175.3	2.2	175.4	2.1	0.1	1	94	61 ^{167}Ir^{m}	Ara		97Da07	
		188.0	9.2			−1.4	U			Jyp		05Sc22	*
*^{167}Lu−u	$M-A$=–57501(28) keV for mixture gs+m at 33.7(0.4) keV											Nub269	**
*^{167}W^{m}−u	Re-assigned to the iso. by the evaluator											HWJ266	**
*^{167}Re^{m}(α)^{163}Ta	Original assignment to ^{168}Re changed in reference											92Me10	**
*^{167}Re^{m}(α)^{163}Ta	Original assignment to ^{168}Re^{m} changed in reference											92Me10	**
*	original E_α=5250 recalibrated using their ^{168}Os−^{170}Os results											GAu	**
*^{167}Ir^{m}(p)^{166}Os	Replaced by author's value for ^{167}Ir^{m}(IT)^{167}Ir											97Da07	**
*^{167}Ir^{m}(p)^{166}Os	Replaced for ^{167}Ir^{m}(IT)^{167}Ir											05Sc22	**
*^{167}Dy(β^-)^{167}Ho	E_{β^-}=1780(60) to 3/2$^-$ level at 569.69 keV											Ens008	**
*^{167}Yb(β^+)^{167}Tm	E_{β^+}=639(4) to 7/2$^-$ level at 292.820 keV											Ens008	**
*^{167}Lu(β^+)^{167}Yb	E_{β^+}=2060(100) to 5/2$^+$ level at 29.658, 7/2$^-$ at 78.671 keV											Ens008	**
*^{167}Ir^{m}(IT)^{167}Ir	From E_p=1248(7)−E_p=1062(6)											05Sc22	**
^{168}Gd−^{84}Kr$_2$		125290.6	6.0				2			CP2	1.0	24Ra.A	
^{168}Tb−^{136}Xe$_{1.235}$		57927.2	4.5	57964	5	8.2	B			JY1	1.0	20Vi04	*
^{168}Tb−^{133}Cs$_{1.263}$		62788.1	4.9				2			JY2	1.0	24Ja12	
^{168}Dy−^{84}Kr$_2$		114217.2	4.4	114218	3	0.1	1	49	49 ^{168}Dy	CP2	1.0	18Or.A	*
^{168}Dy−^{133}Cs$_{1.263}$		56628.0	4.3	56627	3	−0.1	1	51	51 ^{168}Dy	JY2	1.0	24Ja12	
^{168}Ho−^{84}Kr$_2$		112578.1	3.5	112577.9	2.5	−0.0	1	52	52 ^{168}Ho	CP2	1.0	24Ra.A	
^{168}Ho^{m}−^{84}Kr$_2$		112641.1	3.5	112641.3	2.5	0.0	1	52	52 ^{168}Ho^{m}	CP2	1.0	24Ra.A	
C$_{13}$ H$_{12}$−^{168}Er		161543.3	5.1	161522.44	0.27	−1.0	U			M23	4.0	79Ha32	
C$_{12}$ H$_{10}$ N−^{168}Er		148884	44	148946.38	0.27	0.4	U			R04	4.0	64De15	
C$_{11}$ ^{13}C H$_9$ N−^{168}Er		144524	29	144476.19	0.27	−0.4	U			R04	4.0	64De15	
C$_{12}$ H$_{10}$ N−^{168}Yb		147010	100	147433.381	0.028	1.1	U			R04	4.0	64De15	
^{168}Lu−u		−61194	62	−61240	40	−0.8	1	49	49 ^{168}Lu	GS2	1.0	05Li24	*
^{168}Lu^{m}−^{133}Cs$_{1.263}$		58320.7	6.2				2			MA8	1.0	19Hu15	
^{168}Hf−u		−59560	104	−59430	30	1.2	U			GS1	1.0	00Ra23	
		−59432	30				2			GS2	1.0	05Li24	
^{168}Ta−u		−52020	110	−51950	30	0.6	U			GS1	1.0	00Ra23	
		−51953	30				2			GS2	1.0	05Li24	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{168}W−u		−48181	30	−48196	14	−0.5	1	21	21 ^{168}W	GS2	1.0	05Li24	
^{168}Yb $^{35}Cl_2$−^{164}Dy $^{37}Cl_2$		10612.8	8.7	10610.7	0.7	−0.1	U			H27	2.5	74Ba90	
^{168}Yb−^{172}Yb$_{.977}$		−3958.822	0.025	−3958.823	0.025	−0.0	1	100	100 ^{168}Yb	Hep	1.0	25Do.1	
^{168}Er ^{35}Cl−^{166}Er ^{37}Cl		5037	50	5027.34	0.24	−0.1	U			R08	1.5	69De19	
		5026	3			0.2	U			H23	2.5	70Wh01	
		5028.9	1.5			−0.4	U			H25	2.5	72Ba08	
^{168}Yb−^{170}Yb$_{.988}$		−1658.726	0.100	−1659.092	0.028	−3.7	B			JY2	1.0	20Ne.1	
^{168}Yb−^{168}Er		1512.91	0.27	1513.00	0.26	0.3	1	96	95 ^{168}Er	SH1	1.0	11El04	
^{168}Er−^{167}Er		284	31	322.09	0.13	0.3	U			R04	4.0	64De15	
		320.9	4.3			0.1	U			M24	2.5	79Ha32	
^{168}W(α)^{164}Hf		4506.5	12.	4500	11	−0.5	1	87	69 ^{164}Hf			91Me05	
^{168}Re(α)^{164}Ta		5063	13	5200#	100#	10.5	D			Bea		92Me10	*
^{168}Os(α)^{164}W		5819.0	3.	5815.8	2.7	−1.0	–			Ora		82De11	Z
		5800.4	8.			1.9	–					84Sc06	
		5812.7	8.			0.4	–					95Hi02	
	ave.	5816.3	2.7			−0.2	1	99	78 ^{168}Os			average	
^{168}Ir(α)^{164}Re^m		6410.9	5.	6381	9	−5.9	B			Ora		82De11	
		6379.2	15.			0.1	5			Daa		96Pa01	
		6382.2	10.			−0.1	5			Jya		09Ha42	
^{168}Ir^m(α)^{164}Re		6477.5	8.	6476	6	−0.1	5			Daa		96Pa01	
		6474.4	10.			0.2	5			Jya		09Ha42	*
^{168}Pt(α)^{164}Os		6990.8	20.	6990	3	−0.1	U			GSa		81Ho10	
		6998.9	10.			−0.9	U			ORa		96Bi07	
		6986.7	8.			0.4	o			Jya		04Ke06	
		6989.7	3.1				2			Jya		09Go16	
^{168}Er(p,t)^{166}Er		−5723	6	−5725.93	0.21	−0.5	U			Min		73Oo01	
^{168}Yb(p,t)^{166}Yb		−7647	7				2			Min		73Oo01	
^{167}Er(n,γ)^{168}Er		7771.43	0.40	7771.29	0.12	−0.3	–					70Mi01	Z
		7771.05	0.20			1.2	–			ILn		79Br25	Z
		7771.6	1.0			−0.3	U			ORn		84Ka22	
		7771.0	0.5			0.6	U					85Va.A	
		7771.45	0.16			−1.0	–			Bdn		06Fi.A	
^{167}Er(d,p)^{168}Er		5541	6	5546.73	0.12	1.0	U			Tal		67Ha25	
^{168}Er(d,t)^{167}Er		−1523	10	−1514.06	0.12	0.9	U			Kop		69Tj01	
^{167}Er(n,γ)^{168}Er	ave.	7771.31	0.12	7771.29	0.12	−0.1	1	99	96 ^{167}Er			average	
^{167}Er(α,t)^{168}Tm−^{168}Er()^{169}Tm		−262.3	1.5	−262.3	1.5	0.0	1	100	100 ^{168}Tm	McM		75Bu02	
^{168}Yb(d,t)^{167}Yb		−2797	12	−2804	4	−0.6	1	11	11 ^{167}Yb	Kop		66Bu16	
^{168}Ho^m(IT)^{168}Ho		59	1	59.0	1.0	−0.0	1	96	48 ^{168}Ho			90Ch37	*
^{168}Ho(β^-)^{168}Er		2740	100	2976.5	2.4	2.4	U					73Ka07	*
		2930	30			1.6	U					90Ch37	
^{168}Lu(β^+)^{168}Yb		4475	80	4530	40	0.7	o					70Ch28	*
		4475	80			0.7	–					72Ch44	*
		4500	80			0.4	–			IRS		83Vi.A	
	ave.	4490	60			0.8	1	51	51 ^{168}Lu			average	
^{168}Lu^m(β^+)^{168}Yb		4696	100	4672	6	−0.2	U					72Ch44	*
*^{168}Tb−^{136}Xe$_{1.235}$	Ramsey spectrum contains contamination											24Ja12	**
*^{168}Dy−$^{84}Kr_2$	this datum was missed in Ame2020											18Or.A	**
*^{168}Lu−u	$M-A$=–56922(28) keV for mixture gs+m at 160(40) keV											Nub269	**
*^{168}Re(α)^{164}Ta	E_α=4833(13) to level at 111.8 keV											92Me10	**
*^{168}Re(α)^{164}Ta	Trends in Mass Surface TMS suggest ^{168}Re 140 keV less bound											HWJ261	**
*^{168}Ir^m(α)^{164}Re	E_α=6320(10), 6260(10) to ground state and level at 69 keV											09Ha42	**
*^{168}Ho^m(IT)^{168}Ho	below detection limit, estimated											90Ch37	**
*^{168}Ho(β^-)^{168}Er	E_{β^-}=1900(100) to 2^+ level at 821.17 and 3^+ at 895.79 keV											Ens108	**
*^{168}Lu(β^+)^{168}Yb	E_{β^+}=1230(80) to 2222.37 level											Ens108	**
*^{168}Lu^m(β^+)^{168}Yb	E_{β^+}=1470(100) to 4^+ level at 2203.84 keV											Ens108	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{169}Tb−^{133}Cs$_{1.271}$	66098	10				2			JY2	1.0	24Ja12	
^{169}Dy−^{133}Cs$_{1.271}$	60564.1	6.1	60570.9	1.6	1.1	1	7	7 ^{169}Dy	JY2	1.0	24Ja12	
^{169}Dy−^{84}Kr$_{2.012}$	118467.4	1.7	118466.9	1.6	−0.3	1	93	93 ^{169}Dy	CP2	1.0	24Ra.A	
^{169}Ho−^{133}Cs$_{1.271}$	57024.0	5.1	57058	4	6.6	B			JY2	1.0	24Ja12	
^{169}Ho−^{85}Rb$_{1.988}$	112249.2	4.2				2			CP2	1.0	24Ra.A	
C_{12} H_{11} N−^{169}Tm	154920	60	154930.5	0.8	0.0	U			R04	4.0	64De15	
^{169}Lu−u	−62362	31	−62355	3	0.2	U			GS2	1.0	05Li24	*
^{169}Hf−u	−58741	30				2			GS2	1.0	05Li24	
^{169}Ta−u	−53960	110	−53990	30	−0.3	U			GS1	1.0	00Ra23	
	−53989	30				2			GS2	1.0	05Li24	
^{169}W−u	−48195	30	−48225	17	−1.0	1	31	31 ^{169}W	GS2	1.0	05Li24	
^{169}Re−u	−41209	66	−41242	11	−0.5	U			GS2	1.0	05Li24	*
^{169}Tm $^{35}Cl_2$−^{165}Ho $^{37}Cl_2$	9793.0	1.1	9790.3	1.1	−1.0	1	16	8 ^{165}Ho	H25	2.5	72Ba08	
^{169}Tm ^{35}Cl−^{167}Er ^{37}Cl	5107	3	5113.1	0.8	0.8	U			H23	2.5	70Wh01	
	5113.2	1.1			−0.0	1	9	8 ^{169}Tm	H25	2.5	72Ba08	
^{169}Re(α)^{165}Ta^m	4989.3	12.				4			Bea		92Me10	*
^{169}Re^m(α)^{165}Ta	5189.1	3.	5189	3	−0.1	1	99	78 ^{165}Ta	Ora		82De11	
	5191.1	10.			−0.2	U			ChR		84Sc06	*
	5184.0	10.			0.5	U			Bea		92Me10	
^{169}Os(α)^{165}W	5717.6	4.	5713	3	−1.0	2			Ora		82De11	
	5699.2	8.			1.7	2					84Sc06	*
	5713	8			0.1	2					95Hi02	*
	5711.5	8.			0.2	2			Daa		96Pa01	
^{169}Ir(α)^{165}Re	6150.8	8.	6162	4	1.4	5			Ara		99Po09	
	6138.5	4.			5.8	o			Jya		05Sc22	
	6165	14			−0.2	U			Jya		12Th13	
	6165.2	4.1			−0.7	5			Jya		14ThZZ	
^{169}Ir^m(α)^{165}Re^m	6276.0	3.	6267.1	2.4	−3.0	B			Ora		82De11	Z
	6258.4	10.			0.9	U			GSa		84Sc.A	
	6267.6	9.			−0.1	–			Daa		96Pa01	
	6254.3	5.			2.6	B			Ara		99Po09	
	6265.6	3.			0.5	–			Jya		05Sc22	
	6268	14			−0.1	U			Jya		12Th13	
	6268.6	4.1			−0.4	–			Jya		14ThZZ	
ave.	6266.7	2.4			0.1	1	99	58 ^{169}Ir^m			average	
^{169}Pt(α)^{165}Os	6840.2	15.	6858	5	1.2	U			GSa		81Ho10	
	6860.7	23.			−0.1	U			Daa		96Pa01	
	6853.5	8.2			0.5	o			Jya		04Ke06	
	6857.6	5.1				6			Jya		09Go16	
^{168}Er(n,γ)^{169}Er	6002.5	0.7	6003.25	0.15	1.1	U					70Bo29	Z
	6003.5	0.3			−0.8	2					70Mu15	Z
	6003.16	0.18			0.5	2			Bdn		06Fi.A	
^{168}Er(d,p)^{169}Er	3773	12	3778.68	0.15	0.5	U			Tal		68Ha10	
	3781	10			−0.2	U			Kop		69Tj01	
^{168}Er(α,t)^{169}Tm	−14244.8	10.	−14239.7	0.8	0.5	U			McM		75Bu02	
^{169}Tm(γ,n)^{168}Tm	−8110	50	−8033.6	1.5	1.5	U			Phi		60Ge01	
^{169}Tm(d,t)^{168}Tm	−1775	6	−1776.4	1.5	−0.2	U			Pit		73Ko06	
^{168}Yb(n,γ)^{169}Yb	6866.8	0.4	6866.98	0.15	0.4	2					68Mi08	Z
	6867.2	0.4			−0.6	2					68Sh12	Z
	6866.97	0.18			0.0	2			Bdn		06Fi.A	
^{168}Yb(d,p)^{169}Yb	4636	12	4642.41	0.15	0.5	U			Kop		66Bu16	
^{169}Au^m(p)^{168}Pt	2195	28				3			Jvp		20Hi.B	
^{169}Dy(β^-)^{169}Ho	3200	300	3272	4	0.2	U			LBL		90Ch34	
^{169}Ho(β^-)^{169}Er	2070	100	2132	4	0.6	U					63Mi17	*
^{169}Er(β^-)^{169}Tm	343.8	3.	353.3	0.8	3.2	B					56Bi30	*
	347.8	5.			1.1	U					65Du02	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{169}Yb(ε)^{169}Tm		913	12	898.9	0.8	−1.2	U					86Ad07	*
		900	100			−0.0	U					87Sa53	*
^{169}Lu(β^+)^{169}Yb		2293	3				3					77Bo31	
^{169}Hf(β^+)^{169}Lu		3365	200	3366	28	0.0	U					69Ar23	*
		3250	90			1.3	U					73Me09	*
*^{169}Lu−u	$M-A$=–58075(28) keV for mixture gs+m at 29.0 keV											Nub269	**
*^{169}Re−u	$M-A$=–38293(29) keV for mixture gs+m at 185(11) keV											Nub269	**
*^{169}Re(α)^{165}Ta^m	E_α=4871(12), and a stronger E_α=4700(12)											92Me10	**
*^{169}Re^m(α)^{165}Ta	Original E_α=5050 recalibrated using their ^{168}Os−^{170}Os results											GAu	**
*^{169}Os(α)^{165}W	Used to recalibrate other results in same reference											GAu	**
*^{169}Os(α)^{165}W	E_α=5578(8), 5536(10) to ground state, (3/2$^-$) level at 43 keV											Ens066	**
*^{169}Ho(β^-)^{169}Er	E_{β^-}=1200(100) to 5/2$^-$ level at 853.0 and 7/2$^-$ at 941.04 keV											Ens089	**
*^{169}Er(β^-)^{169}Tm	E_{β^-}=340(2) 344(4) respectively, 55% to ground state, 45% to 3/2$^+$ level at 8.41 keV											Ens089	**
*^{169}Yb(ε)^{169}Tm	From decay rates to (5/2)$^+$ level at 781.796, (7/2$^+$) 878.35 of same band											Ens089	**
*^{169}Yb(ε)^{169}Tm	pK=0.812(0.029) to 9/2$^-$ level at 472.88 keV											Ens089	**
*^{169}Hf(β^+)^{169}Lu	E_{β^+}=1850(200) to 7/2$^-$ level at 492.88 keV											Ens089	**
*^{169}Hf(β^+)^{169}Lu	K/β^+=5.2(1.0) to 7/2$^-$ level at 492.88 keV											Ens089	**
C_{12} H_{12} N−^{170}Er		161210	70	161501.7	1.2	1.0	U			R04	4.0	64De15	
C_{12} H_{12} N−^{170}Yb		161831	43	162207.148	0.011	2.2	U			R04	4.0	64De15	
C_{11} H_8 O N−^{170}Yb		125370	150	125821.639	0.011	0.8	U			R04	4.0	64De15	
C_{11} ^{13}C H_{11} N−^{170}Yb		157320	210	157736.951	0.011	0.5	U			R04	4.0	64De15	
^{170}Yb−^{129}Xe$_{1.318}$		60266.078	0.012	60266.068	0.009	−0.8	1	53	48 ^{170}Yb	FS1	1.0	12Ra34	
^{170}Yb−^{132}Xe$_{1.288}$		58215.483	0.013	58215.491	0.009	0.6	1	45	43 ^{170}Yb	FS1	1.0	12Ra34	
^{170}Lu−u		−61529	42	−61521	18	0.2	R			GS2	1.0	05Li24	*
^{170}Hf−u		−60400	104	−60390	30	0.1	U			GS1	1.0	00Ra23	
		−60391	30				2			GS2	1.0	05Li24	
^{170}Ta−u		−53810	104	−53830	30	−0.1	U			GS1	1.0	00Ra23	
		−53825	30				2			GS2	1.0	05Li24	
^{170}W−u		−50710	110	−50768	14	−0.5	U			GS1	1.0	00Ra23	
		−50755	30			−0.4	1	22	22 ^{170}W	GS2	1.0	05Li24	
^{170}Re−u		−41782	30	−41764	12	0.6	–			GS2	1.0	05Li24	
	ave.	−41780	28			0.6	1	17	17 ^{170}Re			average	
^{170}Os−u		−36454	31	−36422	10	1.0	1	11	11 ^{170}Os	GS2	1.0	05Li24	
^{170}Er ^{35}Cl−^{168}Er ^{37}Cl		6073	31	6044.8	1.2	−0.6	U			R08	1.5	69De19	
		6040	3			0.6	U			H23	2.5	70Wh01	
		6046.9	1.8			−0.5	1	7	7 ^{170}Er	H25	2.5	72Ba08	
^{170}Yb ^{35}Cl−^{168}Yb ^{37}Cl		3806.0	7.6	3826.43	0.07	1.1	U			H27	2.5	74Ba90	
^{170}Yb−^{172}Yb$_{.988}$		−2382.796	0.025	−2382.780	0.012	0.6	1	24	14 ^{172}Yb	Hep	1.0	25Do.1	
^{170}Dy−^{170}Yb		7567.7	7.0				2			JY2	1.0	24Ja12	
^{170}Ho^m−^{170}Ho		162.0	5.8				3			JY2	1.0	24Ja12	
^{170}Ho−^{170}Yb		4831.1	8.3				2			JY2	1.0	24Ja12	
^{170}Er−^{168}Er		3450	70	3094.7	1.2	−1.3	U			R04	4.0	64De15	
^{170}Yb−^{168}Yb		910	200	876.297	0.028	−0.0	U			R04	4.0	64De15	
^{170}Os(α)^{166}W		5533.5	10.	5536.9	2.7	0.3	–			ORa		72To06	Z
		5541.8	4.1			−1.2	–			Ora		82De11	Z
		5523.2	8.			1.7	–					84Sc06	*
		5533.4	8.			0.4	–					95Hi02	
		5537.5	5.			−0.1	–			Bka		02Ro17	
	ave.	5537.1	2.7			−0.1	1	99	89 ^{170}Os			average	
^{170}Ir(α)^{166}Re		6175.4	10.	6272	10	9.6	B					78Sc26	*
		6172.7	5.			19.8	B			Ora		82De11	*
		6147.9	10.			12.4	B			Daa		96Pa01	*
		6229.9	11.			3.8	B			Daa		96Pa01	*
		6235.0	5.			7.3	B			Ara		04GoZZ	*
		6271.7	10.2				6			Jya		07Ha45	*
^{170}Ir^m(α)^{166}Re^m		5955.4	10.	5955	5	−0.0	7			Bka		02Ro17	
		5955.4	5.1			−0.0	7			Ara		04GoZZ	
^{170}Pt(α)^{166}Os		6703.0	8.	6707	3	0.5	–			GSa		81Ho10	
		6705.0	10.			0.2	–					82En03	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{170}Pt(α)^{166}Os		6708.1	6.	6707	3	−0.1	–				ORa		96Bi07	
		6711.2	11.			−0.3	–				Jya		97Uu01	
		6723.5	14.			−1.1	–				Bka		01Ro.B	
		6707.1	7.			0.0	–				Jya		04Ke06	
	ave.	6708	3			−0.1	1	84	81	^{170}Pt			average	
^{170}Au(α)^{166}Ir		7174.1	11.	7177	15	0.3	o				Jya		02Ke.C	
		7170.0	12.			0.6	U				Jya		04Ke06	
^{170}Au^m(α)^{166}Ir^m		7277.5	6.	7285	12	0.2	o				Jya		02Ke.C	
		7226.3	15.			1.1	U				Ara		02Ma61	
		7278.5	9.			0.1	U				Jya		04Ke06	
^{170}Hg(α)^{166}Pt		7773.2	30.7				6				Jya		19Hi06	
^{170}Er(p,α)^{167}Ho		7036	5				2				NDm		83Ta.A	
^{170}Er(^{18}O,^{20}Ne)^{168}Dy		4710	140	4638	3	−0.5	U						98Lu08	
^{170}Er(p,t)^{168}Er		−4785	5	−4778.1	1.1	1.4	U				Min		73Oo01	
^{170}Yb(p,t)^{168}Yb		−6861	6	−6844.574	0.026	2.7	B				Min		73Oo01	
^{170}Er(d,^{3}He)^{169}Ho		−3107	20	−3113	4	−0.3	U						76Su.A	
^{170}Er(d,t)^{169}Er		−1010	10	−999.5	1.1	1.1	U				Kop		69Tj01	
^{169}Tm(n,γ)^{170}Tm		6595.	2.5	6591.98	0.17	−1.2	U						66Sh03	
		6592.1	1.5			−0.1	U						70Or.A	
		6591.7	0.9			0.3	U				BNn		96Ho12	Z
		6591.95	0.17			0.1	1	99	66	^{169}Tm	Bdn		06Fi.A	
^{169}Tm(d,p)^{170}Tm		4420	20	4367.41	0.17	−2.6	U				CIT		66Ry01	
		4369	15			−0.1	U				Tal		66Sh03	
^{170}Yb(d,t)^{169}Yb		−2211	12	−2202.16	0.15	0.7	U				Kop		66Bu16	
^{170}Au(p)^{169}Pt		1473.8	15.	1472	12	−0.1	o				Jyp		02Ke.C	
		1471.7	12.				7				Jyp		04Ke06	
^{170}Au^m(p)^{169}Pt		1749.5	8.	1751	5	0.2	o				Jyp		02Ke.C	
		1745.4	10.			0.6	7				Arp		02Ma61	
		1753.5	6.			−0.4	7				Jyp		04Ke06	
^{170}Ho(β^-)^{170}Er		3870	50	3843	8	−0.5	U						78Tu04	
^{170}Ho^m(β^-)^{170}Er		3970	60	3994	9	0.4	U						78Tu04	
^{170}Tm(β^-)^{170}Yb		970	2	968.5	0.7	−0.7	–						54Po26	
		967.3	1.			1.2	–						69Va17	*
	ave.	967.8	0.9			0.8	1	67	67	^{170}Tm			average	
^{170}Lu(β^+)^{170}Yb		3467	20	3458	17	−0.5	2						60Dz02	
		3410	50			1.0	2						65Ha30	
*^{170}Lu−u	$M-A$=–57267(29) keV for mixture gs+m at 92.91 keV												Nub269	**
*^{170}Os(α)^{166}W	Used to recalibrate other results in same reference												GAu	**
*^{170}Ir(α)^{166}Re	E_α=6029.8(10,Z) 6027.2(5,Z) 6003(10) most probably to low levels in ^{166}Re												GAu	**
*^{170}Ir(α)^{166}Re	Correlated with ^{166}Re E_α=5533 keV												96Pa01	**
*^{170}Ir(α)^{166}Re	e$^-$ summed line; energy not trusted												FKo258	**
*^{170}Ir(α)^{166}Re	Correlated with E_α=6618,6547,6471,6431 keV from high-spin level ^{174}Au												04GoZZ	**
*^{170}Ir(α)^{166}Re	e$^-$ summed line; energy not trusted												FKo258	**
*^{170}Ir(α)^{166}Re	E_α=6121(10) 5951(10) to gs. 175, 6007(10) to 122, 6053(10) to 75 keV												07Ha45	**
*^{170}Ir(α)^{166}Re	E_α determined by gating on E(γ) - no e$^-$ coin summing												FKo258	*
*^{170}Tm(β^-)^{170}Yb	E_{β^-}=883(1) to 2$^+$ level at 84.25468 keV												Ens02b	**
C_{11} ^{13}C H_{12} N−^{171}Yb		164140	80	163997.710	0.014	−0.4	U				R04	4.0	64De15	
C_{10} H_7 O N_2−^{171}Yb		119640	270	119506.338	0.014	−0.1	U				R04	4.0	64De15	
^{171}Yb−^{129}Xe$_{1.326}$		62592.096	0.012	62592.096	0.012	−0.0	1	100	100	^{171}Yb	FS1	1.0	12Ra34	
^{171}Lu−u		−62132	41	−62081.8	1.8	1.2	U				GS2	1.0	05Li24	*
^{171}Hf−u		−59570	104	−59510	30	0.6	U				GS1	1.0	00Ra23	*
		−59508	31				2				GS2	1.0	05Li24	*
^{171}Ta−u		−55550	104	−55520	30	0.2	U				GS1	1.0	00Ra23	
		−55524	30				2				GS2	1.0	05Li24	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{171}W−u		−50650	110	−50550	30	0.9	U			GS1	1.0	00Ra23	
		−50549	30				2			GS2	1.0	05Li24	
^{171}Re−u		−44284	30				2			GS2	1.0	05Li24	
^{171}Os^{m}−u		−36796	30	−36760	18	1.2	1	34	34 ^{171}Os^{m}	GS2	1.0	05Li24	*
^{171}Yb ^{35}Cl$_2$−^{167}Er ^{37}Cl$_2$		10178.0	1.7	10175.9	0.3	−0.5	U			H27	2.5	74Ba90	
^{171}Yb ^{35}Cl−^{169}Tm ^{37}Cl		5055	3	5062.8	0.8	1.0	U			H23	2.5	70Wh01	
		5061.9	1.7			0.2	U			H27	2.5	74Ba90	
^{171}Dy−^{171}Yb		9804	12				2			JY2	1.0	24Ja12	
^{171}Ho−^{171}Yb		5043.6	7.9				2			JY2	1.0	24Ja12	
^{171}Yb−^{170}Yb		1220	60	1564.273	0.015	1.4	U			R04	4.0	64De15	
^{171}Os(α)^{167}W		5365.8	10.	5372.3	2.4	0.6	–			ORa		72To06	
		5365.8	10.			0.6	–					78Sc26	
		5393.4	15.			−1.4	–			ISa		79Ha10	
		5367.9	8.			0.5	–					95Hi02	*
		5374.0	9.			−0.2	–			Daa		96Pa01	
		5372.3	4.1			−0.0	–			Jya		23Zh03	*
	ave.	5372	3			0.2	1	65	39 ^{167}W			average	
^{171}Os^{m}(α)^{167}W		5560.4	4.1	5558.6	2.4	−0.4	1	35	21 ^{167}W	Jya		23Zh03	*
^{171}Ir(α)^{167}Re^{m}		5854.2	10.	5866	5	1.1	5			Bka		02Ro17	*
		5865.4	8.			0.0	5			Ara		11Ko.B	*
		5871.6	7.2			−0.8	5			Anv		13An10	
^{171}Ir^{m}(α)^{167}Re		6159.2	3.	6159.4	2.0	0.1	11			Ora		82De11	*
		6159	5			0.1	11					92Sc16	*
		6180	11			−1.9	11			Daa		96Pa01	*
		6159.2	8.			0.0	11			Anv		10An01	*
		6172.4	8.			−1.6	11			Ara		11Ko.B	*
		6153.8	4.1			1.4	11			Jya		23Zh03	*
^{171}Pt(α)^{167}Os		6608.1	4.	6607	3	−0.2	5			Ora		81De22	Z
		6606.8	5.			0.1	5			GSa		81Ho10	Z
		6604.8	11.			0.2	5			Jya		97Uu01	
		6600.6	15.			0.5	U			Anv		10An01	
^{171}Au^{m}(α)^{167}Ir^{m}		7163.9	6.	7164	4	0.1	–			Ara		97Da07	
		7162.9	8.			0.2	–			Jya		04Ke06	
	ave.	7164	5			0.2	1	69	43 ^{171}Au^{m}			average	
^{171}Hg(α)^{167}Pt		7667.7	15.				6			Jya		04Ke06	
^{171}Yb(p,t)^{169}Yb		−6599	5	−6591.80	0.15	1.4	U			Min		73Oo01	
^{170}Er(n,γ)^{171}Er		5681.5	0.5	5681.6	0.3	0.2	–					71Al01	
		5681.6	0.5			0.0	–			Bdn		06Fi.A	
^{170}Er(d,p)^{171}Er		3450	10	3457.1	0.3	0.7	U			Tal		68Ha10	
		3458	10			−0.1	U			Kop		69Tj01	
^{170}Er(n,γ)^{171}Er	ave.	5681.6	0.4	5681.6	0.3	0.2	1	98	71 ^{171}Er			average	
^{170}Er(α,t)^{171}Tm−^{168}Er()^{169}Tm		817.9	1.0	817.3	0.9	−0.6	1	80	66 ^{170}Er	McM		75Bu02	
^{170}Yb(n,γ)^{171}Yb		6614.3	0.6	6614.207	0.014	−0.2	U					72Wa10	Z
		6616.6	0.4			−6.0	C			Bdn		06Fi.A	
^{170}Yb(d,p)^{171}Yb		4390	12	4389.641	0.014	−0.0	U			Kop		66Bu16	
^{171}Yb(d,t)^{170}Yb		−359	12	−356.977	0.014	0.2	U			Kop		66Bu16	
^{170}Yb(α,t)^{171}Lu−^{174}Yb()^{175}Lu		−1156.2	2.0	−1156.8	1.7	−0.3	1	70	68 ^{171}Lu	McM		75Bu02	
^{171}Au(p)^{170}Pt		1452.6	17.	1448	10	−0.3	2			Arp		99Po09	
		1445.6	12.			0.2	2			Jyp		04Ke06	
^{171}Au^{m}(p)^{170}Pt		1702.1	6.	1702	4	0.1	–					97Da07	
		1704.1	6.			−0.3	–			Jyp		04Ke06	
	ave.	1703	4			−0.2	1	77	57 ^{171}Au^{m}			average	
^{171}Ho(β^-)^{171}Er		3200	600	3108	7	−0.2	U			LBL		90Ch34	
^{171}Er(β^-)^{171}Tm		1490	2	1492.2	1.1	1.1	1	29	29 ^{171}Er			61Ar15	*
^{171}Tm(β^-)^{171}Yb		96.5	1.0	97.5	0.4	1.0	–					57Sm73	
		97.60	0.38			−0.4	–					23Ju01	
	ave.	97.5	0.4			−0.0	1	99	99 ^{171}Tm			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{171}Lu(β^+)^{171}Yb	1479.3	3.	1478.0	1.7	−0.4	1	32	32 ^{171}Lu			77Bo32	*
^{171}Re(β^+)^{171}W	5670	200	5840	40	0.8	U			Got		87Ru05	
^{171}Os^m(IT)^{171}Os	186.32	0.13	186.32	0.13	0.0	1	100	52 ^{171}Os^m			Ens186	
^{171}Au^m(IT)^{171}Au	250	16	255	10	0.3	o					99Po09	*
*^{171}Lu−u	$M-A$=–57840(33) keV for mixture gs+m at 71.13 keV										Nub269	**
*^{171}Hf−u	$M-A$=–55480(100) keV for mixture gs+m at 21.93 keV										Nub269	**
*^{171}Hf−u	$M-A$=–55420(28) keV for mixture gs+m at 21.93 keV										Nub269	**
*^{171}Os(α)^{167}W	E_α=5241(8), 5166(8) to ground state and level at 79 keV										95Hi02	**
*^{171}Os(α)^{167}W	E_α=5248(4),5168(4),5115(4) to gs., 79.2(0.3), 134.2(0.3) keV levels										23Zh03	**
*^{171}Os^m(α)^{167}W	E_α=5306(4) to 13/2$^+$ level at 127.1(1.7) keV										23Zh03	**
*^{171}Ir(α)^{167}Re^m	Correlated with E_α=6412 of ^{175}Au										02Ro17	**
*^{171}Ir(α)^{167}Re^m	Correlated with E_α=6430(8) of ^{175}Au and 6556(8) of ^{179}Tl										11Ko.B	**
*^{171}Ir^m(α)^{167}Re	E_α=5925.2(3,Z) 5925(5) 5945(11) 5925(8) respectively, to (11/2$^-$) level at 92 keV										92Sc16	**
*	E_α=5920 correlated with ^{175}Au E_α=6438 keV										02Ro17	**
*^{171}Ir^m(α)^{167}Re	E_α=5938(8) to 92 level; correlated with E_α=6431(8) of ^{175}Au^m										11Ko.B	**
*	and 7194(8) of ^{179}Tl^m										11Ko.B	**
*^{171}Ir^m(α)^{167}Re	E_α=6011(5),5919(4) to gs. and 92 level										23Zh03	**
*^{171}Er(β^-)^{171}Tm	E_{β^-}=1065(2) to 7/2$^-$ level at 424.95 keV										Ens029	**
*^{171}Lu(β^+)^{171}Yb	E_{β^+}=362(3) to 7/2$^+$ level at 95.28 keV										Ens029	**
*^{171}Au^m(IT)^{171}Au	Redundant; use only their Q_p										GAu	**
C_{10} H_6 O_2 N−^{172}Yb	103560	60	103466.776	0.012	−0.4	U			R04	4.0	64De15	
^{172}Yb−^{132}Xe$_{1.303}$	61272.578	0.013	61272.582	0.010	0.3	1	61	60 ^{172}Yb	FS1	1.0	12Ra34	
^{172}Hf−u	−60555	30	−60550	26	0.2	2			GS2	1.0	05Li24	
^{172}Ta−u	−55105	30				2			GS2	1.0	05Li24	
^{172}W−u	−52770	110	−52710	30	0.6	U			GS1	1.0	00Ra23	
	−52708	30				2			GS2	1.0	05Li24	
^{172}Re−u	−44760	220	−44610	40	0.7	U			GS1	1.0	00Ra23	*
	−44647	53			0.6	1	52	52 ^{172}Re	GS2	1.0	05Li24	*
^{172}Yb ^{35}Cl$_2$−^{168}Er ^{37}Cl$_2$	9906.7	1.7	9908.98	0.30	0.5	U			H27	2.5	74Ba90	
^{172}Yb ^{35}Cl−^{170}Yb ^{37}Cl	4568.5	2.0	4569.55	0.07	0.2	U			H27	2.5	74Ba90	
^{172}Yb−^{170}Yb$_{1.012}$	2402.251	0.085	2402.209	0.012	−0.5	U			JY2	1.0	20Ne.1	
^{172}Yb−^{171}Yb	−50	230	55.145	0.016	0.1	U			R04	4.0	64De15	
^{172}Os(α)^{168}W	5226.8	10.	5226	4	−0.1	–					71Bo06	
	5227.8	10.			−0.2	–			Daa		96Pa01	
	5226.8	4.1			−0.2	–			Jya		23Zh03	*
ave.	5227	4			−0.2	1	98	62 ^{168}W			average	
^{172}Ir(α)^{168}Re	5990.6	10.				4					92Sc16	*
^{172}Ir(α)^{168}Re^m	5777.7	10.2	5777	5	−0.0	5			ISa		21Ha32	*
	5777.1	5.1			0.0	5			Jya		23Zh03	*
^{172}Ir^m(α)^{168}Re	6129.3	3.	6126.2	2.1	−1.0	4			Ora		82De11	*
	6161	20			−1.7	F			GSa		84Sc.A	*
	6129.1	5.			−0.6	4					92Sc16	*
	6123.0	12.			0.3	U			Daa		96Pa01	*
	6125.0	10.2			0.1	4			ISa		21Ha32	*
	6118.8	4.1			1.8	4			Jya		23Zh03	*
^{172}Pt(α)^{168}Os	6464.8	4.	6464	4	−0.2	1	97	75 ^{172}Pt	Ora		81De22	Z
	6474.8	15.			−0.7	U			Anv		09An20	
^{172}Au(α)^{168}Ir	6923.3	10.2				6			Jya		09Ha42	
^{172}Au^m(α)^{168}Ir^m	7023.6	10.	7034	6	1.0	6					93Se09	
	7042.1	9.			−0.9	6			Daa		96Pa01	
	7033.8	10.			0.0	6			Jya		09Ha42	*
^{172}Hg(α)^{168}Pt	7525.3	12.	7524	6	−0.1	3					99Se14	
	7536.5	16.			−0.8	o			Jya		04Ke06	
	7523.3	7.2			0.1	3			Jya		09Sa27	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{170}Er(t,p)^{172}Er	4034	4	4036	4	0.5	1	89	88 ^{172}Er			80Sh14	
^{172}Yb(p,t)^{170}Yb	−6161	5	−6152.361	0.011	1.7	U			Min		73Oo01	
^{171}Yb(n,γ)^{172}Yb	8020.3	0.7	8019.951	0.015	−0.5	U					71Al14	Z
	8020.1	0.5			−0.3	U					75Gr32	
	8019.67	0.35			0.8	U			ILn		85Ge02	Z
	8019.27	0.17			4.0	C			Bdn		06Fi.A	
^{171}Yb(d,p)^{172}Yb	5797	12	5795.384	0.015	−0.1	U			Kop		66Bu16	
	5789	5			1.3	U			Tal		66Sh14	
^{172}Yb(d,t)^{171}Yb	−1772	12	−1762.721	0.015	0.8	U			Kop		66Bu16	
^{171}Yb(^{3}He,d)^{172}Lu	−792	34	−774.7	2.0	0.5	U			Roc		76El11	
^{171}Yb(α,t)^{172}Lu−^{174}Yb()^{175}Lu	−791.9	2.0	−791.9	2.0	−0.0	1	100	100 ^{172}Lu	McM		75Bu02	
^{172}Er(β^-)^{172}Tm	888	5	891	5	0.6	1	82	70 ^{172}Tm			62Gu03	*
^{172}Tm(β^-)^{172}Yb	1870	10	1882	5	1.2	1	30	30 ^{172}Tm			66Ha15	
^{172}Hf(ε)^{172}Lu	350	50	334	25	−0.3	R					79To18	
^{172}Ta(β^+)^{172}Hf	4920	180	5070	40	0.8	U					73Ca10	*
^{172}W(β^+)^{172}Ta	3210	100	2230	40	−9.8	C					74Ca.A	*
*^{172}Re−u	$M-A$=–41640(200) keV for mixture gs+m at 110#50 keV										Nub269	**
*^{172}Re−u	$M-A$=–41533(28) keV for mixture gs+m at 110#50 keV										Nub269	**
*^{172}Os(α)^{168}W	E_α=5106(4),4909(5) to gs. and 2^+ level at 199.2(0.3) keV										23Zh03	**
*^{172}Ir(α)^{168}Re	E_α=5510(10) to 89.7+123.2+136.3 level										92Sc16	**
*^{172}Ir(α)^{168}Re^m	E_α=5510(10) to 136.3-keV level above ^{168}Re^m										21Ha32	**
*^{172}Ir(α)^{168}Re^m	E_α=5547(5),5537(5),5517(5),5505(5) to levels at 89.4(0.3),102.8(0.3)										23Zh03	**
*	123.0(0.2) 136.0(0.2) keV above ^{168}Re^m										23Zh03	**
*^{172}Ir^m(α)^{168}Re	E_α=5736 followed by XK(Re), 128 M2 and 161 M3 γ's										84Sc.A	**
*^{172}Ir^m(α)^{168}Re	F : first assigned to ^{173}Ir(α); seen in neither nuclide in reference										92Sc16	**
*^{172}Ir^m(α)^{168}Re	E_α=5828.2(3,Z) 5828(5) 5822(12) respectively, to (8^+) level at 162.1 keV										Ens108	**
*^{172}Ir^m(α)^{168}Re	E_α=5983(15), 5824(10) to ground state and 8^+ level at 162.1										21Ha32	**
*^{172}Ir^m(α)^{168}Re	E_α=5818(4) to 162.1 keV level										23Zh03	**
*^{172}Au^m(α)^{168}Ir^m	E_α=6870(10) 6800(10) to ground state and 70 keV level										09Ha42	**
*^{172}Er(β^-)^{172}Tm	E_{β^-}=278(5) to 1^+ level at 610.062 keV										Ens15c	**
*^{172}Ta(β^+)^{172}Hf	E_{β^+}=2480(180) to 4^- level at 1418.55 keV										Ens959	**
*^{172}W(β^+)^{172}Ta	E_{β^+}=1600(100) in coinc. with 459.2 keV γ from 586.3 level										Ens15c	**
C_{14} H_5−^{173}Yb	101030	70	100908.949	0.012	−0.4	U			R04	4.0	64De15	
C_{10} H_7 O_2 N−^{173}Yb	109810	60	109462.255	0.012	−1.4	U			R04	4.0	64De15	
^{173}Yb−^{129}Xe$_{1.341}$	65905.075	0.013	65905.080	0.010	0.3	1	60	56 ^{173}Yb	FS1	1.0	12Ra34	
^{173}Yb−^{132}Xe$_{1.311}$	63868.901	0.015	63868.895	0.010	−0.4	1	46	44 ^{173}Yb	FS1	1.0	12Ra34	
^{173}Hf−u	−59487	30	−59464	3	0.8	U			GS2	1.0	05Li24	
^{173}Ta−u	−56270	104	−56250	30	0.2	U			GS1	1.0	00Ra23	
	−56250	30				2			GS2	1.0	05Li24	
^{173}W−u	−52340	104	−52310	30	0.3	U			GS1	1.0	00Ra23	
	−52311	30				2			GS2	1.0	05Li24	
^{173}Re−u	−46910	110	−46760	30	1.4	U			GS1	1.0	00Ra23	
	−46757	30				2			GS2	1.0	05Li24	
^{173}Os−u	−40169	30	−40195	16	−0.9	1	29	29 ^{173}Os	GS2	1.0	05Li24	
^{173}Ir−u	−32440	100	−32492	11	−0.5	U			GS2	1.0	05Li24	*
^{173}Yb ^{35}Cl$_2$−^{169}Tm ^{37}Cl$_2$	9898.3	1.2	9897.6	0.8	−0.2	U			H27	2.5	74Ba90	
^{173}Yb ^{35}Cl−^{171}Yb ^{37}Cl	4827	4	4834.83	0.07	0.8	U			H23	2.5	70Wh01	
	4835.3	1.6			−0.1	U			H27	2.5	74Ba90	
^{173}Hf−^{173}Yb	2319.3	3.3				2			JY2	1.0	23Ja10	
^{173}Yb−^{172}Yb	1970	120	1829.553	0.014	−0.3	U			R04	4.0	64De15	
^{173}Os(α)^{169}W	5057.2	10.	5055	6	−0.3	–					71Bo06	
	5055.2	7.			−0.1	–			GSa		84Sc.A	
ave.	5056	6			−0.2	1	97	69 ^{169}W			average	
^{173}Ir(α)^{169}Re^m	5544.4	10.	5541	10	−0.3	1	90	79 ^{169}Re^m			92Sc16	
^{173}Ir^m(α)^{169}Re	5930.4	5.	5942.6	2.3	2.4	3					67Si02	*
	5947.1	4.			−1.1	3			Ora		82De11	*
	5937	10			0.6	3			GSa		84Sc.A	*
	5944.8	5.			−0.4	3					92Sc16	*
	5951.9	13.			−0.7	3			Daa		96Pa01	*
	5927.3	20.			0.8	U			Ara		01Ko.B	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{173}Ir$^{m}(\alpha)^{169}$Re		5945.7	5.1	5942.6	2.3	−0.6	3			ISa		21Ha32	*
^{173}Pt$(\alpha)^{169}$Os		6359.3	8.2	6383	4	0.5	B			ISa		79Ha10	Z
		6352.3	3.			0.6	B			Ora		81De22	Z
		6382.9	10.			−0.0	o			GSa		84Sc.A	
		6372.6	9.			0.2	3			Daa		96Pa01	
		6387.9	15.			−0.1	U			Anv		09An20	
		6382.9	4.1			−0.0	3			Jya		14ThZZ	*
		6383.0	8.2			−0.0	o			Jya		14Pe.A	
^{173}Au$(\alpha)^{169}$Ir		6830.2	6.	6842	3	2.0	4			Ara		99Po09	
		6847.6	8.			−0.6	4			Ara		01Ko44	
		6846.6	14.			−0.3	U			Ara		12Th13	
		6846.6	4.1			−1.0	4			Jya		14ThZZ	
^{173}Au$^{m}(\alpha)^{169}$Ir^{m}		6896.8	10.	6897.5	2.4	0.1	–			GSa		84Sc.A	
		6909.1	9.			−1.2	–			Daa		96Pa01	
		6891.6	4.			1.4	–			Ara		99Po09	
		6900.8	6.			−0.5	–			Ara		01Ko44	
		6899	15			−0.1	U			Jya		12Th13	
		6898.8	4.1			−0.3	–			Jya		14ThZZ	
	ave.	6897.2	2.4			0.1	1	99	57 ^{173}Au^{m}			average	
^{173}Hg$(\alpha)^{169}$Pt		7382.0	11.3	7378	4	−0.4	7					99Se14	
		7362.5	15.4			1.0	7			Jya		04Ke06	
		7378.9	5.			−0.2	7					12Od01	
^{173}Yb(p,t)^{171}Yb		−5913	5	−5905.255	0.015	1.5	U			Min		73Oo01	
^{172}Yb(n,γ)^{173}Yb		6367.3	0.4	6367.101	0.013	−0.5	U					71Al01	Z
		6367.2	0.6			−0.2	U			Bdn		06Fi.A	
^{173}Yb(γ,n)^{172}Yb		−6500	80	−6367.101	0.013	1.7	U			Phi		60Ge01	
^{172}Yb(d,p)^{173}Yb		4145	12	4142.534	0.013	−0.2	U			Kop		66Bu16	
^{173}Yb(d,t)^{172}Yb		−114	12	−109.871	0.013	0.3	U			Kop		66Bu16	
^{172}Yb(α,t)^{173}Lu−^{174}Yb()^{175}Lu		−595.6	1.0	−595.6	1.0	−0.0	1	100	100 ^{173}Lu	McM		75Bu02	
^{173}Tm$(\beta^{-})^{173}$Yb		1260	50	1295	4	0.7	U					63Ku22	
		1320	40			−0.6	U					63Or01	
^{173}Lu$(\varepsilon)^{173}$Yb		675	20	669.7	1.1	−0.3	U					73Ko13	
^{173}Ta$(\beta^{+})^{173}$Hf		3670	200	2994	28	−3.4	B					73Re03	
^{173}W$(\beta^{+})^{173}$Ta		4000	300	3670	40	−1.1	U					80Vi.A	
*^{173}Ir−u	$M-A$=–30113(70) keV for mixture gs+m at 216(4) keV											Nub269	**
*^{173}Ir$^{m}(\alpha)^{169}$Re	E_α=5660.0(5,Z) 5676.2(4,Z) 5666(10) 5674(5) 5681(13) respectively,											AHW	**
*	to (11/2^{-}) level at 136.40 keV											Ens155	**
*^{173}Ir$^{m}(\alpha)^{169}$Re	E_α=5675(5) to (11/2^{-}) level at 136.2 keV; Also tentatively assign											21Ha32	**
*^{173}Ir$^{m}(\alpha)^{169}$Re	E_α=5815(15) to ground state											21Ha32	**
*^{173}Pt$(\alpha)^{169}$Os	E_α=6201(4) to 34.84 keV; quoted E_α=6235(4) is e^{-} summed peak											FKo25b	**
*^{173}Pt$(\alpha)^{169}$Os	E_α=6201(8) to 34.84 keV											14Pe.A	**
C_{14} H_6−^{174}Yb		108308	38	108082.647	0.011	−1.5	U			R04	4.0	64De15	
^{174}Yb−^{129}Xe$_{1.349}$		67318.179	0.011	67318.168	0.009	−1.0	1	66	62 ^{174}Yb	FS1	1.0	12Ra34	
^{174}Yb−^{132}Xe$_{1.318}$		65191.113	0.017	65191.140	0.009	1.6	1	30	28 ^{174}Yb	FS1	1.0	12Ra34	
^{174}Ta−u		−55546	30				2			GS2	1.0	05Li24	
^{174}W−u		−53940	104	−53920	30	0.2	U			GS1	1.0	00Ra23	
		−53921	30				2			GS2	1.0	05Li24	
^{174}Re−u		−46930	104	−46890	30	0.4	U			GS1	1.0	00Ra23	
		−46885	30				2			GS2	1.0	05Li24	
^{174}Os−u		−42880	110	−42936	11	−0.5	U			GS1	1.0	00Ra23	
		−42919	30			−0.6	1	13	13 ^{174}Os	GS2	1.0	05Li24	
^{174}Ir−u		−33090	55	−33051	12	0.7	R			GS2	1.0	05Li24	*
^{174}Yb ^{35}Cl−^{172}Yb ^{37}Cl		5420	4	5431.02	0.07	1.1	U			H23	2.5	70Wh01	
		5430.3	1.1			0.3	U			H27	2.5	74Ba90	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{174}Yb−^{172}Yb$_{1.012}$	3244.248	0.026	3244.244	0.013	−0.1	1	24	14 ^{172}Yb	Hep	1.0	25Do.1	
^{174}Yb−^{173}Yb	700	50	651.334	0.013	−0.2	U			R04	4.0	64De15	
^{174}Hf(α)^{170}Yb	2558.9	30.	2494.7	1.8	−2.1	U					61Ma05	
^{174}Os(α)^{170}W	4872.2	10.	4871	10	−0.2	1	90	78 ^{170}W			71Bo06	
^{174}Ir(α)^{170}Re^{m}	5624.1	10.	5616	4	−0.8	4					92Sc16	*
	5619.4	5.1			−0.7	o			Jya		19Mo.B	*
	5614.3	4.1			0.3	4			Jya		23Zh03	*
^{174}Ir^{m}(α)^{170}Re	5817.6	6.	5817	3	−0.1	–					67Si02	*
	5816.4	5.			0.1	–					92Sc16	*
	5819.2	13.3			−0.2	o			Jya		19Mo.B	*
	5816.4	5.1			0.1	–			Jya		23Zh03	*
ave.	5817	3			0.1	1	99	83 ^{170}Re			average	
^{174}Pt(α)^{170}Os	6176.3	10.	6183	3	0.7	2			ISa		79Ha10	Z
	6185.7	5.			−0.5	2			Ora		81De22	Z
	6182.4	5.1			0.2	2			Ara		04Go38	
^{174}Au(α)^{170}Ir	6701.3	5.	6650#	50#	−9.5	F			Ara		04GoZZ	*
^{174}Au(α)^{170}Ir^{m}	6683.9	20.5	6690	9	0.3	8			GSa		83Sc24	*
	6698.3	10.			−0.8	F			Daa		96Pa01	*
	6692.1	10.2			−0.2	8			Bka		02Ro17	*
	6701.3	5.			−2.2	F			Ara		04GoZZ	*
^{174}Au^{m}(α)^{170}Ir	6782.2	10.2	6779	4	−0.4	7			GSa		84Sc.A	*
	6793.5	13.			−1.1	7			Daa		96Pa01	*
	6775.4	5.1			0.6	7			Ara		04GoZZ	*
^{174}Hg(α)^{170}Pt	7235.6	11.	7233	6	−0.2	2			Jya		97Uu01	
	7232.5	8.2			0.1	2					99Se14	
	7231.5	14.3			0.1	2			Bka		01Ro.B	
^{174}Yb(p,t)^{172}Yb	−5359	5	−5349.909	0.012	1.8	U			Min		73Oo01	
^{173}Yb(n,γ)^{174}Yb	7464.63	0.06	7464.604	0.012	−0.4	U			MMn		82Is05	Z
	7464.58	0.35			0.1	U			ILn		87Ge01	Z
	7465.5	0.4			−2.2	U			Bdn		06Fi.A	
^{173}Yb(d,p)^{174}Yb	5239	12	5240.038	0.012	0.1	U			Kop		66Bu16	
	5229	5			2.2	U			Tal		66Sh14	
^{174}Yb(d,t)^{173}Yb	−1218	12	−1207.374	0.012	0.9	U			Kop		66Bu16	
^{173}Yb(α,t)^{174}Lu−^{174}Yb()^{175}Lu	−202.1	1.0	−202.1	1.0	−0.0	1	100	100 ^{174}Lu	McM		75Bu02	
^{174}Tm(β^-)^{174}Yb	3080	100	3080	40	−0.0	2					64Ka16	*
	3080	50			−0.0	2					67Gu12	*
^{174}Lu(β^+)^{174}Yb	1402	5	1373.7	1.1	−5.7	B					68Kl08	
^{174}Lu(ε)^{174}Yb	1370	7			0.5	U					68Li01	*
^{174}Ta(β^+)^{174}Hf	3845	80	4104	28	3.2	B					71Ch26	*
*^{174}Ir−u	$M-A$=–30761(36) keV for mixture gs+m at 126(15) keV										Nub269	**
*^{174}Ir(α)^{170}Re^{m}	E_α=5275(10) to (3$^+$) level at 224.7 keV above ^{170}Re^{m}										Ens02b	**
*^{174}Ir(α)^{170}Re^{m}	E_α=5271(5) to 224.2(0.3) keV level above ^{170}Re^{m}										19Mo.B	**
*^{174}Ir(α)^{170}Re^{m}	E_α=5292(8),5268(5) to 193.1(0.4),224.0(0.6) keV levels										23Zh03	**
*^{174}Ir^{m}(α)^{170}Re	E_α=5478(6) to (7$^+$) level at 210.32 keV										Ens02b	**
*^{174}Ir^{m}(α)^{170}Re	E_α=5478(5) 5316(10) to (7$^+$) at 210.32 and 370.1 level										Ens02b	**
*^{174}Ir^{m}(α)^{170}Re	E_α=5480(13) to 210.1(0.1) keV level above ^{170}Re										19Mo.B	**
*^{174}Ir^{m}(α)^{170}Re	E_α=5499(8),5476(5) to 191(1),210.1(0.3) keV levels										23Zh03	**
*^{174}Au(α)^{170}Ir	overlaps with stronger E_α from ^{174}Au^{m}; not used										FKo258	**
*^{174}Au(α)^{170}Ir^{m}	associated with ^{174}Au because of the short T$_1$/2=120(20) ms										FKo258	**
*^{174}Au(α)^{170}Ir^{m}	overlaps with stronger E_α from ^{174}Au^{m}; not used										FKo258	**
*^{174}Au(α)^{170}Ir^{m}	Correlated with E_α=5817 keV ^{170}Ir^{m} (low-spin)										02Ro17	**
*	and only this with ^{178}Tl α's (no decays via ^{174}Au^{m})										02Ro17	**
*^{174}Au(α)^{170}Ir^{m}	overlaps with stronger E_α from ^{174}Au^{m}; not used										FKo258	**
*^{174}Au^{m}(α)^{170}Ir	also E_α=6546(10) to 73 keV level										84Sc.A	**
*^{174}Au^{m}(α)^{170}Ir	also E_α=6544(10) to 73 keV level										96Pa01	**
*^{174}Au^{m}(α)^{170}Ir	E_α=6471(5) to 153.2(0.5) keV and 6431(5) to 191.2(0.5) keV;										04GoZZ	**
*^{174}Au^{m}(α)^{170}Ir	others E_α=6547(5), 6618(15). All αs correlated										04GoZZ	**
*^{174}Au^{m}(α)^{170}Ir	with E_α=6088 keV (affected by e$^-$ summing) in ^{170}Ir decay										04GoZZ	**
*^{174}Tm(β^-)^{174}Yb	E_{β^-}=1200(100) 1200(50) respectively, to 5$^-$ level at 1884.674 keV, and other $E_{\beta-}$										Ens998	**
*^{174}Lu(ε)^{174}Yb	No K capture to 2$^-$ level at 1318.361 keV, Q <1380; and L capture										Ens998	**
*	of ^{174}Lu^{m} at 170.83 to ^{174}Yb^{m} at 1518.148 keV → Q_{gs}>1357 keV										Nub269	**
*^{174}Ta(β^+)^{174}Hf	E_{β^+}=2525(80) to 4$^+$ level at 297.38 keV										Ens04a	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{175}Lu ^{37}Cl−^{142}Nd ^{35}Cl$_2$		61249.5	2.5	61246.7	1.2	−0.5	U			H31	2.5	77So02	
C_{14} H_7−^{175}Lu		114121	37	113998.6	0.5	−0.8	U			R04	4.0	64De15	
C_{13} ^{13}C H_6−^{175}Lu		109763	36	109528.4	0.5	−1.6	U			R04	4.0	64De15	
^{175}Ta−u		−56350	120	−56260	30	0.7	U			GS1	1.0	00Ra23	
		−56263	30				2			GS2	1.0	05Li24	
^{175}W−u		−53290	104	−53280	30	0.1	U			GS1	1.0	00Ra23	
		−53283	30				2			GS2	1.0	05Li24	
^{175}Re−u		−48630	104	−48620	30	0.1	U			GS1	1.0	00Ra23	
		−48619	30				2			GS2	1.0	05Li24	
^{175}Os−u		−43120	110	−43050	11	0.6	U			GS1	1.0	00Ra23	
		−43024	30			−0.9	1	13	13 ^{175}Os	GS2	1.0	05Li24	
^{175}Ir−u		−34353	1288	−35848	12	−0.5	U				2.5	91Br17	
		−35828	30			−0.7	1	15	15 ^{175}Ir	GS2	1.0	05Li24	
^{175}Lu ^{35}Cl−^{173}Yb ^{37}Cl		5503	4	5510.5	0.5	0.8	U			H23	2.5	70Wh01	
		5507.3	1.4			0.9	U			H27	2.5	74Ba90	
^{175}Lu O−C_{16}		−64316.3	4.5	−64308.8	0.5	1.1	U			TG1	1.5	11Ke03	
^{175}Ir(α)^{171}Re		5709.0	5.	5433	30	−55.2	B					67Si02	*
		5709.2	5.			−55.3	B					92Sc16	*
^{175}Pt(α)^{171}Os		6166	10	6178.0	2.2	1.2	o			Ora		66Si08	*
		6176	10			0.2	o			Ora		73Ga08	*
		6179.8	4.4			−0.4	–			ISa		79Ha10	*
		6175.3	3.1			0.9	–			Ora		82De11	*
		6163	4			3.7	B			Jya		14Pe02	*
		6182.4	6.1			−0.7	–			Lza		24Hu14	*
		6181.4	8.2			−0.4	–			Ara		25Ko.B	*
	ave.	6177.9	2.3			0.0	1	100	73 ^{175}Pt			average	
^{175}Au(α)^{171}Ir		6562.3	15.	6583.4	2.7	1.4	U			Bka		02Ro17	*
		6580.7	8.2			0.3	6			Ara		11Ko.B	*
		6583.7	4.			−0.1	6			Anv		13An10	
		6583.7	4.			−0.1	6			Jya		17Ba46	
^{175}Au^m(α)^{171}Ir^m		6590.9	10.	6583.5	2.5	−0.7	10			Ora		75Ca06	
		6775.8	10.			−19.2	F			GSa		84Sc.A	*
		6588.8	9.			−0.6	10			Daa		96Pa01	
		6579.6	6.			0.6	10			Ara		01Ko44	
		6582.7	5.			0.1	10			Anv		10An01	
		6581.7	8.2			0.2	10			Ara		11Ko.B	*
		6583.7	4.			−0.1	10			Jya		17Ba46	
^{175}Hg(α)^{171}Pt		7020.7	20.	7072	5	2.5	o			GSa		83Sc24	
		7039.2	20.			1.7	U			GSa		84Sc.A	
		7071.0	24.			0.1	o			Daa		96Pa01	
		7058.7	11.			1.2	6			Jya		97Uu01	
		7075	5			−0.5	6			Daa		09Od01	
		7082.1	20.			−0.5	U			Anv		10An01	
^{174}Yb(n,γ)^{175}Yb		5822.35	0.07				2			MMn		82Is05	Z
		5822.5	0.4	5822.35	0.07	−0.4	U			Bdn		06Fi.A	
^{174}Yb(d,p)^{175}Yb		3595	12	3597.78	0.07	0.2	U			Kop		66Bu16	
^{174}Yb(α,t)^{175}Lu		−14303	10	−14303.2	0.4	−0.0	U			McM		75Bu02	
^{175}Lu(γ,n)^{174}Lu		−7880	80	−7666.7	1.0	2.7	U			Phi		60Ge01	
^{175}Lu(d,t)^{174}Lu		−1400	10	−1409.5	1.0	−0.9	U			Tal		70Jo08	
^{174}Hf(n,γ)^{175}Hf		6708.4	0.5	6708.5	0.4	0.2	–					71Al01	Z
		6708.8	0.6			−0.5	–			Bdn		06Fi.A	
	ave.	6708.6	0.4			−0.2	1	99	88 ^{175}Hf			average	
^{175}Tm(β^-)^{175}Yb		2385	50				3					66Wi04	*
^{175}Yb(β^-)^{175}Lu		466	3	470.7	0.4	1.6	U					55De18	
		468	5			0.5	U					55Mi90	
		471	3			−0.1	U					56Co13	
		467	3			1.2	U					62Ba32	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{175}Hf(ε)^{175}Lu	628	9	684.8	1.8	6.3	B					68Ja11	*
	650	20			1.7	U					69Jo16	*
	630	3			18.3	B					88Si22	*
*^{175}Ir(α)^{171}Re	E_α=5392.8(5,Z) to 189.8 level										95Hi02	**
*^{175}Ir(α)^{171}Re	E_α=5393(5) to 189.8 level										95Hi02	**
*^{175}Pt(α)^{171}Os	E_α=5950(10) to 76.70(0.07) level										66Si08	**
*^{175}Pt(α)^{171}Os	E_α=5960(10) to 76.70(0.07) level										73Ga08	**
*^{175}Pt(α)^{171}Os	E_α=6038(10), 5964(5) to ground state, 76.70(0.07) level										79Ha10	**
*^{175}Pt(α)^{171}Os	E_α=5959(3) to 76.70(0.07) level										82De11	**
*^{175}Pt(α)^{171}Os	E_α=6021(4),5948(4),5819(4),5814(4) to ground state,76.70(0.07),										14Pe02	**
*^{175}Pt(α)^{171}Os	207.57(0.07),211.0(0.4) level										14Pe02	**
*^{175}Pt(α)^{171}Os	E_α=5966(6) to 76.70(0.07) keV level										24Hu14	**
*^{175}Pt(α)^{171}Os	E_α=5965(8) to 76.70(0.07) level										25Ko.B	**
*^{175}Au(α)^{171}Ir	Analysis by AHW of data in Fig. 3 of reference										02Ro17	**
*^{175}Au(α)^{171}Ir	Correlated with E_α=6556(8) of ^{179}Tl and 5728(8) of ^{171}Ir										11Ko.B	**
*^{175}Au^m(α)^{171}Ir^m	E_α=6435(10) and 6470(20) to 190.0 and 152.7 levels										84Sc.A	**
*^{175}Au^m(α)^{171}Ir^m	F : reassigned to ^{174}Au										01Ko.B	**
*^{175}Au^m(α)^{171}Ir^m	Correlated with E_α=7194(8) of ^{179}Tl^m and 5958(8) of ^{171}Ir^m										11Ko.B	**
*	different method and different detectors as compared to 01Ko44										11Ko.B	**
*^{175}Tm(β^-)^{175}Yb	E_{β^-}=1870(50) to 1/2$^-$ level at 514.866 keV										Ens04a	**
*^{175}Hf(ε)^{175}Lu	pK=0.712(0.008) 0.740(0.015) 0.714(0.002) respectively,										AHW	**
*	to 7/2$^+$ level at 432.74 keV, and other capture ratios, recalculated										Ens04a	**
C_{14} H_8 −^{176}Yb	119980	46	120025.552	0.014	0.2	U			R04	4.0	64De15	
C_{13} H_6 N−^{176}Yb	107190	110	107449.492	0.014	0.6	U			R04	4.0	64De15	
^{176}Yb−^{129}Xe$_{1.364}$	72453.619	0.016	72453.613	0.012	−0.4	1	61	59 ^{176}Yb	FS1	1.0	12Ra34	
^{176}Yb−^{132}Xe$_{1.333}$	70335.958	0.027	70335.972	0.013	0.5	1	22	22 ^{176}Yb	FS1	1.0	12Ra34	
^{176}Lu ^{37}Cl−^{143}Nd ^{35}Cl$_2$	61067.2	1.4	61070.2	1.2	0.9	1	12	11 ^{143}Nd	H31	2.5	77So02	
C_{14} H_8 −^{176}Lu	119962	49	119909.1	0.4	−0.3	U			R04	4.0	64De15	
^{176}Lu O−C_{16}	−62394.1	7.6	−62394.2	0.4	−0.0	U			TG1	1.5	11Ke03	
^{176}Hf O−C_{16}	−63668.5	9.8	−63675.2	0.4	−0.5	U			TG1	1.5	11Ke03	
^{176}Ta−u	−55143	33				2			GS2	1.0	05Li24	
^{176}W−u	−54420	104	−54370	30	0.5	U			GS1	1.0	00Ra23	
	−54366	30				2			GS2	1.0	05Li24	
^{176}Re−u	−48380	110	−48380	30	0.0	U			GS1	1.0	00Ra23	
	−48377	30				2			GS2	1.0	05Li24	
^{176}Os−u	−45150	110	−45230	12	−0.7	U			GS1	1.0	00Ra23	
	−45194	30			−1.2	1	15	15 ^{176}Os	GS2	1.0	05Li24	
^{176}Ir−u	−36328	30	−36354	9	−0.9	1	9	9 ^{176}Ir	GS2	1.0	05Li24	
^{176}Yb ^{35}Cl$_2$−^{172}Yb ^{37}Cl$_2$	12088.9	2.4	12088.30	0.14	−0.1	U			H27	2.5	74Ba90	
^{176}Yb−^{172}Yb$_{1.023}$	7651.151	0.026	7651.153	0.015	0.1	1	32	20 ^{176}Yb	Hep	1.0	25Do.1	
^{176}Yb ^{35}Cl−^{174}Yb ^{37}Cl	6652	3	6657.29	0.07	0.7	U			H23	2.5	70Wh01	
	6656.3	1.4			0.3	U			H27	2.5	74Ba90	
^{176}Hf ^{35}Cl−^{174}Hf ^{37}Cl	4106	16	4311.6	1.9	5.1	B			H24	2.5	73Ba40	
	4314.21	0.86			−1.2	1	76	76 ^{174}Hf	H37	2.5	77Sh12	
^{176}Lu−^{176}Yb	116.91	0.83	116.4	0.4	−0.6	1	28	28 ^{176}Lu	MS1	1.0	23Qu02	
^{176}Yb−^{176}Hf	1164.67	0.46	1164.5	0.4	−0.4	1	73	73 ^{176}Hf	MS1	1.0	23Qu02	
^{176}Lu−^{176}Hf	1279.96	0.72	1280.9	0.4	1.3	1	32	20 ^{176}Lu	MS1	1.0	23Qu02	
^{176}Lu−^{175}Lu	1980	60	1914.51	0.16	−0.3	U			R04	4.0	64De15	
^{176}Yb−^{174}Yb	4000	50	3707.159	0.015	−1.5	U			R04	4.0	64De15	
^{176}Ir(α)^{172}Re	5237.3	8.	5270	40	0.6	1	49	48 ^{172}Re			67Si02	
^{176}Pt(α)^{172}Os	5890.1	5.	5883.9	1.8	−1.2	–			ISa		79Ha10	Z
	5881.4	4.			0.6	–			Bka		82Bo04	Z
	5887.3	3.			−1.1	–			Ora		82De11	Z
	5874.8	8.			1.1	–			Daa		96Pa01	
	5880.9	5.1			0.6	–			Ara		04GoZZ	
	5881.9	7.			0.3	–			ISa		17An16	
	5876.8	8.2			0.9	–			Jya		21Gi08	
	5883.9	6.1			−0.0	–			Lza		24Hu14	
ave.	5884.1	1.8			−0.1	1	100	64 ^{172}Os			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{176}Au(α)^{172}Ir	6574.2	10.	6432	4	−14.2	B			Ora		75Ca06	*
	6541.5	10.			−10.9	C			GSa		84Sc.A	*
	6433.4	7.			−0.2	5			Anv		14An10	*
	6431.8	5.1			0.1	5			ISa		21Ha32	*
^{176}Au^m(α)^{172}Ir^m	6436.6	10.	6433	3	−0.3	5			Ora		75Ca06	*
	6428.4	10.			0.5	5			GSa		84Sc.A	*
	6433.4	6.			0.0	5			Ara		01Ko44	*
	6434.4	7.			−0.1	5			Anv		14An10	*
	6433.5	5.1			−0.0	5			ISa		21Ha32	*
^{176}Hg(α)^{172}Pt	6907.2	20.	6898	6	−0.5	o			GSa		83Sc24	
	6924.7	10.			−2.7	C			GSa		84Sc.A	
	6907.3	20.			−0.5	U			Daa		96Pa01	
	6897.0	6.			0.1	–			Ara		99Po09	
	6917.5	15.			−1.3	–			Anv		09An20	
ave.	6900	6			−0.3	1	94	70 ^{176}Hg			average	
^{176}Yb(p,α)^{173}Tm	7628.8	4.4				2			NDm		78Ta10	
^{176}Yb(p,t)^{174}Yb	−4216	5	−4207.643	0.014	1.7	U			Min		73Oo01	
^{176}Hf(p,t)^{174}Hf	−6397	5	−6392.6	1.7	0.9	1	12	12 ^{174}Hf	Min		73Oo01	
^{176}Yb(d,t)^{175}Yb	−621	12	−609.86	0.07	0.9	U			Kop		66Bu16	
^{175}Lu(n,γ)^{176}Lu	6293.2	1.2	6287.96	0.15	−4.4	B					70Wa20	
	6287.96	0.15			0.0	1	100	99 ^{175}Lu	ILn		91Kl02	Z
	6289.78	0.24			−7.6	C			Bdn		06Fi.A	
^{175}Lu(d,p)^{176}Lu	4070	8	4063.40	0.15	−0.8	U			Tal		67St14	
^{176}Lu(d,t)^{175}Lu	−25	15	−30.73	0.15	−0.4	U			Tal		71Mi01	
^{176}Hf(d,t)^{175}Hf	−1925	8	−1908.7	1.8	2.0	U			Tal		73Za08	
^{176}Tl(p)^{175}Hg	1265.2	18.				7			Jyp		04Ke06	
^{176}Tm(β^-)^{176}Yb	4120	100				2					67Gu11	*
^{176}Lu(β^+)^{176}Yb	108.9	0.8	108.5	0.4	−0.6	1	26	26 ^{176}Lu			23Qu02	
^{176}Lu(β^-)^{176}Hf	1162	25	1193.2	0.4	1.2	U					69Pr11	*
	1194.1	1.0			−0.9	–					73Va11	*
	1193.1	0.9			0.1	–					23Qu02	*
ave.	1193.5	0.7			−0.6	1	33	20 ^{176}Lu			average	
^{176}Ta(β^+)^{176}Hf	3100	90	3210	30	1.2	U					71Be10	*
*^{176}Au(α)^{172}Ir	E_α=6260(10) coinc. with E(γ)=168.4(0.5) keV										75Ca06	**
*^{176}Au(α)^{172}Ir	E_α=6228(10) to 168.4(0.5) γ										84Sc.A	**
*^{176}Au(α)^{172}Ir	E_α=6260 correlated with ^{172}Ir E_α=5510 keV										02Ro17	**
*^{176}Au(α)^{172}Ir	and E_α=6157(20),6138(15),6054(20),5798(20) to 126.3, 151.5, 236.6,500.0										14An10	**
*^{176}Au(α)^{172}Ir	E_α=6260(5) to (2$^-$,3$^-$) level at 25 keV; Also E_α=6052(15),										21Ha32	**
*^{176}Au(α)^{172}Ir	6138(10) to levels at 236.0(0.2), 151.5(0.3) keV										21Ha32	**
*^{176}Au^m(α)^{172}Ir^m	E_α=6286 correlated with ^{172}Ir^m E_α=5828 keV										02Ro17	**
*^{176}Au^m(α)^{172}Ir^m	E_α=6119+E(γ)=175.1 was misassigned to ^{177}Au										01Ko44	**
*^{176}Au^m(α)^{172}Ir^m	E_α=6115(6) coinc. with 175.1 γ of reference										84Sc.A	**
*^{176}Au^m(α)^{172}Ir^m	E_α=6287(7), 6117(7), 6082(7) to 0, 175.2, 211.6 above ^{172}Ir^m										14An10	**
*^{176}Au^m(α)^{172}Ir^m	E_α=6280(10), 6114(5), 6080(5) to 0, 175.2, 211.6 above ^{172}Ir^m										21Ha32	**
*^{176}Tm(β^-)^{176}Yb	E_{β^-}=2000(100), 1150(100) to (3$^+$,4$^+$) level at 2053.34, (3$^+$,4$^+$,5$^+$) 3052.2										Ens062	**
*^{176}Lu(β^-)^{176}Hf	E_{β^-}=565(25) to 6$^+$ level at 596.82 keV										Ens062	**
*^{176}Lu(β^-)^{176}Hf	Q_{β^-}=1317(1) from ^{176}Lu^m at 122.845 keV										Nub269	**
*^{176}Lu(β^-)^{176}Hf	average Q_{β^-}=596.1(0.9),195.7(2.5) to 6$^+$ level at 597,										23Qu02	**
*	8$^+$ level at 998.0 keV										23Qu02	**
*^{176}Ta(β^+)^{176}Hf	KLM/β^+=119(50) to 2$^-$ level at 1247.70 keV, 1$^+$ level at 2994 keV										Ens062	**
^{177}Ta−u	−55559	30	−55518	3	1.4	U			GS2	1.0	05Li24	
^{177}W−u	−53420	110	−53360	30	0.6	U			GS1	1.0	00Ra23	
	−53357	30				2			GS2	1.0	05Li24	
^{177}Re−u	−49620	104	−49670	30	−0.5	U			GS1	1.0	00Ra23	
	−49672	30				2			GS2	1.0	05Li24	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{177}Os−u		−45020	104	−45042	16	−0.2	U			GS1	1.0	00Ra23	
		−45012	30			−1.0	R			GS2	1.0	05Li24	
^{177}Ir−u		−38810	110	−38699	21	1.0	U			GS1	1.0	00Ra23	
		−38699	30			0.0	2			GS2	1.0	05Li24	
^{177}Pt−u		−31545	30	−31526	16	0.6	1	29	29 ^{177}Pt	GS2	1.0	05Li24	
^{177}Hf O−C_{16}		−61845.2	7.2	−61855.5	0.9	−1.0	U			TG1	1.5	11Ke03	
^{177}Ir(α)^{173}Re		5127.1	10.	5080	30	−0.9	F					67Si02	*
^{177}Pt(α)^{173}Os		5654.6	6.	5650	4	−0.8	–			ISa		79Ha10	Z
		5640.5	3.1			3.0	B			Bka		82Bo04	Z
		5648.9	5.			0.2	–			Ara		04GoZZ	
		5647.8	10.2			0.2	–			ISa		21Ha32	
	ave.	5651	4			−0.2	1	99	55 ^{177}Pt			average	
^{177}Au(α)^{173}Ir		6292.5	10.	6297.9	2.9	0.5	–			Daa		75Ca06	
		6292.5	20.			0.3	U			GSa		84Sc.A	
		6296.5	10.			0.1	–			Daa		96Pa01	
		6298.6	6.			−0.1	–			Ara		01Ko44	
		6303.7	7.			−0.8	o			Anv		09An14	
		6301.6	7.2			−0.5	–			Anv		18Cu04	
		6298.6	5.1			−0.1	–			ISa		21Ha32	
		6295.8	9.2			0.2	–			Jya		25Sp01	*
	ave.	6298.2	2.9			−0.1	1	99	89 ^{173}Ir			average	
^{177}Au^{m}(α)^{173}Ir		6489.6	15.3	6479.8	3.0	−0.6	U			Jya		25Sp01	*
^{177}Au^{m}(α)^{173}Ir^{m}		6251.5	10.	6263.3	2.8	1.2	2			Ora		75Ca06	
		6260.8	10.			0.3	2			GSa		84Sc.A	*
		6259.7	9.			0.4	2			Daa		96Pa01	*
		6263.8	6.			−0.1	2			Ara		01Ko44	
		6265.8	7.			−0.4	2			Anv		09An14	
		6266.9	5.1			−0.7	2			ISa		21Ha32	*
		6261.8	9.2			0.2	2			Jya		25Sp01	*
^{177}Hg(α)^{173}Pt		6732.4	8.	6740	60	0.1	4			ISa		79Ha10	
		6747.8	10.			−0.2	4					91Ko.A	
		6729.4	9.2			0.1	4			Daa		96Pa01	
		6734.5	15.			0.0	4			Anv		09An20	
^{177}Tl(α)^{173}Au		7067.0	7.				3			Ara		99Po09	
^{177}Tl^{m}(α)^{173}Au^{m}		7660.4	13.	7658	9	−0.2	–			Ara		99Po09	
		7645.1	13.			1.0	–			Jya		04Ke06	
	ave.	7653	9			0.6	1	84	43 ^{173}Au^{m}			average	
^{177}Hf(p,t)^{175}Hf		−6071	5	−6060.4	1.9	2.1	1	15	12 ^{175}Hf	Min		73Oo01	
^{176}Yb(n,γ)^{177}Yb		5565.1	1.0	5566.40	0.22	1.3	U					72Al19	Z
		5566.40	0.22				2			Bdn		06Fi.A	
^{176}Yb(d,p)^{177}Yb		3340	16	3341.83	0.22	0.1	U			Tal		63Ve09	
		3337	12			0.4	U			Kop		66Bu16	
^{176}Yb(α,t)^{177}Lu−^{174}Yb()^{175}Lu		674.1	1.0	671.41	0.22	−2.7	U			McM		75Bu02	
^{176}Lu(n,γ)^{177}Lu		7071.2	0.4	7072.88	0.16	4.2	B					71Ma45	Z
		7073.1	0.4			−0.5	–					72Mi16	Z
		7072.85	0.17			0.2	–			Bdn		06Fi.A	
^{176}Lu(d,p)^{177}Lu		4843	10	4848.32	0.16	0.5	U			Tal		71Mi01	
^{176}Lu(n,γ)^{177}Lu	ave.	7072.89	0.16	7072.88	0.16	−0.0	1	99	96 ^{177}Lu			average	
^{176}Hf(n,γ)^{177}Hf		6385.8	0.8	6376.3	0.8	−11.8	C			Bdn		06Fi.A	
^{177}Hf(γ,n)^{176}Hf		−6400	30	−6376.3	0.8	0.8	U			Phi		60Ge01	
^{176}Hf(d,p)^{177}Hf		4150	7	4151.8	0.8	0.3	U			Tal		68Ri07	
^{177}Hf(d,t)^{176}Hf		−127	11	−119.1	0.8	0.7	U			Tal		72Za04	
^{177}Tl(p)^{176}Hg		1162.6	20.	1157	19	−0.3	o			Arp		99Po09	*
^{177}Tl^{m}(p)^{176}Hg		1969.2	10.	1964	7	−0.5	–			Arp		99Po09	
		1965.2	12.			−0.1	–			Jyp		04Ke06	
	ave.	1968	8			−0.5	1	89	59 ^{177}Tl^{m}			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{177}Yb(β^-)^{177}Lu		1400	20	1398.0	0.5	−0.1	U					64Jo03	
^{177}Lu(β^-)^{177}Hf		497	2	496.6	0.8	−0.2	–					55Ma12	
		496.4	1.0			0.2	–					62El02	*
	ave.	496.5	0.9			0.1	1	74	70 ^{177}Hf			average	
^{177}Ta(β^+)^{177}Hf		1166	3				2					61We11	
^{177}Au^m(IT)^{177}Au		210	30	181.9	0.4	−0.9	o					01Ko44	*
		181.9	0.4			0.1	1	100	63 ^{177}Au^m			20Ve03	
^{177}Tl^m(IT)^{177}Tl		807	18				2					99Po09	
*^{177}Ir(α)^{173}Re	F : final state uncertain: possibly to 5/2$^-$ level at 214.7 keV											95Hi02	**
*^{177}Au(α)^{173}Ir	E_α=6156(9),5998(9) to gs. and 3/2$^+$ level at 156.1 keV											25Sp01	**
*^{177}Au^m(α)^{173}Ir	E_α=5928(15) to (9/2$^-$) level at 424.4 keV											25Sp01	**
*^{177}Au^m(α)^{173}Ir^m	Followed by a 175.1(0.5) γ											84Sc.A	**
*	γ associated with E_α=6116 keV from ^{176}Au											01Ko44	**
*	yet E_α=6118 correlated with E_α=5672 of ^{173}Ir^m											02Ro17	**
*^{177}Au^m(α)^{173}Ir^m	E_α correlated with ^{173}Ir E_α=5681(13); also with ^{181}Tl E_α=6180 keV											96Pa01	**
*	evaluator doubts about correctness of latter remark											AHW	**
*^{177}Au^m(α)^{173}Ir^m	Also E_α=5932(12) to 9/2$^-$ level at 424.4 keV											21Ha32	**
*^{177}Au^m(α)^{173}Ir^m	Also E_α=5928(15) to (9/2$^-$) level at 424.4 keV											25Sp01	**
*^{177}Tl(p)^{176}Hg	Replaced by ^{177}Tl^m(IT)											AHW	**
*^{177}Lu(β^-)^{177}Hf	E_{β^-}=384(2), 175(1) to 112.95, 321.32 levels											Ens035	**
*^{177}Au^m(IT)^{177}Au	Authors say 157.9+x, x estimated by evaluator											AHW	**
*	x is better known from ^{181}Tl^m(IT)^{181}Tl combined with Q_α											09An14	**
^{178}Yb−^{85}Rb$_{2.094}$		131396.8	9.4	131382	7	−1.6	1	57	57 ^{178}Yb	MA8	1.0	19Hu15	
^{178}Yb−^{140}Ce ^{19}F$_2$		44429	34	44415	7	−0.4	U			TR1	1.0	26Br.A	
^{178}W−u		−54152	30	−54115	16	1.2	U			GS2	1.0	05Li24	
^{178}Re−u		−48800	110	−49002	22	−1.8	U			GS1	1.0	00Ra23	
		−49011	30			0.3	1	53	53 ^{178}Re	GS2	1.0	05Li24	
^{178}Os−u		−46790	104	−46747	15	0.4	U			GS1	1.0	00Ra23	
		−46710	30			−1.2	1	24	24 ^{178}Os	GS2	1.0	05Li24	
^{178}Ir−u		−38950	110	−38922	20	0.3	U			GS1	1.0	00Ra23	
		−38888	30			−1.1	1	45	45 ^{178}Ir	GS2	1.0	05Li24	
^{178}Pt−u		−34783	1181	−34350	11	0.1	U				2.5	91Br17	
		−34300	110			−0.5	U			GS1	1.0	00Ra23	
		−34333	30			−0.6	1	13	13 ^{178}Pt	GS2	1.0	05Li24	
^{178}Au−^{133}Cs$_{1.338}$		102562	11				2			MA8	1.0	20Cu04	
^{178}Au^n−^{133}Cs$_{1.338}$		102764	11	102761	10	−0.2	1	87	87 ^{178}Au^n	MA8	1.0	20Cu04	
^{178}Hf ^{35}Cl−^{176}Hf ^{37}Cl		5236	5	5247.9	0.9	1.0	U			H23	2.5	70Wh01	
		5239.5	1.3			2.6	U			H27	2.5	74Ba90	
^{178}Re−^{178}Hf		7300	32	7290	22	−0.3	1	47	47 ^{178}Re	RI1	1.0	26Ya.1	
^{178}Hf O−C$_{16}$		−61364.8	7.9	−61377.4	0.9	−1.1	U			TG1	1.5	11Ke03	
^{178}Pt−^{175}Ir		−472	1052	1498	16	0.7	U				2.5	91Br17	
^{178}Pt(α)^{174}Os		5583.3	5.	5573.0	2.1	−2.0	–			ISa		79Ha10	Z
		5569.9	3.			1.0	–			Bka		82Bo04	Z
		5568.4	13.			0.4	U			Lvn		94Wa23	
		5572.4	4.			0.1	–			Ara		00Ko16	*
		5572.4	8.2			0.1	–			Lza		24Hu14	
	ave.	5573.1	2.1			−0.1	1	99	75 ^{174}Os			average	
^{178}Au(α)^{174}Ir		6056.4	10.	6059	4	0.3	3					68Si01	*
		6058.4	4.1			0.2	o			Jya		19Mo.B	*
		6058.4	5.1			0.1	3			ISa		20Cu04	*
		6064.5	10.2			−0.5	3			Jya		21Gi08	
^{178}Au^n(α)^{174}Ir^m		6117.7	20.	6119	5	0.0	–			GSa		86Ke03	
		6116.4	4.0			0.5	o			Jya		19Mo.B	*
		6119.0	7.0			−0.1	–			ISa		20Cu04	*
		6116.5	8.2			0.2	–			Jya		21Gi08	*
	ave.	6118	5			0.1	1	97	84 ^{174}Ir^m			average	
^{178}Hg(α)^{174}Pt		6578.1	6.	6577.1	2.8	−0.2	3			ISa		79Ha10	
		6576.1	9.			0.1	3			Daa		96Pa01	
		6577.1	4.			−0.0	3			Ara		00Ko48	
		6578.1	8.			−0.1	3			Anv		09An14	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{178}Hg(α)^{174}Pt		6575.0	8.2	6577.1	2.8	0.2	3			Lza		24Hu14	
^{178}Tl(α)^{174}Au		7017.0	5.	7020	10	0.6	o			Bka		02Ro17	*
		7020.0	10.				9			Bka		13Li49	*
^{178}Pb(α)^{174}Hg		7790.4	14.	7789	13	−0.1	3			Bka		01Ro.B	
		7785.2	30.7			0.1	3			Jya		16Ba60	
^{178}Pt(p,α)^{175}Ir		4420	980	6260	15	1.9	U					91Br17	
^{176}Yb(t,p)^{178}Yb		3865	10	3847	7	−1.8	1	43	43 ^{178}Yb	Phi		82Zu02	
^{176}Lu(t,p)^{178}Lu^{m}		4482	5	4492.6	2.9	2.1	1	34	34 ^{178}Lu^{m}	LAl		81Gi01	
^{178}Hf(p,t)^{176}Hf		−5531	5	−5520.5	0.8	2.1	U			Min		73Oo01	
^{177}Hf(n,γ)^{178}Hf		7625	1	7625.93	0.18	0.9	U					69Fa01	
		7624.4	1.5			1.0	U					77St10	
		7626.2	0.3			−0.9	–			ILn		86Ha22	Z
		7625.80	0.22			0.6	–			Bdn		06Fi.A	
^{178}Hf(d,t)^{177}Hf		−1364	9	−1368.70	0.18	−0.5	U			Tal		68Ri07	
^{177}Hf(n,γ)^{178}Hf	ave.	7625.94	0.18	7625.93	0.18	−0.1	1	99	71 ^{178}Hf			average	
^{178}Yb(β^-)^{178}Lu		641	30	661	7	0.7	U					73Or03	*
^{178}Lu^{m}(IT)^{178}Lu		120	3	123.8	2.6	1.3	1	76	66 ^{178}Lu^{m}	McM		93Bu02	
^{178}Lu(β^-)^{178}Hf		2046	50	2097.3	2.0	1.0	U					73Or03	*
		2117	30			−0.7	U					75Ka15	*
^{178}Ta^{m}(β^+)^{178}Hf		1937	15				2					61Ga05	*
^{178}W(ε)^{178}Ta^{m}		91.3	2.				3					67Ni02	
^{178}Re(β^+)^{178}W		4660	180	4762	25	0.6	U					70Go20	*
*^{178}Pt(α)^{174}Os	Also E_α=5289(8) keV to 2$^+$ 158.601 level (not used)											GAu	**
*^{178}Au(α)^{174}Ir	others E_α=5840(10),5812(4),5758(8) to 90.3,115.9,											19Mo.B	**
*^{178}Au(α)^{174}Ir	175.1 keV levels above ^{174}Ir											19Mo.B	**
*^{178}Au(α)^{174}Ir	others E_α=5840(10),5811(10),5750(10) to 90.0,115.7,											20Cu04	**
*^{178}Au(α)^{174}Ir	174.8 keV levels above ^{174}Ir											20Cu04	**
*^{178}Au^{n}(α)^{174}Ir^{m}	others E_α=5978(8), 5923(7), 5573(12), 5516(7) to											19Mo.B	**
*^{178}Au^{n}(α)^{174}Ir^{m}	0, 56.8, 421.4, 472.1 levels above ^{174}Ir^{m}											19Mo.B	**
*^{178}Au^{n}(α)^{174}Ir^{m}	E_α=5977(10), 5925(7), 5571(7), 5521(7) to											20Cu04	**
*^{178}Au^{n}(α)^{174}Ir^{m}	0, 56.8, 421.4, 472.1 levels above ^{174}Ir^{m}											20Cu04	**
*^{178}Au^{n}(α)^{174}Ir^{m}	E_α=5976(8), 5926(8) to 0, 56.8 keV level above ^{174}Ir^{m}											21Gi08	**
*^{178}Tl(α)^{174}Au	And a stronger E_α=6704; both correlated with ^{174}Au E_α=6538 keV											02Ro17	**
*^{178}Tl(α)^{174}Au	Also 6693(10) and 6595(10) to 173.0, 273.0 levels											13Li49	**
*^{178}Yb(β^-)^{178}Lu	E_{β^-}=250(30) to 1$^+$ level at 390.8 keV											Ens097	**
*^{178}Lu(β^-)^{178}Hf	E_{β^-}=2000(50) to ground state and 50% to 2$^+$ level at 93.18 keV											Ens097	**
*^{178}Lu(β^-)^{178}Hf	E_{β^-}=2050(50) to ground state and 50% to 2$^+$ level at 93.18 keV and											Ens097	**
*	E_{β^-}=770(30) from ^{178}Lu^{m} at 123.8(2.6) to 8$^-$ level 1479.025 keV											Nub269	**
*^{178}Ta^{m}(β^+)^{178}Hf	E_{β^+}=890(10) to ground state and 2$^+$ level at 93.18 keV, ratio 2.7 to 1											Ens097	**
*^{178}Re(β^+)^{178}W	E_{β^+}=3300(180) to 4$^+$ level at 342.74 keV											Ens097	**
^{179}Yb−^{160}Dy ^{19}F		26162	29				2			TR1	1.0	26Br.A	
^{179}Lu−^{85}Rb$_{2.106}$		132997	21	133103	5	5.1	C			MA8	1.0	13Ro.A	
C$_{14}$ H$_{11}$−^{179}Hf		140260.3	1.8	140250.0	0.9	−1.4	U			M23	4.0	79Ha32	
^{179}W−u		−52964	76	−52921	16	0.6	U			GS2	1.0	05Li24	*
^{179}Re−u		−50010	30	−50010	26	−0.0	1	78	78 ^{179}Re	GS2	1.0	05Li24	
^{179}Os−u		−46220	104	−46184	17	0.3	U			GS1	1.0	00Ra23	
		−46176	30			−0.3	1	31	31 ^{179}Os	GS2	1.0	05Li24	
^{179}Ir−u		−40910	104	−40882	11	0.3	U			GS1	1.0	00Ra23	
		−40852	30			−1.0	1	12	12 ^{179}Ir	GS2	1.0	05Li24	
^{179}Pt−u		−34710	110	−34641	8	0.6	U			GS1	1.0	00Ra23	
		−34625	30			−0.5	U			GS2	1.0	05Li24	
^{179}Au−u		−26811	31	−26824	11	−0.4	1	13	13 ^{179}Au	GS2	1.0	05Li24	
^{179}Hg−^{208}Pb$_{.861}$		1900	34	1885	18	−0.5	1	27	27 ^{179}Hg	MA6	1.0	01Sc41	
^{179}Hf ^{35}Cl−^{177}Hf ^{37}Cl		5539	3	5545.66	0.22	0.9	U			H23	2.5	70Wh01	
		5544.4	0.7			0.7	U			H27	2.5	74Ba90	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{179}Hf O−C_{16}		−59261.8	6.5	−59260.0	0.9	0.2	U			TG1	1.5	11Ke03	
^{179}Pt(α)^{175}Os		5370	10	5407	7	3.7	F					66Si08	*
		5416	10			−0.9	o			ISa		79Ha10	*
		5382	3			8.5	F			Bka		82Bo04	*
		5409.0	7.2			−0.2	1	94	87 ^{175}Os	ISa		21Ha32	*
^{179}Au(α)^{175}Ir		5981.8	5.	5982	3	−0.0	–					68Si01	
		5986.9	15.			−0.4	U			Jya		04Ra28	*
		5981.9	4.1			−0.1	–			ISa		21Ha32	*
	ave.	5982	3			−0.1	1	99	85 ^{175}Ir			average	
^{179}Hg(α)^{175}Pt		6431.0	5.	6428.9	2.6	−0.4	–			ISa		79Ha10	Z
		6418.7	9.			1.1	–			Daa		96Pa01	
		6430.0	4.			−0.3	–			Ara		02Ko09	*
		6426.8	6.1			0.3	–			Lza		24Hu14	
		6430	9			−0.1	–			Jya		25Sp01	*
	ave.	6428.8	2.6			0.0	1	100	73 ^{179}Hg			average	
^{179}Tl(α)^{175}Au		6710.2	20.	6709.1	2.6	−0.1	U			GSa		83Sc24	
		6718.4	18.			−0.5	U			Daa		96Pa01	
		6719.4	10.			−1.0	7			Ara		98To14	
		6706.1	8.			0.4	7			Ara		11Ko.B	
		6710.2	4.			−0.3	7			Anv		13An10	
		6707.2	4.1			0.5	7			Jya		17Ba46	
^{179}Tl^m(α)^{175}Au^m		7364.5	20.	7369.7	3.0	0.3	U			GSa		83Sc24	
		7366.0	20.			0.2	U			Daa		96Pa01	
		7378.1	10.			−0.8	o			Ara		98To14	
		7372.0	5.1			−0.4	9			Anv		10An01	
		7358.7	8.2			1.3	9			Ara		11Ko.B	
		7371.0	4.1			−0.3	9			Jya		17Ba46	
^{179}Pb(α)^{175}Hg		7598.3	20.	7596	5	−0.1	7			Anv		10An01	*
		7596	5			0.0	7			Jya		17Ba46	*
^{179}Hf(p,t)^{177}Hf		−5249	5	−5243.12	0.19	1.2	U			Min		73Oo01	
^{179}Hf(t,α)^{178}Lu−^{178}Hf()^{177}Lu		−72	2	−73.7	1.9	−0.8	1	89	89 ^{178}Lu	McM		93Bu02	
^{178}Hf(n,γ)^{179}Hf		6099.02	0.10	6098.99	0.08	−0.3	–			ILn		89Ri03	Z
		6098.95	0.12			0.3	–			Bdn		06Fi.A	
^{179}Hf(γ,n)^{178}Hf		−6000	70	−6098.99	0.08	−1.4	U			Phi		60Ge01	
^{178}Hf(d,p)^{179}Hf		3877	14	3874.42	0.08	−0.2	U			Tal		63Ve09	
^{178}Hf(n,γ)^{179}Hf	ave.	6098.99	0.08	6098.99	0.08	−0.1	1	100	71 ^{179}Hf			average	
^{179}Lu(β^-)^{179}Hf		1350	50	1404	5	1.1	U					61Ku10	
		1380	70			0.3	U					63St06	
^{179}Ta(ε)^{179}Hf		129	16	105.6	0.4	−1.5	U					61Jo15	*
		105.61	0.41			−0.1	1	99	97 ^{179}Ta			01Hi06	
^{179}Re(β^+)^{179}W		2710	50	2711	27	0.0	1	29	22 ^{179}Re			75Me20	*
*^{179}W−u	$M-A$=–49225(29) keV for mixture gs+m at 221.91 keV											Nub269	**
*^{179}Pt(α)^{175}Os	F : part of double line (with ^{180}Pt)											AHW	**
*^{179}Pt(α)^{175}Os	E_α=5150(10) 5195(10) 5161(3) respectively, to 1/2$^-$ level at 102.3 keV, recalibrated											Ens092	**
*^{179}Pt(α)^{175}Os	E_α=5188(7), 5116(15) to 1/2$^-$, 3/2$^-$ levels at 102.3, 175.6 keV											21Ha32	**
*^{179}Au(α)^{175}Ir	E_α=5853(15), 5810(15) to ground state, 49 keV level											04Ra28	**
*^{179}Au(α)^{175}Ir	Also E_α=5600(10), 5705(15), 5718(10) to 260.3, 146.0(0.7),											21Ha32	**
*	131.9(0.4) kev level above 26.1 level											21Ha32	**
*^{179}Hg(α)^{175}Pt	no γs in coincidence with E_α=8286 → ground state to ground state decay											02Ko09	**
*^{179}Hg(α)^{175}Pt	Also E_α=6156(10) keV to 9/2$^-$ level at 131.3(0.5) keV											25Sp01	**
*^{179}Pb(α)^{175}Hg	E_α=7350(20) to 80 keV level											10An01	**
*^{179}Pb(α)^{175}Hg	E_α=7348(5) to 80 keV level											17Ba46	**
*^{179}Ta(ε)^{179}Hf	As corrected in reference											76He03	**
*^{179}Re(β^+)^{179}W	E_{β^+}=950(50) to 3/2$^+$ level at 720.18 and 5/2$^+$ at 773.65 keV											Ens092	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{180}Yb−^{156}Gd ^{12}C$_2$		29763	27				2				TR1	1.0	26Br.A	
C$_{14}$ H$_{12}$−^{180}Hf		147356.6	4.8	147341.2	0.9	−0.8	U				M23	4.0	79Ha32	
^{180}W−u		−53299	30	−53287.0	0.9	0.4	U				GS2	1.0	05Li24	
^{180}Re−u		−49209	30	−49209	22	0.0	–				GS2	1.0	05Li24	
	ave.	−49208	23			−0.0	1	91	91	^{180}Re			average	
^{180}Re−^{85}Rb$_{2.118}$		137618	75	137621	22	0.0	1	9	9	^{180}Re	RI1	1.0	26Ya.1	
^{180}Os−u		−47650	104	−47618	17	0.3	U				GS1	1.0	00Ra23	
		−47626	30			0.3	1	32	32	^{180}Os	GS2	1.0	05Li24	
^{180}Ir−u		−40800	104	−40771	23	0.3	U				GS1	1.0	00Ra23	
		−40765	30			−0.2	2				GS2	1.0	05Li24	
^{180}Pt−u		−36900	104	−36963	11	−0.6	U				GS1	1.0	00Ra23	
		−36918	30			−1.5	1	13	13	^{180}Pt	GS2	1.0	05Li24	
^{180}Au−u		−27496	30	−27511	5	−0.5	U				GS2	1.0	05Li24	
^{180}Au−^{133}Cs$_{1.353}$		100411.5	5.3	100412	5	0.2	1	93	93	^{180}Au	MA8	1.0	17Ma29	
^{180}Hg−^{208}Pb$_{.865}$		−1569	22	−1543	13	1.2	1	37	37	^{180}Hg	MA6	1.0	01Sc41	
^{180}Hf ^{35}Cl$_2$−^{176}Hf ^{37}Cl$_2$		11036.1	3.0	11049.3	0.9	1.8	U				H27	2.5	74Ba90	
^{180}Hf ^{35}Cl−^{178}Hf ^{37}Cl		5797	3	5801.35	0.19	0.6	U				H23	2.5	70Wh01	
		5798.4	0.7			1.7	U				H27	2.5	74Ba90	
^{180}W−^{180}Hf		153.73	0.30	153.79	0.30	0.2	1	98	73	^{180}W	SH1	1.0	12Dr01	
^{180}Ta^m−^{180}Ta		82.43	0.60	83.0	0.4	0.9	1	41	41	^{180}Ta	JY2	1.0	22Ne10	
^{180}Hf−^{179}Hf		730.8	4.7	733.84	0.16	0.3	U				M24	2.5	79Ha32	
^{180}Hf O−C$_{16}$		−58524.5	6.5	−58526.2	0.9	−0.2	U				TG1	1.5	11Ke03	
^{180}W(α)^{176}Hf		2516.4	1.6	2514.6	0.9	−1.1	1	29	27	^{180}W			04Co26	
^{180}Pt(α)^{176}Os		5257.1	10.	5276	5	1.9	F						66Si08	*
		5279	3			−0.9	F				Bka		82Bo04	*
		5277.5	5.1			−0.2	1	97	85	^{176}Os			20Cu02	*
^{180}Au(α)^{176}Ir		5814.4	10.2	5812	7	−0.2	F				GSa		86Ke03	*
		5857	30			−1.5	F				Lvn		93Wa03	*
		5813.4	7.2			−0.1	1	93	90	^{176}Ir	ISa		20Ha24	*
^{180}Hg(α)^{176}Pt		6258.3	5.	6258.6	2.1	0.0	–				ISa		79Ha10	Z
		6259.5	5.			−0.2	–				Lvn		93Wa03	*
		6258.3	4.			0.1	–				Ara		00Ko48	
		6259.3	5.			−0.2	–				Anv		03An27	
		6258.3	7.			0.0	–				ISa		17An16	
		6259.3	6.1			−0.1	–				Lza		24Hu14	
	ave.	6258.8	2.1			−0.1	1	99	65	^{176}Pt			average	
^{180}Tl(α)^{176}Au		6709.4	10.	6704	6	−0.5	o				Ara		98To14	*
		6707.3	10.2			−0.3	6				Ara		13KoZR	
		6703.0	7.			0.2	6				Isa		17An16	*
^{180}Pb(α)^{176}Hg		7394.6	40.	7419	5	0.6	U				ORa		96To08	
		7415.1	15.			0.2	2				Ara		99To11	
		7419.2	10.			−0.1	2				Anv		09An20	
		7419.2	7.			−0.1	2				Jya		10Ra12	
^{180}Hf(p,t)^{178}Hf		−5011	5	−5004.94	0.17	1.2	U				Min		73Oo01	
^{180}Hf(t,α)^{179}Lu−^{178}Hf()^{177}Lu		−669	5	−669	5	−0.0	1	100	100	^{179}Lu	McM		92Bu12	
^{179}Hf(n,γ)^{180}Hf		7387.3	0.4	7387.75	0.15	1.1	–						74Bu22	Z
		7387.8	0.6			−0.1	–						90Bo52	Z
		7387.85	0.17			−0.6	–				Bdn		06Fi.A	
^{180}Hf(γ,n)^{179}Hf		−7470	110	−7387.75	0.15	0.7	U				Phi		60Ge01	
^{179}Hf(d,p)^{180}Hf		5167	7	5163.19	0.15	−0.5	U				Tal		72Za04	
^{180}Hf(d,t)^{179}Hf		−1112	4	−1130.52	0.15	−4.6	B				Tal		68Ri07	
^{179}Hf(n,γ)^{180}Hf	ave.	7387.77	0.15	7387.75	0.15	−0.1	1	99	75	^{180}Hf			average	
^{180}W(d,t)^{179}W		−2155	15	−2155	15	−0.0	1	94	94	^{179}W	Kop		72Ca01	
^{180}Lu(β^-)^{180}Hf		3148	100	3100	70	−0.4	2						71Gu02	*
		3058	100			0.5	2						71Sw01	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{180}Ta^{m}(IT)^{180}Ta		78.0	0.5	77.3	0.4	−1.4	1	52	51 ^{180}Ta			02We01	*
^{180}Ta(β^-)^{180}W		705	15	700.5	1.7	−0.3	U					51Br87	
		712	15			−0.8	U					62Ga07	
^{180}Re(β^+)^{180}W		3830	60	3799	20	−0.5	R					67Go22	*
		3790	40			0.2	R					67Ho12	*
*^{180}Pt(α)^{176}Os	F : part of double line (with ^{179}Pt); E_α=5140(10) keV											AHW	**
*^{180}Pt(α)^{176}Os	F : part of double line (with ^{179}Pt)											AHW	**
*^{180}Pt(α)^{176}Os	Also E_α=5028(7) keV to 2^+ level at 135.1 keV											20Ha24	**
*^{180}Au(α)^{176}Ir	E_α=5685(10) to 40(30) level, former treatment not supported by 20Ha24											93Wa03	**
*^{180}Au(α)^{176}Ir	E_α=5647(10,Z) to 80(30) level, not supported by 20Ha24											93Wa03	**
*^{180}Au(α)^{176}Ir	E_α=5639(7) to 46.0(0.8) keV level											20Ha24	**
*^{180}Hg(α)^{176}Pt	E_α=6120 5862 5689(5) to ground state, 2^+ level at 264.0, 0^+ at 443 keV											Ens062	**
*^{180}Hg(α)^{176}Pt	supersedes reference											93Wa.A	**
*^{180}Tl(α)^{176}Au	Highest E_α; not necessarily ground state to ground state											98To14	**
*^{180}Tl(α)^{176}Au	E_α=6553(7),6354(7),6245(7),6199(7) to levels 0,204..8,317.1,361.7											17An16	**
*^{180}Lu(β^-)^{180}Hf	E_{β^-}=1540(100) 1450(100) respectively, to 4^+ level at 1607.67 keV											Ens156	**
*^{180}Ta^{m}(IT)^{180}Ta	Deduced from two separate level energy differences in ^{181}Ta(p,d)											FGk24c	*W
*^{180}Ta^{m}(IT)^{180}Ta	as 78.7(0.5) keV and 77.4(0.5) keV in 02We01											FGk24c	*W
*^{180}Re(β^+)^{180}W	E_{β^+}=1800(60) 1760(40) respectively, to 2^- level 1006.381 keV											Ens156	**
^{181}Lu−u		−48092	135				2			GS3	1.0	13Sh30	
^{181}Ta O−^{202}Tl$_{.975}$		−29891.4	2.5	−29892.3	1.9	−0.4	1	56	31 ^{202}Tl	MA8	1.0	17We09	
^{181}Re−u		−49915	30	−49939	13	−0.8	R			GS2	1.0	05Li24	
^{181}Os−u		−46670	110	−46753	27	−0.8	U			GS1	1.0	00Ra23	*
		−46756	34			0.1	1	64	64 ^{181}Os	GS2	1.0	05Li24	*
^{181}Ir−u		−42330	104	−42378	9	−0.5	U			GS1	1.0	00Ra23	
		−42372	30			−0.2	U			GS2	1.0	05Li24	
^{181}Pt−u		−36880	104	−36910	15	−0.3	U			GS1	1.0	00Ra23	
		−36900	30			−0.3	–			GS2	1.0	05Li24	
	ave.	−36890	21			−1.0	1	48	48 ^{181}Pt			average	
^{181}Au−u		−30030	110	−29921	21	1.0	U			GS1	1.0	00Ra23	
		−29920	30			−0.0	R			GS2	1.0	05Li24	
^{181}Hg−^{208}Pb$_{.870}$		−1929	40	−1863	17	1.7	1	17	17 ^{181}Hg	MA6	1.0	01Sc41	
^{181}Tl−^{133}Cs$_{1.361}$		114936	11	114940	10	0.3	1	79	79 ^{181}Tl	MA8	1.0	08We02	
^{181}Ta ^{35}Cl−^{179}Hf ^{37}Cl		5128.6	2.1	5122.8	1.8	−1.1	1	12	9 ^{181}Ta	H35	2.5	80Sh06	
^{181}Ta ^{17}O ^{35}Cl−^{180}Ta^{m} O ^{37}Cl		7572	21	7617.29	0.21	0.9	U			H35	2.5	80Sh06	
^{181}Yb−^{181}Ta		7450	27				2			TR1	1.0	26Br.A	
^{181}Pt(α)^{177}Os		5133.7	20.	5150	5	0.8	U					66Si08	
		5150.1	5.				2			ORa		95Bi01	
^{181}Au(α)^{177}Ir		5750.1	5.	5751.4	2.9	0.2	3					68Si01	Z
		5751.9	5.			−0.1	3			ISa		79Ha10	Z
		5735	4			4.1	F			IRa		92Sa03	*
		5752	5			−0.1	3			ORa		95Bi01	*
^{181}Hg(α)^{177}Pt		6288	5	6284	4	−0.7	–			ISa		79Ha10	*
		6283	10			0.1	–			GSa		86Ke03	*
		6269.3	13.			1.2	–			Daa		96Pa01	*
	ave.	6285	4			−0.2	1	99	83 ^{181}Hg			average	
^{181}Tl(α)^{177}Au		6319.9	20.	6319	3	−0.0	U					92Bo.D	
		6326.1	10.			−0.7	–			Ara		98To14	
		6320.9	7.			−0.3	–			Anv		09An14	
		6323.0	7.2			−0.5	–			Anv		18Cu04	
	ave.	6323	5			−0.8	1	58	53 ^{177}Au			average	
^{181}Tl(α)^{177}Au^{m}		6120.3	20.	6137	3	0.8	U			GSa		84Sc.A	*
		6132.6	10.			0.5	–			Ara		98To14	*
		6133.1	6.4			0.6	–			Anv		09An14	*
	ave.	6133	5			0.8	1	41	37 ^{177}Au^{m}			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{181}Pb(α)^{177}Hg		7374.3	10.	7251	8	−12.3	F			GSa		86Ke03	*
		7203.5	15.			3.2	F			ORa		89To01	*
		7224.9	20.			1.3	F			Ara		96To01	*
		7250.7	10.			0.0	5			Ara		05Ca.A	*
		7252.0	15.			−0.1	5			Anv		09An20	*
^{181}Ta(p,t)^{179}Ta		−5738	5	−5742.5	1.7	−0.9	1	12	9 ^{181}Ta	Min		73Oo01	
^{180}Hf(n,γ)^{181}Hf		5695.2	0.6	5694.80	0.07	−0.7	U					71Al22	
		5694.80	0.07				2			Prn		02Bo41	
		5695.58	0.20			−3.9	C			Bdn		06Fi.A	
^{180}Hf(d,p)^{181}Hf		3440	25	3470.23	0.07	1.2	U			Sac		66Ga06	
		3475	10			−0.5	U			Tal		68Ri07	
^{181}Ta(γ,n)^{180}Ta		−7717	25	−7574.8	0.4	5.7	B			Phi		60Ge01	*
		−7852	26			10.7	B			Phi		60Ge01	
		−7580	5			1.0	U			McM		79Ba06	
		−7579	2			2.1	–			McM		81Co17	
^{181}Ta(d,t)^{180}Ta		−1317.7	1.8	−1317.6	0.4	0.1	–			NDm		79Ta.B	
^{181}Ta(γ,n)^{180}Ta	ave.	−7576.8	1.3	−7574.8	0.4	1.4	1	9	9 ^{180}Ta			average	
^{180}Ta^{m}(n,γ)^{181}Ta		7651.8	0.5	7652.12	0.18	0.6	–			MMn		81Co17	Z
		7652.13	0.20			−0.0	–			ILn		84Fo.A	Z
	ave.	7652.08	0.19			0.2	1	98	98 ^{180}Ta^{m}			average	
^{180}W(n,γ)^{181}W		6669.02	0.16				2			Bdn		15Hu07	
^{180}W(d,p)^{181}W		4468	15	4444.45	0.16	−1.6	U			Kop		72Ca01	
^{181}Hg(εp)^{180}Pt		6150	200	6484	18	1.7	F					72Ho19	*
^{181}Hf(β^-)^{181}Ta		1023	8	1036.2	1.7	1.7	U					52Fa14	*
		1020	5			3.2	B					53Ba81	*
^{181}W(ε)^{181}Ta		184	12	205.2	1.7	1.8	U					66Ra03	
		190	6			2.5	U					83Se17	
^{181}Os(β^+)^{181}Re		2990	200	2967	28	−0.1	U					67Go25	*
^{181}Au^{p}(IT)^{181}Au		1.8	0.1				4					20Se13	
^{181}Hg^{m}(IT)^{181}Hg		212	50				2					09An17	*
*^{181}Os−u	$M-A$=–43450(100) keV for mixture gs+m at 49.20 keV											Nub269	**
*^{181}Os−u	$M-A$=–43529(28) keV for mixture gs+m at 49.20 keV											Nub269	**
*^{181}Au(α)^{177}Ir	E_α=5609(8), 5462(4) to ground state and (3/2$^-$) level at 148.00 keV											Ens035	**
*	F : all lines in ^{181}Au and ^{183}Au shifted by 16−20keV											GAu	**
*^{181}Au(α)^{177}Ir	E_α=5626(5) to gs; favored 5479(5) to (3/2$^-$) level at 148.00 keV											Ens035	**
*^{181}Hg(α)^{177}Pt	E_α=6147.0(10,Z), 6005.0(5,Z) to ground state and 1/2$^-$ isomer at 147.5 keV											Nub269	**
*^{181}Hg(α)^{177}Pt	E_α=6136.6(10,Z), 6005.6(10,Z) to ground state and 1/2$^-$ isomer at 147.5 keV											Nub269	**
*^{181}Hg(α)^{177}Pt	E_α=5986(13) to 1/2$^-$ isomer at 147.5 keV											Nub269	**
*^{181}Tl(α)^{177}Au^{m}	E_α=6566(20) Q_α=6956.2 from ^{181}Tl^{m} at 835.9 to 241.5 above ^{177}Au^{m}											Nub269	**
*^{181}Tl(α)^{177}Au^{m}	E_α=6578(10) Q_α=6968.5 from ^{181}Tl^{m} at 835.9 to 241.5 above ^{177}Au^{m}											Nub269	**
*^{181}Tl(α)^{177}Au^{m}	E_α=6818(15), 6578(7) from ^{181}Tl^{m} to ^{177}Au^{m} and level 241.5 above ^{177}Au^{m}											09An14	**
*^{181}Pb(α)^{177}Hg	F : This α-line not found in same reaction											96To01	**
*^{181}Pb(α)^{177}Hg	E_α=7044(5) is broad (Figure 3); ce summing from the 77.2-keV level											FKo25c	**
*^{181}Pb(α)^{177}Hg	Seen in correlation with ^{177}Hg E_α=6580 keV											96To01	**
*^{181}Pb(α)^{177}Hg	summed with ce from the 77.2-keV level											FKo25c	**
*^{181}Pb(α)^{177}Hg	E_α=7015(10) to level at 77.2 keV											09An20	**
*^{181}Pb(α)^{177}Hg	E_α=7016(15) to level at 77.2 keV											09An20	**
*^{181}Ta(γ,n)^{180}Ta	Q =–7640(25) to ^{180}Ta^{m} at 77.3(0.4) keV											Nub269	**
*^{181}Hg(εp)^{180}Pt	F : retracted by authors in PrvCom											AHW	**
*^{181}Hf(β^-)^{181}Ta	E_{β^-}=408(8) 405(5) respectively, to ^{181}Ta^{n} at 615.19 keV											Nub269	**
*^{181}Os(β^+)^{181}Re	E_{β^+}=1750(200) from ^{181}Os^{m} at 49.20 to ^{181}Re^{m} at 262.91 keV											Nub269	**
*^{181}Hg^{m}(IT)^{181}Hg	From cascade x+90.3+71.4, with x estimated 50#											09An17	**
^{182}Yb−^{163}Dy ^{19}F		30937	36	30934	28	−0.1	1	61	61 ^{182}Yb	TR1	1.0	26Br.A	
^{182}Re−u		−48311	65	−48790	110	−7.3	C			GS2	1.0	03Li.A	*
^{182}Os−u		−47883	30	−47890	23	−0.2	1	61	61 ^{182}Os	GS2	1.0	05Li24	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{182}Ir−u		−41942	30	−41924	23	0.6	1	56	56 ^{182}Ir	GS2	1.0	05Li24	
^{182}Pt−u		−38870	104	−38829	14	0.4	U			GS1	1.0	00Ra23	
		−38860	30			1.0	1	22	22 ^{182}Pt	GS2	1.0	05Li24	
^{182}Au−u		−30420	110	−30388	20	0.3	U			GS1	1.0	00Ra23	
		−30412	30			0.8	1	44	44 ^{182}Au	GS2	1.0	05Li24	
^{182}Hg−u		−25297	30	−25312	10	−0.5	1	12	12 ^{182}Hg	GS2	1.0	05Li24	
^{182}Hg−^{208}Pb$_{.875}$		−4893	19	−4881	10	0.6	–			MA6	1.0	01Sc41	
		−4898	21			0.8	–			MA6	1.0	01Sc41	
	ave.	−4895	14			1.0	1	55	55 ^{182}Hg			average	
^{182}Yb−^{182}W		9863	45	9868	28	0.1	1	39	39 ^{182}Yb	TR1	1.0	26Br.A	
^{182}Pt(α)^{178}Os		4928.5	30.	4951	5	0.7	U					63Gr08	
		4948.9	20.			0.1	U					66Si08	
		4952.0	5.			−0.2	1	97	76 ^{178}Os	ORa		95Bi01	
^{182}Au(α)^{178}Ir		5529	10	5524	4	−0.5	o			ISa		79Ha10	*
		5525.5	5.			−0.3	–			ORa		95Bi01	*
		5523.6	5.1			0.1	–			ISa		25Mi06	*
	ave.	5525	4			−0.1	1	99	55 ^{178}Ir			average	
^{182}Hg(α)^{178}Pt		5998.1	5.	5994	4	−0.9	–			ISa		79Ha10	Z
		5999.1	10.2			−0.5	o			Lvn		93Wa.A	
		5989.9	13.3			0.3	–			Lvn		94Wa23	
		5988.9	7.2			0.7	–			Lza		24Hu14	
	ave.	5995	4			−0.2	1	96	64 ^{178}Pt			average	
^{182}Tl(α)^{178}Au		6550.2	10.	6547	5	−0.4	F			GSa		86Ke03	*
		6593.1	15.			−3.1	B					04Ra28	*
		6546.7	5.1				3			ISa		16Va01	*
^{182}Tl^{m}(α)^{178}Au^{q}		6186.2	20.				5					92Bo.D	
^{182}Pb(α)^{178}Hg		7076.8	10.	7063	5	−1.4	4			GSa		86Ke03	
		7074.8	15.			−0.8	4			ORa		87To09	
		7050.2	10.			1.2	4			Ara		99To11	
		7066.6	10.			−0.4	4			Jya		00Je09	
		7056.3	8.2			0.8	4			Lza		24Hu14	
^{180}Hf(t,p)^{182}Hf		3931	6				2			McM		83Bu03	
^{180}W(t,p)^{182}W		6265	5	6270.5	1.1	1.1	U			LAl		76Ca10	*
^{182}W(p,t)^{180}W		−6261	10	−6270.5	1.1	−0.9	U			Min		73Oo01	
^{181}Ta(n,γ)^{182}Ta		6063.0	0.4	6062.94	0.11	−0.2	–					71He13	Z
		6063.1	0.5			−0.3	–					77St15	Z
		6063.1	0.5			−0.3	–			MMn		81Co17	Z
		6062.95	0.2			−0.1	–			ILn		83Fo.B	
		6062.89	0.14			0.3	–			Bdn		06Fi.A	
^{181}Ta(d,p)^{182}Ta		3832	8	3838.37	0.11	0.8	U			MIT		64Er02	
^{181}Ta(n,γ)^{182}Ta	ave.	6062.93	0.11	6062.94	0.11	0.1	1	100	69 ^{182}Ta			average	
^{182}W(d,t)^{181}W		−1809	10	−1826.0	1.1	−1.7	U			Kop		72Ca01	
^{182}Hf(β^-)^{182}Ta		431	50	381	6	−1.0	U					74Wa14	*
^{182}Ta(β^-)^{182}W		1809	5	1815.1	1.5	1.2	–					64Da15	*
		1813	3			0.7	–					67Ba01	*
	ave.	1811.9	2.6			1.2	1	34	31 ^{182}Ta			average	
^{182}Re^{m}(β^+)^{182}W		2860	20				2					63Ba37	*
^{182}Re^{m}(IT)^{182}Re		60	100				3					63Ba37	
^{182}Os(ε)^{182}Re^{m}		848	15	777	30	−4.7	B					70Ak02	*
^{182}Ir(β^+)^{182}Os		5700	200	5560	30	−0.7	U					72We.A	
^{182}Pt(β^+)^{182}Ir		2900	200	2883	25	−0.1	U					72We.A	
^{182}Au(β^+)^{182}Pt		6850	200	7863	23	5.1	C					72We.A	
^{182}Hg(β^+)^{182}Au		4950	200	4728	21	−1.1	U					72We.A	
*^{182}Re−u	$M-A$=–44972(29) keV for mixture gs+m at 60(100) keV											Nub269	**
*^{182}Au(α)^{178}Ir	E_α=5353(10) to 2^+ level at 54.4(0.5) keV											Ens097	**
*^{182}Au(α)^{178}Ir	E_α=5403(5), 5352(5) to ground state, 54.4 level											95Bi01	**
*^{182}Au(α)^{178}Ir	Also E_α=5350(5),5293(8) 5282(5) to 55.0(0.2),114.7(0.5)											25Mi06	**
*^{182}Au(α)^{178}Ir	127.5(0.7) keV levels											25Mi06	**
*^{182}Tl(α)^{178}Au	F : identification from excitation function assuming 100% α decay											WgM118	**
*^{182}Tl(α)^{178}Au	E_α=6403(15) in coincidence with 46 keV γ											04Ra28	**
*^{182}Tl(α)^{178}Au	E_α=6046(5),6165(6) in coincidence with 361.5(0.1),247.2(0.5) keV γs											16Va01	**
*^{180}W(t,p)^{182}W	$Q-Q(^{170}\mathrm{Y(t,p)})$=112(5,Ca), Q(170)=–6153(4) keV											AHW	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{182}$Hf(β^-)^{182}Ta		E_{β^-}=970(70) 480(50) from ^{182}Hf^m to 651.215 (4)$^-$, 1115.96 (7$^-$) levels										Ens15c	**
$*^{182}$Ta(β^-)^{182}W		E_{β^-}=520(5) to 2$^-$ level at 1289.1498 keV										Ens15c	**
$*^{182}$Ta(β^-)^{182}W		E_{β^-}=1713(3) to 2$^+$ level at 100.10598 keV										Ens15c	**
$*^{182}$Re^m(β^+)^{182}W		E_{β^+}=1740(20), 550(20) to 2$^+$ level at 100.10598, 2$^-$ at 1289.1498 keV										Ens15c	**
$*^{182}$Os(ε)^{182}Re^m		pK=0.47(0.07) to 1$^+$ level at 726.97 keV above ^{182}Re^m, recalculated Q										Ens15c	**
^{183}Lu−u		−42637	86				2			GS3	1.0	13Sh30	
^{183}W O−C$_2$ ^{35}Cl$_5$		100858.0	2.7	100875.5	0.8	2.6	U			H29	2.5	77Sh04	
		100873.6	0.8			1.0	1	16	16 ^{183}W	H48	2.5	03Ba49	
^{183}Re−u		−49151	30	−49179	9	−0.9	U			GS2	1.0	05Li24	
^{183}Os−u		−46879	61	−46880	50	0.0	1	76	76 ^{183}Os	GS2	1.0	05Li24	*
^{183}Ir−u		−43160	104	−43168	22	−0.1	U			GS1	1.0	00Ra23	
		−43145	30			−0.8	1	54	54 ^{183}Ir	GS2	1.0	05Li24	
^{183}Pt−u		−38440	107	−38404	15	0.3	U			GS1	1.0	00Ra23	
		−38400	32			−0.1	1	23	23 ^{183}Pt	GS2	1.0	05Li24	*
^{183}Au−u		−32440	104	−32411	10	0.3	U			GS1	1.0	00Ra23	
		−32371	30			−1.3	1	11	11 ^{183}Au	GS2	1.0	05Li24	
^{183}Hg−u		−25537	35	−25556	8	−0.5	U			GS2	1.0	05Li24	
^{183}Hg−^{208}Pb$_{.880}$		−5009	19	−5009	8	0.0	–			MA6	1.0	01Sc41	
		−5002	19			−0.3	–			MA6	1.0	01Sc41	
	ave.	−5006	13			−0.2	1	32	32 ^{183}Hg			average	
^{183}Tl−^{133}Cs$_{1.376}$		112286	11	112290	10	0.4	1	81	81 ^{183}Tl	MA8	1.0	08We02	
^{183}W O$_2$−^{178}Hf ^{37}Cl		30455.7	5.0	30443.1	1.2	−1.0	U			H35	2.5	80Sh06	
^{183}W O$_2$−^{180}W ^{35}Cl		24421	9	24487.9	1.2	3.0	B			H24	2.5	73Ba40	
		24509	6			−1.4	U			H28	2.5	77Sh04	
^{183}W ^{35}Cl−^{181}Ta ^{37}Cl		5177.2	1.2	5176.4	1.6	−0.3	1	29	26 ^{181}Ta	H35	2.5	80Sh06	
^{183}W O$_2$ ^{37}Cl−^{182}W ^{35}Cl$_2$		20045.6	1.8	20045.19	0.11	−0.1	U			H28	2.5	77Sh04	
^{183}Yb−^{183}W		12078	27				2			TR1	1.0	26Br.A	
^{183}Ir−^{183}W		6585	40	6607	22	0.6	1	31	30 ^{183}Ir	RI1	1.0	26Ya.1	
^{183}Pt(α)^{179}Os		4846.1	30.	4822	9	−0.8	U					63Gr08	
		4835.9	20.0			−0.7	–					66Si08	
		4819.6	10.2			0.2	–			ORa		95Bi01	
	ave.	4823	9			−0.1	1	93	69 ^{179}Os			average	
^{183}Au(α)^{179}Ir		5462.6	5.	5465.3	2.9	0.5	–					68Si01	Z
		5465.5	5.			−0.0	–			Bka		82Bo04	Z
		5449.3	10.			1.6	F					84Br.A	*
		5468.8	5.			−0.7	–			ORa		95Bi01	
	ave.	5465.6	3.0			−0.1	1	99	88 ^{179}Ir			average	
^{183}Hg(α)^{179}Pt		6043.4	6.	6038.0	2.8	−0.9	–			ORa		76To06	
		6036.2	5.			0.3	–			ISa		79Ha10	Z
		6037.2	4.1			0.2	–			Ara		22Hu09	
	ave.	6038.2	2.8			−0.1	1	99	93 ^{179}Pt			average	
^{183}Tl^m(α)^{179}Au		6593.4	15.3	6602	6	0.6	–			GSa		80Sc09	*
		6600.5	15.3			0.1	–			Jya		04Ra28	*
		6609.5	10.			−0.8	–			Anv		11Ve01	*
		6602.6	15.3			−0.0	–			ISa		22Ve01	*
	ave.	6604	7			−0.3	1	92	73 ^{179}Au			average	
^{183}Pb(α)^{179}Hg		6929	7	6931	5	0.4	2			Anv		02Je09	*
		6944	10			−1.3	2			GSa		86Ke03	*
		6950	26			−0.7	U			GSa		80Sc09	*
		6938	15			−0.4	2			ORa		89To01	*
		6921	10			1.0	2			Lza		24Hu14	*
^{183}Pb^m(α)^{179}Hg		7037	20	7020	4	−0.8	U			GSa		80Sc09	*
		7029	20			−0.4	U			GSa		84Sc.A	*
		7026.9	10.2			−0.7	2			GSa		86Ke03	
		7040	10			−2.0	o			ORa		87To09	*
		7033	10			−1.3	2			ORa		89To01	*
		7018	5			0.4	2			Anv		02Je09	*
		7014.1	7.2			0.8	2			Lza		24Hu14	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{183}W(p,t)^{181}W		−5810	10	−5792.3	1.1	1.8	U			Min		73Oo01	
^{182}Ta(n,γ)^{183}Ta		6934.18	0.20				2			ILn		83Fo.B	
^{182}W(n,γ)^{183}W		6191.6	2.0	6190.84	0.04	−0.4	U					67Sp03	Z
		6190.1	1.5			0.5	U					70Or.A	
		6190.76	0.12			0.6	–			Ltn		97Pr02	
		6190.89	0.13			−0.4	o			Bdn		06Fi.A	
		6190.81	0.06			0.4	–			ILn		11Bo09	
		6190.88	0.06			−0.7	–			Bdn		14Hu02	
^{183}W(γ,n)^{182}W		−6290	50	−6190.84	0.04	2.0	U			Phi		60Ge01	
^{182}W(d,p)^{183}W		3967	5	3966.27	0.04	−0.1	U			ANL		65Er03	
		3979	10			−1.3	U			Kop		72Ca01	
^{183}W(d,t)^{182}W		57	15	66.39	0.04	0.6	U			Kop		72Ca01	
^{182}W(n,γ)^{183}W	ave.	6190.84	0.04	6190.84	0.04	0.0	1	100	100 ^{182}W			average	
^{182}W(^{3}He,d)^{183}Re		−610	40	−641	8	−0.8	U			Roc		71Lu01	
^{183}Hg(εp)^{182}Pt		5000	200	5075	15	0.4	F					72Ho19	
^{183}Hf(β^-)^{183}Ta		2010	30				3					67Mo13	*
^{183}Ta(β^-)^{183}W		1068	10	1071.7	1.5	0.4	U					55Mu19	*
^{183}Re(ε)^{183}W		556	8				2					69Ku03	*
^{183}Ir(β^+)^{183}Os		3450	100	3450	50	0.0	1	27	24 ^{183}Os			70Be.A	*
^{183}Tl^m(IT)^{183}Tl		628.7	0.3	628.70	0.30	0.0	1	100	81 ^{183}Tl^m			22Ve01	
*^{183}Os−u	$M-A$=–43582(28) keV for mixture gs+m at 170.73 keV											Nub269	**
*^{183}Pt−u	$M-A$=–35752(28) keV for mixture gs+m at 34.74 keV											Nub269	**
*^{183}Au(α)^{179}Ir	F : all lines in ^{181}Au and ^{183}Au shifted by 16−20keV											GAu	**
*^{183}Tl^m(α)^{179}Au	E_α=6400(10), 6378(10) summed with e$^-$, in coinc. with 62.4, 89.5 γ											GAu	**
*^{183}Tl^m(α)^{179}Au	also E_α=6333(9), 6058(15) to levels at 127(17), 407(17) keV											22Ve01	**
*^{183}Pb(α)^{179}Hg	E_α=6775(7), 6570(10) to ground state, 217 level											02Je09	**
*^{183}Pb(α)^{179}Hg	E_α=6580(10) to 217 level											FKo255	**
*^{183}Pb(α)^{179}Hg	E_α=6798(25) to ground state											FKo255	**
*^{183}Pb(α)^{179}Hg	E_α=6781(15),6579(15) to ground state, 217 level											FKo255	**
*^{183}Pb(α)^{179}Hg	E_α=6765(10),6562(10) to ground state, 217 level											24Hu14	**
*^{183}Pb^m(α)^{179}Hg	E_α=6715(20) to 171.4(0.4) isomer											FKo255	**
*^{183}Pb^m(α)^{179}Hg	E_α=6868(20), 6715(20) to ground state, 171.4(0.4) isomer											02Je09	**
*	original assignment to ^{182}Pb changed											AHW	**
*^{183}Pb^m(α)^{179}Hg	E_α=6718(10) to 171.4(0.4) isomer											FKo255	**
*^{183}Pb^m(α)^{179}Hg	E_α=6874(15), 6712(10) to ground state, 171.4(0.4) isomer											Nub269	**
*^{183}Pb^m(α)^{179}Hg	E_α=6860(11), 6698(5) to ground state, 171.4 isomer											Nub269	**
*^{183}Pb^m(α)^{179}Hg	probably decays to the 171.4 keV level											HWJ253	**
*^{183}Hg(εp)^{182}Pt	F : retracted by authors in PrvCom											AHW	**
*^{183}Hf(β^-)^{183}Ta	E_{β^-}=1540(30) to (5/2$^+$) level at 459.062 keV, and other E_{β^-}											Ens164	**
*^{183}Ta(β^-)^{183}W	E_{β^-}=615(10) to 7/2$^-$ level at 453.0695 keV											Ens164	**
*^{183}Re(ε)^{183}W	pK=0.40(0.07) to 7/2$^-$ level at 453.0695 keV											Ens164	**
*^{183}Ir(β^+)^{183}Os	Q_{β^+}=3190(100) mainly to 3/2$^-$ level at 258.34 keV											Ens164	**
^{184}W−u		−49066.76	0.99	−49066.9	0.8	−0.1	1	28	28 ^{184}W	TG1	1.5	12Sm07	
^{184}Os−u		−47504.3	1.2	−47507.1	0.9	−1.6	1	24	24 ^{184}Os	TG1	1.5	12Sm07	
^{184}Ir−u		−42460	110	−42549	12	−0.8	U			GS1	1.0	00Ra23	
		−42524	30			−0.8	1	16	16 ^{184}Ir	GS2	1.0	05Li24	
^{184}Pt−u		−40120	104	−40078	16	0.4	U			GS1	1.0	00Ra23	
		−40068	30			−0.3	1	28	28 ^{184}Pt	GS2	1.0	05Li24	
^{184}Au−u		−32540	104	−32548	24	−0.1	U			GS1	1.0	00Ra23	*
		−32557	37			0.2	R			GS2	1.0	05Li24	*
^{184}Hg−u		−28230	110	−28283	10	−0.5	U			GS1	1.0	00Ra23	
		−28296	30			0.4	–			GS2	1.0	05Li24	
	ave.	−28301	21			0.8	1	23	23 ^{184}Hg			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{184}Hg−^{204}Pb$_{.902}$		−3986	20	−3967	10	1.0	1	26	26 ^{184}Hg	MA6	1.0	01Sc41	
^{184}Hg−^{208}Pb$_{.885}$		−7620	19	−7619	10	0.1	1	29	29 ^{184}Hg	MA6	1.0	01Sc41	
^{184}Tl−u		−18115	112	−18122	11	−0.1	U			GS2	1.0	05Li24	*
^{184}Tl−^{133}Cs$_{1.383}$		112645.4	23.2	112638	11	−0.3	1	21	21 ^{184}Tl	MA8	1.0	14Bo26	*
^{184}W O$_2$−^{181}Ta ^{35}Cl		23917.5	2.8	23911.6	1.6	−0.8	U			H35	2.5	80Sh06	
^{184}W ^{35}Cl−^{182}W ^{37}Cl		5675	3	5677.68	0.16	0.4	U			H22	2.5	70Mc03	
		5676.3	2.2			0.3	U			H28	2.5	77Sh04	
^{184}W O$_2$ ^{37}Cl−^{183}W ^{35}Cl$_2$		18734.7	3.0	18735.18	0.17	0.1	U			H28	2.5	77Sh04	
^{184}Yb−^{184}W		14372	40				2			TR1	1.0	26Br.A	
^{184}Ta−^{184}W		3143.5	7.7				2			RI1	1.0	26Ya.1	
^{184}Os−^{184}W		1560.59	0.70	1559.8	0.7	−0.8	1	46	31 ^{184}Os	TG1	1.5	12Sm07	
^{184}Ir−^{184}W		6513	13	6518	12	0.4	1	84	84 ^{184}Ir	RI1	1.0	26Ya.1	
^{184}Pt(α)^{180}Os		4579.8	20.	4599	8	0.9	–					63Gr08	
		4600.2	20.			−0.1	–					66Si08	
		4602.2	10.			−0.3	–			ORa		95Bi01	
	ave.	4598	8			0.1	1	94	68 ^{180}Os			average	
^{184}Au^m(α)^{180}Ir		5287.1	15.	5302	5	1.0	U			ISa		70Ha18	
		5302.5	5.				3			ORa		95Bi01	
^{184}Hg(α)^{180}Pt		5658.2	15.	5660	4	0.1	–			ISa		70Ha18	
		5662.3	5.1			−0.4	–			ORa		76To06	
		5658.2	10.2			0.2	o			Lvn		93Wa.A	
		5662.3	10.2			−0.2	–			Lvn		93Wa03	Z
	ave.	5662	4			−0.4	1	96	75 ^{180}Pt			average	
^{184}Tl(α)^{180}Au		6299.4	5.	6321	9	0.4	U			ORa		76To06	Z
		6292.9	10.			0.5	U			GSa		80Sc09	Z
		6319	10			0.2	1	83	79 ^{184}Tl	ISa		16Va01	*
^{184}Pb(α)^{180}Hg		6765.4	10.	6774	3	0.8	–					80Du02	
		6779.6	10.			−0.6	–			GSa		80Sc09	
		6773.6	10.			0.0	–			GSa		84Sc.A	
		6781.7	10.2			−0.8	–			ORa		87To09	
		6773.6	6.			0.1	–			Jya		98Co27	
		6772.5	10.			0.1	–			Ara		99To11	
		6773.6	6.			0.1	–			Anv		04An07	
	ave.	6774	3			−0.0	1	99	70 ^{184}Pb			average	
^{184}Bi(α)^{180}Tl		7830	15	8040#	100#	14.0	D			Anv		03An27	*
^{184}Bi^m(α)^{180}Tl^p		7354.2	20.4				8			Anv		03An27	
^{182}W(t,p)^{184}W		5127	7	5120.15	0.14	−1.0	U			LAl		76Ca10	
^{184}W(p,t)^{182}W		−5124	5	−5120.15	0.14	0.8	U			Min		73Oo01	
^{183}W(n,γ)^{184}W		7411.2	0.5	7411.11	0.13	−0.2	U					74Gr11	Z
		7411.8	0.3			−2.3	B					75Bu01	Z
		7411.15	0.16			−0.3	o			Bdn		06Fi.A	
		7411.11	0.13			−0.0	1	99	73 ^{183}W	Bdn		14Hu02	
^{183}W(d,p)^{184}W		5187	15	5186.54	0.13	−0.0	U			Kop		72Ca01	
^{184}W(d,t)^{183}W		−1154	10	−1153.88	0.13	0.0	U			Kop		72Ca01	
^{184}Hf(β^-)^{184}Ta		1340	30				3					73Wa18	*
^{184}Ta(β^-)^{184}W		2866	26	2928	7	2.4	U					73Ya02	*
^{184}Ir(β^+)^{184}Os		5100	250	4618	11	−1.9	U					70Be.A	*
		4300	100			3.2	B					73Ho09	*
		4285	70			4.8	B					89Po09	*
^{184}Au(β^+)^{184}Pt		6380	50	7014	27	12.7	C					84Da.A	*
^{184}Au^m(IT)^{184}Au		68.46	0.04				4					Ens106	
^{184}Hg(β^+)^{184}Au		3760	30	3973	24	7.1	C					84Da.A	
*^{184}Au−u	$M-A$=–30280(100) keV for mixture gs+m at 68.46 keV											Nub269	**
*^{184}Au−u	$M-A$=–30292(28) keV for mixture gs+m at 68.46 keV											Nub269	**
*^{184}Tl−u	$M-A$=–16899(102) keV for mixture gs+m at –50(30) keV											Nub211	**
*^{184}Tl−^{133}Cs$_{1.383}$	D_M=112618.6(6.2) μu for mixture gs+m at –50(30) keV; $M-A$=–16898.5(5.8) keV											Nub211	**
*	transition to ground state in ^{180}Ir											95Bi01	**
*^{184}Tl(α)^{180}Au	E_α=6161(10),5988(12),5964(13) to levels at 17.1(3),201.3(0.3),											16Va01	**
*	224.3(0.3) keV											16Va01	**
*^{184}Bi(α)^{180}Tl	E_α=7220(15) to level at 449(1) keV											03An27	**
*^{184}Bi(α)^{180}Tl	Trends in Mass Surface TMS suggest ^{184}Bi 210 keV less bound											HWJ266	**
*^{184}Hf(β^-)^{184}Ta	E_{β^-}=1100(30) to 1$^-$ level at 228.4 keV , and other E_{β^-}											Ens102	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
$*^{184}$Ta(β^-)^{184}W	E_{β^-}=1165(26) to 6^+ level at 1746.03 keV, and other E_{β^-}												Ens102	**
$*^{184}$Ir(β^+)^{184}Os	Q_{β^+}=4720(250) to 4^+ level at 383.68 keV												Ens102	**
$*^{184}$Ir(β^+)^{184}Os	E_{β^+}=2900(100) to 4^+ level at 383.68 keV												Ens102	**
$*^{184}$Ir(β^+)^{184}Os	E_{β^+}=2320(70) to 2^+ level at 942.86 keV												Ens102	**
$*^{184}$Au(β^+)^{184}Pt	Q_{β^+}=6450(50) from ^{184}Au^m at 68.6(0.1) keV												94Ib01	**
^{185}Hf−u		−41138	69				2				GS3	1.0	13Sh30	
^{185}Os−u		−46037	31	−45954.0	0.9	2.7	F				GS2	1.0	03Li.A	*
^{185}Ir−u		−43340	110	−43300	30	0.3	U				GS1	1.0	00Ra23	
		−43302	30				2				GS2	1.0	05Li24	
^{185}Pt−u		−39334	112	−39386	28	−0.5	U				GS1	1.0	00Ra23	*
		−39381	44			−0.1	1	40	40	^{185}Pt	GS2	1.0	05Li24	*
^{185}Au−u		−34173	97	−34214	8	−0.4	o				GS1	1.0	00Ra23	*
		−34183	31			−1.0	U				GS2	1.0	05Li24	*
^{185}Au−^{133}Cs$_{1.391}$		97302.5	7.9				2				MA8	1.0	17Ma29	*
^{185}Hg−u		−28070	107	−28110	15	−0.4	U				GS1	1.0	00Ra23	
		−28088	44			−0.5	1	11	11	^{185}Hg	GS2	1.0	05Li24	*
^{185}Hg−^{208}Pb$_{.889}$		−7373	29	−7352	15	0.7	1	25	25	^{185}Hg	MA6	1.0	01Sc41	*
^{185}Tl−u		−21354	145	−21209	22	1.0	U				GS2	1.0	05Li24	*
^{185}Re O_2−^{182}W ^{35}Cl		25731	6	25729.2	0.7	−0.1	U				H22	2.5	70Mc03	
^{185}Re ^{35}Cl−^{183}W ^{37}Cl		5695	3	5684.1	0.7	−1.5	U				H22	2.5	70Mc03	
		5678.7	1.0			2.1	U				H28	2.5	77Sh04	
^{185}Re(α,^{8}He)^{181}Re		−26480	14	−26486	13	−0.5	2				INS		90Ka19	
^{185}Pt(α)^{181}Os		4436.6	10.2	4437	10	0.0	1	96	60	^{185}Pt	ORa		91Bi04	*
^{185}Au(α)^{181}Ir		5180.2	5.	5180	5	−0.0	3						68Si01	Z
		5182.9	15.			−0.2	U				ISa		70Ha18	Z
		5179	10			0.1	3				ORa		91Bi04	*
^{185}Hg(α)^{181}Pt		5777	15	5773	4	−0.3	–				ISa		70Ha18	*
		5775	5			−0.4	–				ORa		76To06	*
		5761	15			0.8	–						76Gr.A	*
	ave.	5774	5			−0.2	1	97	52	^{181}Pt			average	
^{185}Tl^m(α)^{181}Au^p		6112.6	7.	6143	5	4.4	C						75Co.A	Z
		6143.3	5.				5				ORa		76To06	*
		6145.6	15.			−0.2	U				GSa		80Sc09	Z
^{185}Pb(α)^{181}Hg		6693	15	6695	5	0.1	U				GSa		80Sc09	*
		6555.0	15.			2.7	B				ORa		87To09	
		6695	5				2				Anv		02An15	*
^{185}Pb^m(α)^{181}Hg^m		6622.9	20.	6550	5	−3.7	B				Ora		75Ca06	
		6679.7	20.			−6.5	B				GSa		80Sc09	
		6549.8	5.				3				Anv		02An15	
^{185}Bi(α)^{181}Tl		8258.9	30.	8218	12	−1.3	–				Ara		01Po05	*
		8207.8	15.3			0.7	–				Anv		04An07	
	ave.	8218	14			0.0	1	75	63	^{185}Bi			average	
^{184}W(n,γ)^{185}W		5753.7	0.3	5753.74	0.05	0.1	U				BNn		87Br05	Z
		5754.62	0.24			−3.7	C				Bdn		06Fi.A	
		5753.74	0.05			−0.1	1	100	85	^{185}W	Bdn		14Hu02	
^{184}W(d,p)^{185}W		3524	5	3529.17	0.05	1.0	U				ANL		65Er03	
		3533	10			−0.4	U				Kop		72Ca01	
^{184}W(^{3}He,d)^{185}Re		−98	40	−90.9	0.7	0.2	U				Roc		71Lu01	
^{185}Re(d,t)^{184}Re−^{187}Re()^{186}Re		−310	4	−310	4	0.0	1	100	100	^{184}Re	Roc		76El12	
^{184}Os(n,γ)^{185}Os		6625.4	0.9	6624.66	0.27	−0.8	U						74Pr15	
		6624.52	0.28			0.5	1	95	51	^{185}Os	Bdn		06Fi.A	
^{185}Bi(p)^{184}Pb		1669	50	1607	13	−1.2	o						95Da.A	*
		1606.8	16.			−0.0	1	67	37	^{185}Bi	Ara		01Po05	*
		1568.6	50.			0.8	o						03An27	*
		1591.7	5.			3.0	F						04An07	*
^{185}Ta(β^-)^{185}W		2013	20	1993	14	−1.0	2						69Ku07	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{185}W(β^-)^{185}Re		432.6	1.0	431.2	0.7	−1.4	1	44	28 ^{185}Re			67Wi19	
^{185}Os(ε)^{185}Re		1012.7	1.0	1013.1	0.4	0.4	–					67Sc15	*
		1012.8	0.5			0.7	–					70Sc06	*
	ave.	1012.8	0.4			0.8	1	88	49 ^{185}Os			average	
^{185}Au(β^+)^{185}Pt		4707	40	4818	27	2.8	F					86Da.A	*
^{185}Tl^m(IT)^{185}Tl		454.8	1.5	454.70	0.22	−0.1	U					Ens061	
		454.70	0.22				6					13Sa43	*
*^{185}Os−u	F : contaminated by isomeric state AND by other nuclides											03Li.B	**
*^{185}Pt−u	$M-A$=–36590(100) keV for mixture gs+m at 103.41 keV											Nub269	**
*^{185}Pt−u	$M-A$=–36631(28) keV for mixture gs+m at 103.41 keV											Nub269	**
*^{185}Au−u	$M-A$=–31820(90) keV for mixture gs+m at 23.6(0.1) keV											Nub269	**
*^{185}Au−u	$M-A$=–31829(28) keV for mixture gs+m at 23.6(0.1) keV											Nub269	**
*^{185}Au−^{133}Cs$_{1.391}$	D_M=97315.2(2.8) μu for mixture g+m at 23.6(0.1) keV;											Nub269	**
*^{185}Au−^{133}Cs$_{1.391}$	$M-A$=–31858.2(2.6) keV											Nub269	**
*^{185}Hg−u	$M-A$=–26112(28) keV for mixture gs+m at 103.7 keV											Nub269	**
*^{185}Hg−^{208}Pb$_{.889}$	Original error (17keV) increased by 20 due to isomer+ground state lines in trap											01Sc41	**
*^{185}Tl−u	$M-A$=–19664(31) keV for mixture gs+m at 454.70(0.22) keV											Nub269	**
*^{185}Pt(α)^{181}Os	E_α=4444(10) assumed from (1/2$^-$) isomer at 103.41 keV											Nub269	**
*^{185}Au(α)^{181}Ir	E_α=5069(10), 4826(10) to ground state, 243.3 level											91Bi04	**
*	unhindered E_α=5069(10) to ground state or very low level; from coinc.											95Bi01	**
*^{185}Hg(α)^{181}Pt	E_α=5653.4(15,Z), 5576.4(15,Z) to ground state, 3/2$^-$ level at 79.41 keV											Ens061	**
*	and E_α=5376.4(15,Z) from ^{185}Hg^m at 103.8 to 13/2$^+$ level at 380.92 keV											Ens061	**
*^{185}Hg(α)^{181}Pt	E_α=5653(5), 5569(5) to ground state, 3/2$^-$ level at 79.41 keV											Ens061	**
*	and 5371(10) from ^{185}Hg^m at 103.8 to 13/2$^+$ level at 380.92 keV											Ens061	**
*^{185}Hg(α)^{181}Pt	E_α=5365(15) from ^{185}Hg^m at 103.8 to 13/2$^+$ level at 380.92 keV											Ens061	**
*^{185}Tl^m(α)^{181}Au^p	E_α=6010.2(5,Z); also an E_α=5975.2(5,Z), 4 times stronger branch											76To06	**
*^{185}Tl^m(α)^{181}Au^p	E_α=6012.5(15,Z); also an E_α=5970.5(15,Z), 4 times stronger branch											80Sc09	**
*^{185}Pb(α)^{181}Hg	E_α=6485(15) to 64 level											02An15	**
*^{185}Pb(α)^{181}Hg	E_α=6486(5),6288(5) to 64, 269 levels											02An15	**
*^{185}Bi(α)^{181}Tl	E_α=8030, by same authors, from only one event											96Da06	**
*^{185}Bi(p)^{184}Pb	Read from graph											AHW	**
*^{185}Bi(p)^{184}Pb	Average by authors of E_p=1618(11), and 1585(9) in reference											96Da06	**
*^{185}Bi(p)^{184}Pb	As read from graph											GAu	**
*^{185}Bi(p)^{184}Pb	F : rejected by authors: no dedicated calibration with known proton activity											04An07	**
*^{185}Ta(β^-)^{185}W	E_{β^-}=1770(20) to 7/2$^-$ level at 243.62 keV											Ens061	**
*^{185}Os(ε)^{185}Re	L/K=0.600(0.006) to 3/2$^+$ level at 874.81 and 1/2$^+$ at 880.33 keV											Ens061	**
*^{185}Os(ε)^{185}Re	pK=0.109(0.005) to 3/2$^+$ level at 931.06 keV, and other pK, recalculated											Ens061	**
*^{185}Au(β^+)^{185}Pt	F : insufficient information											GAu	**
*^{185}Tl^m(IT)^{185}Tl	E(γ)=169.2(0.1) 285.5(0.2) keV											13Sa43	**
^{186}Hf−u		−39103	55				2			GS3	1.0	13Sh30	
^{186}W O−C ^{13}C ^{35}Cl$_4$ ^{37}Cl		104592.7	3.2	104611.6	1.3	2.4	U			H29	2.5	77Sh04	
^{186}Ir−u		−42063	30	−42053	18	0.3	2			GS2	1.0	05Li24	*
^{186}Pt−u		−40656	30	−40649	23	0.2	1	61	61 ^{186}Pt	GS2	1.0	05Li24	
^{186}Au−u		−34029	30	−34047	23	−0.6	1	56	56 ^{186}Au	GS2	1.0	05Li24	
^{186}Hg−u		−30660	104	−30639	12	0.2	U			GS1	1.0	00Ra23	
		−30630	30			−0.3	1	17	17 ^{186}Hg	GS2	1.0	05Li24	
^{186}Hg−^{204}Pb$_{.912}$		−6065	20	−6053	12	0.6	–			MA6	1.0	01Sc41	
	ave.	−6058	17			0.3	1	56	56 ^{186}Hg			average	
^{186}Tl−u		−21670	220	−21361	16	1.4	o			GS1	1.0	00Ra23	*
		−21387	35			0.7	U			GS2	1.0	05Li24	*
^{186}Tl−^{133}Cs$_{1.398}$		110819	24	110817	16	−0.1	o			MA8	1.0	08We02	*
		110817	18			−0.0	1	76	76 ^{186}Tl	MA8	1.0	14Bo26	*
^{186}Tl^n−^{133}Cs$_{1.398}$		111254	34	111267	20	0.4	1	34	34 ^{186}Tl^n	MA8	1.0	14Bo26	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{186}W O_2−^{183}W ^{35}Cl	25122	5	25117.3	1.3	−0.4	U			H28	2.5	77Sh04	
^{186}W ^{35}Cl−^{184}W ^{37}Cl	6374	3	6382.1	1.2	1.1	U			H22	2.5	70Mc03	
	6382.0	1.4			0.0	1	13	11 ^{186}W	H28	2.5	77Sh04	
^{186}Os(α)^{182}W	2820.6	50.	2821.2	0.9	0.0	U					75Vi01	
^{186}Pt(α)^{182}Os	4323.2	20.	4320	18	−0.2	1	79	39 ^{182}Os			63Gr08	
^{186}Au(α)^{182}Ir	4907	15	4912	14	0.3	1	87	44 ^{182}Ir	ORa		90Ak04	*
^{186}Hg(α)^{182}Pt	5206.2	15.	5204	10	−0.1	–			ISa		70Ha18	
	5204.2	15.			0.0	–					96Ri12	
ave.	5205	11			−0.1	1	83	57 ^{182}Pt			average	
^{186}Tl(α)^{182}Au	6016.2	52.1	5984	23	−0.6	1	19	12 ^{182}Au	ISa		20St11	*
^{186}Tl^m(α)^{182}Au^m	5891.9	7.	5892	5	−0.0	3					75Co.A	
	5891.9	7.			−0.0	3			ORa		77Ij01	*
^{186}Pb(α)^{182}Hg	6458.2	20.	6471	5	0.6	2			Ora		74Le02	Z
	6470.1	10.			0.1	2			GSa		80Sc09	Z
	6474.7	10.			−0.4	2			ORa		84To09	Z
	6474.5	10.2			−0.3	2			Lvn		93Wa.A	
	6476.5	15.			−0.4	2			ORa		97Ba25	
	6459.2	15.			0.8	2			Anv		97An09	
^{186}Bi(α)^{182}Tl	7759.6	20.5	7756	12	−0.2	4			Ara		97Ba21	*
	7754.7	15.4			0.1	4			Anv		03An27	*
^{186}Bi^m(α)^{182}Tl^p	7349.3	25.	7423	5	2.9	B			GSa		84Sc.A	
	7420.9	20.4			0.1	5			Ara		97Ba21	
	7422.9	5.1			−0.0	5			Anv		03An27	
^{186}Po(α)^{182}Pb	8493	30	8503	15	0.3	o					05Hu.A	
	8503.2	15.3				5					13An13	
^{186}W(p,t)^{184}W	−4474	5	−4464.0	1.2	2.0	U			Min		73Oo01	
^{186}W(p,t)^{184}W−^{184}W()^{182}W	660.1	1.6	656.2	1.2	−2.4	o					09Le03	
	657.0	1.8			−0.5	1	42	35 ^{186}W			09Le.A	
^{186}W(t,α)^{185}Ta	11430	20	11411	14	−1.0	R			LAl		80Lo10	
^{186}W(γ,n)^{185}W	−7120	60	−7192.0	1.2	−1.2	U			Phi		60Ge01	
^{186}W(d,t)^{185}W	−939	10	−934.8	1.2	0.4	U			Kop		72Ca01	
^{185}Re(n,γ)^{186}Re	6179.8	0.8	6179.59	0.05	−0.3	U			Tal		69La11	Z
	6178.6	1.5			0.7	U					70Or.A	
	6179.34	0.18			1.4	o			Bdn		06Fi.A	
	6179.59	0.05			0.0	1	100	72 ^{186}Re	Bdn		16Ma35	
^{185}Re(d,p)^{186}Re	3939	25	3955.02	0.05	0.6	U			Tal		69La11	
^{186}Ta(β^-)^{186}W	3901	60				2					69Mo16	*
^{186}Re(β^-)^{186}Os	1064	2	1072.7	0.8	4.3	B					56Jo05	
	1071.5	1.3			0.9	–					56Po28	
	1076	3			−1.1	–					64Ma36	
	1064	3			2.9	B					68An11	
ave.	1072.2	1.2			0.4	1	49	28 ^{186}Re			average	
^{186}Ir(β^+)^{186}Os	3760	200	3828	17	0.3	U					62Bo22	*
	3831	20			−0.2	R					63Em02	
^{186}Au(β^+)^{186}Pt	5950	200	6150	30	1.0	U					72We.A	
^{186}Hg(β^+)^{186}Au	3250	200	3175	24	−0.4	U					72We.A	
^{186}Tl^n(IT)^{186}Tl^m	373.9	0.5	374.20	0.10	0.6	o			Lvn		91Va04	
	374.0	0.2			1.0	o					Ens036	
	374.2	0.1				2			ISa		20St11	
*^{186}Ir−u	$M-A$=–39181(28) keV for mixture gs+m at 0.8 keV										Nub269	**
*^{186}Tl−u	$M-A$=–20030(180) keV for mixture gs+m+n at 44(15) and 418(15) keV										Nub269	**
*^{186}Tl−u	$M-A$=–19900(29) keV for mixture gs+m at 44(15) keV,										Nub269	**
*	n state can be resolved										Nub269	**
*^{186}Tl−^{133}Cs$_{1.398}$	D_M=110842.1(9.2) μu for mixture gs+m at 44(15) keV; $M-A$=–19874.4(8.6) keV										Nub269	**
*^{186}Tl−^{133}Cs$_{1.398}$	D_M=110840.4(8.6) μu for mixture gs+m at 20(40) keV; $M-A$=–19876.0(8.0) keV										Nub269	**
*^{186}Au(α)^{182}Ir	E_α=4653(15) to 3^- level at 152.3 keV										95Sa42	**
*^{186}Tl(α)^{182}Au	E_α=5760(51) to 129.4(0.1) keV										20St11	**
*^{186}Tl^m(α)^{182}Au^m	scattered events around E_α=5.76 Mev and $T_1/2>20$ s										77Ij01	**
*^{186}Bi(α)^{182}Tl	E_α=7158(20) followed by E(γ)=444(1) keV										03An27	**
*^{186}Bi(α)^{182}Tl	E_α=7152(15), 7085(15) followed by E(γ)=444(1), 520(1) keV										03An27	**
*^{186}Ta(β^-)^{186}W	E_{β^-}=2240(60) to (2^-,3^-) level at 1661.3817 keV										Ens036	**
*^{186}Ir(β^+)^{186}Os	E_{β^+}=2600(200) assumed to 2^+ level at 137.159 keV, also other E_{β^+}										Ens036	**
*^{186}Ir(β^+)^{186}Os	E_{β^+}=1940(20) to 6^+ level at 868.94 keV										Ens036	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{187}Ta−u		−39609	60				2			GS3	1.0	13Sh30	
^{187}Ir−u		−42458	30				2			GS2	1.0	05Li24	
^{187}Pt−u		−39500	110	−39384	26	1.1	U			GS1	1.0	00Ra23	
		−39413	30			1.0	1	74	74 ^{187}Pt	GS2	1.0	05Li24	
^{187}Au−u		−35470	114	−35459	24	0.1	U			GS1	1.0	00Ra23	*
		−35441	30			−0.6	1	64	64 ^{187}Au	GS2	1.0	05Li24	*
^{187}Hg−u		−30190	110	−30187	14	0.0	U			GS1	1.0	00Ra23	*
		−30155	36			−0.9	1	15	15 ^{187}Hg	GS2	1.0	05Li24	*
^{187}Hg−^{208}Pb$_{.899}$		−9210	20	−9196	14	0.7	1	48	48 ^{187}Hg	MA6	1.0	01Sc41	
^{187}Hg^{m}−^{208}Pb$_{.899}$		−9152	19	−9133	16	1.0	o			MA6	1.0	01Sc41	*
^{187}Tl−u		−24120	107	−24097	9	0.2	U			GS1	1.0	00Ra23	
		−23928	109			−1.6	U			GS2	1.0	05Li24	*
^{187}Tl^{m}−^{133}Cs$_{1.406}$		109151	24	109199	8	2.0	F			MA8	1.0	08We02	*
^{187}Pb−u		−16076	45	−16089	5	−0.3	U			GS2	1.0	05Li24	*
^{187}Pb−^{133}Cs$_{1.406}$		116843.5	5.9	116845	5	0.3	1	86	86 ^{187}Pb	MA8	1.0	05We11	*
^{187}Pb^{m}−^{133}Cs$_{1.406}$		116871.6	5.6	116866	12	−1.0	o			MA8	1.0	05We11	*
^{187}Re O$_2$−^{184}W ^{35}Cl		25797.4	3.5	25795.6	0.9	−0.2	U			H28	2.5	77Sh04	
^{187}Re ^{35}Cl−^{185}Re ^{37}Cl		5737	3	5744.1	0.9	0.9	U			H22	2.5	70Mc03	
		5744.2	1.2			−0.0	1	10	6 ^{185}Re	H28	2.5	77Sh04	
^{187}Re−^{187}Os		2.676	0.036	2.6508	0.0011	−0.7	U			SH1	1.0	14Ne15	
		2.6526	0.0014			−1.3	1	58	52 ^{187}Re	Hep	1.0	21Fi03	*
^{187}Au(α)^{183}Ir		4792.7	20.	4756	28	−0.7	1	27	15 ^{187}Au			68Si01	*
^{187}Hg(α)^{183}Pt		5229.9	20.	5229	14	−0.0	1	49	29 ^{183}Pt	ISa		70Ha18	*
^{187}Hg^{m}(α)^{183}Pt		5293.4	20.	5288	14	−0.3	1	52	27 ^{187}Hg^{m}	ISa		70Ha18	*
^{187}Tl^{m}(α)^{183}Au		5643	20	5656	6	0.6	–			ORa		76To06	*
		5661.5	10.			−0.6	–			GSa		80Sc09	*
		5645.1	12.			0.9	o			Lvn		85Co06	*
		5661.5	10.			−0.6	–			Lvn		91Wa21	*
	ave.	5659	7			−0.6	1	91	77 ^{183}Au			average	
^{187}Pb(α)^{183}Hg		6393.0	10.	6393	6	0.0	–			Ora		75Ca06	*
		6395.0	19.			−0.1	o			GSa		80Sc09	
		6398.4	10.			−0.5	–			GSa		81Mi12	*
	ave.	6396	7			−0.3	1	76	62 ^{183}Hg			average	
^{187}Pb^{m}(α)^{183}Hg^{m}		6213.1	20.	6212	3	−0.1	o			Ora		74Le02	
		6213.1	10.			−0.1	2			Ora		75Ca06	
		6223.3	10.			−1.1	o			GSa		80Sc09	
		6206.0	10.2			0.6	2			GSa		81Mi12	
		6202.9	15.			0.6	2			Anv		99An36	
		6213.2	4.1			−0.3	2			Ara		22Hu09	
^{187}Bi(α)^{183}Tl^{m}		7139.0	10.	7150	4	1.1	2			GSa		84Sc.A	
		7153.3	8.			−0.4	2			ORa		99Ba45	*
		7158.4	10.			−0.8	o			Anv		03An27	
		7147.2	8.			0.4	2			Jya		03Ke08	*
		7153.3	10.			−0.3	o			Anv		04An07	
		7153.3	5.			−0.6	2			Anv		06An11	*
^{187}Bi^{m}(α)^{183}Tl		7749.1	10.	7887	7	13.8	F			GSa		84Sc.A	*
		7890.1	15.			−0.2	2			ORa		99Ba45	
		7882.9	11.			0.4	2			Jya		03Ke08	
		7890.1	10.			−0.3	2			Anv		06An11	
^{187}Po(α)^{183}Pb		7978.9	15.				3			Anv		06An11	*
^{187}Po^{m}(α)^{183}Pb^{m}		7889.1	20.				3			Anv		06An11	
^{186}W(n,γ)^{187}W		5466.3	0.3	5466.76	0.04	1.5	U			BNn		87Br05	Z
		5467.22	0.3			−1.5	U			Ltn		92Be17	*
		5466.59	0.12			1.4	o			Bdn		06Fi.A	
		5466.83	0.05			−1.4	–			Prn		08Bo26	
		5466.62	0.07			2.0	–			Bdn		14Hu02	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{186}W(d,p)^{187}W		3236	5	3242.19	0.04	1.2	U			ANL		65Er03	
		3240	10			0.2	U			Kop		72Ca01	
^{186}W(n,γ)^{187}W	ave.	5466.76	0.04	5466.76	0.04	0.0	1	100	55 ^{186}W			average	
^{186}W(^{3}He,d)^{187}Re		530	40	503.4	1.1	−0.7	U			Roc		71Lu01	
^{187}Re(γ,n)^{186}Re		−7180	80	−7360.5	0.9	−2.3	U			Phi		60Ge01	
^{187}Re(d,t)^{186}Re		−1055	25	−1103.3	0.9	−1.9	U			Tal		69La11	
^{186}Os(n,γ)^{187}Os		6291.1	1.0	6290.3	0.5	−0.8	–					74Pr15	Z
		6289.4	0.8			1.1	–			Bdn		06Fi.A	
	ave.	6290.1	0.6			0.4	1	70	39 ^{186}Os			average	
^{187}W(β^-)^{187}Re		1314	2	1312.5	1.1	−0.8	–					69Na03	
		1310	2			1.2	–					70He14	
	ave.	1312.0	1.4			0.3	1	63	55 ^{187}W			average	
^{187}Re(β^-)^{187}Os		2.62	0.09	2.4692	0.0010	−1.7	U					65Br12	
		2.64	0.05			−3.4	B					67Hu05	
		2.667	0.020			−9.9	B					92Co23	
		2.70	0.09			−2.6	U					93As02	
		2.460	0.011			0.8	U					99Al20	
		2.470	0.004			−0.2	–					01Ga01	*
		2.4661	0.0017			1.8	–					03Ar36	
	ave.	2.4667	0.0016			1.6	1	40	36 ^{187}Re			average	
^{187}Ir(β^+)^{187}Os		1550	200	1670	28	0.6	U					71Ma24	*
^{187}Os(^{3}He,t)^{187}Ir		−1521	6	−1688	28	−27.9	B			INS		90Ka27	
^{187}Au(β^+)^{187}Pt		3600	40	3656	27	1.4	1	47	26 ^{187}Pt			83Gn01	*
^{187}Hg^m(IT)^{187}Hg		54	21	58	14	0.2	1	46	28 ^{187}Hg^m	MA6		01Sc41	*
^{187}Pb^m(IT)^{187}Pb		33	13	19	10	−1.1	1	61	61 ^{187}Pb^m	MA8		05We11	
*^{187}Au−u	$M-A$=–32980(100) keV for mixture gs+m at 120.33 keV											Nub269	**
*^{187}Au−u	Observed for 10 minutes: no isomeric contamination											03Li.A	**
*^{187}Hg−u	$M-A$=–28090(100) keV for mixture gs+m at 58(14) keV											Nub269	**
*^{187}Hg−u	$M-A$=–28060(28) keV for mixture gs+m at 58(14) keV											Nub269	**
*^{187}Hg^m−^{208}Pb$_{.899}$	Use instead their difference between ground state and ^{187}Hg^m lines											GAu	**
*^{187}Tl−u	$M-A$=–22121(28) keV for mixture gs+m at 337(4) keV											Nub269	**
*^{187}Tl^m−^{133}Cs$_{1.406}$	F : contamination from ground state not resolved											08We02	**
*^{187}Pb−u	$M-A$=–14965(41) keV for mixture gs+m at 19(10) keV											Nub269	**
*^{187}Pb−^{133}Cs$_{1.406}$	D_M=116851.3(4.3) μu for mixture gs+m at 19(10) keV with R=0.62(0.02);											Nub269	**
*	$M-A$=–14981.5(4.0) keV											GAu	**
*^{187}Pb^m−^{133}Cs$_{1.406}$	D_M=116869.5(5.5) for mixture gs+m at 19(10) R=8.7(0.7); $M-A$=–14964.5(5.1) keV											Nub269	**
*	used are only the equations for the ^{187}Pb doublet and ^{187}Pb^m(IT)^{187}Pb											GAu	**
*^{187}Au(α)^{183}Ir	Assignment uncertain											Ens095	**
*^{187}Hg(α)^{183}Pt	E_α=5035(20) to 3/2$^-$ level at 84.73 keV											Ens164	**
*^{187}Hg^m(α)^{183}Pt	E_α=4870(20) to (13/2$^+$) level at 316.9 level											Ens164	**
*^{187}Tl^m(α)^{183}Au	E_α=5510(20) 5528(10) 5512(12) 5528(10) respectively, to (9/2)$^-$ at 12.4(0.4)keV											Ens164	**
*^{187}Pb(α)^{183}Hg	E_α=6190(10) to 3/2$^-$ level at 67.16 keV											Ens164	**
*^{187}Pb(α)^{183}Hg	E_α=6194(10) 5993(10) to 3/2$^-$ levels at 67.16 and 275.33 keV											Ens164	**
*^{187}Bi(α)^{183}Tl^m	Also E_α=7612(15) keV to ground state											99Ba45	**
*^{187}Bi(α)^{183}Tl^m	Also E_α=7605(16) keV to ground state											03Ke08	**
*^{187}Bi(α)^{183}Tl^m	Also E_α=7612(5), 7342(15) keV to ground state, 273(1) keV											06An11	**
*^{187}Bi^m(α)^{183}Tl	F : for T=700 μs instead of Nubase=370(20) μs											Nub269	**
*^{187}Po(α)^{183}Pb	E_α=7528(15) to 286(1) keV level; also 1 event E_α=7796(15) to ground state											06An11	**
*^{186}W(n,γ)^{187}W	Only statistical error 0.04 keV given; Z recalibrated											GAu	**
*^{187}Re(β^-)^{187}Os	supersedes 2.481(0.006) of same group											98Ga50	**
*^{187}Ir(β^+)^{187}Os	p$^+$ <0.15(0.05), resulting Q <1550 keV											Ens095	**
*^{187}Au(β^+)^{187}Pt	K/β^+=31.6(2.8) to 1/2$^+$ level at 1341.07 keV, recalculated											Ens095	**
*^{187}Hg^m(IT)^{187}Hg	Original error (7 keV) increased by 20 due to isomer+ground state lines in trap											01Sc41	**
^{188}Ta−u		−36084	59	−36400#	220#	−5.4	D			GS3	1.0	13Sh30	*
^{188}Au−u		−34750	104	−34752.0	2.9	−0.0	U			GS1	1.0	00Ra23	
		−34674	30			−2.6	B			GS2	1.0	05Li24	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{188}Au−^{133}Cs$_{1.414}$		98938.9	2.9				2			MA8	1.0	17Ma29	
^{188}Hg−u		−32500	104	−32420	7	0.8	U			GS1	1.0	00Ra23	
		−32428	30			0.3	1	6	6 ^{188}Hg	GS2	1.0	05Li24	
^{188}Hg−^{208}Pb$_{.904}$		−11330	20	−11312	7	0.9	–			MA6	1.0	01Sc41	
	ave.	−11317	17			0.3	1	19	19 ^{188}Hg			average	
^{188}Tl−u		−23827	110	−23980	30	−1.4	U			GS1	1.0	00Ra23	*
		−23994	38			0.4	2			GS2	1.0	05Li24	*
^{188}Pb−u		−19070	110	−19122	11	−0.5	U			GS1	1.0	00Ra23	
		−19144	30			0.7	R			GS2	1.0	05Li24	
^{188}W−^{190}Os$_{.989}$		−484	76	−414	3	0.9	U			RKM	1.0	25Mu03	
^{188}Re−^{190}Os$_{.989}$		−814	25	−788.9	0.5	1.0	U			RKM	1.0	25Mu03	
^{188}Os ^{35}Cl−^{186}W ^{37}Cl		4426	3	4422.3	1.2	−0.5	U			H22	2.5	70Mc03	
^{188}Os−^{190}Os$_{.989}$		−3067	53	−3065.3	0.5	0.0	U			RKM	1.0	25Mu03	
^{188}Au−^{190}Os$_{.989}$		6298	87	6345.4	3.0	0.5	U			RKM	1.0	25Mu03	
^{188}Pt(α)^{184}Os		4015.7	10.	4007	5	−0.9	–					63Gr08	
		4000.3	10.			0.6	–			ORa		78El11	
		3990.1	15.			1.1	–			ISa		79Ha10	
	ave.	4005	7			0.3	1	65	65 ^{188}Pt			average	
^{188}Hg(α)^{184}Pt		4710.4	20.	4709	15	−0.1	1	52	47 ^{184}Pt	ISa		79Ha10	
^{188}Pb(α)^{184}Hg		6110.3	10.	6109	3	−0.1	2			Ora		74Le02	Z
		6109.2	10.			−0.0	2			Ora		77De32	Z
		6120.5	15.			−0.8	2			GSa		80Sc09	Z
		6110.5	5.			−0.3	2			ORa		81To02	Z
		6110.3	10.3			−0.1	o			Lvn		93Wa.A	
		6109.3	10.			−0.0	2			Lvn		93Wa03	Z
		6100.0	8.			1.1	2			Jya		03Ke04	
^{188}Bi(α)^{184}Tl		7274.5	25.	7262	3	−0.5	o			GSa		80Sc09	*
		7279.7	10.			−1.8	2			GSa		84Sc.A	*
		7255.2	7.			1.0	2			Lvn		97Wa05	*
		7259.3	5.			0.5	o			Anv		03An26	*
		7264.8	10.			−0.3	2			Anv		06An04	*
		7260.7	4.1			0.3	2			Ara		22Zh46	*
^{188}Bi^n(α)^{184}Tl^m		7462.9	5.				5			Anv		03An26	*
^{188}Bi^n(α)^{184}Tl^n		6968.5	20.	6964	3	−0.2	o			GSa		80Sc09	
		6968.5	10.			−0.5	5			GSa		84Sc.A	
		6963.5	6.			0.1	5			Lvn		97Wa05	
		6961.3	5.			0.5	o			Anv		03An26	
		6963.5	10.2			0.0	5			Anv		06An04	
		6963.5	4.1			0.1	5			Ara		22Zh46	
^{188}Po(α)^{184}Pb		8087.4	25.	8077	11	−0.4	o			Anv		99An52	
		8080.2	15.			−0.2	o			Anv		01Va.B	
		8082.3	15.			−0.3	2			Anv		03Va16	
		8071.1	16.3			0.4	2			Ara		22Zh46	
^{188}Os(p,t)^{186}Os		−5802	5	−5798.1	0.5	0.8	U			Min		73Oo01	
		−5803	4			1.2	U			McM		75Th04	
^{187}Re(n,γ)^{188}Re		5871.77	0.3	5871.65	0.04	−0.4	U					72Sh13	Z
		5871.75	0.13			−0.8	U			Bdn		06Fi.A	
		5871.65	0.04				2			Prn		10Ba48	
^{188}Os(t,α)^{187}Re		12604	10	12604.14	0.15	0.0	U			McM		76Hi08	
^{187}Os(n,γ)^{188}Os		7989.6	0.3	7989.60	0.15	0.0	–					83Fe06	Z
		7989.58	0.17			0.1	–			Bdn		06Fi.A	
	ave.	7989.58	0.15			0.1	1	98	57 ^{187}Os			average	
^{188}At(p)^{187}Po		1510	40	2080#	300#	14.2	D			Jyp		25Ko13	*
^{188}W(β^-)^{188}Re		349	3				3					64Bu10	
^{188}Re(β^-)^{188}Os		2116	2	2120.42	0.15	2.2	U					56Jo05	
		2111	3			3.1	B					68An11	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{188}Ir(β^+)^{188}Os		2833	10	2792	9	−4.1	B					62Wa20	*
		2781	20			0.6	–					69Ya02	*
		2827	30			−1.2	–					70Ag03	*
	ave.	2795	17			−0.2	1	32	32 ^{188}Ir			average	
^{188}Pt(ε)^{188}Ir		525	10	524	9	−0.1	1	75	68 ^{188}Ir	ORa		78El11	*
^{188}Au(β^+)^{188}Pt		5520	30	5450	6	−2.3	U					84Da.A	
^{188}Hg(β^+)^{188}Au		2040	20	2173	7	6.6	B					84Da.A	
*^{188}Ta−u	Trends in Mass Surface TMS suggest ^{188}Ta 300 keV more bound											GAu	**
*^{188}Tl−u	$M-A$=–22180(100) keV for mixture gs+m at 30(40) keV											GAu	**
*^{188}Tl−u	$M-A$=–22335(28) keV for mixture gs+m at 30(40) keV											GAu	**
*^{188}Bi(α)^{184}Tl	E_α=7005(25) 7010(10) 6987(6) respectively, to (3^+) level at 117.5(0.5) keV											Ens102	**
*	E_α=7029(7) 3 times weaker exists too, possible mixture in older results											97Wa05	**
*^{188}Bi(α)^{184}Tl	E_α=7106(5), 6992(5), 6889(10) to ground state, 117.5, 216 levels											03An26	**
*^{188}Bi(α)^{184}Tl	E_α=6995(10) to 117.5 level											06An04	**
*^{188}Bi(α)^{184}Tl	E_α=6991(4) to 117.5 level											22Zh46	**
*^{188}Bi^n(α)^{184}Tl^m	E_α=7302(5), 7232(10), 6995(15) to ground state, 70.5, 320 levels											03An26	**
*^{188}At(p)^{187}Po	Trends in Mass Surface TMS suggest ^{188}At 570 keV less bound											HWJ261	**
*^{188}Ir(β^+)^{188}Os	E_{β^+}=1656(10) 1605(20) 1650(30) respectively, to 2^+ level at 155.021 keV											Ens024	**
*^{188}Pt(ε)^{188}Ir	pL=0.67(0.05) to 1^+ level at 478.17 keV											Ens024	**
^{189}W−u		−38237	43	−38260	40	−0.6	1	67	67 ^{189}W	GS3	1.0	13Sh30	
C_{14} H_{21}−^{189}Os		206188.3	6.2	206179.7	0.7	−0.3	U			M23	4.0	79Ha32	
^{189}Au−u		−36080	140	−36052	22	0.2	U			GS1	1.0	00Ra23	*
		−36045	31			−0.2	2			GS2	1.0	05Li24	
		−36058	30			0.2	2			GS2	1.0	05Li24	*
^{189}Hg−u		−31788	111	−31810	30	−0.2	U			GS1	1.0	00Ra23	*
		−31791	42			−0.4	1	65	65 ^{189}Hg	GS2	1.0	05Li24	*
^{189}Hg^m−^{208}Pb$_{.909}$		−10501	20	−10498	19	0.2	1	92	92 ^{189}Hg^m	MA6	1.0	01Sc41	
^{189}Tl−u		−26497	139	−26426	9	0.5	U			GS1	1.0	00Ra23	*
		−26313	93			−1.2	U			GS2	1.0	05Li24	*
^{189}Pb−u		−19206	97	−19157	15	0.5	U			GS1	1.0	00Ra23	*
		−19193	34			1.1	1	20	20 ^{189}Pb	GS2	1.0	05Li24	*
^{189}Os ^{35}Cl−^{187}Re ^{37}Cl		5341	3	5343.9	0.5	0.4	U			H22	2.5	70Mc03	
^{189}W−^{190}Os$_{.995}$		3030	61	3080	40	0.9	1	33	33 ^{189}W	RKM	1.0	25Mu03	
^{189}Re−^{190}Os$_{.995}$		587	32	575	9	−0.4	U			RKM	1.0	25Mu03	
^{189}Os−^{190}Os$_{.995}$		−551	74	−507.27	0.20	0.6	U			RKM	1.0	25Mu03	
^{189}Ir−^{190}Os$_{.995}$		51	27	65	12	0.5	1	20	20 ^{189}Ir	RKM	1.0	25Mu03	
^{189}Pt−^{190}Os$_{.995}$		2181	34	2193	10	0.4	1	9	9 ^{189}Pt	RKM	1.0	25Mu03	
^{189}Pb(α)^{185}Hg		5954.2	10.	5915	4	−3.9	B			Ora		72Ga27	*
		5943.9	10.			−2.9	B			Ora		74Le02	*
		5915	10			−0.0	–					05Fr.A	*
		5914.8	5.4			−0.0	–			ISa		13Sa43	*
	ave.	5915	5			−0.0	1	82	67 ^{189}Pb			average	
^{189}Pb^m(α)^{185}Hg		5958	10	5955	5	−0.3	1	28	25 ^{189}Pb^m			05Fr.A	*
		5955	5			0.0	o			ISa		13Sa43	*
^{189}Bi(α)^{185}Tl		7269.2	10.	7268.7	2.8	−0.0	7			Ora		74Le02	*
		7274.2	10.2			−0.5	7			GSa		84Sc.A	*
		7270.7	5.1			−0.4	7			Lvn		85Co06	*
		7271.8	15.			−0.2	U			Anv		97An09	*
		7268.1	6.1			0.1	7			Lvn		97Wa05	*
		7266.5	15.3			0.1	7			Anv		99An52	*
		7266.0	5.			0.5	o			Jya		02Hu14	*
		7266.7	4.1			0.5	o			Jya		03Ke08	*
		7266	5			0.5	7			Jya		07DoZW	*
^{189}Bi^m(α)^{185}Tl		7362.1	20.	7448	4	1.6	U			GSa		84Sc.A	*
		7499.0	30.			−1.7	U			Dbb		93An19	
		7458.2	40.			−0.3	U			ORa		95Ba75	
		7458.2	15.			−0.7	o			Anv		97An09	
		7450.0	6.			−0.3	7			Ara		97Wa05	
		7456.1	10.2			−0.8	7			Anv		99An52	
		7453.1	6.			−1.0	o			Jya		03Ke08	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{189}Bi$^{m}(\alpha)^{185}$Tl	7444.9	5.1	7448	4	0.6	7			Jya		07DoZW	
^{189}Po$(\alpha)^{185}$Pb	7699.4	15.	7694	15	−0.3	o			Anv		99An52	*
	7694.3	15.				3			Anv		05Va04	*
^{189}Os(p,t)^{187}Os	−5431	5	−5428.6	0.5	0.5	U			Min		73Oo01	
	−5432	4			0.8	U			McM		75Th04	
^{188}Os(n,γ)^{189}Os	5920.8	2.	5920.8	0.4	0.0	U					76Be50	
	5920.6	0.5			0.4	1	80	59 ^{188}Os	ILn		92Br17	
	5922.0	0.4			−3.0	C			Bdn		06Fi.A	
^{188}Os(d,p)^{189}Os	3689	10	3696.2	0.4	0.7	U			Kop		75Mo29	
^{189}Os(d,t)^{188}Os	335	15	336.4	0.4	0.1	U			Tal		75Th06	
^{189}W$(\beta^-)^{189}$Re	2500	200	2340	30	−0.8	U					65Ka07	
^{189}Re$(\beta^-)^{189}$Os	1000	20	1008	8	0.4	R					63Cr06	
	1015	20			−0.4	R					65Bl06	
^{189}Pt$(\beta^+)^{189}$Ir	1950	20	1982	13	1.6	1	41	25 ^{189}Ir			71Pl08	*
^{189}Au$(\beta^+)^{189}$Pt	3160	300	2889	22	−0.9	U					75Un.A	
^{189}Hg$(\beta^+)^{189}$Au	4200	200	3960	40	−1.2	U					75Un.A	
^{189}Hg^{m}(IT)^{189}Hg	100	50	80	30	−0.4	1	43	35 ^{189}Hg	MA6		01Sc41	
^{189}Tl$^{m}(\beta^+)^{189}$Hg	5460	200	5300	30	−0.8	U					75Un.A	*
^{189}Pb^{m}(IT)^{189}Pb	40	4	40	4	0.1	1	88	75 ^{189}Pb^{m}	ISa		13Sa43	
*^{189}Au−u	$M-A$=–33490(100) keV for mixture gs+m at 247.25 keV										Nub269	**
*^{189}Au−u	$M-A$=–33341(28) keV for ^{189}Au^{m} at 247.25 keV										Nub269	**
*^{189}Hg−u	$M-A$=–29570(100) keV for mixture gs+m at 80(30) keV										Nub269	**
*^{189}Hg−u	$M-A$=–29573(28) keV for mixture gs+m at 80(30) keV										Nub269	**
*^{189}Tl−u	$M-A$=–24540(100) keV for mixture gs+m at 285(6) keV										Nub269	**
*^{189}Tl−u	$M-A$=–24369(28) keV for mixture gs+m at 285(6) keV										Nub269	**
*^{189}Pb−u	$M-A$=–17870(90) keV for mixture gs+m at 40(4) keV										Nub269	**
*^{189}Pb−u	$M-A$=–17858(29) keV for mixture gs+m at 40(4) keV										Nub269	**
*^{189}Pb$(\alpha)^{185}$Hg	E_α=5730.1(10,Z) possibly from ground state, to 3/2$^-$ level at 26.1 keV										Ens061	**
*^{189}Pb$(\alpha)^{185}$Hg	E_α=5720(10) possibly from ground state, to 3/2$^-$ level at 26.1 keV										Ens061	**
*^{189}Pb$(\alpha)^{185}$Hg	E_α=5761 to 3/2$^-$ level at 26.1 and E_α=5623 to 173.8 level										05Fr.A	**
*^{189}Pb$(\alpha)^{185}$Hg	E_α=5764(3)(5) to 3/2$^-$ level at 26.02 and E_α=5619(2)(5) to 173.9 level										13Sa43	**
*^{189}Pb$^{m}(\alpha)^{185}$Hg	E_α=5730 to 103.8 level										05Fr.A	**
*^{189}Pb$^{m}(\alpha)^{185}$Hg	E_α=5727(2)(5) to 103.7(0.4) level										13Sa43	**
*^{189}Pb$^{m}(\alpha)^{185}$Hg	use instead ^{189}Pb^{m}(IT)^{189}Pb where systematic errors cancel										GAu	**
*^{189}Bi$(\alpha)^{185}$Tl	E_α=6670.1(10,Z) to ^{185}Tl^{m} at 454.70(0.22) keV										Nub269	**
*^{189}Bi$(\alpha)^{185}$Tl	E_α=6675(10) to ^{185}Tl^{m} at 454.70(0.22) keV										Nub269	**
*^{189}Bi$(\alpha)^{185}$Tl	E_α=7115.6(15,Z) and 6671.6(5,Z) to ground state and ^{185}Tl^{m} at 454.70(0.22) keV										Nub269	**
*^{189}Bi$(\alpha)^{185}$Tl	E_α=7120(15) and 6670(15) to ground state and ^{185}Tl^{m} at 454.70(0.22) keV										Nub269	**
*^{189}Bi$(\alpha)^{185}$Tl	E_α=7114(6) and 6833(7) to ground state and 284.5(0.2) keV level in ^{185}Tl										Fgk261	**
*^{189}Bi$(\alpha)^{185}$Tl	E_α=6834(15) to 284.5(0.2) keV level in ^{185}Tl										Fgk261	**
*^{189}Bi$(\alpha)^{185}$Tl	E_α=6667(4) to ^{185}Tl^{m} at 454.70(0.22) keV										Nub269	**
*^{189}Bi$(\alpha)^{185}$Tl	E_α=7114(6) and 6667(4) to ground state and ^{185}Tl^{m} at 454.70(0.22) keV										Nub269	**
*^{189}Bi$(\alpha)^{185}$Tl	E_α=7107(5) and 6672(5) to ground state and ^{185}Tl^{m} at 454.70(0.22) keV										Nub269	**
*^{189}Bi$^{m}(\alpha)^{185}$Tl	Only one event; not seen in reference										93An19	**
*^{189}Po$(\alpha)^{185}$Pb	E_α=7264(15) to 278(1) level										99An52	**
*^{189}Po$(\alpha)^{185}$Pb	E_α=7259(15) to 278(1) level										05Va04	**
*^{189}Pt$(\beta^+)^{189}$Ir	E_{β^+}=885(10) to ground state, 1/2$^+$ level at 94.34 and 3/2$^+$ at 176.53 keV										Ens039	**
*^{189}Tl$^{m}(\beta^+)^{189}$Hg	E_{β^+}=4140(200) to several levels around 300 keV										75Un.A	**
^{190}W−u	−36917	44	−36920	30	−0.2	1	56	56 ^{190}W	GS3	1.0	13Sh30	
^{190}Ir−u	−39479	188	−39456.6	1.5	0.1	U			GS1	1.0	97Sc.B	*
^{190}Au−u	−35300	130	−35248	4	0.4	U			GS2	1.0	05Li24	*
^{190}Au−^{133}Cs$_{1.429}$	99860.9	3.7				2			MA8	1.0	17Ma29	
^{190}Hg−u	−33670	107	−33680	17	−0.1	U			GS1	1.0	00Ra23	
^{190}Hg−^{208}Pb$_{.913}$	−12361	20	−12362	17	−0.0	1	71	71 ^{190}Hg	MA6	1.0	01Sc41	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$^{190}Tl^{m}-u$		−26055	107	−26083	6	−0.3	o			GS1	1.0	00Ra23	*
		−26048	30			−1.2	U			GS2	1.0	05Li24	*
$^{190}Tl^{m}-^{133}Cs_{1.429}$		109033.5	6.9	109026	6	−1.1	1	64	64 $^{190}Tl^{m}$	MA8	1.0	14Bo26	
$^{190}Pb-u$		−21940	104	−21919	13	0.2	U			GS1	1.0	00Ra23	
		−21905	30			−0.5	R			GS2	1.0	05Li24	
$^{190}Bi^{m}-^{133}Cs_{1.429}$		123800	27	123879	20	2.9	F			MA8	1.0	08We02	*
$^{186}Os-^{190}Pt_{.979}$		−6953.13	0.86	−6953.3	0.6	−0.2	1	53	39 ^{186}Os	MS1	1.0	16Ei01	
$^{190}Os\ ^{35}Cl-^{188}Os\ ^{37}Cl$		5557	3	5558.3	0.5	0.2	U			H22	2.5	70Mc03	
$^{190}W-^{190}Os$		4549	65	4630	30	1.3	1	25	25 ^{190}W	RKM	1.0	25Mu03	
$^{190}Re-^{190}Os$		3395	49	3355	5	−0.8	U			RKM	1.0	25Mu03	
$^{190}Ir-^{190}Os$		2093	19	2097.9	1.3	0.3	U			RKM	1.0	25Mu03	
$^{190}Os-^{190}Pt$		−1504.31	0.59	−1504.4	0.5	−0.1	1	62	33 ^{190}Pt	MS1	1.0	16Ei01	
$^{190}Pt-^{190}Os$		1504	74	1504.4	0.5	0.0	U			RKM	1.0	25Mu03	
$^{190}Au-^{190}Os$		6280	27	6306	4	1.0	U			RKM	1.0	25Mu03	
$^{190}Os-C_{14}\ H_{21}$		−205897.8	5.8	−205880.2	0.7	0.8	U			M23	4.0	79Ha32	
$^{190}Os-^{189}Os$		285.2	5.2	299.49	0.20	1.1	U			M24	2.5	79Ha32	
$^{190}Pt(\alpha)^{186}Os$		3238.3	20.	3268.6	0.6	1.5	U					61Pe23	
		3248.5	20.			1.0	U					63Gr08	
$^{190}Pb(\alpha)^{186}Hg$		5699.8	10.	5698	5	−0.2	2			Ora		74Le02	Z
		5697.0	5.			0.1	2			ORa		81El03	Z
$^{190}Bi(\alpha)^{186}Tl$		6862.2	5.	6862	3	−0.0	–			Lvn		91Va04	*
		6863.3	5.			−0.3	–			Anv		03An26	*
		6860.3	6.			0.3	–			Jya		13Ny01	*
	ave.	6862	3			−0.0	1	99	83 ^{190}Bi			average	
$^{190}Bi^{m}(\alpha)^{186}Tl^{n}$		6589	10	6592.8	2.8	0.4	–			Ora		74Le02	
		6595.1	5.1			−0.4	–			Lvn		91Va04	*
		6595.1	5.1			−0.4	–			Anv		03An26	*
		6589	5			0.8	–			Jya		13Ny01	
	ave.	6592.7	2.8			0.0	1	99	66 $^{186}Tl^{n}$			average	
$^{190}Po(\alpha)^{186}Pb$		7643.2	20.	7693	7	2.5	C			GSa		88Qu.A	
		7651.4	40.			1.0	U			ORa		96Ba35	
		7691.2	10.			0.2	3			ORa		97Ba25	
		7695.3	10.			−0.2	3			Anv		00An14	*
$^{190}At(\alpha)^{186}Bi$		7917	20	7910	60	−0.0	5			Jya		23Ko10	*
		7912.9	10.2			0.0	5			Anv		23An09	*
$^{190}At^{m}(\alpha)^{186}Bi^{m}$		7909.8	10.2	7910	60	0.1	6			Anv		23An09	*
		7917.0	20.4			−0.1	6			Jya		23Ko10	*
$^{190}Os(p,t)^{188}Os$		−5234	5	−5231.4	0.5	0.5	U			Min		73Oo01	
		−5237	4			1.4	U			McM		75Th04	
$^{190}Pt(p,t)^{188}Pt$		−7150	10	−7146	5	0.4	1	28	28 ^{188}Pt	Ors		78Ve10	
$^{190}Os(t,\alpha)^{189}Re$		11796	10	11796	8	0.0	2			McM		76Hi08	
$^{189}Os(n,\gamma)^{190}Os$		7791.8	1.0	7792.34	0.19	0.5	U			BNn		79Ca02	Z
		7792.31	0.19			0.2	1	97	79 ^{189}Os	Bdn		06Fi.A	
$^{190}Os(d,t)^{189}Os$		−1541	10	−1535.11	0.19	0.6	U			Kop		75Mo29	
		−1530	4			−1.3	U			Tal		76Be50	
$^{190}Pt(p,d)^{189}Pt$		−6693	11	−6682	10	1.0	1	75	75 ^{189}Pt	Ors		80Ka19	
$^{190}W(\beta^{-})^{190}Re$		1270	70	1190	30	−1.2	1	19	19 ^{190}W			76Ha39	*
$^{190}Re(\beta^{-})^{190}Os$		3090	300	3125	5	0.1	U					55At21	*
		3190	300			−0.2	U					69Ha44	*
		3146	200			−0.1	U					64Fl02	*
$^{190}Ir(\beta^{+})^{190}Os$		2000	200	1954.2	1.2	−0.2	U					60Ka14	*
$^{190}Au(\beta^{+})^{190}Pt$		4442	15	4473	4	2.1	U					73Jo11	
		4380	55			1.7	U					74Di.A	
		4380	200			0.5	U					75Un.A	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{190}Hg(β^+)^{190}Au	2105	80	1461	16	−8.1	C						74Di.A	
^{190}Tl(β^+)^{190}Hg	7000	400	7006	17	0.0	U						75Un.A	*
^{190}Tl^m(β^+)^{190}Hg	6975	300	7076	16	0.3	U						76Bi09	*
^{190}Bi(β^+)^{190}Pb	8700	500	9806	19	2.2	F						76Bi09	*
*^{190}Ir−u	$M-A$=–36761(175) keV for mixture gs+m at 26.1 keV											Nub269	**
*^{190}Au−u	$M-A$=–32701(28) keV for mixture gs+m at 358#100 keV											Nub269	**
*^{190}Tl^m−u	Assumed by evaluator to be the 7^+ excited isomer											GAu	**
*^{190}Bi$^m-^{133}$Cs$_{1.429}$	F : contamination due to ground state not resolved											08We02	**
*^{190}Bi(α)^{186}Tl	E_α=6716(5), 6507(5), 6431(5) to ground state, 215.2, 293.7 levels											91Va04	**
*^{190}Bi(α)^{186}Tl	E_α=6431(5) to 293.7 level											03An26	**
*^{190}Bi(α)^{186}Tl	E_α=6428(6) to 293.7 level											13Ny01	**
*^{190}Po(α)^{186}Pb	E_α=7545(15) same dataset as in reference 2000An14											97An09	**
*^{190}W(β^-)^{190}Re	E_{β^-}=950(70) to 1^+ level at 319.7 keV											Ens036	**
*^{190}Re(β^-)^{190}Os	E_{β^-}=1700(300) 1800(300) respectively, to 3^- level at 1387.00 keV											Ens036	**
*^{190}Re(β^-)^{190}Os	E_{β^-}=1600(200) from isomer at 204(10) to several levels around 1750 keV											Nub269	**
*^{190}Ir(β^+)^{190}Os	p^+ =6(1)×10^{-5} to 4^+ levels at 1163.19 and 955.37 keV, level at 1872.15 keV fed											Ens036	**
*^{190}Tl(β^+)^{190}Hg	E_{β^+}=5700(400) to ground state and 2^+ level at 416.32 keV											Ens036	**
*^{190}Tl^m(β^+)^{190}Hg	E_{β^+}=4180(300) to 6^+ level at 1772.94 keV											Ens036	**
*^{190}Bi(β^+)^{190}Pb	F : E_{β^+}=5700(300) to a level around 2000 at least											AHW	**
^{191}W−u	−33469	45				2				GS3	1.0	13Sh30	
^{191}Au−u	−36180	88	−36284	5	−1.2	U				GS2	1.0	05Li24	*
^{191}Au−^{133}Cs$_{1.436}$	99487.3	5.3	99487	5	0.0	1	100	100	^{191}Au	MA8	1.0	13Kr15	
^{191}Hg−u	−32811	51	−32843	24	−0.6	1	22	22	^{191}Hg	GS2	1.0	05Li24	*
^{191}Hg−^{208}Pb$_{.918}$	−11414	29	−11408	24	0.2	1	68	68	^{191}Hg	MA6	1.0	01Sc41	*
^{191}Tl−u	−28340	130	−28216	8	1.0	U				GS1	1.0	00Ra23	*
	−28234	30			0.6	U				GS2	1.0	05Li24	
	−28192	31			−0.8	U				GS2	1.0	05Li24	*
^{191}Pb−u	−21770	110	−21785	7	−0.1	U				GS1	1.0	00Ra23	*
	−21744	35			−1.2	U				GS2	1.0	05Li24	*
^{191}Bi−^{133}Cs$_{1.436}$	121552.1	8.6	121558	8	0.7	1	87	87	^{191}Bi	MA8	1.0	08We02	
^{191}Pb^m(α)^{187}Hg^m	5403.4	20.	5402	15	−0.1	1	51	44	^{187}Hg^m	Ora		74Le02	
^{191}Bi(α)^{187}Tl	6780.8	5.	6782	4	0.3	2				Lvn		85Co06	Z
	6785.3	10.2			−0.3	2				ORa		98Bi.A	
	6782.3	10.2			−0.0	2				Anv		99An36	
	6783.3	7.2			−0.2	2				Jya		13Ny01	
^{191}Bi(α)^{187}Tl^m	6440.0	5.	6445.6	2.7	1.1	–						67Tr06	Z
	6455.0	10.			−0.9	U				Ora		74Le02	Z
	6445.9	5.			−0.1	–				Lvn		85Co06	Z
	6447	10			−0.1	U				ORa		98Bi.A	
	6458.5	20.			−0.6	U				RIa		99Ta20	
	6445	10			0.1	U				Anv		99An36	
	6443.2	3.			0.8	o				Jya		03Ke04	
	6445.2	10.			0.0	U				Jya		13Uu01	
	6450.3	4.			−1.2	–				Jya		13Ny01	
ave.	6446.2	2.7			−0.2	1	99	86	^{187}Tl^m			average	
^{191}Bi^m(α)^{187}Tl	7022.8	5.	7023	3	−0.0	3				Lvn		85Co06	Z
	7023.4	10.			−0.1	U				ORa		98Bi.A	
	7016.2	20.			0.3	U				RIa		99Ta20	
	7017.2	3.			1.7	o				Jya		03Ke04	
	7031.0	15.			−0.6	U				Jya		13Uu01	*
	7022.4	4.			0.0	3				Jya		13Ny01	
	6963	34			1.8	U				Lza		24Wa14	
^{191}Po(α)^{187}Pb	7470.8	20.	7493	5	1.1	F				GSa		93Qu03	*
	7487.1	15.			0.4	U				ORa		97Ba25	
	7491.2	5.			0.4	1	94	94	^{191}Po	Anv		02An19	*
^{191}Po(α)^{187}Pb^m	7493.2	15.	7474	10	−1.2	1	45	39	^{187}Pb^m	Anv		02An19	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{191}Po$^m(\alpha)^{187}$Pb^m		7535	5				2			Anv		02An19	*
^{191}At$(\alpha)^{187}$Bi^m		7713.9	11.				3			Jya		03Ke08	
^{191}At$^m(\alpha)^{187}$Bi		7880.4	15.				3			Jya		03Ke08	
^{191}Ir(p,t)^{189}Ir		−5903	15	−5916	11	−0.9	1	56	55 ^{189}Ir	McM		78Lo07	
^{190}Os(n,γ)^{191}Os		5758.2	2.	5758.73	0.11	0.3	U					77Be15	
		5759.1	1.5			−0.2	U					77Ca19	
		5758.67	0.16			0.4	–			ILn		91Bo35	Z
		5758.81	0.15			−0.5	–			Bdn		06Fi.A	
	ave.	5758.74	0.11			−0.1	1	100	99 ^{191}Os			average	
^{190}Os$(\alpha$,t)^{191}Ir		−14569	15	−14523.9	1.1	3.0	B			McM		71Pr13	
^{191}Ir(d,t)^{190}Ir		−1769.3	0.4				2					95Ga04	*
^{191}Os$(\beta^-)^{191}$Ir		313.3	3.	313.6	1.1	0.1	–					48Sa18	*
		314.3	2.			−0.4	–					51Ko17	*
		316.3	3.			−0.9	–					58Na15	*
		314.3	3.			−0.2	–					60Fe03	*
		318.3	3.			−1.6	–					63Pl01	*
	ave.	315.1	1.2			−1.3	1	90	90 ^{191}Ir			average	
^{191}Pt$(\varepsilon)^{191}$Ir		1000	15	1011	4	0.7	U					70Sc20	*
^{191}Au$(\beta^+)^{191}$Pt		1830	50	1900	6	1.4	U					76Vi.A	*
^{191}Hg$(\beta^+)^{191}$Au		3430	200	3205	23	−1.1	U					75Un.A	*
		3180	70			0.4	1	11	10 ^{191}Hg			76Vi.A	*
^{191}Tl$^m(\beta^+)^{191}$Hg		5178	200	4607	23	−2.9	C					75Un.A	*
*^{191}Au−u	$M-A$=–33568(28) keV for mixture gs+m at 266.1 keV											Nub269	**
*^{191}Hg−u	$M-A$=–30499(28) keV for mixture gs+m at 128(22) keV											Nub269	**
*^{191}Hg−^{208}Pb$_{.918}$	Original error (19keV) increased by 20 due to isomer+ground state lines in trap											01Sc41	**
*^{191}Tl−u	$M-A$=–26250(90) keV for mixture gs+m at 297(7) keV											Nub269	**
*^{191}Tl−u	$M-A$=–25964(28) keV for ^{191}Tl^m at 297(7) keV											Nub269	**
*^{191}Pb−u	Possible isomeric contamination											00Ra23	**
*^{191}Pb−u	$M-A$=–20226(28) keV for mixture gs+m at 59(10) keV											Nub269	**
*^{191}Bi$^m(\alpha)^{187}$Tl	average 6882(20) 6885(15)											13Uu01	**
*^{191}Po$(\alpha)^{187}$Pb	F : probably mainly ^{189}Bi^m											97Ba25	**
*^{191}Po$(\alpha)^{187}$Pb	E_α=7334(10), 6960(15) to ground state, 375(1) superseded by 02An19											99An10	**
*^{191}Po$^m(\alpha)^{187}$Pb^m	E_α=7376(5), 6888(5) to ^{187}Pb^m and 494(1) above											02An19	**
*^{191}Po$^m(\alpha)^{187}$Pb^m	E_α=7378(10), 6888(15) superseded by 02An19											99An10	**
*^{191}Ir(d,t)^{190}Ir	Feeds ground state											96Ga30	**
*^{191}Os$(\beta^-)^{191}$Ir	E_{β^-}=142(3) 143(2) 145(3) 143(3) 147(3) respectively, to 11/2$^-$ level at 171.29 keV											Ens07a	**
*^{191}Pt$(\varepsilon)^{191}$Ir	pL=0.73(0.12) to (1/2$^+$,3/2,5/2$^+$) at 935.46 keV , no K capture											Ens07a	**
*^{191}Au$(\beta^+)^{191}$Pt	E_{β^+}=850(30) to ground state and (5/2$^-$,7/2$^-$) level at 9.547 keV; also E_{β^+}=470(60) to											Ens07a	**
*	(3/2$^-$,5/2$^-$) level at 277.88 and 5/2$^-$ level at 293.458 keV											Ens07a	**
*^{191}Hg$(\beta^+)^{191}$Au	Reassigned by evaluator to mainly ground state, partly 3/2$^+$ 207.9 level											Ens07a	**
*^{191}Tl$^m(\beta^+)^{191}$Hg	E_{β^+}=3820(200) to level(5/2$^-$) at 336.32 keV											Ens07a	**
^{192}Re−u		−33912	76	−34000	50	−1.2	1	39	39 ^{192}Re	GS3	1.0	13Sh30	
^{192}Hg−u		−34440	104	−34367	17	0.7	U			GS1	1.0	00Ra23	
		−34342	30			−0.8	R			GS2	1.0	05Li24	
^{192}Hg−^{208}Pb$_{.923}$		−12826	20	−12816	17	0.5	2			MA6	1.0	01Sc41	
^{192}Tl−u		−27830	123	−27780	30	0.4	U			GS1	1.0	00Ra23	*
		−27775	34				2			GS2	1.0	05Li24	
^{192}Pb−u		−24280	104	−24211	6	0.7	U			GS1	1.0	00Ra23	
		−24185	30			−0.9	R			GS2	1.0	05Li24	
^{192}Bi−u		−14783	128	−14530	30	2.0	U			GS1	1.0	00Ra23	*
		−14489	60			−0.7	R			GS2	1.0	05Li24	*
^{192}Bi^m−^{133}Cs$_{1.444}$		122143.5	9.6				2			MA8	1.0	08We02	
^{192}Os O$_2$−^{189}Os ^{35}Cl		24301	6	24309.3	2.4	0.6	U			H22	2.5	70Mc03	
^{192}Re−^{190}Os$_{1.011}$		7947	61	8010	50	1.0	1	61	61 ^{192}Re	RKM	1.0	25Mu03	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{192}Os ^{35}Cl−^{190}Os ^{37}Cl	5984	3	5983.4	2.4	−0.1	U			H22	2.5	70Mc03	
^{192}Os−^{190}Os$_{1.011}$	3447	55	3490.4	2.4	0.8	U			RKM	1.0	25Mu03	
^{192}Ir−^{190}Os$_{1.011}$	4688	78	4614.1	1.2	−0.9	U			RKM	1.0	25Mu03	
^{192}Pt−^{190}Os$_{1.011}$	2900	230	3054.5	2.7	0.7	U			RKM	1.0	25Mu03	
^{192}Au−^{190}Os$_{1.011}$	6866	34	6837	15	−0.9	1	20	20 ^{192}Au	RKM	1.0	25Mu03	
^{192}Pb(α)^{188}Hg	5221.0	5.	5222	5	0.1	1	88	69 ^{188}Hg	ORa		79To06	Z
^{192}Bi(α)^{188}Tl	6376.0	5.	6377	4	0.2	3			Lvn		91Va04	*
	6377.9	5.			−0.2	3			Jya		13Ny01	*
^{192}Bi^m(α)^{188}Tl^m	6481.4	5.	6485	3	0.7	3					67Tr06	*
	6491.6	10.			−0.7	3			Ora		74Le02	*
	6483.3	5.			0.3	3			Lvn		91Va04	*
	6494	8			−1.1	3			Jya		03Ke04	*
^{192}Po(α)^{188}Pb	7322.8	20.	7320	3	−0.2	U					81Le23	
	7319.8	7.			−0.0	3			Lvn		93Wa04	
	7364.6	35.			−1.3	U			RIa		95Mo14	
	7349.4	30.			−1.0	U			RIa		97Pu01	
	7319.8	11.			−0.0	o			Jya		01Ke06	
	7318.8	8.			0.1	3			Jya		03Ke04	
	7319.8	4.			−0.0	3			Anv		03Va16	
^{192}At(α)^{188}Bi	7695.6	25.				3			Anv		06An04	
^{192}At(α)^{188}Bi^m	7629.3	15.				4			Anv		06An04	*
^{192}At^m(α)^{188}Bi	7695.6	25.				3			Anv		06An04	
^{192}At^m(α)^{188}Bi^n	7542.4	15.				4			Anv		06An04	*
^{192}Os(p,t)^{190}Os	−4835	5	−4835.3	2.2	−0.1	–			Min		73Oo01	
	−4837	4			0.4	–			McM		75Th04	
ave.	−4836	3			0.3	1	51	51 ^{192}Os			average	
^{192}Pt(p,t)^{190}Pt	−6629	7	−6642.7	2.5	−2.0	1	13	13 ^{192}Pt	Ors		80Ka19	
^{192}Os(t,α)^{191}Re	10993	10				2			McM		76Hi08	
^{192}Os(d,t)^{191}Os	−1265	15	−1301.2	2.2	−2.4	U			Tal		77Be15	
^{191}Ir(n,γ)^{192}Ir	6197.7	0.3	6198.12	0.11	1.4	o			ILn		87Ke.A	
	6198.1	0.2			0.1	–			ILn		91Ke10	
	6198.14	0.13			−0.1	–			Bdn		06Fi.A	
ave.	6198.13	0.11			−0.1	1	100	91 ^{192}Ir			average	
^{192}Pt(p,d)^{191}Pt	−6448	6	−6436.9	2.9	1.8	1	23	26 ^{191}Pt	Ors		80Ka19	
^{192}Pt(p,d)^{191}Pt−^{194}Pt()^{193}Pt	−307	3	−309.8	2.7	−0.9	1	81	74 ^{191}Pt	Ors		78Be09	
^{192}Ir(β^+)^{192}Os	1468	10	1046.7	2.4	−42.1	B					60An04	*
^{192}Ir(β^-)^{192}Pt	1456.7	4.	1452.8	2.3	−1.0	–					65Jo04	*
	1453.3	3.			−0.2	–					77Ra17	*
ave.	1454.5	2.4			−0.7	1	89	87 ^{192}Pt			average	
^{192}Au(β^+)^{192}Pt	3514	20	3523	14	0.5	–					66Ny01	
	3520	25			0.1	–					74Di.A	
ave.	3516	16			0.4	1	81	80 ^{192}Au			average	
^{192}Hg(β^+)^{192}Au	1745	30	753	21	−33.1	F					74Di.A	*
^{192}Tl(β^+)^{192}Hg	6380	200	6140	40	−1.2	U					75Un.A	*
*^{192}Tl−u	$M-A$=–25830(100) keV for mixture gs+m at 196(7) keV										Nub269	**
*^{192}Bi−u	$M-A$=–13700(110) keV for mixture gs+m at 140(30) keV										Nub269	**
*^{192}Bi−u	$M-A$=–13426(31) keV for mixture gs+m at 140(30) keV										Nub269	**
*^{192}Bi(α)^{188}Tl	E_α=6245(5), 6060(5) to ground state, 184.6 level										91Va04	**
*^{192}Bi(α)^{188}Tl	E_α=6064(5) to 184.6 level										Ens024	**
*^{192}Bi^m(α)^{188}Tl^m	E_α=6050(5) to (10$^-$) level, 302.4 above ^{188}Tl^m										91Va04	**
*^{192}Bi^m(α)^{188}Tl^m	E_α=6060(10) to (10$^-$) level, 302.4 above ^{188}Tl^m										91Va04	**
*^{192}Bi^m(α)^{188}Tl^m	E_α=6348(5), 6253(5), 6081(10), 6052(5) to ^{188}Tl^m and										91Va04	**
*	to levels 103.2, 268.8, 302.4 above ^{188}Tl^m										91Va04	**
*^{192}Bi^m(α)^{188}Tl^m	E_α=6062(5) to level 302.4 above ^{188}Tl^m										03Ke04	**
*^{192}At(α)^{188}Bi^m	Also E_α=7435(15) keV followed by 36 keV γ										06An04	**
*^{192}At^m(α)^{188}Bi^n	Also E_α=7224(15), 7195(15) keV followed by 165 and 188 keV γ										06An04	**
*^{192}Ir(β^+)^{192}Os	E_{β^+}=240(10) to 2$^+$ level at 205.7944 keV										Ens129	**
*^{192}Ir(β^-)^{192}Pt	E_{β^-}=672(4) 666(2) respectively, to 4$^+$ level at 784.5759, and other E_{β^-}										Ens129	**
*^{192}Hg(β^+)^{192}Au	F : most probably due to backscattering of 2.5 MeV Au positons										AHW	**
*^{192}Tl(β^+)^{192}Hg	E_{β^+}=4940(200) to 2$^+$ level at 422.79 keV										Ens129	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{193}Re−u	−32455	42				2			GS3	1.0	13Sh30	
^{193}Au−u	−35736	96	−35862	9	−1.3	U			GS2	1.0	05Li24	*
^{193}Hg−u	−33288	53	−33347	17	−1.1	U			GS2	1.0	05Li24	*
^{193}Hg−^{208}Pb$_{.928}$	−11673	29	−11679	17	−0.2	1	33	33 ^{193}Hg	MA6	1.0	01Sc41	*
^{193}Tl−u	−29690	158	−29497	7	1.2	U			GS1	1.0	00Ra23	*
	−29329	120			−1.4	U			GS2	1.0	05Li24	*
^{193}Tl−^{133}Cs$_{1.451}$	107692.1	7.2				2			MA8	1.0	14Bo26	*
^{193}Pb−u	−23840	110	−23864	11	−0.2	o			GS1	1.0	00Ra23	*
	−23826	43			−0.9	–			GS2	1.0	05Li24	*
ave.	−23820	40			−1.1	1	8	8 ^{193}Pb			average	
^{193}Bi−u	−16980	110	−17053	8	−0.7	U			GS1	1.0	00Ra23	
	−17025	30			−0.9	R			GS2	1.0	05Li24	
^{193}Bi−^{133}Cs$_{1.451}$	120147	11	120136	8	−1.0	–			MA8	1.0	08We02	
ave.	120149	10			−1.2	1	62	62 ^{193}Bi			average	
^{193}Bi(α)^{189}Tl	6304.5	5.	6307	5	0.4	1	92	70 ^{189}Tl	Lvn		85Co06	Z
^{193}Bi(α)^{189}Tl^m	6017.8	5.	6021	3	0.7	2					67Tr06	Z
	6024.6	10.			−0.3	2			Ora		74Le02	Z
	6023.7	5.			−0.5	2			Lvn		85Co06	Z
^{193}Bi^m(α)^{189}Tl	6617.4	10.	6611	4	−0.6	–			Ora		74Le02	
	6611.9	5.			−0.1	–			Lvn		85Co06	Z
	6618.4	14.			−0.5	U			Jya		05Uu02	
ave.	6613	5			−0.4	1	94	64 ^{193}Bi^m			average	
^{193}Po(α)^{189}Pb	7128.1	20.	7094	4	−1.7	U					67Si09	
	7087.1	20.			0.3	U			Ora		77De32	
	7096.4	5.			−0.5	2			Lvn		93Wa04	
	7093.3	30.			0.0	U			RIa		95Mo14	
	7089.2	6.			0.7	2			Jya		96En02	
	7096.4	10.			−0.3	2			Anv		02Va13	
^{193}Po^m(α)^{189}Pb^m	7143.3	10.	7154	3	1.0	2			Ora		77De32	
	7148.4	20.			0.3	U					81Le23	
	7152.5	5.			0.2	2			Lvn		93Wa04	
	7139.2	30.			0.5	U			RIa		95Mo14	
	7159.7	6.			−1.0	2			Jya		96En02	
	7152.5	10.			0.1	2			Anv		02Va13	
^{193}At(α)^{189}Bi^m	7388.5	5.				8			Jya		03Ke08	
^{193}At^m(α)^{189}Bi	7556.9	20.	7580	5	1.1	o			Jya		95Le15	
	7490	6			15.1	C			Jya		98En.A	
	7580.4	5.				8			Jya		03Ke08	*
^{193}At^n(α)^{189}Bi	7614.3	5.				8			Jya		03Ke08	*
^{193}Rn(α)^{189}Po	8040.0	12.				4			Anv		06An36	*
^{193}Ir(p,t)^{191}Ir	−5490	15	−5488.32	0.23	0.1	U			McM		78Lo07	
^{192}Os(n,γ)^{193}Os	5583.5	2.	5583.42	0.20	−0.0	U					78Be22	
	5583.40	0.20			0.1	1	100	81 ^{193}Os			79Wa04	
	5584.01	0.16			−3.7	C			Bdn		06Fi.A	
^{192}Os(α,t)^{193}Ir	−13923	15	−13870.9	2.4	3.5	B			McM		71Pr13	
^{193}Ir(t,α)^{192}Os−^{191}Ir()^{190}Os	−661	4	−653.0	2.2	2.0	1	31	31 ^{192}Os	LAl		82La22	
^{192}Ir(n,γ)^{193}Ir	7772.0	0.2	7771.99	0.20	−0.0	1	100	94 ^{193}Ir			85Co.B	Z
^{193}Ir(γ,n)^{192}Ir	−7790	50	−7771.99	0.20	0.4	U			Phi		60Ge01	
^{192}Pt(n,γ)^{193}Pt	6247	3	6262.6	2.3	5.2	B					68Sa13	
^{193}Os(β^-)^{193}Ir	1132	5	1141.9	2.4	2.0	1	23	19 ^{193}Os			58Na15	
^{193}Pt(ε)^{193}Ir	56.6	0.3	56.63	0.30	0.1	1	100	96 ^{193}Pt			83Jo04	
^{193}Au(β^+)^{193}Pt	1355	20	1075	9	−14.0	B					76Di15	*
^{193}Hg(β^+)^{193}Au	2341	30	2342	14	0.0	–					58Br88	*
	2340	20			0.1	–					76Di15	*
ave.	2340	17			0.1	1	75	67 ^{193}Hg			average	
*^{193}Au−u	$M-A$=–33143(29) keV for mixture gs+m at 290.20 keV										Nub269	**
*^{193}Hg−u	$M-A$=–30937(28) keV for mixture gs+m at 140.76 keV										Nub269	**
*^{193}Hg−^{208}Pb$_{.928}$	Original error (18keV) increased by 20 due to isomer+ground state lines in trap										01Sc41	**
*^{193}Tl−u	$M-A$=–27470(100) keV for mixture gs+m at 372(4) keV										Nub269	**
*^{193}Tl−u	$M-A$=–27134(28) keV for mixture gs+m at 372(4) keV										Nub269	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{193}$Tl−^{133}Cs$_{1.451}$		D_M=108091.5(5.6) μu for ^{193}Tl^m at 372(4) keV; $M-A$=–27104.3(5.2) keV										Nub269	**
$*^{193}$Pb−u		$M-A$=–22160(100) keV for mixture gs+m at 93(12) keV										Nub269	**
$*^{193}$Pb−u		$M-A$=–22147(28) keV for mixture gs+m at 93(12) keV										Nub269	**
$*^{193}$At$^m(\alpha)^{189}$Bi		E_α=7423(5), 7325(5) to ground state and 99.6(5) level										03Ke08	**
$*^{193}$At$^n(\alpha)^{189}$Bi		E_α=7106(5) to 357.6(0.5) ^{189}Bi^n level										03Ke08	**
$*^{193}$Rn$(\alpha)^{189}$Po		E_α=7875(20), 7685(15) to ground state and 194 level										06An36	**
$*^{193}$Au$(\beta^+)^{193}$Pt		E_{β^+}=153(15) to 3/2$^-$ level at 187.81 keV, and other E_{β^+}										Ens062	**
$*^{193}$Hg$(\beta^+)^{193}$Au		E_{β^-}=1170(30) from ^{193}Hg^m at 140.76 to 11/2$^-$ level at 290.19 keV										Ens062	**
$*^{193}$Hg$(\beta^+)^{193}$Au		E_{β^+}=1287(15) reinterpreted by AHW as going to ground state and 1/2$^+$ level at 38.22keV										Ens062	**
^{194}Pt−u		−37315.72	0.67	−37316.5	0.5	−1.1	1	63	63 ^{194}Pt	MS1	1.0	16Ei01	
^{194}Au−u		−34768	114	−34580.9	2.3	1.6	U			GS2	1.0	05Li24	*
^{194}Hg−u		−34527	30	−34551	3	−0.8	U			GS2	1.0	05Li24	
^{194}Hg−^{133}Cs$_{1.459}$		103394.7	3.1				2			MA8	1.0	10El11	
^{194}Hg−^{208}Pb$_{.933}$		−12766	19	−12766	3	0.0	U			MA6	1.0	01Sc41	
^{194}Tl−u		−28803	135	−28919	15	−0.9	o			GS1	1.0	00Ra23	*
		−28778	87			−1.6	U			GS2	1.0	05Li24	*
^{194}Tl−^{133}Cs$_{1.459}$		109027	15				2			MA8	1.0	14Bo26	
^{194}Tl^m−^{133}Cs$_{1.459}$		109306.4	4.1	109306	4	0.0	o			MA8	1.0	14Bo26	
		109306.4	4.1				2			MA8	1.0	13St25	
^{194}Pb−u		−25980	104	−25991	18	−0.1	U			GS1	1.0	00Ra23	
^{194}Bi−u		−17160	120	−17201	6	−0.3	o			GS1	1.0	00Ra23	*
		−17165	59			−0.6	U			GS2	1.0	05Li24	*
^{194}Bi^m−^{133}Cs$_{1.459}$		120900	54				2			MA8	1.0	08We02	*
^{190}Os−^{194}Pt$_{.979}$		−5022.45	0.68	−5021.7	0.5	1.1	1	57	52 ^{190}Os	MS1	1.0	16Ei01	
^{190}Pt−^{194}Pt$_{.979}$		−3516.95	0.68	−3517.3	0.5	−0.5	1	58	53 ^{190}Pt	MS1	1.0	16Ei01	
^{194}Pt−^{197}Au$_{.985}$		−4396.4	3.2	−4388.0	0.6	2.6	U			CP1	1.0	05Sh52	
^{194}Au−^{197}Au$_{.985}$		−1652.5	2.2				2			MA8	1.0	10El11	
^{194}Pb$(\alpha)^{190}$Hg		4737.9	20.	4737	16	−0.0	1	65	35 ^{194}Pb	ORa		87El09	
^{194}Bi$(\alpha)^{190}$Tl		5918.3	5.				3			Lvn		91Va04	*
^{194}Bi$^n(\alpha)^{190}$Tl^m		6015.7	5.	6012	4	−0.8	1	78	42 ^{194}Bi^n	Lvn		91Va04	*
^{194}Po$(\alpha)^{190}$Pb		6991.5	10.	6987	3	−0.4	3					67Si09	Z
		6990.9	7.			−0.5	3					67Tr06	Z
		6984.4	5.			0.5	3			Ora		77De32	Z
		6990.0	5.			−0.6	o			Lvn		85Va03	Z
		6986.3	6.			0.1	3			Lvn		93Wa04	
		6993.4	4.			−1.6	o			Jya		96En02	
		6987.3	14.			−0.0	3			Jya		05Uu02	
^{194}At$(\alpha)^{190}$Bi		7462.5	15.	7460	14	−0.2	1	85	68 ^{194}At	Anv		09An11	*
^{194}At$(\alpha)^{190}$Bi^m		7362.1	20.	7310	5	−2.6	o					80Ya.A	
		7351.9	20.			−2.1	U					84Ya.A	
		7341.7	20.			−1.6	o			Jya		95Le15	
		7329	15			−1.3	–			Anv		09An11	*
		7306.9	5.1			0.5	–			Jya		13Ny01	
	ave.	7309	5			0.1	1	98	66 ^{190}Bi^m			average	
^{194}At$^m(\alpha)^{190}$Bi		7412.5	20.	7456	11	2.2	o			Jya		95Le15	*
		7464	15			−0.5	2			Anv		09An11	*
		7446	17			0.6	2			Jya		13Ny01	*
^{194}Rn$(\alpha)^{190}$Po		7862.5	10.				4			Anv		06An36	
^{193}Ir(n,γ)^{194}Ir		6067.0	0.4	6066.79	0.11	−0.5	2					82Ra.A	
		6066.9	0.2			−0.6	2					98Ba85	
		6066.71	0.14			0.6	2			Bdn		06Fi.A	
^{194}Pt(t,α)^{193}Ir		12286	20	12301.1	1.3	0.8	U			Tal		78Ya07	
^{194}Pt(d,t)^{193}Pt		−2126	20	−2094.5	1.3	1.6	U			Pit		64Co11	
^{194}Pt(p,d)^{193}Pt−^{196}Pt()^{195}Pt		−445	3	−429.8	1.3	5.1	B			Ors		78Be09	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{194}Os(β^-)^{194}Ir	96.6	2.				3					64Wi07	*
^{194}Ir(β^-)^{194}Pt	2254	4	2228.3	1.3	−6.4	B					76Ra33	
^{194}Ir^n(β^-)^{194}Pt	2560	70				2					68Su02	*
^{194}Pt(d,α)^{192}Ir−^{192}Os()^{190}Re	375	4	375	4	−0.1	1	100	100 ^{190}Re			20Gr08	
^{194}Au(β^+)^{194}Pt	2465	20	2548.1	2.1	4.2	B					56Th11	
	2509	15			2.6	U					60Ba17	
	2485	30			2.1	U					70Ag03	*
^{194}Hg(ε)^{194}Au	40	20	28	4	−0.6	U					81Ho18	
*^{194}Au−u	$M-A$=–32192(29) keV for mixture gs+m+n at 107.4 and 475.7 keV										Nub269	**
*^{194}Tl−u	$M-A$=–26700(100) keV for mixture gs+m at 260(14) keV										Nub269	**
*^{194}Tl−u	$M-A$=–26677(28) keV for mixture gs+m at 260(14) keV										Nub269	**
*^{194}Bi−u	$M-A$=–15870(100) keV for mixture gs+m+n at 150(50) and 163(4) keV										Nub269	**
*^{194}Bi−u	$M-A$=–15885(28) keV for mixture gs+m+n at 150(50) and 163(4) keV										Nub269	**
*^{194}Bi^m−^{133}Cs$_{1.459}$	Original error 16 μu increased to include possible 3^+ and 10^- contam.										08We02	**
*^{194}Bi(α)^{190}Tl	E_α=5799(5), 5645(5) to ground state, 151.3 level										91Va04	**
*^{194}Bi^n(α)^{190}Tl^m	E_α=5892(5), 5781(5) to levels 0, 112.2 above ^{190}Tl^m										91Va04	**
*^{194}At(α)^{190}Bi	E_α=7190(15) to 121(15); further E_α: 7310(15), 7266(15), 7145(15) keV										09An11	**
*^{194}At(α)^{190}Bi^m	E_α=7178(15), 7135(15), 7087(15), 6908(15) to levels										09An11	**
*^{194}At(α)^{190}Bi^m	0, 40.3(0.5), 96.7(0.5), 274(1) keV above ^{190}Bi^m										09An11	**
*^{194}At^m(α)^{190}Bi	E_α=7140(20) to 121(15)										09An11	**
*^{194}At^m(α)^{190}Bi	E_α=7310(15), 7266(15), 7190(15), 7145(15) to levels										09An11	**
*^{194}At^m(α)^{190}Bi	at 0, 45(15), 121(15), 168(15) keV										09An11	**
*^{194}At^m(α)^{190}Bi	E_α=7174(8) to 121(15) keV										13Ny01	**
*^{194}Os(β^-)^{194}Ir	E_{β^-}=54.5(2.0) to 0^- level at 43.119 keV, and other E_{β^-}										Ens066	**
*^{194}Ir^n(β^-)^{194}Pt	from E_{β^-} <250 to 10^+ level at 2438.41 keV										68Su02	**
*^{194}Au(β^+)^{194}Pt	E_{β^+}=1230(30) to 2^+ level at 328.464 keV, and other E^+										Ens066	**
^{195}Os−u	−31682	60				2			GS3	1.0	13Sh30	
^{195}Hg−u	−33283	62	−33295	25	−0.2	U			GS2	1.0	05Li24	*
^{195}Hg−^{208}Pb$_{.938}$	−11381	28	−11393	25	−0.4	1	79	79 ^{195}Hg	MA6	1.0	01Sc41	*
^{195}Tl−u	−30320	200	−30226	12	0.5	U			GS1	1.0	00Ra23	*
	−30209	40			−0.4	–			GS2	1.0	05Li24	
	−30264	33			1.1	–			GS2	1.0	05Li24	*
ave.	−30242	25			0.6	1	22	22 ^{195}Tl			average	
^{195}Tl−^{133}Cs$_{1.466}$	108375	27	108381	12	0.2	–			MA8	1.0	14Bo26	
	108472	79			−1.1	–			MA8	1.0	14Bo26	*
ave.	108385	26			−0.2	1	22	22 ^{195}Tl			average	
^{195}Pb−u	−25423	150	−25486	6	−0.4	o			GS1	1.0	00Ra23	*
	−25461	70			−0.4	U			GS2	1.0	05Li24	*
	−25451	38			−0.9	U			1.0	1.0	199Po-C	
	−25462	38			−0.6	U			1.0	1.0	199Po-C	
^{195}Bi−u	−19320	100	−19351	6	−0.3	U			GS1	1.0	00Ra23	
	−19537	128			1.5	U			GS2	1.0	05Li24	*
^{195}Bi−^{133}Cs$_{1.466}$	119258.2	6.0	119256	6	−0.3	1	89	89 ^{195}Bi	MA8	1.0	08We02	
^{195}Po−^{133}Cs$_{1.466}$	126671.8	6.6				2			MA8	1.0	17Al34	
^{195}Po^m−^{133}Cs$_{1.466}$	126832.4	7.8	126833	8	0.0	1	94	94 ^{195}Po^m	MA8	1.0	17Al34	
^{195}Pt−^{197}Au$_{.990}$	−2119.9	3.2	−2110.1	0.6	3.1	B			CP1	1.0	05Sh52	
^{195}Bi(α)^{191}Tl	5832.5	5.				2			Lvn		85Co06	Z
^{195}Bi(α)^{191}Tl^m	5542.9	10.	5535	5	−0.8	2			Ora		74Le02	Z
	5533.3	5.			0.4	2			Lvn		85Co06	Z
^{195}Bi^m(α)^{191}Tl	6228.1	5.	6232	3	0.7	3					67Tr06	Z
	6238.4	10.			−0.6	3			Ora		74Le02	Z
	6233.7	5.			−0.4	3			Lvn		85Co06	Z
^{195}Po(α)^{191}Pb	6763.1	8.	6749.9	2.8	−1.6	3					67Si09	Z
	6747.4	5.			0.5	3					67Tr06	Z
	6744.6	5.			1.0	3			Lvn		93Wa04	
	6752.8	14.			−0.2	o			Jya		96Le09	
	6744.6	10.			0.5	3			Anv		02Va13	
	6755.9	6.			−1.0	3			Jya		05Uu02	
^{195}Po^m(α)^{191}Pb^m	6850.8	10.	6840.6	2.8	−1.0	–					67Si09	
	6839.4	5.			0.2	–					67Tr06	Z

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{195}Po$^m(\alpha)^{191}$Pb^m		6839.6	5.	6840.6	2.8	0.2	–				Lvn		93Wa04	
		6852.8	10.			−1.2	o				Jya		96Le09	
		6839.6	10.			0.1	–				Anv		02Va13	
		6840.6	6.			0.0	–				Jya		05Uu02	
	ave.	6840.6	2.9			−0.0	1	99	93	^{191}Pb^m			average	
^{195}At$(\alpha)^{191}$Bi^m		7095.8	20.	7102	4	0.3	U				Jya		95Le15	
		7105	20			−0.2	U				RIa		99Ta20	
		7098.9	3.			0.9	o				Jya		03Ke04	*
		7113.2	10.			−1.1	o				Jya		13Uu01	
		7101.9	4.			−0.1	4				Jya		13Ny01	
		7101	8			0.1	4				Anv		13Ka16	
		7118	34			−0.5	U				Lza		24Wa14	
^{195}At$^m(\alpha)^{191}$Bi		7340.9	30.	7373	4	1.1	o				Jya		83Le.A	*
		7371.5	30.			0.1	o				Jya		95Le.A	
		7403	30			−1.0	U				RIa		99Ta20	*
		7372.5	4.0			0.2	o				Jya		03Ke04	*
		7373.5	4.0				2				Jya		13Ny01	*
^{195}Rn$(\alpha)^{191}$Po		7694.1	11.				2				Jya		01Ke06	
^{195}Rn$^m(\alpha)^{191}$Po^m		7713.5	11.				3				Jya		01Ke06	
^{194}Ir(n,γ)^{195}Ir		7231.92	0.11	7231.86	0.06	−0.5	o				ILn		87Ci.A	
		7231.86	0.06				3				ILn		87Co08	Z
^{194}Pt(n,γ)^{195}Pt		6105.06	0.12	6105.09	0.12	0.3	1	99	72	^{195}Pt	ILn		81Ho.B	Z
		6109.17	0.13			−31.4	F				Bdn		06Fi.A	
^{195}Pt(γ,n)^{194}Pt		−6205	44	−6105.09	0.12	2.3	U				Phi		60Ge01	
^{194}Pt(d,p)^{195}Pt		3908	20	3880.53	0.12	−1.4	U				Pit		64Co11	
^{195}Pt(d,t)^{194}Pt		140	20	152.14	0.12	0.6	U				Pit		64Co11	
^{195}Os$(\beta^-)^{195}$Ir		2000	500	2180	60	0.4	U						57Ba08	
^{195}Ir$(\beta^-)^{195}$Pt		1116	20	1101.6	1.3	−0.7	U						73Ja10	*
^{195}Au$(\varepsilon)^{195}$Pt		226.8	1.0	226.8	1.0	0.0	1	100	100	^{195}Au			Averag	*
^{195}Hg$(\beta^+)^{195}$Au		1510	50	1553	23	0.9	1	21	21	^{195}Hg			71Fr03	*
^{195}Tl$(\beta^+)^{195}$Hg		3000	300	2859	26	−0.5	U						78Go15	*
^{195}Pb^m(IT)^{195}Pb		202.9	0.7	202.9	0.7	−0.0	1	100	99	^{195}Pb	Oak		91Gr12	
^{195}Bi$(\beta^+)^{195}$Pb		4850	550	5715	7	1.6	U				Oak		91Gr12	
*^{195}Hg−u	$M-A$=–30914(28) keV for mixture gs+m at 176.07 keV												Nub269	**
*^{195}Hg−^{208}Pb$_{.938}$	Corrected 40(20) keV for isomeric mixture R=0.3(0.2) E=176.07 keV												Nub269	**
*^{195}Tl−u	$M-A$=–28000(100) keV for mixture gs+m at 482.63 keV												Nub269	**
*^{195}Tl−u	$M-A$=–27708(31) keV for ^{195}Tl^m at 482.63 keV												Nub269	**
*^{195}Tl−^{133}Cs$_{1.466}$	D_M=108990(79) μu for ^{195}Tl^m at 482.63 keV; $M-A$=–27589(73) keV												Nub269	**
*^{195}Pb−u	$M-A$=–23580(100) keV for mixture gs+m at 202.9 keV												Nub269	**
*^{195}Pb−u	$M-A$=–23615(28) keV for mixture gs+m at 202.9 keV												Nub269	**
*^{195}Bi−u	$M-A$=–17999(28) keV for mixture gs+m at 399(6) keV												Nub269	**
*^{195}At$(\alpha)^{191}$Bi^m	Correlated with E_α=6313 of ^{191}Bi^m												03Ke04	**
*^{195}At$^m(\alpha)^{191}$Bi	E_α=7190(30) to 148.7(0.5) level												03Ke04	**
*	correlated with α of 12 s ^{191}Bi ground state												95Le15	**
*^{195}At$^m(\alpha)^{191}$Bi	E_α=7105(30) to 148.7(0.5) level												03Ke04	**
*^{195}At$^m(\alpha)^{191}$Bi	E_α=7221(4) and 7075(4) to 148.7(0.5) level												03Ke04	**
*^{195}At$^m(\alpha)^{191}$Bi	E_α=7222(4) and 7076(5) to 148.7(0.5) level												13Ny01	**
*^{195}Ir$(\beta^-)^{195}$Pt	E_{β^-}=980(30) to 3/2$^-$ level at 98.880 keV and 5/2$^-$ level at 129.772 keV,												Ens148	**
*	and E_{β^-}=410(20) from ^{195}Ir^m at 100(5) to 9/2$^-$ at 814.50, and other E_{β^-}												Ens148	**
*^{195}Au$(\varepsilon)^{195}$Pt	Average pK=0.179(0.006) to 5/2$^-$ level at 129.772 from the following references:												Ens148	**
*	pK=0.195(0.015) to 129.78 level												65De20	**
*	pK=0.166(0.020) to 129.78 level												68Ja11	**
*	pK=0.160(0.017) to 129.78 level												73Go05	**
*	pK=0.183(0.009) to 129.78 level												80Sa11	**
*	pK=0.176(0.012) to 129.78 level												82Be.A	**
*^{195}Hg$(\beta^+)^{195}$Au	Assuming 511 γ is annihilation of β^+ to ground state and 1/2$^+$ level at 61.434												Ens148	**
*^{195}Tl$(\beta^+)^{195}$Hg	K/β^+=6(1) to ground state and 3/2$^-$ level at 37.083 keV												Ens148	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{196}Hg−u		−34161.8	16.3	−34166	3	−0.3	U			MA8	1.0	20Ku19	*
^{196}Hg−^{208}Pb$_{.942}$		−12178	20	−12171	3	0.4	U			MA6	1.0	01Sc41	
^{196}Tl−u		−29188	126	−29519	13	−2.6	U			GS2	1.0	05Li24	*
^{196}Tl−^{133}Cs$_{1.474}$		109845	13				2			MA8	1.0	08We02	*
^{196}Pb−^{208}Pb$_{.942}$		−5228	22	−5213	9	0.7	–			MA6	1.0	01Sc41	
	ave.	−5231	18			1.0	1	23	23 ^{196}Pb			average	
^{196}Pb−u		−27200	104	−27209	9	−0.1	U			GS1	1.0	00Ra23	
		−27232	30			0.8	R			GS2	1.0	05Li24	
^{196}Bi−u		−19320	120	−19325.7	1.0	−0.0	o			GS1	1.0	00Ra23	*
		−19325	30			−0.0	–			GS2	1.0	05Li24	
		−19361	54			0.7	–			MA8	1.0	08We02	*
		−19324.9	1.1			−0.7	–			SH2	1.0	22Mo.A	
	ave.	−19324.9	1.1			−0.7	1	80	80 ^{196}Bi			average	
^{196}Bi^{m}−u		−19147.8	1.8	−19147.9	1.3	−0.1	1	55	55 ^{196}Bi^{m}	SH2	1.0	22Mo.A	
^{196}Bi^{n}−u		−19040.73	0.89				2			SH2	1.0	22Mo.A	
^{196}Po−^{133}Cs$_{1.474}$		124905.4	6.3	124904	6	−0.2	1	84	84 ^{196}Po	MA8	1.0	17Al34	
^{196}Pt−^{197}Au$_{.995}$		−1781.1	3.0	−1782.6	0.6	−0.5	U			CP1	1.0	05Sh52	
^{196}Bi(α)^{192}Tl^{p}		5260.6	5.				2			Lvn		91Va04	
^{196}Po(α)^{192}Pb		6662.3	8.2	6658.2	2.4	−0.5	–					67Si09	Z
		6653.7	5.			0.9	–					67Tr06	Z
		6658.4	8.			−0.0	–					71Ho01	Z
		6656.7	5.			0.3	o			Lvn		85Va03	Z
		6656.7	5.			0.3	–			Lvn		93Wa04	
		6653.1	18.			0.3	o			Ara		95Le04	
		6657.1	10.			0.1	o			Jya		96Le09	
		6654.0	5.0			0.8	–			Ara		96Ta18	*
		6669.4	6.			−1.8	–			Jya		05Uu02	
		6658.2	25.			0.0	U			Anv		10He25	
	ave.	6658.1	2.4			0.1	1	97	81 ^{192}Pb			average	
^{196}At(α)^{192}Bi		7202.3	7.	7196.4	2.7	−0.8	4					67Tr06	
		7187.0	25.			0.4	U			Jya		95Le15	
		7200.2	30.			−0.1	U			RIa		95Mo14	
		7191.0	7.			0.8	o			Jya		96En01	
		7195.1	5.			0.3	o			Jya		00Sm06	
		7202.3	12.			−0.5	o			Anv		05De01	
		7195.1	12.			0.1	o			Jya		13Uu01	*
		7194.1	5.			0.5	4			Jya		13Ny01	
		7192.1	5.			0.9	4			Anv		14Ka23	
		7200.2	5.			−0.7	4			Anv		16Tr07	
		7201	15			−0.3	U			Lza		22Hu12	
^{196}At^{m}(α)^{192}Bi^{m}		7023.6	15.				3			Jya		96En01	*
^{196}Rn(α)^{192}Po		7583.1	35.	7617	9	0.9	o			RIa		95Mo14	
		7648.4	30.			−1.1	U			RIa		97Pu01	
		7616.7	9.				4			Jya		01Ke06	
^{196}Pt(t,α)^{195}Ir		11565	20	11572.7	1.3	0.4	U			Tal		78Ya07	
		11545	20			1.4	U			LAl		81Fl.A	
^{195}Pt(n,γ)^{196}Pt		7921.96	0.20	7921.98	0.13	0.1	–			ILn		81Ho.B	Z
		7921.92	0.17			0.3	–			Bdn		06Fi.A	
^{196}Pt(γ,n)^{195}Pt		−8290	140	−7921.98	0.13	2.6	U			Phi		60Ge01	
^{195}Pt(d,p)^{196}Pt		5712	25	5697.41	0.13	−0.6	U			Pit		64Co11	
^{196}Pt(d,t)^{195}Pt		−1686	20	−1664.75	0.13	1.1	U			Pit		64Co11	
^{195}Pt(n,γ)^{196}Pt	ave.	7921.94	0.13	7921.98	0.13	0.3	1	99	71 ^{196}Pt			average	
^{196}Os(β^-)^{196}Ir		900	40	1160	60	6.5	B					77Ha32	*
^{196}Ir(β^-)^{196}Pt		3150	60	3210	40	1.0	2					66Vo05	*
		3250	50			−0.8	2					67Mo10	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{196}Ir$^{m}(\beta^-)^{196}$Pt	3418	20				2					65Bi04	*
	3630	100	3418	20	−2.1	U					68Ja06	*
^{196}Au$(\beta^+)^{196}$Pt	1498	7	1507.1	2.6	1.3	1	13	13 ^{196}Au			63Ik01	*
^{196}Au$(\varepsilon)^{196}$Pt	1490	10			1.7	U					62Wa16	*
^{196}Au$(\beta^-)^{196}$Hg	685	4	688.0	3.0	0.7	1	56	33 ^{196}Hg			62Li03	*
*^{196}Hg−u	From frequency ratio $^{196}Hg^+/(^{39}K)^+$ =2.514744076(209)										20Ku19	**
*^{196}Tl−u	$M-A$=–26991(28) keV for mixture gs+m at 394.2 keV										Nub269	**
*^{196}Tl−^{133}Cs$_{1.474}$	Q =110268(13) μu $M-A$=–27103(12) keV for ^{196}Tl^{m} at 394.2 keV										Nub269	**
*^{196}Bi−u	$M-A$=–17850(100) keV for mixture gs+m+n at 166.4(2.9),272(3) keV										Nub269	**
*^{196}Bi−u	Q =120182(15) μu for ^{196}Bi^{m}−^{133}Cs$_{1.474}$, $M-A$(^{196}Bi^{m})=–17868(14) keV at										08We02	**
*	167(3) keV; error increased to include possible 3^+ and 10^- contam.										08We02	**
*^{196}Po$(\alpha)^{192}$Pb	Including systematic uncertainty 5 keV										96Ta18	**
*^{196}At$(\alpha)^{192}$Bi	Same group, but much less events than in 00Sm06										WgM151	**
*^{196}At$^{m}(\alpha)^{192}$Bi^{m}	Correlated with E_α=7550 of ^{200}Fr(α)										96En01	**
*^{196}Os$(\beta^-)^{196}$Ir	E_{β^-}=435(20) to $(0,1)^+$ levels at 407.88, 522.37 keV										Ens076	**
*^{196}Ir$(\beta^-)^{196}$Pt	Original value 3170(60) recalibrated using ^{62}Cu										AHW	**
*^{196}Ir$^{m}(\beta^-)^{196}$Pt	E_{β^-}=950(20) to $(10^-,11^-)$ level at 2468.0 keV										Ens076	**
*^{196}Ir$^{m}(\beta^-)^{196}$Pt	E_{β^-}=1160(100) to $(10^-,11^-)$ level at 2468.0 keV										Ens076	**
*^{196}Au$(\beta^+)^{196}$Pt	KL/β^+=2.0(0.4)×10^{+6} to 2^+ level at 355.68 keV, recalculated										Ens076	**
*^{196}Au$(\varepsilon)^{196}$Pt	pL=0.64(0.06) to 3^- level at 1447.043 keV										Ens076	**
*^{196}Au$(\beta^-)^{196}$Hg	E_{β^-}=259(4) to 2^+ level at 425.98 keV										Ens076	**
^{197}Ir−u	−30279	15	−30300	12	−1.4	1	67	67 ^{197}Ir	GS5	1.0	26Fe.B	
^{197}Hg−u	−32766	30	−32786	3	−0.7	U			GS2	1.0	05Li24	
	−32765	30			−0.7	U			GS2	1.0	05Li24	*
^{197}Hg−^{208}Pb$_{.947}$	−10664	30	−10674	3	−0.3	U			MA6	1.0	01Sc41	
^{197}Tl−u	−30450	30	−30440	15	0.3	R			GS2	1.0	05Li24	
^{197}Pb−u	−26520	110	−26566	5	−0.4	U			GS1	1.0	00Ra23	
	−26609	30			1.4	U			GS2	1.0	05Li24	
	−26543	30			−0.8	U			GS2	1.0	05Li24	*
^{197}Pb^{m}−^{133}Cs$_{1.481}$	113799.6	6.0	113803	5	0.5	1	73	73 ^{197}Pb^{m}	MA8	1.0	08We02	
^{197}Bi−^{208}Pb$_{.947}$	982	22	981.0	2.0	−0.0	R			MA6	1.0	01Sc41	
^{197}Bi−u	−21389	190	−21130.9	2.0	1.4	U			GS1	1.0	00Ra23	*
	−21187	31			1.8	U			GS2	1.0	05Li24	
^{197}Bi−^{133}Cs$_{1.481}$	118870	26	118894.7	2.0	1.0	R			MA8	1.0	08We02	*
^{197}Po−u	−14420	130	−14378	11	0.3	o			GS1	1.0	00Ra23	*
	−14288	71			−1.3	R			GS2	1.0	05Li24	*
^{197}Po−^{133}Cs$_{1.481}$	125644	11	125648	11	0.3	1	93	93 ^{197}Po	MA8	1.0	17Al34	
^{197}Po^{m}−^{133}Cs$_{1.481}$	125858.0	7.0				2			MA8	1.0	17Al34	
^{197}At−^{133}Cs$_{1.481}$	133186	20	133203	9	0.8	1	18	18 ^{197}At	MA8	1.0	17Ma29	
^{197}At^{m}−^{133}Cs$_{1.481}$	133234	15	133251	10	1.1	1	42	42 ^{197}At^{m}	MA8	1.0	17Ma29	
^{197}Au−C$_{16}$	−33432.5	7.3	−33429.9	0.6	0.2	o			TG1	1.5	09Ke.A	
	−33432.9	5.4			0.4	U			TG1	1.5	10Ke09	
^{197}Au$(\alpha,^{8}$He$)^{193}$Au	−26919	9	−26920	9	−0.1	1	93	93 ^{193}Au			89Ka04	
^{197}Bi$^{m}(\alpha)^{193}$Tl	5890.8	10.	5898	5	0.7	o			Ora		72Ga27	
	5889.7	10.			0.8	3			Ora		74Le02	Z
	5899.6	5.			−0.4	3			Lvn		85Co06	Z
^{197}Po$(\alpha)^{193}$Pb	6420.7	10.	6411	3	−0.9	–					67Si09	Z
	6410.1	5.			0.2	–					67Tr06	Z
	6409.4	9.			0.2	–					71Ho01	Z
	6411.4	5.0			−0.0	–			Ara		96Ta18	*
ave.	6412	3			−0.1	1	99	92 ^{193}Pb			average	
^{197}Po$^{m}(\alpha)^{193}$Pb^{m}	6510.1	5.	6514.7	2.1	0.9	3					67Tr06	Z
	6511.4	9.			0.4	U					71Ho01	Z
	6518.0	3.			−1.1	3			Bka		82Bo04	Z
	6512.4	5.0			0.4	3			Ara		96Ta18	*
	6517.6	10.			−0.3	o			Anv		02Va13	
	6516.6	30.			−0.1	U			Anv		10He25	
	6513.0	4.6			0.4	3			Tex		12Fo09	
^{197}At$(\alpha)^{193}$Bi	7103.0	5.	7104	3	0.3	–					67Tr06	Z
	7100.5	5.			0.8	o			Jya		96En01	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{197}At(α)^{193}Bi		7104.5	5.	7104	3	−0.0	o			Jya		99Sm07	
		7103.5	6.			0.1	–			Jya		05Uu02	
		7107.5	5.			−0.6	–			Anv		14Ka23	
	ave.	7105	3			−0.1	1	98	82 ^{197}At			average	
^{197}At^{m}(α)^{193}Bi^{m}		6846.2	10.	6844	4	−0.2	o			Lvn		86Co12	
		6846.2	5.			−0.4	–			Jya		99Sm07	
		6845.2	9.			−0.1	–			Jya		05Uu02	
		6837.0	16.			0.5	U			Anv		14Ka23	
	ave.	6846	4			−0.4	1	94	58 ^{197}At^{m}			average	
^{197}Rn(α)^{193}Po		7410.8	20.	7411	7	−0.0	o			RIa		95No.A	
		7411.8	30.			−0.0	U			RIa		95Mo14	
		7410.8	7.				3			Jya		96En02	
^{197}Rn^{m}(α)^{193}Po^{m}		7523.1	30.	7509	6	−0.5	U			RIa		95Mo14	
		7508.7	7.			0.1	3			Jya		96En02	
		7510.7	14.			−0.1	3			Jya		05Uu02	
^{197}Fr(α)^{193}At^{m}		7888.4	15.				9			Anv		13Ka16	
^{196}Pt(n,γ)^{197}Pt		5846.4	0.4	5846.55	0.26	0.4	–					78Ya07	Z
		5846.0	0.9			0.6	–			ILn		81Ho.B	Z
		5846.6	0.5			−0.1	–			BNn		83Ca04	Z
		5846.0	0.7			0.8	–			Bdn		06Fi.A	
^{196}Pt(d,p)^{197}Pt		3627	20	3621.99	0.26	−0.3	U			Pit		64Co11	
		3606	20			0.8	U			Tal		78Ya07	
^{196}Pt(n,γ)^{197}Pt	ave.	5846.36	0.27	5846.55	0.26	0.7	1	94	65 ^{197}Pt			average	
^{197}Au(γ,n)^{196}Au		−8057	22	−8073.7	2.5	−0.8	U			Phi		60Ge01	
		−8080	5			1.3	–			McM		79Ba06	
		−8072	7			−0.2	–					79Be.A	
^{197}Au(d,t)^{196}Au		−1820	30	−1816.4	2.5	0.1	U			Pit		64Co11	
^{197}Au(γ,n)^{196}Au	ave.	−8077	4	−8073.7	2.5	0.9	1	39	38 ^{196}Au			average	
^{196}Hg(n,γ)^{197}Hg		6785.3	1.5	6785.6	1.5	0.2	1	97	85 ^{197}Hg	BNn		78Zg.A	Z
^{197}Ir(β^-)^{197}Pt		2000	200	2196	11	1.0	U					61Ho10	
^{197}Pt(β^-)^{197}Au		719.0	0.6	720.0	0.5	1.7	1	70	36 ^{197}Au			71Pr03	
^{197}Hg(ε)^{197}Au		415	20	600	3	9.3	B					65De20	*
		610	100			−0.1	U					92Da14	*
^{197}Tl(β^+)^{197}Hg		2220	100	2185	14	−0.4	U					61Ju05	
^{197}Pb^{m}(IT)^{197}Pb		319.31	0.11	319.31	0.11	−0.0	1	100	86 ^{197}Pb			Ens053	
*^{197}Hg−u	$M-A$=–30221(28) keV for ^{197}Hg^{m} at 298.93 keV											Nub269	**
*^{197}Pb−u	$M-A$=–24405(28) keV for ^{197}Pb^{m} at 319.31 keV											Nub269	**
*^{197}Bi−u	$M-A$=–19650(90) keV for mixture gs+m at 529(8) keV											Nub269	**
*^{197}Bi−^{133}Cs$_{1.481}$	Q =118887(12) μu $M-A$=–19690(11) keV corrected by –16(22) keV due to possible											08We02	**
*	contamination from ^{197}Bi^{m}											08We02	**
*^{197}Po−u	$M-A$=–13330(110) keV for mixture gs+m at 196(12) keV											Nub269	**
*^{197}Po−u	$M-A$=–13210(32) keV for mixture gs+m at 196(12) keV											Nub269	**
*^{197}Po(α)^{193}Pb	Also E_α=6283(5) keV from uncorrelated decays											96Ta18	**
*^{197}Po^{m}(α)^{193}Pb^{m}	Also E_α=6381(5) keV from uncorrelated decays											96Ta18	**
*^{197}Hg(ε)^{197}Au	pK=0.54(0.06) to 3/2$^+$ level at 268.788 keV											Ens053	**
*^{197}Hg(ε)^{197}Au	pK=0.746(0.033) to 268.75 level → Q =574(+139–62) keV											Ens053	**
^{198}Os−u		−25415	30				2			GS5	1.0	26Fe.B	
^{198}Ir−u		−27654	27				2			GS5	1.0	26Fe.B	
^{198}Hg−^{161}Dy ^{37}Cl		74130	60	73927.5	0.9	−0.8	U			R04	4.0	64De15	
^{198}Hg−^{163}Dy ^{35}Cl		68979	37	69179.6	0.9	1.4	U			R04	4.0	64De15	
^{198}Hg−u		−33231.6	0.6	−33230.8	0.5	1.3	1	67	67 ^{198}Hg	ST2	1.0	02Bf02	
^{198}Tl−^{133}Cs$_{1.489}$		111228.7	8.1				2			MA8	1.0	14Bo26	*
^{198}Pb−^{208}Pb$_{.952}$		−5748	23	−5757	9	−0.4	–			MA6	1.0	01Sc41	
	ave.	−5739	18			−1.0	1	26	26 ^{198}Pb			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{198}Pb−u		−27990	104	−27986	9	0.0	U				GS1	1.0	00Ra23	
		−27951	30			−1.2	R				GS2	1.0	05Li24	
^{198}Bi−u		−20900	120	−20780.9	1.5	1.0	o				GS1	1.0	00Ra23	*
		−20794	30			0.4	U				GS2	1.0	05Li24	
		−20780.9	1.5			−0.0	1	100	100	^{198}Bi	SH2	1.0	22Mo.A	
^{198}Bi^{m}−u		−20652.4	1.7				2				SH2	1.0	22Mo.A	
^{198}Bi^{n}−u		−20222	30	−20514.1	1.6	−9.7	B				GS2	1.0	05Li24	
		−20507.47	0.99			−6.7	B				SH2	1.0	22Mo.A	*
^{198}Po−^{208}Pb$_{.952}$		5616	24	5614	18	−0.1	1	54	54	^{198}Po	MA6	1.0	01Sc41	
^{198}Po−u		−16600	104	−16615	18	−0.1	U				GS1	1.0	00Ra23	
^{198}At−^{133}Cs$_{1.489}$		133570.0	7.3	133580	5	1.4	o				MA8	1.0	13Ma.A	
		133573.7	6.3			1.0	1	70	70	^{198}At	MA8	1.0	13St25	
^{198}At^{m}−^{133}Cs$_{1.489}$		133898	39	133866	6	−0.8	U				MA8	1.0	13St25	
^{198}Hg ^{35}Cl−^{196}Hg ^{37}Cl		3885.91	1.66	3885	3	−0.2	1	54	54	^{196}Hg	H33	2.5	80Ko25	
^{198}Pt−^{197}Au$_{1.005}$		1494.7	3.0	1494.2	2.2	−0.2	1	54	53	^{198}Pt	CP1	1.0	05Sh52	
^{198}Po(α)^{194}Pb		6312.8	5.	6309.7	1.4	−0.6	U						67Si09	Z
		6305.7	5.			0.8	U						67Tr06	Z
		6301.2	8.			1.1	U						71Ho01	Z
		6311.1	3.			−0.5	–				Bka		82Bo04	Z
		6307.7	5.			0.4	U				Lvn		93Wa04	
		6309.7	5.0			−0.0	U				Ara		96Ta18	*
		6309.3	1.7			0.2	–				Tex		12Fo09	
	ave.	6309.7	1.4			−0.0	1	100	65	^{194}Pb			average	
^{198}At(α)^{194}Bi		6887.5	5.	6889.4	1.9	0.4	2						67Tr06	Z
		6904.9	7.			−2.2	U				Ora		75Ba.B	Z
		6889.4	15.			−0.0	U						80Ew03	Z
		6893.3	3.5			−1.1	2				Lvn		92Hu04	*
		6892.5	4.			−0.8	o				Jya		96En01	
		6887.4	6.			0.3	2				Jya		05Uu02	
		6888.4	3.6			0.3	2				Tex		12Fo09	
		6886.4	5.			0.6	2				Anv		14Ka23	
		6888.9	8.2			0.1	2				ISa		19Gh11	*
		6890	15			−0.0	U				Lza		25Zh34	
^{198}At^{m}(α)^{194}Bi^{n}		6990.0	5.	6992.8	2.3	0.6	–						67Tr06	Z
		6997.5	10.			−0.5	–						80Ew03	Z
		6997.6	4.			−1.2	–				Lvn		92Hu04	
		6996.6	4.			−0.9	o				Jya		96En01	
		6991.5	6.			0.2	–				Jya		05Uu02	
		6990.5	5.			0.5	–				Anv		14Ka23	
		6996.9	8.2			−0.5	–				ISa		19Gh11	*
	ave.	6993.7	2.3			−0.4	1	95	58	^{194}Bi^{n}			average	
^{198}Rn(α)^{194}Po		7344.7	10.	7349	4	0.5	4						84Ca32	
		7353.8	5.			−0.9	4				Lvn		95Bi17	
		7344.7	6.			0.8	4				Jya		96En02	
^{198}Fr(α)^{194}At		7869.2	20.4				2				Anv		13Ka16	*
^{198}Fr^{m}(α)^{194}At^{m}		7889.6	20.4				3				Anv		13Ka16	
^{198}Pt(^{14}C,^{16}O)^{196}Os		6130	40				2				BNL		83Bo29	
^{198}Hg(d,α)^{196}Au		10890.0	5.0	10893.9	2.5	0.8	1	26	26	^{196}Au	Mun		03WiZV	
^{198}Pt(t,α)^{197}Ir		10885	20	10845	12	−2.0	1	33	33	^{197}Ir	LAl		83Ci01	
^{198}Pt(p,d)^{197}Pt		−5332	3	−5330.6	2.1	0.5	1	47	46	^{198}Pt	Ors		78Be09	*
^{198}Pt(d,t)^{197}Pt		−1305	20	−1297.9	2.1	0.4	U				Pit		64Co11	
		−1311	20			0.7	U				Tal		78Ya07	
^{197}Au(n,γ)^{198}Au		6512.35	0.11	6512.36	0.09	0.1	–				ILn		79Br26	Z
		6512.32	0.16			0.3	–				Bdn		06Fi.A	
^{197}Au(d,p)^{198}Au		4282	30	4287.79	0.09	0.2	U				Pit		64Co11	
		4298	5			−2.0	U				MIT		67Sp09	
^{197}Au(n,γ)^{198}Au	ave.	6512.34	0.09	6512.36	0.09	0.2	1	99	63	^{197}Au			average	
^{198}Au(β^{-})^{198}Hg		1372.3	0.7	1373.6	0.5	1.8	–						65Ke04	*
		1372.8	1.2			0.6	–						65Pa08	*
	ave.	1372.4	0.6			1.9	1	66	44	^{198}Au			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{198}Tl(β^+)^{198}Hg		3460	80	3426	8	−0.4	U					61Gu02	
^{198}Bi^n(IT)^{198}Bi		248.5	0.5	248.5	0.5	−0.0	1	99	99 ^{198}Bi^n	Lvn		92Hu04	
^{198}At^m(IT)^{198}At		265	3	266.6	2.7	0.5	1	84	57 ^{198}At^m	ISa		19Gh11	*
*^{198}Tl−^{133}Cs$_{1.489}$	D_M=111812.3(8.1) μu for ^{198}Tl^m at 543.6(0.4) keV; $M-A$=–26985.1(7.5) keV											Nub269	**
*^{198}Bi−u	$M-A$=–19350(100) keV for mixture gs+m+n at 119.7(2.1) and											Nub269	**
*^{198}Bi−u	248.5(0.5) keV											Nub269	**
*^{198}Bi^n−u	at variance with ME of gs. + 248.5(0.5) keV Eg from 10^- to 7^+											Fgk25a	**
*^{198}Po(α)^{194}Pb	Also E_α=6182(5) keV from uncorrelated decays											96Ta18	**
*^{198}At(α)^{194}Bi	E_α=6755(4), 6539(10), 6360(10) to ground state, 218, 396 levels											92Hu04	**
*^{198}At(α)^{194}Bi	E_α=6358(8) keV to 399.7(0.2) keV; Also E_α=6275(8), 6361(9),											19Gh11	**
*^{198}At(α)^{194}Bi	6535(8) to 485.6(0.7), 382.4(0.1), 218.2(0.1) keV											19Gh11	**
*^{198}At^m(α)^{194}Bi^n	E_α=6753(8) keV to level at 104.5(0.2) keV; Also E_α=6322(12),											19Gh11	**
*^{198}At^m(α)^{194}Bi^n	6338(8) keV to levels at 538.3(0.4), 525.4(0.2) keV											19Gh11	**
*^{198}Fr(α)^{194}At	Evaluator's interpretation of Fig. 3d, no 210 keV γ											GAu	**
*^{198}Pt(p,d)^{197}Pt	$Q-Q$(^{196}Pt(p,d))=365(3) keV											AHW	**
*^{198}Au(β^-)^{198}Hg	E_{β^-}=960.5(0.7) 961.0(1.2) respectively, to 2^+ level at 411.80251 keV											Ens163	**
*^{198}At^m(IT)^{198}At	E(γ)=511(3)+130.0(0.1)+150.8(0.1) - E(γ)=527.3(0.2)											19Gh11	**
^{199}Ir−u		−26194	29				2			GS5	1.0	26Fe.B	
^{199}Pt−u		−29365	25	−29402.5	2.3	−1.5	U			GS5	1.0	26Fe.B	
^{199}Hg−C$_2$ ^{35}Cl$_5$		124023.43	0.53	124017.4	0.6	−4.5	B			H34	2.5	80Ko25	
		124017.21	0.37			0.2	1	36	33 ^{199}Hg	H48	2.5	03Ba49	
^{199}Hg−^{183}W O		23144.4	0.9	23141.9	0.9	−1.1	1	16	11 ^{183}W	H48	2.5	03Ba49	
^{199}Hg−^{162}Dy ^{37}Cl		75661	41	75574.2	0.9	−0.5	U			R04	4.0	64De15	
^{199}Hg−^{164}Dy ^{35}Cl		70087	31	70247.7	0.9	1.3	U			R04	4.0	64De15	
^{199}Hg−^{164}Er ^{35}Cl		70310	80	70220.8	0.9	−0.3	U			R04	4.0	64De15	
^{199}Tl−u		−30123	30				2			GS2	1.0	05Li24	
^{199}Pb−u		−27028	137	−27087	7	−0.4	U			GS2	1.0	05Li24	*
^{199}Bi−u		−22328	31	−22327	11	0.0	–			GS2	1.0	05Li24	
		−22263	30			−2.1	–			GS2	1.0	05Li24	*
	ave.	−22294	22			−1.5	1	28	28 ^{199}Bi			average	
^{199}Po−u		−16249	144	−16362	6	−0.8	U			GS1	1.0	00Ra23	*
		−16327	38			−0.9	R			GS2	1.0	05Li24	
		−16339	38			−0.6	R			GS2	1.0	05Li24	*
^{199}Po^m−^{133}Cs$_{1.496}$		125416.5	5.3	125417	5	0.0	1	99	99 ^{199}Po^m	MA8	1.0	17Al34	
^{199}Bi^m(α)^{195}Tl		5598.7	6.	5599	6	0.1	1	93	56 ^{195}Tl			66Ma51	
^{199}Po(α)^{195}Pb		6074.1	2.	6074.3	1.9	0.1	–			Dba		68Go.B	Z
		6075.3	5.0			−0.2	–			Ara		96Ta18	
	ave.	6074.3	1.9			0.0	1	100	99 ^{199}Po			average	
^{199}Po^m(α)^{195}Pb^m		6190.7	5.	6183.3	1.7	−1.5	–					67Si09	Z
		6177.6	5.1			1.1	–					67Tr06	Z
		6182.3	3.1			0.3	–			Dba		68Go.B	Z
		6183.5	3.			−0.1	–			Bka		82Bo04	Z
		6183.5	5.0			−0.1	–			Ara		96Ta18	*
		6173.3	3.6			2.8	B			Tex		12Fo09	
	ave.	6183.3	1.7			−0.0	1	100	99 ^{195}Pb^m			average	
^{199}At(α)^{195}Bi		6775.1	5.	6777.3	1.2	0.4	–					67Tr06	Z
		6781.3	3.			−1.3	–			Ora		75Ba.B	Z
		6775.4	5.0			0.4	–			Ara		96Ta18	
		6779.4	6.			−0.4	U			Jya		05Uu02	
		6776.8	1.5			0.3	–			Tex		12Fo09	
	ave.	6777.4	1.2			−0.1	1	100	89 ^{199}At			average	
^{199}Rn(α)^{195}Po		7133.7	15.	7132	4	−0.1	3					80Di07	
		7132.7	10.			−0.1	3					82Hi14	
		7138.8	10.			−0.7	3					84Ca32	
		7112.2	15.			1.3	o			Jya		96Le09	
		7132.6	6.			−0.1	3			Jya		05Uu02	
		7121.4	10.2			1.0	3			Anv		14Ka23	
^{199}Rn^m(α)^{195}Po^m		7205.1	15.	7203	4	−0.1	2					80Di07	
		7205.1	10.			−0.2	2					82Hi14	
		7204.1	10.			−0.1	2					84Ca32	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{199}Rn$^{m}(\alpha)^{195}$Po^{m}		7205.1	15.	7203	4	−0.1	o			Jya		96Le09	
		7205.1	6.			−0.3	2			Jya		05Uu02	
		7194.9	10.2			0.8	2			Anv		14Ka23	
^{199}Fr$(\alpha)^{195}$At		7812.3	40.	7817	10	0.1	U					99Ta20	
		7821.5	11.			−0.4	5			Anv		13Ka16	
		7801.1	20.			0.8	5			Jya		13Uu01	
		7813	34			0.1	U			Lza		24Wa14	
^{199}Fr$^{m}(\alpha)^{195}$At^{m}		7833.7	6.	7833	6	−0.2	3			Anv		13Ka16	
		7825.6	15.3			0.5	3			Jya		13Uu01	
^{199}Fr$^{n}(\alpha)^{195}$At^{p}		7968.4	20.				4			Jya		13Uu01	
^{199}Hg(p,t)^{197}Hg		−6734	29	−6667	3	2.3	U			Pri		81Ko13	
		−6658	8			−1.2	1	16	15 ^{197}Hg	Ors		82Be21	
^{198}Pt(^{18}O,^{17}F)^{199}Ir		−8240	41	−8239	27	0.0	U					95Zh10	
^{198}Pt(n,γ)^{199}Pt		5602	3	5556.0	0.5	−15.3	B					68Sa13	
		5556.0	0.5				2			BNn		83Ca04	Z
^{198}Pt(d,p)^{199}Pt		3347	20	3331.4	0.5	−0.8	U			Pit		64Co11	
^{198}Au(n,γ)^{199}Au		7584.27	0.15	7584.28	0.06	0.1	o			ILn		79Br26	Z
		7584.28	0.06			−0.0	1	100	80 ^{199}Au	ILn		91Ma65	
^{198}Hg(n,γ)^{199}Hg		6665.2	0.5	6663.1	0.6	−4.1	B			CRn		75Lo03	
^{199}Hg(γ,n)^{198}Hg		−6590	90	−6663.1	0.6	−0.8	U			Phi		60Ge01	
^{199}Pt$(\beta^{-})^{199}$Au		1690	50	1705.5	2.1	0.3	U					64Jo09	
^{199}Au$(\beta^{-})^{199}$Hg		453.0	1.0	452.4	0.6	−0.6	1	37	20 ^{199}Au			68Be06	
^{199}Tl$(\beta^{+})^{199}$Hg		1420	150	1487	28	0.4	U					75Ma05	*
^{199}Pb$(\beta^{+})^{199}$Tl		2870	110	2828	29	−0.4	U					70Do.A	*
^{199}Bi^{m}(IT)^{199}Bi		667	5	667	3	−0.0	–					80Br23	
		667	5			−0.0	–					85St02	
	ave.	667	4			−0.1	1	98	64 ^{199}Bi^{m}			average	
^{199}At^{m}(IT)^{199}At		255	20	244.0	1.0	−0.5	o			Jya		13Ja06	*
		244	1				2			Jya		14Au03	
*^{199}Pb−u	$M-A$=–24961(28) keV for mixture gs+m at 429.5(2.7) keV											Nub269	**
*^{199}Bi−u	$M-A$=–20071(28) keV for ^{199}Bi^{m} at 667(3) keV											Nub269	**
*^{199}Po−u	$M-A$=–14980(100) keV for mixture gs+m at 311.8(2.7) keV											Nub269	**
*^{199}Po−u	$M-A$=–14909(35) keV for ^{199}Po^{m} at 311.8(2.7) keV											Nub269	**
*^{199}Po$^{m}(\alpha)^{195}$Pb^{m}	Also E_α=6059(5) keV from uncorrelated decays											96Ta18	**
*^{199}Tl$(\beta^{+})^{199}$Hg	KL+<500(100) giving Q <1620, (1/2^{-},3/2^{-}) level at 1221.17 fed. Reanalyzed											Ens073	**
*^{199}Pb$(\beta^{+})^{199}$Tl	p^{+} =0.04(0.01) to 3/2^{+} level at 366.89 keV, recalculated											Ens073	**
*^{199}At^{m}(IT)^{199}At	Combining lines 110(20) 160(20) 240(30) as read from Fig. 8a											GAu	**
^{200}Ir−u		−23164	18				2			GS5	1.0	26Fe.B	
^{200}Au−u		−29237	34	−29203	5	1.0	U			GS3	1.0	08ChZI	
^{200}Au^{m}−u		−28135	33	−28163	28	−0.8	1	73	73 ^{200}Au^{m}	GS3	1.0	08ChZI	
^{200}Hg−C ^{13}C ^{35}Cl$_5$		120707.97	1.22	120708.5	0.6	0.2	U			H34	2.5	80Ko25	
^{200}Hg−^{165}Ho ^{35}Cl		69116	33	69145.3	1.0	0.2	U			R04	4.0	64De15	
^{200}Hg−^{163}Dy ^{37}Cl		73527	42	73687.4	0.9	1.0	U			R04	4.0	64De15	
^{200}Pb−u		−28179	30	−28183	11	−0.1	R			GS2	1.0	05Li24	
^{200}Bi−u		−21888	57	−21855	12	0.6	R			GS2	1.0	05Li24	*
^{200}Po−u		−18170	104	−18184	8	−0.1	U			GS1	1.0	00Ra23	
		−18204	30			0.7	U			GS2	1.0	05Li24	
^{200}At−u		−9648.3	1.6	−9642.2	1.5	3.8	B			SH2	1.0	22Mo.A	
^{200}At^{m}−u		−9520.4	1.6	−9520.8	1.2	−0.3	1	54	54 ^{200}At^{m}	SH2	1.0	22Mo.A	
^{200}At^{n}−u		−9245.7	1.6	−9273.0	1.2	−17.0	B			SH2	1.0	22Mo.A	
^{200}Hg−^{208}Pb$_{.962}$		−9205	28	−9210.9	0.6	−0.2	U			MA6	1.0	01Sc41	
^{200}Hg ^{35}Cl−^{198}Hg ^{37}Cl		4525	2	4507.8	0.6	−2.1	U			H17	4.0	64Mc07	
		4508.80	0.48			−0.8	1	28	16 ^{200}Hg	H33	2.5	80Ko25	
^{200}Po$(\alpha)^{196}$Pb		5979.8	5.	5981.7	1.8	0.4	–					67Si09	Z
		5980.0	3.			0.5	–					67Tr06	Z
		5983.4	3.			−0.6	–					70Ra14	Z
		5981.8	5.0			−0.0	–			Ara		96Ta18	*
	ave.	5981.5	1.9			0.1	1	99	77 ^{196}Pb			average	
^{200}At$(\alpha)^{196}$Bi		6594.9	5.	6595.2	1.2	0.1	–					67Tr06	Z
		6596.9	2.			−0.9	–			Ora		75Ba.B	Z

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{200}At(α)^{196}Bi		6593.1	5.	6595.2	1.2	0.4	o			Lvn		87Va09	
		6596.1	2.			−0.5	–			Lvn		92Hu04	
		6593.1	5.0			0.4	–			Ara		96Ta18	
		6599.1	6.			−0.6	U			Jya		05Uu02	
	ave.	6596.2	1.3			−0.8	1	82	72 ^{200}At			average	
^{200}At^{m}(α)^{196}Bi		6708.3	5.	6708.2	1.3	0.0	–			Ora		75Ba.B	Z
		6705.4	5.			0.6	o			Lvn		87Va09	
		6709.5	3.			−0.4	–			Lvn		92Hu04	
	ave.	6709.1	2.6			−0.3	1	26	15 ^{200}At^{m}			average	
^{200}At^{m}(α)^{196}Bi^{m}		6542.8	5.	6542.6	1.1	−0.0	–					67Tr06	Z
		6542.9	2.			−0.1	–			Ora		75Ba.B	Z
		6540.0	5.			0.5	o			Lvn		87Va09	
		6542.1	2.			0.3	–			Lvn		92Hu04	
		6545.1	5.0			−0.5	–			Ara		96Ta18	
		6544.1	6.			−0.2	U			Jya		05Uu02	
	ave.	6542.7	1.3			−0.0	1	71	45 ^{196}Bi^{m}			average	
^{200}At^{m}(α)^{196}Bi^{n}		6439.5	5.	6442.8	1.4	0.7	U					67Tr06	*
		6438.5	5.			0.9	U			Ora		75Ba.B	*
		6433.8	5.			1.8	o			Lvn		87Va09	*
		6439.2	3.			1.2	U			Lvn		92Hu04	*
		6430.5	5.0			2.5	U			Ara		96Ta18	*
		6436.7	6.			1.0	U			Jya		05Uu02	*
^{200}Rn(α)^{196}Po		7020.6	10.	7043.4	2.1	2.3	U					67Va.A	
		7050.3	8.			−0.9	U					71Ho01	
		7040.1	10.			0.3	U			Lvn		84Co.A	
		7043.5	2.5			−0.1	2			Lvn		93Wa04	
		7042.1	12.			0.1	o			Ara		95Le04	
		7039.0	10.			0.4	o			Jya		96Le09	
		7042.1	5.1			0.2	2			Ara		96Ta18	
		7044.1	6.			−0.1	2			Jya		05Uu02	
		7055.4	30.			−0.4	U			Anv		10He25	
^{200}Fr(α)^{196}At		7653.4	30.	7623	4	−1.0	U			RIa		95Mo14	
		7620.7	9.			0.2	5			Jya		96En01	
		7625.8	12.			−0.3	o			Anv		05De01	
		7620.7	15.			0.1	o			Jya		13Uu01	*
		7622.7	5.			−0.0	5			Anv		14Ka23	
		7618.6	12.2			0.3	5			ISa		19Gh11	
		7633	15			−0.7	5			Lza		22Hu12	
^{200}Fr^{m}(α)^{196}At^{m}		7704.4	15.				4			Jya		96En01	*
^{198}Pt(t,p)^{200}Pt		4356	20				2					81Ci01	
^{199}Hg(n,γ)^{200}Hg		8029.1	0.3	8028.52	0.11	−1.9	–			BNn		67Sc30	Z
		8029.6	0.5			−2.2	U			CRn		75Lo03	Z
		8028.51	0.18			0.0	–			ILn		79Br25	Z
		8028.37	0.17			0.9	–			Bdn		06Fi.A	
	ave.	8028.53	0.11			−0.1	1	98	64 ^{200}Hg			average	
^{200}Au(β^-)^{200}Hg		2273	100	2301	5	0.3	U					59Ro53	*
		2200	100			1.0	U					60Gi01	
		2260	70			0.6	U					72He36	*
		2260	100			0.4	o					72Pa24	
^{200}Au^{m}(β^-)^{200}Hg		3202	50	3270	26	1.4	1	27	27 ^{200}Au^{m}			72Cu07	*
^{200}Tl(β^+)^{200}Hg		2450	10	2456	6	0.6	2					57He43	*
		2459	7			−0.4	2					62Va10	*
^{200}At^{n}(IT)^{200}At^{m}		230.9	0.2	230.89	0.20	−0.1	1	100	96 ^{200}At^{n}	Lvn		92Hu04	
*^{200}Bi−u	$M-A$=–20338(28) keV for mixture gs+m at 100#70 keV											Nub269	**
*^{200}Po(α)^{196}Pb	Also E_α=5863(5) keV from uncorrelated decays											96Ta18	**
*^{200}At^{m}(α)^{196}Bi^{n}	E_α=6536.7(5,Z) from ^{200}At^{n} 230.9 above ^{200}At^{m}											92Hu04	**
*^{200}At^{m}(α)^{196}Bi^{n}	E_α=6535.8(5,Z) from ^{200}At^{n} 230.9 above ^{200}At^{m}											92Hu04	**
*^{200}At^{m}(α)^{196}Bi^{n}	E_α=6301(5); 6535(5) from ^{200}At^{n} 230.9 above ^{200}At^{m}											92Hu04	**
*^{200}At^{m}(α)^{196}Bi^{n}	E_α=6306(5); 6538(3) from ^{200}At^{n} 230.9 above ^{200}At^{m}											92Hu04	**
*^{200}At^{m}(α)^{196}Bi^{n}	E_α=6528(5) from ^{200}At^{n} 230.9 above ^{200}At^{m}											92Hu04	**
*^{200}At^{m}(α)^{196}Bi^{n}	E_α=6534(6) from ^{200}At^{n} 230.9 above ^{200}At^{m}											92Hu04	**
*^{200}Fr(α)^{196}At	Same group, but much less events than preceding item											WgM151	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{200}Fr^m(\alpha)^{196}At^m$		Correlated with $^{196}At^m$ E_α=6880(15); two events only										96En01	**
$*^{200}Au(\beta^-)^{200}Hg$		E_{β^-}=2250(200) to ground state, and 700(100) to levels 1^+ at 1570.275,										Ens077	**
*		2^+ at 1573.663, 2^+ at 1593.423 keV										Ens077	**
$*^{200}Au(\beta^-)^{200}Hg$		E_{β^-}=2260(100), 670(70) to ground state, 2^+ level at 1593.423 keV										Ens077	**
$*^{200}Au^m(\beta^-)^{200}Hg$		E_{β^-}=560(50) to 11^- level at 2641.54 keV										Ens077	**
$*^{200}Tl(\beta^+)^{200}Hg$		E_{β^+}=1052(10) 1069(7) respectively, to 2^+ level at 367.943 keV, and other E_{β^+}										Ens077	**
$^{201}Ir-u$		−21549	23				2			GS5	1.0	26Fe.B	
$^{201}Pt-u$		−25572.0	6.1	−25571	6	0.2	1	99	99 ^{201}Pt	GS5	1.0	26Fe.B	
$^{201}Hg-^{185}Re$ O		22440	5	22429.8	1.1	−0.8	U			H48	2.5	03Ba49	
$^{201}Hg-C_2\ ^{35}Cl_4\ ^{37}Cl$		128995.43	0.61	128989.4	0.7	−4.0	B			H34	2.5	80Ko25	
$^{201}Hg-^{164}Dy\ ^{37}Cl$		75086	42	75219.7	1.0	0.8	U			R04	4.0	64De15	
$^{201}Hg-^{166}Er\ ^{35}Cl$		71186	35	71149.3	0.8	−0.3	U			R04	4.0	64De15	
$^{201}Pb-u$		−27418	198	−27131	13	1.4	U			GS2	1.0	05Li24	*
$^{201}Bi-u$		−22935	30	−23006	13	−2.4	R			GS2	1.0	05Li24	
		−22995	30			−0.4	R			GS2	1.0	05Li24	*
$^{201}Po-u$		−17760	190	−17736	5	0.1	U			GS1	1.0	00Ra23	*
		−17649	30			−2.9	U			GS2	1.0	05Li24	
$^{201}Po^m-u$		−17305	30	−17282	5	0.8	U			GS2	1.0	05Li24	
$^{201}At-u$		−11573	31	−11578.8	1.1	−0.2	U			GS2	1.0	05Li24	
		−11578.5	1.2			−0.3	1	86	86 ^{201}At	SH2	1.0	22Mo.A	
$^{201}Hg\ ^{35}Cl-^{199}Hg\ ^{37}Cl$		4981	2	4971.9	0.6	−1.1	U			H17	4.0	64Mc07	
		4972.65	0.37			−0.8	1	43	32 ^{201}Hg	H33	2.5	80Ko25	
		4971.8	1.0			0.1	U			H48	2.5	03Ba49	
$^{201}Bi(\alpha)^{197}Tl$		4500.3	6.				4					66Ma51	*
$^{201}Po(\alpha)^{197}Pb$		5793.9	5.	5799.5	1.2	1.1	–					67Tr06	Z
		5799.4	2.			0.0	–			Dba		68Go.B	Z
		5800.4	4.			−0.2	–					70Ra14	Z
	ave.	5799.0	1.7			0.3	1	50	36 ^{201}Po			average	
$^{201}Po^m(\alpha)^{197}Pb^m$		5899.0	5.1	5903.6	1.2	0.9	–					67Tr06	Z
		5904.5	2.0			−0.5	–			Dba		68Go.B	Z
		5903.9	4.1			−0.1	–					70Ra14	Z
	ave.	5903.8	1.7			−0.1	1	50	37 $^{201}Po^m$			average	
$^{201}At(\alpha)^{197}Bi$		6470.7	3.	6472.8	1.6	0.7	2					67Tr06	Z
		6476.2	5.			−0.7	U					74Ho27	Z
		6474.0	2.			−0.6	2			Ora		75Ba.B	Z
		6471.0	5.0			0.4	U			Ara		96Ta18	
		6472.0	4.			0.2	2			Anv		05De01	
$^{201}Rn(\alpha)^{197}Po$		6862.8	8.	6860.7	2.3	−0.3	U					67Va.A	
		6858.8	8.			0.2	U					71Ho01	
		6866.9	20.			−0.3	U			GSa		87He10	
		6860.5	2.5			0.1	2			Lvn		93Wa04	
		6863.8	7.			−0.4	o			Ara		95Le04	
		6861.8	5.0			−0.2	2			Ara		96Ta18	
$^{201}Rn^m(\alpha)^{197}Po^m$		6906.8	5.	6909.5	2.1	0.5	3					67Va17	Z
		6909.0	8.			0.1	U					71Ho01	Z
		6907.7	20.			0.1	U			GSa		87He10	
		6909.9	2.5			−0.1	3			Lvn		93Wa04	
		6915.9	7.			−0.9	o			Ara		95Le04	
		6910.7	5.0			−0.3	3			Ara		96Ta18	
		6925.1	30.			−0.5	U			Anv		10He25	
$^{201}Fr(\alpha)^{197}At$		7538.0	15.	7519	4	−1.3	U					80Ew03	
		7510.8	7.			1.1	o			Jya		96En01	
		7529.1	7.			−1.4	o			Anv		05De01	
		7519.0	8.			−0.0	2			Jya		05Uu02	
		7519.0	5.			−0.0	2			Anv		14Ka23	
$^{201}Fr^m(\alpha)^{197}At^m$		7605.7	8.	7603	5	−0.3	2			Jya		05Uu02	
		7596.4	8.			0.8	2			Anv		14Ka23	
		7608.7	9.2			−0.6	2			Jya		20Au01	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{201}Ra(α)^{197}Rn		8001.5	12.				4			Anv		14Ka23	
^{201}Ra^{m}(α)^{197}Rn^{m}		8065.8	20.				4			Jya		05Uu02	
^{201}Hg(γ,n)^{200}Hg		−6210	70	−6230.8	0.6	−0.3	U			Phi		60Ge01	
^{201}Pt(β^-)^{201}Au		2660	50	2582	6	−1.6	1	2	1 ^{201}Pt			63Go06	
^{201}Au(β^-)^{201}Hg		1270	100	1262	3	−0.1	U					72Pa24	
^{201}Tl(ε)^{201}Hg		470	70	481	14	0.2	U					60Gu05	*
^{201}Pb(β^+)^{201}Tl		1900	40	1910	18	0.2	1	19	11 ^{201}Tl			79Do09	*
^{201}Po^{m}(IT)^{201}Po		423.41	0.22	423.41	0.22	0.0	1	99	63 ^{201}Po^{m}			Ens229	
*^{201}Pb−u	$M-A$=–25225(28) keV for mixture gs+m at 629.1 keV											Nub269	**
*^{201}Bi−u	$M-A$=–20573(28) keV for ^{201}Bi^{m} at 846.35 keV											Nub269	**
*^{201}Po−u	$M-A$=–16330(100) keV for mixture gs+m at 423.41(0.22) keV											Nub269	**
*^{201}Bi(α)^{197}Tl	E_α=5240(6) from ^{201}Bi^{m} at 846.35 keV											Nub269	**
*^{201}Tl(ε)^{201}Hg	pK=0.70(0.04)to 1/2$^-$ level at 167.47 keV, recalculated											Ens073	**
*^{201}Pb(β^+)^{201}Tl	p$^+$ =10(2)×10^{-3} to 3/2$^+$ level at 331.16 keV											Ens073	**
^{202}Pt−u		−24425	34	−24361	27	1.9	o			GS3	1.0	08ChZI	
		−24361	27				2			GS3	1.0	12Ch19	
^{202}Au−u		−26202	34	−26150	5	1.5	o			GS3	1.0	08ChZI	
		−26144	25			−0.2	U			GS3	1.0	12Ch19	
^{202}Hg−C ^{13}C ^{35}Cl$_4$ ^{37}Cl		125976.01	1.32	125975.3	0.7	−0.2	U			H34	2.5	80Ko25	
C$_{16}$ H$_{10}$−^{202}Hg		107663	40	107606.9	0.7	−0.9	U			R08	1.5	69De19	
C$_{15}$ ^{13}C H$_9$−^{202}Hg		103102	60	103136.7	0.7	0.4	U			R08	1.5	69De19	
^{202}Hg−^{167}Er ^{35}Cl		69740	60	69734.9	0.7	−0.0	U			R04	4.0	64De15	
^{202}Hg−^{165}Ho ^{37}Cl		74470	50	74412.0	1.1	−0.3	U			R04	4.0	64De15	
^{202}Pb−u		−27823	30	−27849	4	−0.9	U			GS2	1.0	05Li24	*
^{202}Pb−^{133}Cs$_{1.519}$		115773.4	3.6	115770	4	−1.0	o			MA8	1.0	10Bo.A	
		115769.2	4.4			0.2	1	86	86 ^{202}Pb	MA8	1.0	14Bo26	
^{202}Bi−u		−22282	30	−22277	15	0.2	1	25	25 ^{202}Bi	GS2	1.0	05Li24	
^{202}Po−u		−19270	104	−19262	9	0.1	U			GS1	1.0	00Ra23	
		−19243	30			−0.6	U			GS2	1.0	05Li24	
^{202}At−u		−11457.7	1.0	−11458.4	2.0	−0.7	F			SH2	1.0	22Mo.A	*
^{202}At^{m}−u		−11252.9	1.9	−11228.0	2.2	13.1	F			SH2	1.0	22Mo.A	*
^{202}At^{n}−u		−11024.8	1.1	−11037.9	2.0	−11.9	F			SH2	1.0	22Mo.A	*
^{202}Hg ^{35}Cl$_2$−^{198}Hg ^{37}Cl$_2$		9774.87	1.06	9774.6	0.8	−0.1	U			H33	2.5	80Ko25	
^{202}Hg ^{35}Cl−^{200}Hg ^{37}Cl		5271	3	5266.7	0.6	−0.4	U			H17	4.0	64Mc07	
		5266.76	0.43			−0.0	1	32	23 ^{202}Hg	H33	2.5	80Ko25	
^{202}Tl−^{203}Tl$_{.995}$		−372.4	2.1	−373.1	1.8	−0.3	1	71	69 ^{202}Tl	MA8	1.0	17We09	
^{202}Po(α)^{198}Pb		5700.9	2.	5701.0	1.3	0.1	–			Dba		68Go.B	Z
		5701.6	3.			−0.2	–					70Ra14	Z
		5701.1	2.			−0.0	–					93Wa04	
	ave.	5701.1	1.3			−0.1	1	100	74 ^{198}Pb			average	
^{202}At(α)^{198}Bi		6259.9	2.	6259.0	1.2	−0.5	–					63Ho18	Z
		6256.8	3.			0.7	–					67Tr06	Z
		6257.2	5.			0.3	–					74Ho27	Z
		6259.0	2.			−0.0	–			Ora		75Ba.B	*
		6260.0	5.0			−0.2	–			Lvn		92Hu04	*
		6257.1	6.0			0.3	–			Ara		96Ta18	
	ave.	6258.9	1.2			0.1	1	95	95 ^{202}At			average	
^{202}At^{m}(α)^{198}Bi^{m}		6355.8	3.	6353.8	1.3	−0.6	3					63Ho18	Z
		6351.7	3.			0.7	3					67Tr06	Z
		6353.2	5.			0.1	3					74Ho27	Z
		6353.9	2.			−0.0	3			Ora		75Ba.B	*
		6354	5			−0.0	3			Lvn		92Hu04	*
		6355.0	6.0			−0.2	3			Ara		96Ta18	
^{202}At^{n}(α)^{198}Bi^{n}		6404.1	5.1	6402.2	1.3	−0.4	1	6	5 ^{202}At^{n}	Lvn		92Hu04	*
^{202}Rn(α)^{198}Po		6771.0	3.	6773.8	1.8	0.9	–					67Va17	Z
		6772.3	10.			0.1	U			GSa		87He10	
		6775.3	2.5			−0.6	–			Lvn		93Wa04	
		6773.4	7.			0.1	o			Ara		95Le04	
		6775.4	5.0			−0.3	–			Ara		96Ta18	
	ave.	6773.8	1.8			−0.0	1	100	89 ^{202}Rn			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{202}Fr(α)^{198}At		7397.7	15.	7385	3	−0.8	2						80Ew03	*
		7382.5	11.			0.2	2				Lvn		92Hu04	*
		7389.6	6.			−0.7	o				Jya		96En01	*
		7387.6	8.			−0.3	2				Jya		05Uu02	
		7384.5	5.1			0.1	2				Anv		14Ka23	
		7383.3	8.2			0.2	2				ISa		19Gh11	*
		7380.4	15.3			0.3	2				Lza		25Zh34	
^{202}Fr^{m}(α)^{198}At		7643.7	8.2	7643	4	−0.1	1	27	24	^{202}Fr^{m}	ISa		19Gh11	*
^{202}Fr^{m}(α)^{198}At^{m}		7382.5	11.	7376	4	−0.6	–				Lvn		92Hu04	*
		7388.6	6.			−2.0	o				Jya		96En01	
		7381.5	8.			−0.7	–				Jya		05Uu02	
		7372.2	5.			0.8	–				Anv		14Ka23	
		7387.3	8.2			−1.4	U				ISa		19Gh11	*
	ave.	7376	4			0.1	1	82	76	^{202}Fr^{m}			average	
^{202}Ra(α)^{198}Rn		8019.1	60.	7880	7	−2.3	o				Jya		96Le09	
		7896.7	20.			−0.8	5				Jya		05Uu02	
		7878.3	7.1			0.3	5				Anv		14Ka23	
^{202}Hg(d,α)^{200}Au		10568.0	5.0				2				Mun		03WiZV	
^{202}Hg(t,α)^{201}Au		11567	15	11581	3	0.9	U				LAl		81Fl05	
^{202}Hg(d,^{3}He)^{201}Au−^{206}Pb()^{205}Tl		−979.9	3.1	−980	3	0.1	1	100	100	^{201}Au			94Gr07	
^{201}Hg(n,γ)^{202}Hg		7754.9	0.5	7753.92	0.12	−2.0	R				BNn		75Br02	Z
		7756.4	0.5			−5.0	B				CRn		75Lo03	Z
		7753.93	0.22			−0.0	–				Bdn		06Fi.A	
		7753.90	0.14			0.2	–				BNn		75Br02	*
^{202}Hg(γ,n)^{201}Hg		−7600	130	−7753.92	0.12	−1.2	U				Phi		60Ge01	
^{201}Hg(n,γ)^{202}Hg	ave.	7753.91	0.12	7753.92	0.12	0.1	1	99	68	^{201}Hg			average	
^{202}Au(β^{-})^{202}Hg		3500	300	2987	5	−1.7	U						67Wa23	
		2700	300			1.0	U						72Bu05	
^{202}Tl(ε)^{202}Hg		1242	20	1364.0	1.8	6.1	B						66Le06	*
		1220	30			4.8	B						65Le04	*
^{202}Pb(ε)^{202}Tl		55	20	41	4	−0.7	U						54Hu61	
^{202}At^{n}(IT)^{202}At		391.7	0.2	391.70	0.20	0.0	1	100	95	^{202}At^{n}	Lvn		92Hu04	
*^{202}Pb−u	$M-A$=–23747(28) keV for ^{202}Pb^{m} at 2169.83 keV												Nub269	**
*^{202}At−u	at variance with ME extracted using ME of ^{198}Bi and ^{202}At(α)^{198}Bi												HWJ25a	**
*^{202}At^{m}−u	at variance with ME extracted using ME of ^{198}Bi^{m} and ^{202}At^{m}(α)^{198}Bi^{m}												HWJ25a	**
*^{202}At^{n}−u	at variance with ME extracted using ME of ^{198}Bi^{n} and ^{202}At^{n}(α)^{198}Bi^{n}												HWJ25a	**
*^{202}At(α)^{198}Bi	Assignment to ^{202}At^{m} in reference; Z recalibrated												92Hu04	**
*^{202}At(α)^{198}Bi	E_α=6135(5); and 6277(5) from ^{202}At^{n}(α)^{198}Bi^{n}, with												92Hu04	**
*	^{202}At^{n}(IT)^{202}At=391.7(0.5) and ^{198}Bi^{n}(IT)^{198}Bi=248.5(0.5) keV												Nub269	**
*^{202}At^{m}(α)^{198}Bi^{m}	E_α=6228(5), 6070(10), 5929(10) to ground state, 164, 303 levels												92Hu04	**
*^{202}Fr(α)^{198}At	E_α=7251(10) has a doublet structure												92Hu04	**
*^{202}Fr(α)^{198}At	E_α=7237(8), is a doublet												92Hu04	**
*^{202}Fr(α)^{198}At	^{202}Fr E_α's in correlation with At daughters												96En01	**
*^{202}Fr(α)^{198}At	E_α=7105(8) keV to level at 130.0(0.1) keV; Also E_α=7086(8) keV												19Gh11	**
*^{202}Fr(α)^{198}At	to level at 154.2(0.3) keV												19Gh11	**
*^{202}Fr^{m}(α)^{198}At	E_α=7217(8) to level at 280.8(0.1) keV												19Gh11	**
*^{202}Fr^{m}(α)^{198}At^{m}	E_α=7237(8), is a doublet												92Hu04	**
*^{202}Fr^{m}(α)^{198}At^{m}	E_α=6724(8) keV to level at 527.3(0.2) keV above ^{198}At^{m} and at												19Gh11	**
*^{202}Fr^{m}(α)^{198}At^{m}	792(3) keV above ^{198}At; not independent												19Gh11	**
*^{201}Hg(n,γ)^{202}Hg	lsq fit to adopted levels and primary γs from 75Br02												FGK227	**
*^{202}Tl(ε)^{202}Hg	average 1235(+26−20), from pL/pK=0.305(0.020) to 2^{+} level at 959.94 keV,												66Le06	**
*^{202}Tl(ε)^{202}Hg	1245(+45−41), from pL/pK=0.196(0.002) to 2^{+} level at 439.512 keV												66Le06	**
*^{202}Tl(ε)^{202}Hg	1214(+40−22), from pK(to 959 keV)/pK(to 439 keV)=0.888(0.027)												65Le04	**
^{203}Pt−u		−21312	20				2				GS5	1.0	26Fe.C	
^{203}Au−u		−24689	28	−24846	3	−5.6	U				GR1	1.0	23Am.A	
C_{16} H_{11}−^{203}Tl		113735	43	113732.8	0.4	−0.0	U				R08	1.5	69De19	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
C_{15} ^{13}C $H_{10}-^{203}Tl$		109216	95	109262.6	0.4	0.3	U			R08	1.5	69De19	
C_{14} N_2 $H_7-^{203}Tl$		88540	48	88580.7	0.4	0.6	U			R08	1.5	69De19	
$^{203}Tl-^{166}Er$ ^{37}Cl		76190	48	76139.2	0.5	−0.7	U			R08	1.5	69De19	
$^{203}Tl-^{168}Er$ ^{35}Cl		71069	36	71111.9	0.5	0.8	U			R08	1.5	69De19	
$^{203}Pb-u$		−26594	30	−26611	7	−0.6	U			GS2	1.0	05Li24	
$^{203}Po-u$		−18581	30	−18584	5	−0.1	U			GS2	1.0	05Li24	
$^{203}Po-^{133}Cs_{1.526}$		125697.0	6.0	125697	5	−0.0	1	71	71 ^{203}Po	MA8	1.0	17Al34	
$^{203}At-u$		−13042	30	−13057	11	−0.5	1	14	14 ^{203}At	GS2	1.0	05Li24	
$^{203}Fr-^{133}Cs_{1.526}$		145205	17	145221	7	1.0	1	15	15 ^{203}Fr	MA8	1.0	08We02	
$^{203}At-^{208}Pb_{.976}$		9690	25	9732	11	1.7	1	21	21 ^{203}At	MA6	1.0	01Sc41	
$^{203}Rn^m-^{208}Pb_{.976}$		16579	30	16537	6	−1.4	U			SH1	1.0	13Dr04	
^{203}Tl $^{35}Cl-^{201}Hg$ ^{37}Cl		4997	3	4989.9	0.8	−0.6	U			H17	4.0	64Mc07	
		4995.23	1.49			−1.4	U			H36	2.5	85De40	
^{202}Hg $H-^{203}Tl$		6154	34	6126.0	0.8	−0.5	U			R08	1.5	69De19	
$^{203}Tl-^{167}Er$ ^{35}Cl		71436	36	71434.0	0.5	−0.0	U			R08	1.5	69De19	
^{167}Er $^{37}Cl-^{203}Tl$		−74404	33	−74384.1	0.5	0.4	U			R08	1.5	69De19	
^{169}Tm $^{35}Cl-^{203}Tl$		−69257	29	−69271.0	0.9	−0.3	U			R08	1.5	69De19	
$^{203}Tl-^{202}Hg$		1722	20	1699.1	0.8	−0.8	U			R08	1.5	69De19	
$^{203}Tl-^{201}Hg$		1999	29	2039.8	0.8	0.9	U			R08	1.5	69De19	
$^{203}Po(\alpha)^{199}Pb$		5496	5				2			Dba		68Go.B	*
$^{203}At(\alpha)^{199}Bi$		6210.3	1.	6210.1	0.8	−0.2	–					63Ho18	Z
		6208.7	3.			0.5	–					67Tr06	Z
		6209.4	2.			0.3	–			Dba		68Go.B	Z
		6211.7	3.			−0.5	–			Ora		75Ba.B	
		6210.6	5.0			−0.1	U			Ara		96Ta18	
	ave.	6210.1	0.8			−0.0	1	100	62 ^{203}At			average	
$^{203}Rn(\alpha)^{199}Po$		6628.6	5.	6629.9	2.1	0.3	–					67Va17	Z
		6630.2	2.5			−0.1	–			Lvn		93Wa04	
		6630	10			−0.0	U			Jya		95Uu01	
		6629.8	5.0			0.0	–			Ara		96Ta18	
	ave.	6629.9	2.1			0.0	1	100	99 ^{203}Rn			average	
$^{203}Rn^m(\alpha)^{199}Po^m$		6679.5	3.	6680.5	1.6	0.3	2					67Va17	Z
		6681.9	10.			−0.1	U			GSa		87He10	
		6680.9	2.5			−0.2	2			Lvn		93Wa04	
		6683.9	7.			−0.5	o			Ara		95Le04	
		6679.8	3.			0.2	2			Jya		96Le09	
		6682.9	5.0			−0.5	2			Ara		96Ta18	
$^{203}Fr(\alpha)^{199}At$		7275.6	5.	7275	4	−0.1	–					67Va20	Z
		7281.7	10.			−0.7	–					80Ew03	Z
		7263.4	25.			0.5	o			Jya		94Le05	
		7273.6	6.			0.2	–			Jya		05Uu02	
	ave.	7276	4			−0.2	1	95	85 ^{203}Fr			average	
$^{203}Fr^m(\alpha)^{199}At^m$		7391.9	5.				3			Jya		13Ja06	
$^{203}Ra(\alpha)^{199}Rn$		7729.6	20.	7736	6	0.3	o			Jya		96Le09	
		7741.8	8.			−0.7	4			Jya		05Uu02	
		7727.6	10.			0.9	4			Anv		14Ka23	
$^{203}Ra^m(\alpha)^{199}Rn^m$		7768.4	20.	7763	6	−0.3	o			Jya		96Le09	
		7765.3	8.			−0.3	3			Jya		05Uu02	
		7760.2	8.2			0.3	3			Anv		14Ka23	
$^{203}Ac(\alpha)^{199}Fr$		8382.5	16.3				6			Lza		24Wa14	
$^{203}Tl(p,t)^{201}Tl$		−6240	15	−6241	14	−0.1	1	89	89 ^{201}Tl	Yal		71Ki01	
$^{202}Hg(d,p)^{203}Hg-^{204}Hg()^{205}Hg$		325	5	327	4	0.4	1	52	48 ^{205}Hg	Pit		72Mo12	
$^{203}Tl(p,d)^{202}Tl$		−5630	20	−5628.1	1.6	0.1	U			Yal		71Ki01	
$^{203}Au(\beta^-)^{203}Hg$		2040	60	2128	3	1.5	U					94We02	
$^{203}Hg(\beta^-)^{203}Tl$		489.2	2.	491.7	1.3	1.3	–					54Th17	*
		493.2	2.			−0.7	–					55Ma40	*
		493.2	3.			−0.5	–					58Ni28	*
	ave.	491.6	1.3			0.1	1	97	97 ^{203}Hg			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{203}Pb(ε)^{203}Tl		940	50	975	6	0.7	U					55Ha.A	*
		980	20			−0.2	1	10	10 ^{203}Pb			65Le07	*
^{203}Bi(β^+)^{203}Pb		3260	50	3262	14	0.0	U					58No30	*
^{203}At(β^+)^{203}Po		5060	200	5148	12	0.4	U					87Se04	
*^{203}Po(α)^{199}Pb	E_α=5383.8(3,Z) to 4(4) level (this is level x<9.3 in Ensdf)											Ens073	**
*^{203}Hg(β^-)^{203}Tl	E_{β^-}=210(2) 214(2) 214(3) respectively, to 3/2$^+$ level at 279.1958 keV											Ens057	**
*^{203}Pb(ε)^{203}Tl	pK=0.36(0.07) 0.71(0.01) respectively, to 5/2$^+$ level at 680.5164 keV											Ens057	**
*^{203}Bi(β^+)^{203}Pb	E_{β^+}=1350(50), 740(50) to levels around 840, 1550 keV											Ens057	**
^{204}Pt−u		−19473.6	10.4				2			GS5	1.0	26Fe.C	
^{204}Au−u		−22156.7	2.8				2			GS5	1.0	26Fe.C	
		−22166	26	−22156.7	2.8	0.4	U			GR1	1.0	23Am.A	
^{204}Hg−C ^{13}C ^{35}Cl$_3$ ^{37}Cl$_2$		131776.05	1.25	131776.0	0.5	−0.0	U			H34	2.5	80Ko25	
^{204}Hg−^{169}Tm ^{35}Cl		70420	100	70422.5	1.0	0.0	U			R04	4.0	64De15	
^{204}Hg−^{167}Er ^{37}Cl		75430	60	75535.6	0.6	0.4	U			R04	4.0	64De15	
^{204}Hg−u		−26505.8	0.6	−26505.9	0.5	−0.2	1	78	78 ^{204}Hg	ST2	1.0	02Bf02	
^{204}Po−u		−19689	30	−19691	11	−0.1	R			GS2	1.0	05Li24	
		−19426	213			−1.2	U			RI1	1.0	17Sc02	*
^{204}At−u		−12748	30	−12735	12	0.4	–			GS2	1.0	05Li24	
		−13099	208			1.8	U			RI1	1.0	17Sc02	*
	ave.	−12752	27			0.7	1	19	19 ^{204}At			average	
^{204}Rn−^{133}Cs$_{1.534}$		136394	60	136484	8	1.5	U			SH1	1.0	13Dr04	
^{204}Rn−u		−8660	86	−8553	8	1.2	U			RI1	1.0	17Sc02	
^{204}Fr−u		600	110	655.4	1.2	0.5	U			RI1	1.0	17Sc02	*
		654.5	1.3			0.7	1	85	85 ^{204}Fr	SH2	1.0	22Mo.A	
^{204}Fr^m−u		704.0	1.4	704.6	1.4	0.5	1	94	94 ^{204}Fr^m	SH2	1.0	22Mo.A	
^{204}Fr^n−u		1010.1	1.3	1000.2	1.7	−7.6	B			SH2	1.0	22Mo.A	
^{204}Hg ^{35}Cl$_2$−^{200}Hg ^{37}Cl$_2$		11066.85	0.55	11067.5	0.7	0.5	1	23	12 ^{200}Hg	H33	2.5	80Ko25	
^{204}Pb−^{208}Pb$_{.981}$		−4047	21	−4052.14	0.18	−0.2	U			MA6	1.0	01Sc41	
^{204}Rn−^{208}Pb$_{.981}$		14358.1	9.4	14353	8	−0.5	1	78	78 ^{204}Rn	SH1	1.0	13Dr04	
		14303	25			2.0	o			SH1	1.0	13Dr04	
^{204}Hg ^{35}Cl−^{202}Hg ^{37}Cl		5807	2	5800.7	0.7	−0.8	U			H17	4.0	64Mc07	
		5800.67	0.53			0.1	1	32	21 ^{202}Hg	H33	2.5	80Ko25	
^{204}Hg−^{203}Tl		1161	25	1151.5	0.7	−0.3	U			R08	1.5	69De19	
^{204}Pb(α)^{200}Hg		2650	100	1967.2	0.5	−6.8	B					58Ri23	
^{204}Pb(α,^{8}He)^{200}Pb		−28043	13	−28044	10	−0.1	–			INS		90Ka10	
	ave.	−28044	11			−0.0	1	84	84 ^{200}Pb			average	
^{204}Po(α)^{200}Pb		5484.6	1.5	5484.8	1.3	0.2	–			Dba		69Go23	*
		5486.3	3.			−0.5	–					70Ra14	Z
		5486.5	5.1			−0.3	–					67Ti04	
		5482.0	5.			0.6	–					70DaZM	
		5487.8	10.2			−0.3	U					70Jo26	
	ave.	5484.8	1.3			0.0	1	100	84 ^{204}Po			average	
^{204}At(α)^{200}Bi		6069.9	3.	6070.2	1.2	0.1	2					63Ho18	Z
		6066.2	3.			1.3	2					67Tr06	Z
		6071.3	2.			−0.5	2					68Go12	
		6067.2	5.1			0.6	2					74Ho27	
		6071.3	3.			−0.4	2			Ora		75Ba.B	
		6072.0	3.			−0.6	2			Dba		81Va27	Z
^{204}Rn(α)^{200}Po		6544.3	3.	6546.2	1.8	0.6	–					67Va17	Z
		6547.5	2.5			−0.5	–			Lvn		93Wa04	
		6537.4	7.			1.2	o			Ara		95Le04	
		6548.6	5.0			−0.5	–			Ara		96Ta18	
		6536.4	8.2			1.2	–					15Ma63	
		6557.8	16.3			−0.7	U			Lza		22Zh45	
	ave.	6546.0	1.8			0.1	1	99	78 ^{200}Po			average	
^{204}Fr(α)^{200}At		7170.4	5.	7167.2	1.5	−0.6	–					67Va20	Z
		7169.4	5.			−0.4	–					74Ho27	Z

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{204}Fr(α)^{200}At		7170.6	5.	7167.2	1.5	−0.7	–				Lvn		92Hu04	*
		7179.0	6.			−1.9	o				Jya		94Le05	
		7167.8	7.			−0.1	–				Ara		95Le04	
		7173.9	6.			−1.1	o				Jya		05Uu02	
		7171.9	5.			−0.9	–				Jya		13Ja06	
		7172.9	14.3			−0.4	–				Lza		22Ya27	
	ave.	7170.4	2.4			−1.3	1	43	28	^{200}At			average	
^{204}Fr^{m}(α)^{200}At		7218.8	8.	7213.1	1.9	−0.7	o				Lvn		92Hu04	
^{204}Fr^{m}(α)^{200}At^{m}		7108.2	5.	7100.1	1.6	−1.6	U						74Ho27	Z
		7105.5	3.			−1.8	U				Bka		82Bo04	Z
		7108.4	5.			−1.7	U				Lvn		92Hu04	*
		7115.6	7.			−2.2	o				Jya		94Le05	*
		7114.7	7.			−2.1	U				Ara		95Le04	
		7117.7	6.			−2.9	o				Jya		05Uu02	*
		7108.7	5.			−1.7	U				Jya		13Ja06	
		7108.7	14.3			−0.6	U				Lza		22Ya27	
^{204}Fr^{n}(α)^{200}At^{n}		7153.5	5.	7144.5	1.9	−1.8	U				Lvn		92Hu04	
		7157.6	6.1			−2.1	o				Jya		05Uu02	
		7153.5	5.			−1.8	1	13	9	^{204}Fr^{n}	Jya		13Ja06	
		7156.5	14.3			−0.8	U				Lza		22Ya27	
^{204}Ra(α)^{200}Rn		7638.1	12.	7637	7	−0.1	3				Ara		95Le04	
		7638.1	25.			−0.1	o				Jya		95Le15	
		7634.0	10.			0.3	o				Jya		96Le09	
		7636.1	8.			0.1	3				Jya		05Uu02	
		7638.1	25.			−0.1	U				Anv		10He25	
^{204}Ac(α)^{200}Fr		8107.3	15.3				6				Lza		22Hu12	
		8116.4	17.3	8107	15	−0.5	o				Lza		24Wa14	
^{204}Hg(d,α)^{202}Au		10379.0	5.0				2				Mun		03WiZV	
^{204}Pb(p,t)^{202}Pb		−6835	10	−6831	4	0.4	1	14	14	^{202}Pb	Yal		71Ki01	
^{204}Hg(t,α)^{203}Au		10962	15	10978	3	1.1	U				LAl		81Fl05	
^{204}Hg(d,^{3}He)^{203}Au−^{206}Pb()^{205}Tl		−1582.0	3.0	−1582.0	3.0	0.0	1	100	100	^{203}Au			94Gr07	
^{204}Hg(d,t)^{203}Hg		−1242	5	−1233.1	1.4	1.8	U				Ald		70An14	
^{203}Tl(n,γ)^{204}Tl		6656.0	0.3	6656.04	0.29	0.1	1	94	93	^{203}Tl	MMn		74Co21	Z
		6654.88	0.14			8.3	C				Bdn		06Fi.A	
^{204}Pb(p,d)^{203}Pb		−6165	10	−6170	6	−0.5	–				Yal		71Ki01	
^{204}Pb(d,t)^{203}Pb		−2160	20	−2138	6	1.1	–				Ald		67Bj01	
^{204}Pb(p,d)^{203}Pb	ave.	−6171	9	−6170	6	0.0	1	52	52	^{203}Pb			average	
^{204}Pt^{m}(IT)^{204}Pt		1995.1	0.7				3						11St21	
^{204}Pt^{n}(IT)^{204}Pt^{m}		44	20				4						11St21	*
^{204}Pt^{p}(IT)^{204}Pt^{n}		1157.5	0.5				5						11St21	
^{204}Au^{n}(IT)^{204}Au^{m}		1815.6	0.7				4						11St21	*
^{204}Au(β^-)^{204}Hg		4500	300	4051.3	2.7	−1.5	F						67Wa23	*
^{204}Tl(ε)^{204}Hg		314	20	342.6	0.5	1.4	U						64Ch17	
		332	20			0.5	U						66Kl02	
		385	20			−2.1	U						73La17	
^{204}Tl(β^-)^{204}Pb		764.24	0.31	763.73	0.18	−1.7	–						67Pa08	
		763.47	0.22			1.2	–						68Wo02	
	ave.	763.73	0.18			−0.0	1	97	96	^{204}Tl			average	
^{204}At(β^+)^{204}Po		6220	160	6480	15	1.6	U						86Ve.B	*
^{204}Fr^{n}(IT)^{204}Fr^{m}		275	1	275.3	1.0	0.3	1	97	91	^{204}Fr^{n}			12Ja01	
*^{204}Po−u	A systematic error 75 keV is added to all data in the same paper.												HWJ182	**
*	freq ratio rounded, not used												HWJ182	**
*^{204}At−u	$M-A$=–11908(94) keV for mixture gs+m at 587.3(0.2) keV												Nub269	**
*^{204}Fr−u	$M-A$=679(79) keV for mixture at gs+m+n 45.9(1.7) and												Nub269	**
*^{204}Fr−u	321.2(1.9) keV												Nub269	**
*^{204}Po(α)^{200}Pb	Printing error in reference: ^{204}Po not ^{206}Po; Z recalibrated												AHW	**
*^{204}Fr(α)^{200}At	E_α=7031(5), 6916(8) to ground state, 113 level												92Hu04	**
*^{204}Fr^{m}(α)^{200}At^{m}	E_α=6969(5); and 7013(5) from ^{204}Fr^{n} 276.1 above ^{204}Fr^{m} to ^{200}At^{n}												95Bi.A	**
*	230.9 above ^{200}At^{m}												92Hu04	**
*^{204}Fr^{m}(α)^{200}At^{m}	E_α=7020(7) from ^{204}Fr^{n} 276.1 above Fr^{m} to ^{200}At^{n} 230.9 above ^{200}At^{m}												95Bi.A	**
*^{204}Fr^{m}(α)^{200}At^{m}	E_α=6976(6); and 7017(6) from ^{204}Fr^{n} 276.1 above ^{204}Fr^{m} to ^{200}At^{n}												GAu	**
*^{204}Pt^{n}(IT)^{204}Pt^{m}	no K X−rays observed, so 10 keV < dE < 78.4 keV above ^{204}Pt^{m}												FGK268	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{204}Au^{n}(IT)^{204}Au^{m}$		839.0(0.5) and 976.6(0.5) γs in a cascade to $^{204}Au^{m}$										FGK268	**
$*^{204}Au(\beta^{-})^{204}Hg$		F : reported 4 s activity does not exist										Ens87a	**
$*^{204}At(\beta^{+})^{204}Po$		E_{β^+}=2950(160) to 8^+ level at 2248.17 keV										Ens102	**
$^{205}Au-u$		−20472	15				2			GS5	1.0	26Fe.C	
		−20469	33	−20472	15	−0.1	U			GR1	1.0	23Am.A	
$C_{16}\ H_{13}-^{205}Tl$		127345	29	127299.7	0.6	−1.0	U			R08	1.5	69De19	
$C_{14}\ N_2\ H_9-^{205}Tl$		102091	36	102147.6	0.6	1.0	U			R08	1.5	69De19	
$^{205}Tl-^{168}Er\ ^{37}Cl$		76198	44	76145.2	0.6	−0.8	U			R08	1.5	69De19	
$^{205}Tl-^{170}Er\ ^{35}Cl$		70034	23	70100.4	1.3	1.9	U			R08	1.5	69De19	
$^{205}Tl-^{133}Cs_{1.541}$		120129	11	120124.3	0.6	−0.4	U			MA8	1.0	08We02	
$^{205}Bi-u$		−22559	30	−22616	5	−1.9	U			GS2	1.0	05Li24	
		−22358	616			−0.4	U			RI1	1.0	17Sc02	
$^{205}Po-u$		−18773	30	−18810	11	−1.2	–			GS2	1.0	05Li24	
		−18811	538			0.0	U			RI1	1.0	17Sc02	*
	ave.	−18790	25			−0.8	1	19	19 ^{205}Po			average	
$^{205}At-u$		−13841	111	−13940	13	−0.9	U			RI1	1.0	17Sc02	
$^{205}Rn-^{133}Cs_{1.541}$		137456	50	137421	5	−0.7	U			SH1	1.0	13Dr04	
		137458	29			−1.3	U			SH1	1.0	13Dr04	
$^{205}Rn-u$		−8494	222	−8277	5	1.0	U			RI1	1.0	17Sc02	*
$^{205}Fr-^{133}Cs_{1.541}$		144293.8	9.7	144295.1	1.3	0.1	U			MA8	1.0	08We02	
		144299	22			−0.2	U			1.0	1.0	197Bi-C	
		144273	26			0.9	U			1.0	1.0	197Bi-x	
$^{205}Fr-u$		−1403.9	1.5	−1403.4	1.3	0.3	1	78	78 ^{205}Fr	SH2	1.0	22Mo.A	
$^{205}Rn-^{208}Pb_{.986}$		14748	11	14746	5	−0.2	–			SH1	1.0	13Dr04	
		14772	20			−1.3	–			SH1	1.0	13Dr04	
	ave.	14754	10			−0.8	1	31	31 ^{205}Rn			average	
$^{205}Tl\ ^{35}Cl-^{203}Tl\ ^{37}Cl$		5040	4	5033.3	0.6	−0.4	U			H17	4.0	64Mc07	
		5031.43	1.07			0.7	–			H36	2.5	85De40	
		5032.88	1.01			0.3	–			H42	1.5	93Si05	
	ave.	5032.5	1.3			0.6	1	19	14 ^{205}Tl			average	
$^{205}Tl-^{167}Er\ ^{37}Cl$		76426	47	76467.3	0.6	0.6	U			R08	1.5	69De19	
$^{205}Tl-^{169}Tm\ ^{35}Cl$		71355	25	71354.2	1.0	−0.0	U			R08	1.5	69De19	
$^{169}Tm\ ^{37}Cl-^{205}Tl$		−74316	32	−74304.3	1.0	0.2	U			R08	1.5	69De19	
$^{205}Tl-^{204}Hg$		938	27	931.7	0.8	−0.2	U			R08	1.5	69De19	
$^{205}Tl-^{203}Tl$		2092	20	2083.2	0.6	−0.3	U			R08	1.5	69De19	
$^{205}Po(\alpha)^{201}Pb$		5324.1	10.	5326	7	0.2	–					67Ti04	
		5328.2	10.2			−0.2	–					70Jo26	
	ave.	5326	7			0.0	1	97	92 ^{201}Pb			average	
$^{205}At(\alpha)^{201}Bi$		6016.3	4.	6019.6	1.7	0.8	3					63Ho18	Z
		6020.5	2.			−0.5	3			Dba		68Go.B	Z
		6018.9	5.			0.1	3					74Ho27	Z
$^{205}Rn(\alpha)^{201}Po$		6386.6	3.	6386.4	1.8	−0.0	–					67Va17	Z
		6386.6	6.			−0.0	–					71Ho01	Z
		6385.7	2.5			0.3	–			Lvn		93Wa04	
	ave.	6386.1	1.9			0.2	1	97	69 ^{205}Rn			average	
$^{205}Fr(\alpha)^{201}At$		7056.5	5.	7053.4	1.4	−0.6	–					67Va20	Z
		7052.2	5.			0.2	–					74Ho27	Z
		7057.3	5.			−0.8	–			ORa		81Ri04	Z
		7052.9	7.			0.1	–			Ara		95Le04	
		7053.9	5.			−0.1	–			Anv		05De01	
	ave.	7054.7	2.4			−0.5	1	36	22 ^{205}Fr			average	
$^{205}Ra(\alpha)^{201}Rn$		7506.7	20.	7486	20	−0.4	F			GSa		87He10	*
		7496.6	25.			−0.4	o			Jya		95Le15	
		7486.4	20.				3			Jya		96Le09	
$^{205}Ra^{m}(\alpha)^{201}Rn^{m}$		7501.7	10.	7505	9	0.3	4			Ara		95Le04	
		7522.1	25.			−0.7	o			Jya		95Le15	
		7517.0	20.			−0.6	4			Jya		96Le09	
		7526.1	30.			−0.7	U			Anv		10He25	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{205}Ac(α)^{201}Fr		8093.2	30.6	8040	15	−0.9	o			Lza		14Zh03	
		8040.2	15.3				3			Lza		22Hu12	
^{204}Hg(d,p)^{205}Hg		3443	5	3445	4	0.4	1	52	52 ^{205}Hg	Ald		70An14	
^{205}Tl(γ,n)^{204}Tl		−7515	29	−7546.1	0.5	−1.1	U			Phi		60Ge01	
		−7548	3			0.6	U			McM		79Ba06	
^{205}Tl(d,t)^{204}Tl		−1288.7	0.6	−1288.9	0.5	−0.3	1	64	61 ^{205}Tl	Mun		90Li40	
^{204}Pb(n,γ)^{205}Pb		6731.53	0.15	6731.66	0.11	0.8	–			ILn		83Hu13	Z
		6731.80	0.16			−0.9	–			Bdn		06Fi.A	
^{204}Pb(d,p)^{205}Pb		4516	20	4507.09	0.11	−0.4	U			Ald		67Bj01	
^{204}Pb(n,γ)^{205}Pb	ave.	6731.66	0.11	6731.66	0.11	0.0	1	99	99 ^{204}Pb			average	
^{205}Au^{m}(IT)^{205}Au		922	10				3					26Fe.C	*
^{205}Hg(β^-)^{205}Tl		1620	200	1534	4	−0.4	U					40Kr08	
		1750	200			−1.1	U					51Ly10	
^{205}Pb(ε)^{205}Tl		41.4	1.1	50.7	0.5	8.5	B					78Pe08	
^{205}Bi(β^+)^{205}Pb		2701.4	10.	2705	5	0.4	–					62Bo25	*
		2715.4	10.			−1.0	–					62Pe08	*
	ave.	2708	7			−0.4	1	46	46 ^{205}Bi			average	
^{205}Po(β^+)^{205}Bi		3390	150	3545	11	1.0	U					69Ho37	*
*^{205}Po−u	$M-A$=–16791(268) keV for mixture at gs+p 1461.2(0.2) keV											Nub269	**
*^{205}Rn−u	$M-A$=–7584(82) keV for mixture at gs+m 657.1(0.5) keV											Nub269	**
*^{205}Ra(α)^{201}Rn	F : possibly mixed with ^{205}Ra^{m}(α)^{201}Rn^{m}											87He10	**
*^{205}Au^{m}(IT)^{205}Au	from ME(^{205}Au^{m})=–18148(14) keV in 26Fe.C and covariance analysis											FGK26a	**
*^{205}Bi(β^+)^{205}Pb	E_{β^+}=976(10) 990(10) respectively, to 7/2$^-$ level at 703.427 keV											Ens044	**
*^{205}Po(β^+)^{205}Bi	p$^+$ =3(1)×10^{-3} to 7/2$^-$ level at 849.84 and 7/2$^-$ at 1001.22 keV											Ens044	**
^{206}Au−u		−15516.0	8.6				2			GS5	1.0	26Fe.B	
^{206}Hg−^{133}Cs$_{1.549}$		123983.4	8.8	123981	8	−0.2	1	86	86 ^{206}Hg	MA8	1.0	21Da16	
^{206}Bi−u		−21429	30	−21503	8	−2.5	U			GS2	1.0	05Li24	
^{206}Po−u		−19471	30	−19527	4	−1.9	U			GS2	1.0	05Li24	
^{206}At−u		−13305	30	−13355	15	−1.7	1	23	23 ^{206}At	GS2	1.0	05Li24	
		−13416	2309			0.0	U			RI1	1.0	17Sc02	
^{206}Rn−u		−9195	649	−9805	9	−0.9	U			RI1	1.0	17Sc02	
^{206}Rn−^{133}Cs$_{1.549}$		136641	15	136650	9	0.6	1	38	38 ^{206}Rn	SH1	1.0	13Dr04	
^{206}Fr−u		−1450	180	−1423	4	0.2	U			RI1	1.0	17Sc02	*
		−1420.6	1.2			−1.7	F			SH2	1.0	22Mo.A	*
^{206}Fr^{n}−u		−872.7	1.3	−853	4	15.5	F			SH2	1.0	22Mo.A	*
^{206}Ra−u		3808.9	53.7	3824	18	0.3	1	11	11 ^{206}Ra	RI1	1.0	21Ni08	
		3714.5	98.8			1.1	U			RI1	1.0	21Ni08	
^{206}Pb ^{35}Cl$_2$−^{202}Hg ^{37}Cl$_2$		9722.09	0.57	9720.5	0.7	−1.1	1	24	23 ^{202}Hg	H36	2.5	85De40	
^{206}Pb ^{35}Cl−^{204}Hg ^{37}Cl		3929	4	3919.8	0.5	−0.6	U			H17	4.0	64Mc07	
^{206}Pb ^{35}Cl−^{204}Pb ^{37}Cl		4378	3	4371.84	0.15	−0.5	U			H17	4.0	64Mc07	
		4370.72	1.17			0.4	U			H36	2.5	85De40	
		4371.29	0.81			0.5	U			H42	1.5	93Si05	
^{206}Pb−^{208}Pb$_{.990}$		−2420.30	0.58	−2420.28	0.12	0.0	U			TG4	1.0	23Ch12	*
^{206}Rn−^{208}Pb$_{.990}$		13307	15	13311	9	0.2	1	38	38 ^{206}Rn	SH1	1.0	13Dr04	
^{206}Pb−^{207}Pb$_{.995}$		−1552.28	0.56	−1552.13	0.10	0.3	U			TG4	1.0	23Ch12	*
^{206}Po(α)^{202}Pb		5327.4	4.	5327.0	1.3	−0.1	2					67Ti04	Z
		5327.4	1.5			−0.3	2			Dba		69Go23	*
		5325.1	3.			0.6	2					70Ra14	Z
^{206}At(α)^{202}Bi		5884.4	3.6	5887	5	0.6	o			Dba		68Go.B	*
		5886.4	5.			0.0	1	98	75 ^{202}Bi	Dba		81Va27	*
^{206}Rn(α)^{202}Po		6381.8	3.	6384.1	1.6	0.7	–					67Va17	Z
		6384.6	3.			−0.2	–			Dba		71Go35	Z
		6384.8	2.5			−0.3	–			Lvn		93Wa04	
		6387.3	5.1			−0.6	–			Tex		15Ma63	
	ave.	6384.2	1.6			−0.1	1	99	74 ^{202}Po			average	
^{206}Fr(α)^{202}At		6925.9	7.	6923	3	−0.4	–					67Va20	*
		6918.9	7.			0.6	–					74Ho27	*
		6924.0	7.			−0.1	–			ORa		81Ri04	*
		6924.8	7.			−0.2	–			Lvn		92Hu04	*
	ave.	6923	4			−0.0	1	100	100 ^{206}Fr			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$^{206}Fr^{m}(\alpha)^{202}At^{m}$		6925.9	7.	6923	4	−0.4	4					67Va20	*
$^{206}Fr^{m}(\alpha)^{202}At^{m}$		6918.9	7.			0.6	4					74Ho27	*
		6924.0	7.			−0.1	4			ORa		81Ri04	*
		6924.8	7.			−0.2	4			Lvn		92Hu04	*
$^{206}Ra(\alpha)^{202}Rn$		7416.3	5.	7415	4	−0.2	–					67Va22	Z
		7414.3	10.			0.1	–			GSa		87He10	
		7412.2	10.			0.3	o			Jya		95Le15	
		7406	15			0.6	o			Jya		95Uu01	
		7407.1	20.4			0.4	o			Jya		95Le41	
		7412.2	10.			0.3	–			Jya		96Le09	
		7438.7	23.5			−1.0	U					21Ni08	
	ave.	7415	4			−0.0	1	99	89 ^{206}Ra			average	
$^{206}Ac(\alpha)^{202}Fr$		7944.6	30.	7942	14	−0.1	3			Jya		98Es02	
		7972.0	30.			−1.0	o			Lza		14Zh03	
		7941.5	15.3			0.0	3			Lza		25Zh34	
$^{206}Ac^{m}(\alpha)^{202}Fr^{m}$		7903.8	30.				2			Jya		98Es02	
$^{204}Pb(t,p)^{206}Pb$		6322	20	6336.52	0.12	0.7	U			Ald		67Ha.A	
$^{204}Pb(\alpha,d)^{206}Bi$		−15798.	11.5	−15792	8	0.5	R			Pit		76Da20	
$^{205}Tl(n,\gamma)^{206}Tl$		6502.91	0.14	6502.92	0.12	0.0	–			ILn		25Ci01	*
		6502.87	0.27			0.2	–			Bdn		06Fi.A	
$^{205}Tl(d,p)^{206}Tl$		4276	5	4278.35	0.12	0.5	U			ANL		65Er02	
$^{205}Tl(n,\gamma)^{206}Tl$	ave.	6502.90	0.12	6502.92	0.12	0.1	1	99	90 ^{206}Tl			average	
$^{205}Tl(^{3}He,d)^{206}Pb$		1761.7	1.4	1760.1	0.5	−1.1	1	13	13 ^{205}Tl	Mun		90Li40	
$^{205}Pb(n,\gamma)^{206}Pb$		8086.66	0.06	8086.66	0.06	−0.0	1	100	100 ^{205}Pb			96Ra16	Z
$^{206}Pb(\gamma,n)^{205}Pb$		−8090	70	−8086.66	0.06	0.0	U			Phi		60Ge01	
		−8087	3			0.1	U			McM		79Ba06	
$^{206}Pb(d,t)^{205}Pb$		−1830	100	−1829.43	0.06	0.0	U			MIT		53Ha66	
		−1831.2	0.5			3.5	B			Mun		90Li40	
$^{206}Hg(\beta^{-})^{206}Tl$		1240	62	1320	8	1.3	U					68Wo09	*
$^{206}Tl(\beta^{-})^{206}Pb$		1534	5	1533.0	0.5	−0.2	U					71Pe23	
		1527	4			1.5	U					72Wi18	
$^{206}Bi(\beta^{+})^{206}Pb$		3683	33	3757	8	2.3	U					62Pe08	*
$^{206}Bi(\varepsilon)^{206}Pb$		3753	10			0.4	2					74Go20	*
$^{206}At(\beta^{+})^{206}Po$		5687	150	5749	14	0.4	U					77Li16	*
$^{206}Fr^{n}(IT)^{206}Fr$		531	1				2			ORa		81Ri04	
$^{206}Fr^{x}(IT)^{206}Fr$		100	100				2					AHW	*
$*^{206}Fr-u$	$M-A$=–1104(131) keV for mixture gs+m+n at 215(6) 531(1) keV											Nub269	**
$*^{206}Fr-u$	at variance with ME extracted using ME of ^{202}At and $^{206}Fr(\alpha)^{202}At$											HWJ25a	**
$*^{206}Fr^{n}-u$	at variance with ME extracted using ME of $^{202}At^{n}$ and $^{206}Fr^{n}(\alpha)^{202}At^{n}$											HWJ25a	**
$*^{206}Pb-^{208}Pb_{.990}$	constructed from PI−ICR frequency ratio											HWJ25a	**
$*^{206}Pb-^{207}Pb_{.995}$	constructed from PI−ICR frequency ratio											HWJ25a	**
$*^{206}Po(\alpha)^{202}Pb$	Printing error in reference: ^{206}Po not ^{211}Po; Z recalibrated											AHW	**
$*^{206}At(\alpha)^{202}Bi$	E_α=5702.8(2,Z) to $(5)^{+}$ level at 68(3) keV											Ens044	**
$*^{206}At(\alpha)^{202}Bi$	E_α=5773.8(5,Z), 5702.8(5,Z) to ground state and $(5)^{+}$ level at 68(3) keV											Ens044	**
$*^{206}Fr(\alpha)^{202}At$	E_α=6793.1(5,Z); correction –2 for being a doublet											AHW	**
$*^{206}Fr(\alpha)^{202}At$	E_α=6786.3(5,Z); correction –2 for being a doublet											AHW	**
$*^{206}Fr(\alpha)^{202}At$	E_α=6791.3(5,Z); correction –2 for being a doublet											AHW	**
$*^{206}Fr(\alpha)^{202}At$	E_α=6792(5); correction –2 for being a doublet											AHW	**
$*^{206}Fr^{m}(\alpha)^{202}At^{m}$	E_α=6793.1(5,Z); correction –2 for being a doublet											FGK25c	**
$*^{206}Fr^{m}(\alpha)^{202}At^{m}$	E_α=6786.3(5,Z); correction –2 for being a doublet											FGK25c	**
$*^{206}Fr^{m}(\alpha)^{202}At^{m}$	E_α=6791.3(5,Z); correction –2 for being a doublet											FGK25c	**
$*^{206}Fr^{m}(\alpha)^{202}At^{m}$	E_α=6792(5); correction –2 for being a doublet											FGK25c	**
$*^{205}Tl(n,\gamma)^{206}Tl$	recalibrated using the latest experimental data											FGK25a	**
$*^{206}Hg(\beta^{-})^{206}Tl$	E_{β^-}=935(62) to 1^{-} level at 304.896 keV											Ens085	**
$*^{206}Bi(\beta^{+})^{206}Pb$	E_{β^+}=977(33) to 4^{+} level at 1683.99 keV											Ens085	**
$*^{206}Bi(\varepsilon)^{206}Pb$	LK=0.509(0.015) to 5^{-} level at 3562.92 keV, original error 22, recalculated											Ens085	**
$*^{206}At(\beta^{+})^{206}Po$	E_{β^+}=3092(150) to 6^{+} level at 1573.38 keV											Ens085	**
$*^{206}Fr^{x}(IT)^{206}Fr$	Assuming a 0.15(0.20)% isomeric mixture											AHW	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{207}Hg−u		−17721	33	−17656	7	2.0	o			GS3	1.0	08ChZI	
		−17700	32			1.4	U			GS3	1.0	12Ch19	
^{207}Hg−^{133}Cs$_{1.556}$		129461.0	6.5				2			MA8	1.0	21Da16	
^{207}Tl−^{133}Cs$_{1.556}$		124538.55	0.72				2			SH2	1.0	22Ka.A	*
^{207}Pb−^{133}Cs$_{1.556}$		123017.38	0.69	123012.07	0.06	−7.7	B			SH2	1.0	22Ka.A	*
^{207}Rn−^{133}Cs$_{1.556}$		137794	28	137847	5	1.9	U			SH1	1.0	13Dr04	
^{207}Fr−^{133}Cs$_{1.556}$		144062	20	144058	19	−0.2	1	88	88 ^{207}Fr	MA8	1.0	14Bo26	
^{207}Ra−u		3798.2	16.1	3796	16	−0.1	1	94	94 ^{207}Ra	RI1	1.0	21Ni08	
^{207}Ra^{m}−u		4390.8	48.3	4371	10	−0.4	U			RI1	1.0	21Ni08	
^{207}Pb ^{35}Cl−^{205}Tl ^{37}Cl		4413	4	4419.7	0.6	0.4	U			H17	4.0	64Mc07	
		4415.60	2.40			0.7	U			H36	2.5	85De40	
		4417.32	1.40			1.1	U			H42	1.5	93Si05	
^{206}Pb H−^{207}Pb		6394.2	1.1	6393.43	0.10	−0.3	U			C4	2.5	71Ke02	
^{207}Pb−^{208}Pb$_{0.995}$		−871.73	0.58	−871.93	0.06	−0.3	U			TG4	1.0	23Ch12	*
^{207}Pb−^{206}Pb$_{1.005}$		1559.47	0.56	1559.29	0.10	−0.3	U			TG4	1.0	23Ch12	*
^{206}Fr^{x}−^{207}Fr$_{.498}$ ^{205}Fr$_{.502}$		930	90	850	100	−0.4	U			P24	2.5	82Au01	
^{207}Po(α)^{203}Pb		5216.0	2.5	5215.9	2.5	−0.0	1	96	59 ^{207}Po	Dba		70Af.A	
^{207}At(α)^{203}Bi		5872.5	3.				3			Dba		69Go23	Z
^{207}Rn(α)^{203}Po		6256.3	3.	6251.2	1.6	−1.6	–					67Va20	Z
		6247.3	3.			1.3	–			Dba		71Go35	Z
		6250.4	2.5			0.3	–			Lvn		93Wa04	
	ave.	6251.2	1.6			−0.0	1	97	68 ^{207}Rn			average	
^{207}Fr(α)^{203}At		6907.8	5.	6889	20	−0.4	–					67Va20	Z
		6895.8	5.			−0.1	–					74Ho27	Z
		6900.9	5.			−0.2	–			ORa		81Ri04	Z
		6895.5	8.2			−0.1	–			Lza		26Zh40	
	ave.	6900	25			−0.2	1	13	9 ^{207}Fr			average	
^{207}Ra(α)^{203}Rn		7273.8	5.	7298	16	0.5	–					67Va22	Z
		7268.7	10.			0.6	–			GSa		87He10	
		7276.9	12.2			0.4	–			Jya		95Uu01	
	ave.	7273	29			0.4	1	7	6 ^{207}Ra			average	
^{207}Ra^{m}(α)^{203}Rn^{m}		7464.5	10.2	7470	8	0.6	3			GSa		87He10	
		7474.7	15.			−0.3	o			Jya		95Le15	
		7475.7	15.			−0.3	3			Jya		96Le09	
		7499.2	29.			−1.0	3					21Ni08	
^{207}Ac(α)^{203}Fr		7864.3	25.	7850	60	−0.3	o			Jya		94Le05	
		7844.9	25.			0.1	2			Jya		98Es02	
		7851.9	14.3			−0.1	2			Lza		22Ya27	
^{207}Th(α)^{203}Ra		8328.2	21.4				5			Lza		22Ya15	
^{205}Tl(t,p)^{207}Tl		4880	15	4870.0	0.8	−0.7	U			Ald		69Ha11	
^{207}Pb(t,α)^{206}Tl		12321	25	12325.4	0.5	0.2	U			Ald		67Ha.A	
^{206}Pb(n,γ)^{207}Pb		6737.85	0.15	6737.78	0.10	−0.4	–			MMn		81Ke11	Z
		6737.72	0.18			0.4	–			ILn		83Hu13	Z
		6737.74	0.17			0.3	–			Bdn		06Fi.A	
^{207}Pb(γ,n)^{206}Pb		−6742	3	−6737.78	0.10	1.4	U			McM		79Ba06	
^{206}Pb(d,p)^{207}Pb		4480	30	4513.22	0.10	1.1	U			MIT		53Ha66	
		4510	20			0.2	U					58Mc64	
		4526	30			−0.4	U			Pit		64Co11	
^{206}Pb(n,γ)^{207}Pb	ave.	6737.78	0.10	6737.78	0.10	0.1	1	99	99 ^{206}Pb			average	
^{207}Hg(β^{-})^{207}Tl		4815	150	4585	6	−1.5	U					81Jo.B	*
^{207}Tl^{m}(IT)^{207}Tl		1348.18	0.16				3					Ens118	
^{207}Tl(β^{-})^{207}Pb		1431	8	1421.9	0.7	−1.1	U					67Da10	
^{207}Bi(ε)^{207}Pb		2392	10	2398.0	2.1	0.6	U					Averag	*
^{207}Po(β^{+})^{207}Bi		2907	10	2909	7	0.2	1	44	41 ^{207}Po			58Ar56	*
^{207}Rn(β^{+})^{207}At		4617	70	4593	13	−0.3	U					75Ze.A	*
*^{207}Tl−^{133}Cs$_{1.556}$	From frequency ratio ^{207}Tl^{++}/(^{133}Cs)$^{+}$ =0.7786632374(27)											HWJ262	**
*^{207}Pb−^{133}Cs$_{1.556}$	From frequency ratio ^{207}Pb^{++}/(^{133}Cs)$^{+}$ =0.7786575146(26)											HWJ262	**
*^{207}Pb−^{208}Pb$_{0.995}$	constructed from PI−ICR frequency ratio											HWJ25a	**
*^{207}Pb−^{206}Pb$_{1.005}$	constructed from PI−ICR frequency ratio											HWJ25a	**
*^{207}Hg(β^{-})^{207}Tl	$E_{\beta-}$=1800(150), 14%, 32%, 16%, 7% to (7/2^{-},9/2) level at 2911.83 keV											81Jo.B	**
*	(9/2)$^{+}$ at 2985.23 and 3104.43, (7/2^{-},9/2,11/2) at 3143.1 keV											Ens112	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{207}$Bi(ε)^{207}Pb	Average pL=0.61(0.05) to 7/2$^-$ level at 2339.921 keV from two references:										Ens112	**
*	pL=0.663(0.014)										64De16	**
*	pL=0.56(0.04); original error 0.08 is 2σ										82Ta18	**
$*^{207}$Po(β^+)^{207}Bi	E_{β^+}=893(10) to 7/2$^-$ level at 992.43 keV, and other E_{β^+}										Ens112	**
$*^{207}$Rn(β^+)^{207}At	E_{β^+}=3250(70) to 7/2$^-$ level at 344.55 keV										Ens112	**
^{208}Hg−u	−14241	33	−14251	18	−0.3	o			GS3	1.0	08ChZI	
	−14241	33			−0.3	2			GS3	1.0	09Ch08	
	−14255	21			0.2	2			MR1	1.0	21Da16	
^{208}Tl−^{220}Rn$_{.945}$	−28749.24	0.52				2			CP2	1.0	23Ya.A	
^{208}Pb−^{133}Cs$_{1.564}$	124532.0	5.6	124523.635	0.016	−1.5	U			MA8	1.0	08We02	
	124524.3	5.5			−0.1	U			MA8	1.0	14Bo26	
^{208}Pb−^{132}Xe$_{1.576}$	127702.083	0.011	127702.082	0.011	−0.1	1	100	100 ^{208}Pb	Hep	1.0	22Kr04	
^{208}Po−u	−18710	31	−18755.8	1.4	−1.5	U			GS2	1.0	05Li24	
^{208}Po−^{133}Cs$_{1.564}$	129125.3	6.0	129117.3	1.4	−1.3	U			MA8	1.0	17Al34	
^{208}Fr−^{133}Cs$_{1.564}$	144984	20	145009	12	1.3	o			MA8	1.0	12Bo.A	
	145030	16			−1.3	o			MA8	1.0	12Bo.A	
	145012	13			−0.2	1	78	78 ^{208}Fr	MA8	1.0	14Bo26	
^{208}Pb ^{35}Cl−^{206}Pb ^{37}Cl	5136	2	5136.92	0.14	0.1	U			H17	4.0	64Mc07	
	5136.23	1.08			0.3	U			H36	2.5	85De40	
	5136.93	0.41			−0.0	U			H42	1.5	93Si05	
^{208}Pb−^{207}Pb$_{1.005}$	875.5	0.6	875.71	0.06	0.3	U			TG4	1.0	23Ch12	*
^{208}Pb−^{206}Pb$_{1.010}$	2442.0	0.6	2442.15	0.12	0.3	U			TG4	1.0	23Ch12	*
^{207}Fr−^{208}Fr$_{.498}$ ^{206}Fr$^x_{.502}$	−890	60	−910	50	−0.1	U			P24	2.5	82Au01	
^{208}Po(α)^{204}Pb	5216.3	2.	5215.4	1.3	−0.5	2			Dba		69Go23	Z
	5214.0	3.			0.5	2					70Ra14	Z
	5215.1	2.			0.1	2					89Ma05	
^{208}At(α)^{204}Bi	5750.6	3.	5751.1	2.2	0.2	3			Dba		69Go23	Z
	5751.6	3.			−0.2	3			Dba		81Va27	Z
^{208}Rn(α)^{204}Po	6269.3	4.	6260.7	1.7	−2.1	–					55Mo69	Z
	6260.0	3.			0.2	–			Dba		71Go35	Z
	6264.7	8.2			−0.5	o			ISa		71Ho01	
	6257.5	5.			0.6	–					74Ho27	
	6258.7	2.5			0.8	–			Lvn		93Wa04	
ave.	6260.7	1.7			0.0	1	100	83 ^{208}Rn			average	
^{208}Fr(α)^{204}At	6778.3	5.	6769.7	2.6	−1.7	–					67Va20	Z
	6767.7	5.			0.4	–					74Ho27	Z
	6767.7	5.			0.4	–			ORa		81Ri04	Z
	6739.8	51.0			0.6	U			GSa		15De22	
	6780.6	18.4			−0.6	U					15Ma63	
	6772.4	18.4			−0.1	U			Lza		19Zh23	
	6762.3	6.1			1.2	–			Lza		26Zh40	
ave.	6769.5	2.7			0.1	1	99	81 ^{204}At			average	
^{208}Ra(α)^{204}Rn	7273.1	5.	7273	5	−0.1	2					67Va22	Z
	7270.1	16.3			0.2	2			Lza		22Zh45	
^{208}Ac(α)^{204}Fr	7720.8	15.	7718	10	−0.2	2			Jya		94Le05	
	7769.7	40.			−1.3	2			JAa		96Ik01	
	7707.5	21.4			0.5	o			Lza		14Ya19	
	7709.5	14.3			0.6	2			Lza		22Ya27	*
^{208}Ac^m(α)^{204}Fr^n	7892.1	20.	7899	10	0.3	2			Dbb		94An01	
	7910.4	20.			−0.6	2			Jya		94Le05	
	7871.7	50.			0.5	o			JAa		96Ik01	
	7899.2	40.8			−0.0	2			JAa		98LuZV	
	7897.2	14.3			0.1	2			Lza		22Ya27	
^{208}Th(α)^{204}Ra	8202.0	30.	8209	16	0.2	4			Anv		10He25	
	8211.2	18.4			−0.1	4			Lza		22Ya15	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{206}Pb(t,p)^{208}Pb		5622	30	5623.86	0.11	0.1	U				Ald		67Ha.A	
^{207}Pb(n,γ)^{208}Pb		7367.95	0.15	7367.87	0.05	−0.5	–				MMn		81Ke11	Z
		7367.96	0.10			−0.9	–						81Su.A	Z
		7367.81	0.11			0.6	–				ILn		83Hu13	Z
		7367.774	0.098			1.0	–						98Be19	Z
		7367.92	0.16			−0.3	–				Bdn		06Fi.A	
^{208}Pb(γ,n)^{207}Pb		−7370	3	−7367.87	0.05	0.7	U				McM		79Ba06	
^{208}Pb(d,t)^{207}Pb		−1114	25	−1110.64	0.05	0.1	U				Pit		64Co11	
^{207}Pb(n,γ)^{208}Pb	ave.	7367.87	0.05	7367.87	0.05	0.0	1	100	100	^{207}Pb			average	
^{208}Tl(β^-)^{208}Pb		4989.7	7.	4994.4	0.5	0.7	U						48Ma29	*
		4997.7	10.			−0.3	U						54El24	*
^{208}Bi(ε)^{208}Pb		2810	4	2878.9	2.0	17.2	B						59Mi19	*
*^{208}Pb−^{207}Pb$_{1.005}$	constructed from PI−ICR frequency ratio												HWJ25a	**
*^{208}Pb−^{206}Pb$_{1.010}$	constructed from PI−ICR frequency ratio												HWJ25a	**
*^{208}Ac(α)^{204}Fr	Also E_α=7483(15) to an unknown excited state												22Ya27	**
*^{208}Tl(β^-)^{208}Pb	E_{β^-}=1792(7) 1800(10) respectively, to 5$^-$ level at 3197.711 keV												Ens077	**
*^{208}Bi(ε)^{208}Pb	pK=0.24(0.01) to 3$^-$ level at 2614.522 keV, recalculated												Ens077	**
^{209}Pb−^{133}Cs$_{1.571}$		129631.85	0.35				2				SH2	1.0	22Ka.A	*
^{209}Bi−^{133}Cs$_{1.571}$		128937.6	4.7	128932.7	0.8	−1.0	U				MA8	1.0	08We02	
^{209}Fr−^{226}Ra$_{.925}$		−27584	36	−27562	12	0.6	–				MA3	1.0	92Bo28	
	ave.	−27550	16			−0.8	1	62	62	^{209}Fr			average	
^{209}Bi ^{35}Cl−^{207}Pb ^{37}Cl		7444	3	7452.5	0.8	0.7	U				H17	4.0	64Mc07	
		7454.13	1.51			−0.4	U				H36	2.5	85De40	
^{208}Fr−^{209}Fr$_{.498}$ ^{207}Fr$_{.502}$		720	60	646	15	−0.5	U				P24	2.5	82Au01	
^{209}Bi(α)^{205}Tl		3137.0	2.2	3137.9	0.8	0.4	1	12	9	^{209}Bi			03De11	
^{209}Po(α)^{205}Pb		4974	5	4979.2	1.4	1.0	2						66Ha29	*
		4980.0	2.			−0.4	2				Dba		69Go23	*
		4979.3	2.			−0.0	2						89Ma05	*
^{209}At(α)^{205}Bi		5757.2	2.	5756.9	2.0	−0.1	1	95	54	^{205}Bi	Dba		69Go23	Z
^{209}Rn(α)^{205}Po		6157.5	3.	6155.4	2.0	−0.7	–				Dba		71Go35	Z
		6154.2	2.5			0.5	–				Lvn		93Wa04	
	ave.	6155.5	2.0			−0.1	1	99	75	^{205}Po			average	
^{209}Fr(α)^{205}At		6777.7	5.	6777	4	−0.0	2						67Va20	Z
		6777.3	5.			0.0	2						74Ho27	Z
^{209}Ra(α)^{205}Rn		7147.0	5.	7143.1	2.7	−0.8	2						67Va22	Z
		7141	5			0.4	2				GSa		03He06	*
		7142.0	4.			0.3	2						08Ha12	
^{209}Ac(α)^{205}Fr		7733.3	15.	7730	60	−0.1	2						68Va04	
		7738.4	20.			−0.2	2				Dbb		94An01	
		7729.2	15.			0.0	2				Jya		94Le05	
		7728.2	40.			0.0	U				JAa		96Ik01	
		7725.1	10.			0.1	2				GSa		00He17	
		7723.0	23.			0.1	o				Lza		14Ya19	
		7724.0	14.3			0.1	2				Lza		22Ya27	
^{209}Th^m(α)^{205}Ra^m		8238.0	50.	8273	23	0.7	5				JAa		96Ik01	*
		8281.8	25.			−0.3	5				Anv		10He25	*
^{207}Pb(t,p)^{209}Pb		2814	12	2815.6	0.3	0.1	U				Ald		68Bj03	
^{209}Bi(p,t)^{207}Bi		−5864.8	2.0	−5864.9	2.0	−0.0	1	98	97	^{207}Bi	MSU		76Be.B	*
^{208}Pb(n,γ)^{209}Pb		3938.0	0.5	3929.5	0.3	−16.9	B						02Bl08	
^{208}Pb(d,p)^{209}Pb		1705	15	1705.0	0.3	−0.0	U				MIT		64Sp12	
		1700	10			0.5	U						67Mu16	
		1718	4			−3.3	B				Pit		72Ko03	*
		1715	10			−1.0	U				Yal		74Ko20	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{209}Bi(t,α)^{208}Pb	16003	25	16015.4	0.8	0.5	U			Ald		68Bj01	
^{209}Bi(γ,n)^{208}Bi	−7432	10	−7459.8	1.9	−2.8	U			Phi		60Ge01	
	−7460	2			0.1	2			McM		79Ba06	
^{209}Bi(d,t)^{208}Bi	−1216	30	−1202.5	1.9	0.4	U			Pit		64Co11	
	−1201	5			−0.3	2			ANL		64Er06	
^{209}Pb(β^-)^{209}Bi	644.6	1.2	651.3	0.8	5.6	B					72Be44	
^{209}Rn(β^+)^{209}At	3928	40	3943	11	0.4	R					74Vy01	*
*^{209}Pb−^{133}Cs$_{1.571}$	From frequency ratio $^{209}Pb^{++}/(^{133}Cs)^{+}$ =0.7862012402(13)										HWJ262	**
*^{209}Po(α)^{205}Pb	E_α=4876.8(5,Z) 4882.8(2,Z) respectively, to (20% ground state + 80% $^{205}Pb^m$ at 2.329 keV)										Ens044	**
*^{209}Po(α)^{205}Pb	E_α=4882.6(2.0) 4622(5) to (ground state + 80% $^{205}Pb^m$ at 2.329), 3/2$^-$ at 262.8keV										Ens044	**
*^{209}Ra(α)^{205}Rn	E_α=7003(10) to ground state, 6625(5) to 387.0 level										03He06	**
*^{209}Th^m(α)^{205}Ra^m	the decay is from ^{209}Th isomer, following $Ea_1-Ea_2-Ea_3-Ea_4$ correlations										FGK141	**
*^{209}Bi(p,t)^{207}Bi	$Q-Q(^{208}$Pb(p,t))=–241(2,Be), Q(Pb)=–5623.82(0.20) keV										AHW	**
*^{208}Pb(d,p)^{209}Pb	$Q-Q(^{209}$Bi(d,p))=–662(4),Q(Bi)=2380.01(0.14) keV										AHW	**
*^{209}Rn(β^+)^{209}At	E_{β^+}=2160(40) to 7/2$^-$ level at 745.81 keV										Ens156	**
^{210}Fr−^{226}Ra$_{.929}$	−27198	24	−27193	14	0.2	1	36	36 ^{210}Fr	MA3	1.0	92Bo28	
^{210}Ra−u	441.3	35.5	474	10	0.9	U			RI1	1.0	18Ro14	
^{210}Ac−u	9260	169	9340	60	0.4	1	13	13 ^{210}Ac	RI1	1.0	18Ro14	
^{209}Fr−^{210}Fr$_{.498}$ ^{208}Fr$_{.502}$	−770	50	−778	14	−0.1	U			P24	2.5	82Au01	
^{210}Pb(α)^{206}Hg	3792.4	20.	3780	8	−0.6	1	14	14 ^{206}Hg			62Ka27	
^{210}Bi(α)^{206}Tl	5042.8	2.	5036.2	0.8	−3.3	B			Orm		60Wa14	*
	5037.3	1.1			−1.0	1	48	37 ^{210}Bi			76Tu.A	*
^{210}Po(α)^{206}Pb	5407.53	0.07	5407.53	0.07	0.0	1	100	100 ^{210}Po			73Go39	Z
	5407.7	2.			−0.1	U					89Ma05	
^{210}At(α)^{206}Bi	5630.9	1.5	5631.2	1.0	0.2	3			Dba		69Go23	*
	5631.4	1.3			−0.2	3			Dba		81Va27	*
^{210}Rn(α)^{206}Po	6162.1	3.	6159.0	2.2	−1.0	3					55Mo69	Z
	6155.9	3.			1.0	3			Dba		71Go35	Z
^{210}Fr(α)^{206}At	6699.9	5.	6671	5	−5.8	B					67Va20	*
	6672.3	5.			−0.3	1	96	54 ^{206}At	GSa		05Ku06	
^{210}Ra(α)^{206}Rn	7156.6	5.	7150	3	−1.2	2					67Va22	Z
	7147	5			0.7	2			GSa		03He06	*
	7145.4	8.2			0.6	2			Jya		07Le14	
	7142.3	14.3			0.6	2			Lza		22Zh45	
^{210}Ac(α)^{206}Fr	7607.2	8.	7600	60	−0.2	–					68Va04	
	7607.2	10.			−0.2	–			GSa		00He17	
ave.	7610	40			−0.2	1	87	87 ^{210}Ac			average	
^{210}Th(α)^{206}Ra	8052.7	17.	8070	5	1.0	2			Jya		95Uu01	
	7962.0	50.			2.2	F			JAa		96Ik01	*
	8071.0	6.			−0.2	2			Anv		10He25	
	8076.1	14.3			−0.4	2			Lza		23Ch24	
^{210}Pa(α)^{206}Ac	8445.2	15.3				4			Lza		25Zh34	
^{208}Pb(t,p)^{210}Pb	628	12	640.3	0.8	1.0	U			Ald		68Bj03	
^{209}Bi(n,γ)^{210}Bi	4604.5	0.3	4604.64	0.08	0.5	–					71Mo03	
	4604.68	0.14			−0.3	–			MMn		83Ts01	Z
	4604.63	0.10			0.1	–			Bdn		06Fi.A	
^{209}Bi(d,p)^{210}Bi	2369	10	2380.07	0.08	1.1	U			MIT		64Sp12	
^{209}Bi(n,γ)^{210}Bi ave.	4604.64	0.08	4604.64	0.08	−0.0	1	100	90 ^{209}Bi			average	
^{210}Tl(β^-)^{210}Pb	5500	100	5481	12	−0.2	U					64We06	*
^{210}Pb(β^-)^{210}Bi	63.5	0.5	63.36	0.19	−0.3	–					67Ha03	*
	63.339	0.20			0.1	–					25Zh.A	*
ave.	63.36	0.19			−0.0	1	100	99 ^{210}Pb			average	
^{210}Bi(β^-)^{210}Po	1160.5	1.5	1161.7	0.8	0.8	–					62Da03	
	1161.5	1.5			0.1	–					67Hs01	
ave.	1161.0	1.1			0.7	1	52	52 ^{210}Bi			average	
^{210}At(ε)^{210}Po	3870	30	3981	8	3.7	B					63Sc15	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value	Adjusted value	v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{210}$Bi(α)^{206}Tl	E_α=4685.3(2,Z), 4648.3(2,Z) to 2^- level at 265.832, 1^- at 304.896 keV								Ens085	**
$*^{210}$Bi(α)^{206}Tl	E_α=4946(1), 4909(1) from ^{210}Bi^m at 271.31 keV								Nub269	**
*	to 2^- level at 265.832, 1^- level at 304.896 keV								Ens085	**
$*^{210}$At(α)^{206}Bi	E_α=5523.8, 5464.8, 5441.8(1.5,Z) to ground state, 4^+ at 59.897, 5^+ at 82.818								Ens085	**
$*^{210}$At(α)^{206}Bi	E_α=5524.1, 5465.3, 5442.8(1.3,Z) to ground state, 4^+ at 59.897, 5^+ at 82.818								Ens085	**
$*^{210}$Fr(α)^{206}At	E_α=6572.0(5,Z) 6542(5,Z) to ground state and level at 31.05 keV								Ens085	**
$*^{210}$Ra(α)^{206}Rn	E_α=7003(10) to ground state, 6447(5) to 574.9 level								03He06	**
$*^{210}$Th(α)^{206}Ra	F : Low energy; may be escape								96Ik01	**
$*^{210}$Tl(β^-)^{210}Pb	E_{β^-}=1870(100) to 3625(19) level, and other E_{β^-}								Ens148	**
$*^{210}$Pb(β^-)^{210}Bi	E_{β^-}=17.0(0.5) to 0^- level at 46.5390 keV								Ens148	**
$*^{210}$Pb(β^-)^{210}Bi	E_{β^-}=16.80(0.20) to 0^- level at 46.5390 keV								Ens148	**
$*^{210}$At(ε)^{210}Po	pK=0.46(0.10) to $(6)^-$ level at 3727.34 keV								Ens148	**
^{211}Tl−u	−6525	45			2			GS3	1.0	08ChZI	
^{211}Pb−^{133}Cs$_{1.586}$	138688.84	0.21	138688.81	0.21	−0.1	1	97	97 ^{211}Pb	SH2	1.0	22Ka.A	*
^{211}Po−u	−13519	147	−13348.3	0.6	1.2	U			GR1	1.0	19An10	
^{211}Fr−^{133}Cs$_{1.586}$	145517	15	145508	13	−0.6	1	74	74 ^{211}Fr	MA8	1.0	09Ko35	
^{211}Fr−u	−4410	43	−4445	13	−0.8	U			GR1	1.0	19An10	
^{211}Fr−^{226}Ra$_{.934}$	−28200	25	−28175	13	1.0	1	27	26 ^{211}Fr	MA3	1.0	92Bo28	
^{211}Ra−^{133}Cs$_{1.586}$	150846.4	8.5	150847	5	0.0	1	41	41 ^{211}Ra	MA8	1.0	09Ko35	
^{211}Ra−u	879	27	894	5	0.5	U			RI1	1.0	18Ro14	
^{211}Ac−u	7522	106	7670	60	1.4	1	27	27 ^{211}Ac	RI1	1.0	18Ro14	
^{211}Po^m−^{211}Po	1580	129	1570	5	−0.0	U				2.5	15Di03	
^{207}Fr−^{211}Fr$_{.327}$ ^{205}Fr$_{.673}$	−930	100	−615	18	1.3	U			P24	2.5	82Au01	
^{208}Fr−^{211}Fr$_{.394}$ ^{206}Fr$^x_{.606}$	−260	50	−290	60	−0.3	U			P24	2.5	82Au01	
^{210}Fr−^{211}Fr$_{.498}$ ^{209}Fr$_{.502}$	580	50	617	16	0.3	U			P24	2.5	82Au01	
^{211}Bi(α)^{207}Tl	6749.5	0.7	6750.4	0.5	1.3	3					61Ry02	Z
	6751.1	0.6			−1.1	3					71Gr17	Z
	6752.6	10.2			−0.2	U			GSt		23Sa03	*
^{211}Po(α)^{207}Pb	7594.5	0.5				2			Orm		62Wa18	Z
	7594.3	3.	7594.6	0.5	0.1	U			Dba		69Go23	Z
	7600.6	2.			−3.0	B					85La17	Z
	7586.0	15.3			0.6	U					15Di03	
	7604.5	30.6			−0.3	U			GSa		18Mi11	
	7594.2	7.1			0.1	U			GRa		25To01	
	7600.4	13.3			−0.4	U			Lza		25Gu12	
^{211}Po^m(α)^{207}Pb	9057.1	5.1				2			Bka		82Bo04	
	9049.0	15.	9057	5	0.5	U					89Ku08	*
	9043.0	15.			0.9	U					15Di03	*
	9047	5			2.0	U			GRa		25Ti01	*
^{211}Po^m(α)^{207}Pb^m	9047.2	5.1	*			U			GRa		25To01	
^{211}At(α)^{207}Bi	5979.4	2.	5982.4	1.3	1.5	2			Dba		69Go23	Z
	5981.6	3.			0.3	2			Bka		82Bo04	*
	5985.9	2.			−1.7	2					85La17	Z
^{211}Rn(α)^{207}Po	5967.9	2.	5965.5	1.4	−1.2	2					55Mo69	Z
	5963.1	2.			1.2	2			Dba		71Go35	Z
^{211}Fr(α)^{207}At	6660.2	5.1	6662	3	0.4	2					67Va20	Z
	6663.5	4.			−0.3	2			GSa		05Ku06	
^{211}Ra(α)^{207}Rn	7045.3	5.	7041.8	2.9	−0.7	–					67Va22	Z
	7040	5			0.4	–			GSa		03He06	*
	7039.7	6.			0.3	–			Jya		07Le14	
	7033.5	15.3			0.5	U			Lza		19Zh54	
ave.	7042	3			−0.0	1	91	59 ^{211}Ra			average	
^{211}Ac(α)^{207}Fr	7624.8	8.	7570	50	−1.1	–					68Va04	
	7616.7	10.			−1.0	–			GSa		00He17	
	7601.4	7.1			−0.7	–			Lza		26Zh40	
ave.	7614	29			−0.8	1	75	73 ^{211}Ac			average	
^{211}Th(α)^{207}Ra	7942.9	14.	7950	50	0.2	–			Jya		95Uu01	
	7930.6	30.6			0.4	–			Lza		15Ya13	
ave.	7940	40			0.3	1	61	61 ^{211}Th			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$^{211}Th^m(\alpha)^{207}Ra$	8760	62	8730	50	−0.3	1	39	39 $^{211}Th^m$	Jya		21Au03	*
$^{211}Pa(\alpha)^{207}Ac$	8481.0	40.8				3			Jya		20Au04	
$^{211}Pb(\beta^-)^{211}Bi$	1378	8	1363.4	0.8	−1.8	U					65Co06	
$^{211}Th^m(IT)^{211}Th$	778	7	778	7	0.0	1	100	61 $^{211}Th^m$			21Au03	
$*^{211}Pb-^{133}Cs_{1.586}$	From frequency ratio $^{211}Pb^{++}/(^{133}Cs)^+$ =0.7937541548(08)										HWJ262	**
$*^{211}Bi(\alpha)^{207}Tl$	E_α=6280(10) keV to the 351 keV level										23Sa03	**
$*^{211}Po^m(\alpha)^{207}Pb$	E_α=7275(15) to 13/2$^+$ level at 1633.356 keV										Ens112	**
$*^{211}Po^m(\alpha)^{207}Pb$	E_α=7269(15) to 13/2$^+$ level at 1633.356 keV										Ens112	**
$*^{211}Po^m(\alpha)^{207}Pb$	E_α=8877(13),7800(28),7273(5) keV to ground state, 898 keV,1633 keV level										25Ti01	**
$*^{211}At(\alpha)^{207}Bi$	Recalibrated as in reference										91Ry01	**
$*^{211}Ra(\alpha)^{207}Rn$	Average of E_α=6907(5) and several branches to known levels										03He06	**
$*^{211}Th^m(\alpha)^{207}Ra$	E_α=8045(15) keV to 13/2$^+$ level at 560(60) kev										21Au03	**
$^{212}Bi^n-u$	−7127	32				2			GS3	1.0	08ChZI	
$^{212}At-u$	−9234	92	−9263.6	2.3	−0.3	U			GR1	1.0	19An10	
$^{212}Rn-u$	−9242	32	−9298	3	−1.7	U			GR1	1.0	19An10	
$^{212}Fr-^{133}Cs_{1.594}$	146938	10	146935	9	−0.3	1	89	89 ^{212}Fr	MA8	1.0	09Ko35	
$^{212}Fr-u$	−3790	30	−3775	9	0.5	U			GR1	1.0	19An10	
$^{212}Fr-^{226}Ra_{.938}$	−27631	28	−27606	10	0.9	1	12	11 ^{212}Fr	MA3	1.0	92Bo28	
$^{212}Ra-u$	−212.7	26.6	−215	11	−0.1	1	17	17 ^{212}Ra	RI1	1.0	18Ro14	
$^{212}Ac-u$	7840.7	25.4	7835	23	−0.2	1	85	85 ^{212}Ac	RI1	1.0	18Ro14	
$^{212}Pb-^{220}Rn_{.964}$	−19087.45	0.28	−19087.30	0.25	0.5	1	80	81 ^{212}Pb	CP2	1.0	23Ya.A	
$^{209}Fr-^{212}Fr_{.563}\ ^{205}Fr_{.437}$	−1270	70	−1232	13	0.2	U			P24	2.5	82Au01	
$^{206}Fr^x-^{212}Fr_{.139}\ ^{205}Fr_{.861}$	340	130	390	100	0.2	U			P24	2.5	82Au01	
$^{207}Fr-^{212}Fr_{.163}\ ^{206}Fr^x_{.837}$	−1150	70	−1250	90	−0.6	U			P24	2.5	82Au01	
$^{212}Bi(\alpha)^{208}Tl$	6207.12	0.04	6207.262	0.028	3.5	B			BIP		61Ry02	Z
	6207.09	0.08			2.1	o			BIP		69Gr28	*
	6207.262	0.028				3			BIP		72Go.A	*
$^{212}Bi^m(\alpha)^{208}Tl$	6458.1	30.				3					78Ba44	
$^{212}Po(\alpha)^{208}Pb$	8953.6	0.8	8954.20	0.11	0.7	U					61Ry02	Z
	8953.85	0.31			1.1	2					71De52	Z
	8953.3	0.6			1.5	U					71Gr17	Z
	8954.25	0.12			−0.4	2					74Hu15	Z
	8949.2	10.2			0.5	U			GSt		23Sa03	
	8971	13			−1.3	U			GRa		25To01	
$^{212}Po^m(\alpha)^{208}Pb$	11874.6	20.	11877	4	0.1	2					62Pe15	
	11859.3	15.			1.2	o					75Fr.B	
	11884.8	10.2			−0.7	2					76Fr.A	
	11875.6	5.1			0.3	2			GSa		12Ho12	
$^{212}At(\alpha)^{208}Bi$	7829.0	9.	7817.1	0.6	−1.3	U					70Re02	*
	7817.0	0.6				3					76Fr.A	*
	7828.0	10.			−1.1	U					96Li37	*
	7827.0	10.2			−1.0	U			Lza		25Gu12	
$^{212}At^m(\alpha)^{208}Bi$	8049.3	10.	8040.0	0.7	−0.9	U					68Va18	
	8054.3	9.			−1.6	U					70Re02	*
	8040.00	0.61			−0.0	o					76Fr.A	*
	8051.2	10.			−1.1	U					96Li37	*
	8043.2	10.2			−0.3	U			Lza		25Gu12	*
$^{212}Rn(\alpha)^{208}Po$	6392.3	5.	6385.1	2.6	−1.4	3					55Mo69	Z
	6382.5	3.			0.9	3			Dba		71Go35	Z
$^{212}Fr(\alpha)^{208}At$	6531.3	3.	6529.0	1.6	−0.7	2					66Va.A	Z
	6528.0	3.			0.3	2			Dba		81Va27	
	6527.5	3.			0.5	2			Bka		82Bo04	*
	6529.5	4.			−0.1	2			GSa		05Ku06	
$^{212}Ra(\alpha)^{208}Rn$	7030.0	5.	7031.7	1.7	0.3	–					67Va22	Z
	7034.0	5.			−0.4	–					74Ho27	Z
	7032.2	2.			−0.3	–			Bka		82Bo04	Z
	7028	5			0.7	–			GSa		03He06	*
	7040	24			−0.3	U			Lza		14Ya19	
ave.	7031.7	1.7			0.0	1	100	83 ^{212}Ra			average	
$^{212}Ac(\alpha)^{208}Fr$	7521.2	8.	7541	24	0.4	–					68Va04	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{212}Ac(α)^{208}Fr		7515.1	10.	7541	24	0.5	–			GSa		00He17	
		7514.0	18.			0.5	–			Lza		14Ya19	
		7518.0	17.			0.4	o			Lza		19Zh23	
		7502.8	7.1			0.8	–			Lza		26Zh40	
	ave.	7513	26			0.5	1	18	15 ^{212}Ac			average	
^{212}Th(α)^{208}Ra		7952.3	10.	7958	5	0.6	3					80Ve01	
		7959.5	5.			−0.3	3			Anv		10He25	
		7980.8	20.4			−1.1	U			GSa		15De22	
		7963.6	16.3			−0.3	U			Lza		22Zh45	
^{212}Pa(α)^{208}Ac		8429.4	30.	8410	60	−0.3	3			JAa		97Mi03	
		8406.9	20.			0.1	3			Lza		14Ya19	*
		8398.7	20.			0.2	3			Jva		20Au04	
^{210}Pb(t,p)^{212}Pb		515	25	485.4	0.9	−1.2	U					71El05	
^{212}Pb(β^-)^{212}Bi		569.3	2.5	569.4	0.5	0.0	U					48Ma30	*
		576.6	5.			−1.4	U					58Se71	*
^{212}Bi(β^-)^{212}Po		2256	3	2247.4	0.6	−2.9	U					48Fe09	
		2250.5	2.5			−1.2	U					48Ma30	
^{212}At^m(IT)^{212}At		222.9	0.3				4					76FrZO	*
*^{212}Bi(α)^{208}Tl	E_α=6089.86(0.08,Z), 6050.57(0.07,Z) to ground state, 4$^+$ level at 39.858 keV											Ens077	**
*^{212}Bi(α)^{208}Tl	E_α=6089.883(0.037,Z), 6050.837(0.028,Z) to ground state, 4$^+$ level at 39.858 keV											72Go.A	**
*^{212}At(α)^{208}Bi	Original E_α=7679(8); calibration ^{211}Po 7448(1), now 7450.3(0.5) keV											AHW	**
*^{212}At(α)^{208}Bi	Original E_α=7669.0(0.2); calibration ^{211}Po 7450(2), now 7450.3(0.5)											05Ma.A	**
*^{212}At(α)^{208}Bi	E_α=7679(10) to ground state, 7618(10) to 63.3 level											96Li37	**
*	error estimated by the evaluators											GAu	**
*^{212}At^m(α)^{208}Bi	Original E_α=7900(8); calibration ^{211}Po 7448(1), now 7450.3(0.5) keV											GAu	**
*^{212}At^m(α)^{208}Bi	Original E_α=7887.7(0.2); calibration ^{211}Po 7450(2), now 7450.3(0.5)											GAu	**
*^{212}At^m(α)^{208}Bi	E_α=7897(10) to ground state, 7837(10) to 63.3 level											96Li37	**
*	error estimated by the evaluators											GAu	**
*^{212}At^m(α)^{208}Bi	E_α=7879(12) to ground state, 7829(10) to 63.3 level											25Gu12	**
*^{212}Fr(α)^{208}At	E_α=6341(3) (recalibrated as in reference) to 63.70 level											91Ry01	**
*^{212}Ra(α)^{208}Rn	E_α=6898(5) to ground state, 6269(5) to 635.1 level											03He06	**
*^{212}Pa(α)^{208}Ac	E_α=8250(20) keV in reference by combining with ^{97}Mi$_{03}$											14Ya19	**
*^{212}Pb(β^-)^{212}Bi	E_{β^-}=330.7(2.5) 338(5) respectively to 0$^-$ level at 238.632 keV											Ens053	**
*^{212}At^m(IT)^{212}At	from the relative E_α=7887.7(0.2) and 7669.0(0.2) in ^{76}FrZO											FGK219	**
^{213}Tl−u		1893	65	1915	29	0.3	o			GS3	1.0	10Ch19	
		1915	29				2			GS3	1.0	12Ch19	
^{213}Bi−^{133}Cs$_{1.602}$		145847.60	0.32	145847.6	0.3	−0.0	1	98	98 ^{213}Bi	SH2	1.0	22Ka.A	*
^{213}Rn−u		−6159	63	−6116	3	0.7	U			GR1	1.0	19An10	
^{213}Fr−^{133}Cs$_{1.602}$		147649.1	7.4	147650	5	0.1	1	46	46 ^{213}Fr	MA8	1.0	09Ko35	
^{213}Fr−u		−3823.9	13.2	−3816	5	0.6	1	15	15 ^{213}Fr	GR1	1.0	19An10	
^{213}Ra−^{133}Cs$_{1.602}$		151833	12	151837	11	0.3	1	77	77 ^{213}Ra	SH1	1.0	13Dr04	
^{213}Ac−u		6572.2	19.4	6592	13	1.0	1	42	42 ^{213}Ac	RI1	1.0	18Ro14	
^{207}Fr−^{213}Fr$_{.324}$ ^{204}Fr$_{.676}$		−2559	330	−2110	18	0.5	U			P24	2.5	82Au01	*
^{208}Fr−^{213}Fr$_{.279}$ ^{206}Fr$^x_{.721}$		−700	60	−790	70	−0.6	U			P24	2.5	82Au01	
^{209}Fr−^{213}Fr$_{.327}$ ^{207}Fr$_{.673}$		−670	60	−703	17	−0.2	U			P24	2.5	82Au01	
^{209}Fr−^{213}Fr$_{.196}$ ^{208}Fr$_{.804}$		−980	60	−941	14	0.3	U			P24	2.5	82Au01	
^{211}Fr−^{213}Fr$_{.330}$ ^{210}Fr$_{.670}$		−830	60	−727	15	0.7	U			P24	2.5	82Au01	
^{212}Fr−^{213}Fr$_{.498}$ ^{211}Fr$_{.502}$		270	50	333	11	0.5	U			P24	2.5	82Au01	
^{213}Bi(α)^{209}Tl		5982.6	6.	5988	3	0.9	2					64Gr11	
		5990.7	4.			−0.6	2			Gea		13Ma13	
^{213}Po(α)^{209}Pb		8537.1	5.	8536.7	2.6	−0.1	3					64Va20	Z
		8536.5	3.			0.1	3			Bka		82Bo04	Z
		8530.5	10.2			0.6	o			GSt		18Sa45	
		8540.7	10.2			−0.4	U			GSt		23Sa03	
		8548	47			−0.2	U			GRa		25To01	
^{213}At(α)^{209}Bi		9254.2	12.	9254	5	−0.0	2					70Bo13	
		9254.2	5.			−0.0	2			Lvn		88Hu08	
^{213}Rn(α)^{209}Po		8245.1	8.	8245.2	2.9	−0.0	3					67Va20	
		8240.0	10.			0.5	U					70Va13	
		8242	10			0.3	U			GSa		00He17	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{213}Rn(α)^{209}Po		8245.2	3.1	8245.2	2.9	−0.0	3			Jya		01Ku07	
		8218.6	44.			0.6	U					05Li17	
		8204.4	30.6			1.3	U			GSa		18Mi11	
		8245.0	10.			0.0	U			GSa		19Mi08	*
		8255.3	15.3			−0.7	U			Lza		21Hu19	
		8249.2	10.2			−0.4	U			Lza		25Gu12	
^{213}Fr(α)^{209}At		6904.0	5.	6904.7	1.3	0.1	U					67Va20	Z
		6908.0	5.			−0.7	U					74Ho27	Z
		6904.6	2.			0.0	–			Bka		82Bo04	Z
		6904.9	1.7			−0.1	–			GSa		05Ku06	
		6880	20			1.2	U			GSa		15De22	
		6869.2	20.4			1.7	U			GSa		19Mi08	
	ave.	6904.8	1.3			−0.1	1	98	59 ^{209}At			average	
^{213}Ra(α)^{209}Rn		6860.3	5.	6861.7	2.3	0.3	–					67Va22	*
		6862.4	5.			−0.1	–					76Ra37	*
		6862.2	3.			−0.2	–			GSa		06Ku26	*
	ave.	6861.8	2.3			−0.1	1	99	76 ^{209}Rn			average	
^{213}Ra^{m}(α)^{209}Rn		8630.3	5.1	8630	4	−0.1	2					76Ra37	
		8629.3	5.			0.1	2			GSa		06Ku26	*
^{213}Ac(α)^{209}Fr		7505.2	8.	7498	4	−0.8	–					68Va04	*
		7497.0	10.			0.1	o			GSa		00He17	
		7497.0	5.			0.2	–			GSa		02He.A	
	ave.	7499	4			−0.2	1	97	58 ^{213}Ac			average	
^{213}Th(α)^{209}Ra		7837.4	10.	7837	7	−0.1	3					68Va18	*
		7836.5	10.			0.0	3					80Ve01	
^{213}Pa(α)^{209}Ac		8393.9	15.	8384	12	−0.6	3			GSa		00He17	
		8367.4	20.4			0.8	3			Jva		20Au04	
^{213}Bi(β^-)^{213}Po		1430	10	1413.1	2.7	−1.7	U					68Va17	*
*^{213}Bi−^{133}Cs$_{1.602}$	From frequency ratio ^{213}Bi^{++}/(^{133}Cs)$^{+}$ =0.8012995724(12)											HWJ262	**
*^{207}Fr−^{213}Fr$_{.324}$ ^{204}Fr.	D_M=–2470(330) keV for ^{204}Fr mixture gs+m+n at 45.9(1.7) 321.2(1.9) keV											Nub269	**
*^{213}Rn(α)^{209}Po	E_α=8088(10), 7550(15) to ground state, 540.3 level											00He17	**
*^{213}Ra(α)^{209}Rn	E_α=6730.7, 6623.7, 6520.7(3,Z) to ground state, 1/2$^-$ at 110.25, 3/2$^-$ at 214.93											Ens156	**
*^{213}Ra(α)^{209}Rn	E_α=6731.9, 6624.9, 6523.9(5,Z) to ground state, 1/2$^-$ at 110.25, 3/2$^-$ at 214.93											Ens156	**
*^{213}Ra(α)^{209}Rn	E_α=6733(3), 6625(3) to ground state and 1/2$^-$ level at 110.25 keV											Ens156	**
*^{213}Ra^{m}(α)^{209}Rn	E_α=8467 (5) 8358(10) to ground state and 1/2$^-$ level at 110.25 keV											Ens156	**
*^{213}Ac(α)^{209}Fr	Original Q increased by 2 keV, as in reference											91Ry01	**
*^{213}Th(α)^{209}Ra	Original Q decreased by 0.5 keV, as in reference											91Ry01	**
*^{213}Bi(β^-)^{213}Po	E_{β^-}=1420(10) 1018(15) to ground state and 7/2$^+$ level at 440.45 keV											Ens074	**
^{214}Ra−^{133}Cs$_{1.609}$		152235	22	152226	6	−0.4	U			MA8	1.0	08We02	
^{214}Ra−u		63	38	98	6	0.9	U			RI1	1.0	18Ro14	
^{214}Ac−u		6866.8	30.9	6906	15	1.3	1	22	22 ^{214}Ac	RI1	1.0	18Ro14	
^{214}Bi(α)^{210}Tl		5621.3	3.0				2					91Ry01	*
^{214}Po(α)^{210}Pb		7833.54	0.06	7833.54	0.06	−0.0	1	100	99 ^{214}Po			71Gr17	Z
		7853	38			−0.5	U			GRa		25To01	
^{214}At(α)^{210}Bi		8987.2	4.	8988	4	0.1	1	84	84 ^{214}At	Bka		82Bo04	Z
^{214}At^{m}(α)^{210}Bi		9046.4	8.				2					82Ew01	
^{214}At^{n}(α)^{210}Bi		9220.8	5.	9220	4	−0.2	1	76	76 ^{214}At^{n}			82Ew01	*
^{214}Rn(α)^{210}Po		9212.6	20.	9208	9	−0.2	2					70To07	
		9207.5	10.			0.1	2					70Va13	
		9334.8	51.			−2.5	U			GSa		18Mi11	
^{214}Fr(α)^{210}At		8585.5	8.	8589	4	0.4	4					68Va18	*
		8590.9	5.			−0.4	4					70To18	*
		8583.8	10.			0.5	4			Dbb		89An.A	
		8590.8	20.			−0.1	U			GSa		90Ni05	
		8578.7	48.			0.2	U					05Li17	
		8590.9	10.2			−0.2	4			GSa		19Mi08	
		8589.9	15.2			−0.1	4			Lza		21Hu18	
^{214}Fr^{m}(α)^{210}At		8711.7	8.	8710	3	−0.2	4					68Va04	Z
		8711.7	5.			−0.3	4					70To18	*
		8708.1	5.			0.4	4			GSa		05Ku06	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{214}Ra(α)^{210}Rn	7271.7	5.	7271.6	2.4	−0.0	4					67Va22	Z
	7275.6	5.			−0.8	4					74Ho27	Z
	7273.2	10.			−0.2	4			GSa		00He17	*
	7271.2	4.			0.1	4			GSa		06Ku26	*
	7266.1	6.1			0.9	4			Lza		23Lu04	
	7273.2	10.2			−0.2	U			Dba		23Ku17	
^{214}Ra(α)^{210}Rn^m	5563.9	30.				5			GSa		06Ku26	*
^{214}Ac(α)^{210}Fr	7351.7	5.	7351.9	2.5	0.0	–					68Va04	Z
	7347.6	10.			0.4	–			Dbb		89An13	
	7347.6	10.			0.4	o			GSa		00He17	*
	7349.6	5.			0.4	o			GSa		02He.A	
	7352.7	3.			−0.3	–			GSa		04Ku24	*
	7360.6	17.3			−0.5	U			Lza		20Ma27	
ave.	7352.1	2.5			−0.1	1	99	78 ^{214}Ac			average	
^{214}Th(α)^{210}Ra	7828.6	10.	7826	5	−0.3	3					68Va18	
	7823.5	10.			0.2	3					80Ve01	
	7828.6	8.			−0.3	3			Jya		07Le14	
	7814.3	16.3			0.7	3			Lza		22Zh45	
^{214}Pa(α)^{210}Ac	8270.9	15.				2			GSa		00He17	
^{214}U(α)^{210}Th	8695.8	18.3				3			Lza		21Zh22	
^{214}Pb(β^-)^{214}Bi	1024	20	1018	11	−0.3	1	31	31 ^{214}Bi			52Be78	*
^{214}Bi(β^-)^{214}Po	3260	30	3269	11	0.3	–					56Da06	
	3275	15			−0.4	–					60Lu07	
ave.	3272	13			−0.2	1	69	69 ^{214}Bi			average	
*^{214}Bi(α)^{210}Tl	E_α=5516(3) recommended in place of the following E_α:										91Ry01	**
*	E_α=5510.5(1.0) keV										34Le01	**
*	E_α=5515.8(3.0) keV										60Wa14	**
*^{214}At^n(α)^{210}Bi	E_α=8782(5) to 9$^-$ level at 271.31 keV										Ens148	**
*^{214}Fr(α)^{210}At	E_α=8425.5, 8352.5(8,Z) to ground state, (4)$^+$ level at 72.65 keV										Ens148	**
*^{214}Fr(α)^{210}At	E_α=8428.3, 8360.3(5,Z) to ground state, (4)$^+$ level at 72.65 keV										Ens148	**
*^{214}Fr^m(α)^{210}At	E_α=8546.8, 8477.8(5,Z) to ground state, (4)$^+$ level at 72.65 keV										Ens148	**
*^{214}Ra(α)^{210}Rn	E_α=7137(10), 6505(15) to ground state, 641.9 level										00He17	**
*^{214}Ra(α)^{210}Rn	Also E_α=8950(30) keV Q_α=9120.9 keV from ^{214}Ra^n at 1865.2 keV										Nub269	**
*^{214}Ra(α)^{210}Rn^m	E_α=7290(30) Q_α=7429.1 from ^{214}Ra^n at 1865.2 keV										Nub269	**
*^{214}Ac(α)^{210}Fr	E_α=7210(10), 7080(15) to ground state, 138.6 level										00He17	**
*^{214}Ac(α)^{210}Fr	Also E_α=7081(4) keV to 139.0(1) level										04Ku24	**
*^{214}Pb(β^-)^{214}Bi	E_{β^-}=670(20) to (0$^-$,1$^-$) level at 351.9324 keV, and another branch										Ens092	**
^{215}Bi−^{133}Cs$_{1.617}$	154654	16	154633	6	−1.3	1	13	13 ^{215}Bi	MA8	1.0	08We02	
^{215}Po(α)^{211}Pb	7526.45	0.8	7526.0	0.6	−0.6	1	49	46 ^{215}Po			71Gr17	Z
^{215}At(α)^{211}Bi	8178.5	4.	8178	4	−0.2	4			Bka		82Bo04	Z
	8172.3	10.2			0.5	4			GSt		18Sa45	
	8177.4	13.2			0.0	4			Lza		25Gu12	
^{215}Rn(α)^{211}Po	8834.7	20.	8840	5	0.2	3			ORa		69Ha32	
	8839.8	8.			−0.0	3					70Va13	
	8865.3	30.6			−0.8	3			GSa		18Mi11	
	8834.7	10.2			0.5	3			GSt		18Sa45	
	8857.1	19.4			−0.9	3			GRa		25To01	
	8837.8	11.2			0.2	3			Lza		25Gu12	
^{215}Fr(α)^{211}At	9543.0	15.	9540	7	−0.2	3					70Bo13	
	9532.7	10.			0.8	3					74No02	
	9546.9	10.			−0.6	3					84De16	
	9364.4	50.9			3.5	B			GSa		19Mi08	*
^{215}Ra(α)^{211}Rn	8862.7	5.	8862.4	2.3	−0.1	3					68Va18	Z
	8865.5	5.			−0.6	3					70To18	Z
	8865.3	10.			−0.3	U			GSa		00He17	
	8865.3	46.			−0.1	U					05Li17	
	8861.1	3.1			0.4	3			Lza		20Su02	
	8875.3	27.5			−0.5	U			LZa		24Ya27	
^{215}Ac(α)^{211}Fr	7748.4	5.	7746	3	−0.5	2					68Va04	Z
	7746	10			−0.0	o			GSa		00He17	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{215}Ac(α)^{211}Fr	7740.3	5.	7746	3	1.1	o			GSa		02He.A	
	7744.4	4.			0.4	2			GSa		04Ku24	
^{215}Th(α)^{211}Ra	7664.9	8.	7665	4	−0.0	2					68Va18	
	7667.0	10.			−0.2	o			GSa		89He03	
	7664	15			0.0	o			GSa		00He17	*
	7665	5			−0.1	2			GSa		05Ku31	*
	7662.8	10.			0.2	2			Jya		07Le14	*
	7652.2	15.2			0.8	U			Lza		19Zh54	
^{215}Pa(α)^{211}Ac	8238.6	15.	8240	60	−0.0	2					79Sc09	
	8244.7	15.			−0.1	o			GSa		00He17	
	8233.5	20.4			0.1	2			GSa		15De22	
	8240.6	9.2			−0.1	2			Lza		26Zh40	
^{215}U(α)^{211}Th	8588.0	30.6				2			Lza		15Ya13	
*^{215}Ac(α)^{211}Fr	E_α=7602(10) 7026(15) 6960(15) to ground state, 11/2$^-$ at 583.28, 13/2$^-$ at 652.62										Ens136	**
*^{215}Th(α)^{211}Ra	E_α=7520(15), 7387(15), 7336(15) to ground state, 133.6, 192.4 levels										00He17	**
*^{215}Th(α)^{211}Ra	E_α=7523(5), 7392(4), 7335(5), 7236(7) to ground state, 133.9, 194.5, 295.1 levels										05Ku31	**
*^{215}Th(α)^{211}Ra	Also E_α=7399(20) keV to 133.9 level										07Le14	**
^{216}Bi−^{133}Cs$_{1.624}$	159852	12				2			MA8	1.0	08We02	
^{216}Po−^{220}Rn$_{.982}$	−9275.3	1.3	−9274.07	0.11	0.9	U			CP2	1.0	23Ya.A	
^{216}Po(α)^{212}Pb	6906.44	0.5	6906.98	0.25	1.1	1	23	19 ^{212}Pb			71Gr17	Z
^{216}At(α)^{212}Bi	7949.7	3.				4			Bka		82Bo04	Z
	7939	16	7950	3	0.7	U			GRa		25To01	
^{216}At^m(α)^{212}Bi	8110.5	10.				4					71Br13	
^{216}Rn(α)^{212}Po	8199.2	10.	8195	5	−0.4	3					61Ru06	
	8201.2	10.			−0.6	3					70Va13	
	8192.0	10.2			0.3	3					71Br13	
	8202.2	20.4			−0.3	3			GSt		18Sa45	
	8207.3	15.3			−0.8	3			GRa		25To01	
	8173.7	13.2			1.6	3			Lza		25Gu12	
^{216}Fr(α)^{212}At	9175.3	12.	9174	3	−0.1	4					70Bo13	
	9174.1	5.			0.0	4					96Li37	*
	9174.3	5.			0.0	4			GSa		07Ku30	
	9204.8	22.4			−1.4	U			Lza		25Gu12	
^{216}Fr^m(α)^{212}At^m	9170.2	5.				5			GSa		07Ku30	
^{216}Ra(α)^{212}Rn	9525.8	8.	9526	7	−0.0	4					73No09	
	9525.8	17.3			−0.0	4			Lza		17Su18	
^{216}Ac(α)^{212}Fr	9243.3	8.	9240.8	2.9	−0.3	2					70To18	Z
	9223.1	10.			1.8	U			GSa		00He17	
	9241.4	50.9			−0.0	U					05Li17	
	9240.5	3.0			0.1	2			Lza		17Hu08	
	9240.5	6.1			0.1	o			Lza		18Hu13	
	9248.6	22.4			−0.3	o			Lza		21Ma66	
^{216}Ac^m(α)^{212}Fr	9280.0	5.	9279	4	−0.2	2					70To18	Z
	9284	10			−0.5	o			GSa		00He17	*
	9278.2	5.			0.2	o			GSa		02He.A	
	9277.2	7.			0.3	2			GSa		04Ku24	*
^{216}Th(α)^{212}Ra	8070.7	8.	8072	4	0.2	2					68Va18	
	8071	10			0.1	o			GSa		00He17	*
	8073	5			−0.1	2			GSa		05Ku31	*
	8069.7	44.			0.1	U					05Li17	
	8070.7	23.			0.1	U			Lza		14Ya19	
	8068.6	15.3			0.2	U			Lza		19Zh54	
^{216}Th^m(α)^{212}Ra	10099.4	20.	10114	7	0.7	2					83Hi08	
	10107.4	40.			0.2	U			Dbb		93An07	
	10120.8	15.			−0.5	2			GSa		00He17	
	10117.5	10.			−0.4	2					05Ku31	*
	10105.5	15.3			0.5	2			Lza		19Zh54	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{216}Pa(α)^{212}Ac		8013.7	20.	8106	7	4.6	B					79Sc09	
		8110.5	50.			−0.1	U			JAa		98Ik01	
		8097	15			0.6	2			GSa		00He17	*
		8102.3	17.3			0.2	o			Lza		19Zh23	
		8108.5	8.2			−0.3	2			Lza		26Zh40	
^{216}U(α)^{212}Th		8542.5	30.6	8529	16	−0.5	o			Lza		15Ma37	
		8497.6	50.9			0.6	4			GSa		15De22	
		8532.3	17.3			−0.2	4			Lza		21Zh22	
^{216}U^{m}(α)^{212}Th		10782.0	30.6	10738	16	−1.4	o			Lza		15Ma37	
		10738.0	16.3				4			Lza		22Zh45	*
*^{216}Fr(α)^{212}At	E_α=9004(5); and E_α=8933(8) from 133.3 level to 205.6 keV											96Li37	**
*^{216}Ac^{m}(α)^{212}Fr	E_α=9110(10), 9026(15), 8586(15) to ground state, 82.4, 542.2 levels											00He17	**
*^{216}Ac^{m}(α)^{212}Fr	Also E_α=9029(7) keV to 82.6(1) level											04Ku24	**
*^{216}Th(α)^{212}Ra	E_α=7923(10), 7302(15) to ground state, 618.3 level											00He17	**
*^{216}Th(α)^{212}Ra	E_α=7923(5), 7304(4) to ground state, 629.3(1) level											05Ku31	**
*^{216}Th^{m}(α)^{212}Ra	E_α=9930(10), 9312(12) to ground state, 629.3(1) level											05Ku31	**
*^{216}Pa(α)^{212}Ac	E_α=7948(15), 7815(15) to ground state, 133.6 level											00He17	**
*^{216}U^{m}(α)^{212}Th	Also E_α=10163(27) to 2^+ state at 383 keV in ^{212}Th											22Zh45	**
^{217}Bi−u		9420	32	9372	19	−1.5	o			GS3	1.0	08ChZI	
		9372	19				2			GS3	1.0	12Ch19	
^{217}At−u		4759	145	4715.5	0.9	−0.3	U			GR1	1.0	19An10	
^{217}Po(α)^{213}Pb		6660.3	4.	6662.1	2.4	0.4	4			Dba		77Vy02	Z
		6660.0	4.			0.5	4			Orm		97Li23	
		6666.1	4.1			−1.0	4			Anv		03Ku25	
^{217}At(α)^{213}Bi		7200.3	3.	7201.0	0.9	0.2	–					60Vo05	Z
		7200.3	2.			0.4	–			Orm		62Wa28	Z
		7204.6	5.			−0.7	U					64Va20	Z
		7193.1	5.			1.5	o			Dba		77Vy02	Z
		7204.0	2.			−1.5	–			Bka		82Bo04	
		7199.8	1.6			0.7	–			Dba		97Ch19	
	ave.	7201.1	1.0			−0.1	1	71	69 ^{217}At			average	
^{217}Rn(α)^{213}Po		7887.5	4.	7887.2	2.9	−0.1	4					61Ru06	Z
		7886.9	4.			0.1	4			Bka		82Bo04	Z
		7885.6	10.2			0.2	U			GSt		18Sa45	
		7896	26			−0.3	U			GRa		25To01	
		7873.4	12.2			1.1	U			Lza		25Gu12	
^{217}Fr(α)^{213}At		8471.5	8.	8469	4	−0.3	3					70Bo13	
		8468.4	5.			0.2	3			Lvn		88Hu08	
		8469.4	10.2			−0.0	3			GRa		25To01	
^{217}Ra(α)^{213}Rn		9159.1	8.	9161	6	0.2	4					70To07	
		9163.2	10.			−0.2	4					70Va13	
		9159.2	40.8			0.0	U			GSa		19Mi08	*
		9157.1	26.5			0.1	U			Lza		19Ya04	
		9134.7	15.3			1.7	U			Lza		21Hu19	
		9175.5	36.7			−0.4	U			Lza		25Gu12	
^{217}Ac(α)^{213}Fr		9831.6	10.	9832	9	−0.0	1	74	74 ^{217}Ac			73No09	
		9474.9	203.8			1.8	U			GSa		19Mi08	
^{217}Ac^{m}(α)^{213}Fr		11843.8	17.	11844	9	0.0	1	26	26 ^{217}Ac^{m}			85De14	
^{217}Th(α)^{213}Ra		9424.1	10.	9435	4	1.1	2					68Va18	
		9424.1	20.			0.6	U					73Ha32	
		9421.1	15.			0.9	U					00Ni02	
		9442	15			−0.5	o			GSa		00He17	*
		9435.6	5.			−0.1	2			GSa		02He29	*
		9443.5	9.			−0.9	2			GSa		05Ku31	
		9424.1	47.			0.2	U					05Li17	
		9431.0	15.3			0.3	U			Lza		19Zh54	
		9434.3	10.2			0.1	2			Dba		23Ku17	
^{217}Pa(α)^{213}Ac		8486.7	10.	8489	4	0.2	2					68Va18	
		8489.8	15.			−0.1	U					79Sc09	
		8486.7	50.			0.0	U			JAa		98Ik01	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{217}Pa(α)^{213}Ac	8490.8	15.	8489	4	−0.1	U			GSa		00He17	
	8489.3	5.			−0.1	2			GSa		02He29	*
^{217}Pa^{m}(α)^{213}Ac	10351	20	10349	5	−0.1	U					79Sc09	
	10330.8	50.			0.4	U			JAa		98Ik01	
	10346.1	15.			0.2	o			GSa		00He17	
	10349.1	5.				2			GSa		02He29	*
^{217}U(α)^{213}Th^{p}	8155.6	20.	8170	60	0.2	5					00Ma65	
	8175.0	14.3			−0.2	5			Jya		05Le42	*
^{217}Ac^{m}(IT)^{217}Ac	2012.3	0.4	2012.3	0.4	−0.0	1	100	74 ^{217}Ac^{m}			25Sa04	
*^{217}Th(α)^{213}Ra	E_α=9268(15), 8731(15), 8459(15) to ground state, 546.35, 822.7 levels										00He17	**
*^{217}Th(α)^{213}Ra	E_α=9261(5), 8725(5), 8455(5) to ground state, 546.35, 822.7 levels										02He29	**
*^{217}Pa(α)^{213}Ac	E_α=8337(5), 7873(5), 7728(5), 7710(5) to ground state,466.1,612.5,634.3 levels										02He29	**
*^{217}Pa^{m}(α)^{213}Ac	Average of 5 E_α's to known levels										02He29	**
*^{217}U(α)^{213}Th^{p}	Only one event. Not reported in later publication 07Le14										WgM115	**
^{218}Bi−u	14178	34	14188	29	0.3	o			GS3	1.0	08ChZI	
	14188	29				2			GS3	1.0	12Ch19	
^{218}Rn−u	5463	58	5600.1	2.1	2.4	U			GR1	1.0	19An10	
^{218}Po(α)^{214}Pb	6114.76	0.09	6114.75	0.09	−0.0	1	100	100 ^{214}Pb			71Gr17	Z
^{218}At(α)^{214}Bi	6874	3	6876.1	2.6	0.7	2			Orm		58Wa.A	*
	6882.1	5.1			−1.2	2					19Cu02	*
^{218}Rn(α)^{214}Po	7265.0	5.	7262.4	1.8	−0.5	–					56As38	Z
	7262.4	2.			0.0	–			Bka		82Bo04	Z
	7273.7	10.2			−1.1	U			Lza		25Gu12	
ave.	7262.8	1.9			−0.2	1	95	94 ^{218}Rn			average	
^{218}Fr(α)^{214}At	8014.0	2.	8013.7	1.4	−0.2	2			Bka		82Bo04	Z
	8013.3	2.			0.2	2					99Sh03	
^{218}Fr^{m}(α)^{214}At	8099.9	5.	8101	3	0.3	–					82Ew01	Z
	8100.9	5.			0.1	–					99Sh03	
	8106.6	11.2			−0.5	–			Lza		25Gu12	*
ave.	8101	3			0.1	1	89	73 ^{218}Fr^{m}			average	
^{218}Fr^{m}(α)^{214}At^{n}	7870.7	7.1	7869	5	−0.2	1	51	27 ^{218}Fr^{m}			99Sh03	
^{218}Ra(α)^{214}Rn	8549.1	8.	8540	3	−1.1	3					70To07	
	8541.0	10.			−0.1	3					70Va13	
	8577.7	51.			−0.7	U			GSa		18Mi11	
	8537.9	4.1			0.6	3			Jya		19Pa45	
^{218}Ac(α)^{214}Fr	9377.4	15.3	9390	60	0.2	5					70Bo13	
	9391.3	15.			−0.1	o			Lza		17Su18	
	9392.6	30.6			−0.1	5			GSa		19Mi08	
	9376.3	26.			0.2	o			Lza		19Ya04	
	9369	15			0.3	o			Lza		21Hu18	
	9391.5	6.1			−0.1	5			Lza		23Lu04	
^{218}Th(α)^{214}Ra	9861.3	20.4	9862	7	0.0	5					73Ha32	
	9846.1	10.			1.6	5					73No09	
	9851.0	81.5			0.1	U			GSa		15Kh09	
	9902.1	71.3			−0.6	U			Dba		18Br13	
	9878.6	10.2			−1.6	5			Lza		23Lu04	
^{218}Pa(α)^{214}Ac	9794.1	20.	9791	12	−0.1	2					79Sc09	*
	9815	10			−2.4	F			GSa		00He17	*
	9790.0	14.2			0.1	2			Lza		20Zh01	*
	9781.8	17.3			0.6	o			Lza		20Ma27	
^{218}Pa^{m}(α)^{214}Ac	9872.4	15.0				2			Lza		20Zh01	
^{218}U(α)^{214}Th	8786.6	25.	8774	7	−0.5	4			Dbb		92An04	
	8773.2	9.			0.1	4			Jya		07Le14	
	8780.4	35.7			−0.2	U			Lza		18Ya01	
	8773.3	14.4			0.1	4			Lza		21Zh22	
^{218}U^{m}(α)^{214}Th	10878.1	17.	10884	11	0.3	4			Jya		07Le14	
	10901.4	30.6			−0.6	o			Lza		15Ma37	
	10887.8	14.3			−0.3	4			Lza		22Zh45	*
*^{218}At(α)^{214}Bi	E_α=6696.3(3.0,Z) to (2)$^-$ level at 53.2282 keV										Ens092	**
*^{218}At(α)^{214}Bi	E_α=6694(5) to 3$^-$ level at 63 keV										19Cu02	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{218}$At(α)^{214}Bi	also E_α=6655(7) to 4^- level at 100 keV										19Cu02	**
$*^{218}$Fr^m(α)^{214}At	E_α=7819(11),7690(12),7606(15) to 145.1(0.5),278(1) and										Ens216	**
*	343(1) keV levels										Ens216	**
$*^{218}$Pa(α)^{214}Ac	E_α=9614(20)										00He17	**
$*^{218}$Pa(α)^{214}Ac	E_α=9544(10) to 91.8 level; F : from single α-spectra										00He17	**
$*^{218}$Pa(α)^{214}Ac	Also E_α=9524(16) to 91.8(0.8)										20Zh01	**
$*^{218}$U(α)^{214}Th	also reported in ^{22}Zh$_{45}$										21Zh22	*W
$*^{218}$U^m(α)^{214}Th	E_α=10690(14),10073(16) keV to the gs., 2^+ level at 623(1) keV										22Zh45	**
^{219}Po−u	13601	32	13614	17	0.4	o			GS3	1.0	08ChZI	
	13614	17				2			GS3	1.0	12Ch19	
^{219}At−^{133}Cs$_{1.647}$	166879.4	8.4	166881	3	0.2	1	14	14 ^{219}At	MA8	1.0	17Ma29	
^{219}Rn−^{133}Cs$_{1.647}$	165198.87	0.96	165199.2	0.6	0.3	1	39	39 ^{219}Rn	SH2	1.0	22Ka.A	*
^{219}Rn−^{85}Rb$_{2.576}$	236708.1	1.6	236708.2	0.6	0.1	1	14	14 ^{219}Rn	SH2	1.0	22Ka.A	
^{219}Po(α)^{215}Pb	5914.2	5.				3			ISa		15Fi07	
^{219}At(α)^{215}Bi	6390.9	50.	6342	5	−1.0	U					53Hy83	
	6344.0	5.			−0.4	1	90	87 ^{215}Bi	ISa		15Fi07	
^{219}Rn(α)^{215}Po	6946.21	0.3	6946.14	0.29	−0.2	1	93	54 ^{215}Po			71Gr17	Z
	6951.2	11.2			−0.5	U			Lza		25Gu12	
^{219}Fr(α)^{215}At	7448.7	2.0	7448.6	1.8	−0.1	5			Orm		68Ba73	Z
	7448.2	4.			0.1	5			Bka		82Bo04	Z
	7456.5	10.2			−0.8	U			Lza		25Gu12	
^{219}Ra(α)^{215}Rn	8139.0	20.	8138	3	−0.1	U			ORa		69Ha32	
	8128.7	10.			0.9	U					70Va13	
	8128.7	20.			0.5	U			Dbb		89An13	
	8138.0	3.				4					94Sh02	
	8160	31			−0.7	U			GSa		18Mi11	
	8128.8	10.2			0.9	U			GSt		18Sa45	*
	8139	15			−0.1	U			GRa		25To01	*
	8088.0	20.4			2.4	U			Lza		25Gu12	*
^{219}Ac(α)^{215}Fr	8826.5	10.				4					70Bo13	
	9330.6	40.7	8830	50	−7.8	B			GSa		19Mi08	*
^{219}Th(α)^{215}Ra	9514.1	20.	9513	6	−0.1	4					73Ha32	
	9503.9	50.9			0.2	4			GSa		15Kh09	
	9500.9	15.3			0.8	o			Lza		17Su18	
	9503.9	20.			0.4	o			Lza		19Zh54	
	9493.8	24.4			0.8	o			Lza		19Ya04	
	9512.0	16.3			0.1	o			Lza		20Su02	
	9513.1	6.1			−0.0	4			Lza		23Lu04	
	9483	44			0.7	U			Lza		24Ya27	
^{219}Pa(α)^{215}Ac	10084.6	50.	10130	70	0.6	3					87Fa.A	
	10161.8	38.			−0.5	3			Lza		17Su18	
^{219}U(α)^{215}Th	9860.4	40.	9950	12	2.2	U			Dbb		93An07	
	9956.2	18.			−0.4	3			Jya		07Le14	
	9945.0	15.3			0.3	3			Lza		19Zh54	
^{219}Np(α)^{215}Pa	9167.8	203.7	9183	13	0.1	U			GSa		15De22	*
	9207.5	40.7			−0.6	o			Lza		18Ya01	
	9183.1	13.2				3			Lza		26Zh40	
$*^{219}$Rn−^{133}Cs$_{1.647}$	From frequency ratio ^{219}Rn^{++}/(^{133}Cs)$^+$ =0.8239288978(36)										HWJ262	**
$*^{219}$Rn−^{133}Cs$_{1.647}$	From frequency ratio ^{219}Rn^{++}/(^{85}Rb)$^+$ =1.2896312572(93)										HWJ262	**
$*^{219}$Ra(α)^{215}Rn	also E_α=7660(20) to 316-keV level										18Sa45	**
$*^{219}$Ra(α)^{215}Rn	also E_α=7674(13) keV to the 316 keV level										25To01	**
$*^{219}$Ra(α)^{215}Rn	also E_α=7651(21) keV to the 316 keV level										Ens312	**
$*^{219}$Ac(α)^{215}Fr	E_α=8520(40) 9160(40)										19Mi08	**
$*^{219}$Np(α)^{215}Pa	E_α>9000 keV										15De22	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{220}Po−u	16420	32	16386	19	−1.1	o			GS3	1.0	08ChZI	
	16386	19				2			GS3	1.0	12Ch19	
^{220}At−u	15427	32	15433	15	0.2	o			GS3	1.0	08ChZI	
	15433	15				2			GS3	1.0	12Ch19	
^{220}Rn−^{208}Pb$_{1.058}$	36092.14	0.36	36091.52	0.27	−1.7	1	54	54 ^{220}Rn	CP2	1.0	23Ya.A	
^{220}Rn−^{133}Cs$_{1.654}$	167777	11	167770.21	0.27	−0.6	U			MA8	1.0	09Ne03	
	167769.30	0.40			2.3	1	44	44 ^{220}Rn	CP2	1.0	23Ya.A	
^{220}Ra−u	11389	344	11025	7	−1.1	U			GR1	1.0	19An10	
^{210}Fr−^{220}Fr$_{.159}$ ^{208}Fr$_{.841}$	−2930	60	−2925	16	0.0	U			P24	2.5	82Au01	
^{211}Fr−^{220}Fr$_{.240}$ ^{208}Fr$_{.761}$	−4850	70	−4865	15	−0.1	U			P24	2.5	82Au01	
^{212}Fr−^{220}Fr$_{.321}$ ^{208}Fr$_{.679}$	−5450	60	−5389	11	0.4	U			P24	2.5	82Au01	
^{212}Fr−^{220}Fr$_{.263}$ ^{209}Fr$_{.738}$	−3730	60	−3743	12	−0.1	U			P24	2.5	82Au01	
^{213}Fr−^{220}Fr$_{.352}$ ^{209}Fr$_{.649}$	−5170	50	−5140	9	0.2	U			P24	2.5	82Au01	
^{212}Fr−^{220}Fr$_{.193}$ ^{210}Fr$_{.808}$	−3160	60	−3029	14	0.9	U			P24	2.5	82Au01	
^{220}At(α)^{216}Bi^{m}	6053.3	6.				3					89Bu09	
^{220}Rn(α)^{216}Po	6404.75	0.10	6404.77	0.10	0.2	1	97	96 ^{216}Po			71Gr17	Z
	6409.8	10.2			−0.5	U			Lza		25Gu12	
^{220}Fr(α)^{216}At	6799.0	2.	6800.9	1.8	1.0	5			Orm		68Ba73	*
	6811.6	5.			−2.1	5					74Ho27	*
	6807.0	11.2			−0.5	5			Lza		25Gu12	
^{220}Ra(α)^{216}Rn	7593.3	10.	7595	4	0.2	4					61Ru06	
	7595.3	10.			0.0	4					70Va13	
	7598.4	20.4			−0.1	4			Dbb		90An19	
	7587.2	10.			0.8	4			GSa		00He17	
	7598.4	10.2			−0.3	4			GSt		18Sa45	
	7595.4	9.2			0.0	4			GRa		25To01	
	7603.5	11.2			−0.7	4			Lza		25Gu12	
^{220}Ac(α)^{216}Fr	8347.1	10.	8348	4	0.1	5					70Bo13	
	8348	5			−0.0	5					97Sh09	*
^{220}Th(α)^{216}Ra	8953.1	20.	8973	11	1.0	5					73Ha32	
	8981.6	13.2			−0.6	5			Jya		19Pa45	
^{220}Pa(α)^{216}Ac	9829.1	50.	9719	6	−1.6	B					87Fa.A	
	9330	60			5.0	B					87MiZO	
	9696.7	16.3			1.4	o			Lza		17Hu08	
	9718	20			0.0	o			Lza		19Zh54	
	9698.7	25.5			0.8	o			Lza		19Ya04	
	9725.2	30.6			−0.2	o			Lza		20Ma27	
	9725.2	30.6			−0.2	o			Lza		21Ma66	
	9719.1	6.1				3			Lza		23Lu04	
^{220}Pa^{m}(α)^{216}Ac	9730.3	20.4				3			Lza		18Hu13	
	9843.3	40.7	9730	20	−2.8	B			Lza		21Ma66	
^{220}Np(α)^{216}Pa	10226.3	18.3	10260	50	0.7	o			Lza		19Zh23	
	10265.0	16.3				3			Lza		26Zh40	
*^{220}Fr(α)^{216}At	E_α=6675.2, 6631.0, 6570.2(2,Z) to ground state, (2)$^-$ at 44.59, (0)$^-$ at 105.89										Ens075	**
*^{220}Fr(α)^{216}At	E_α=6687.5, 6642.5, 6583.5(2,Z) to ground state, (2)$^-$ at 44.59, (0)$^-$ at 105.89										Ens075	**
*^{220}Ac(α)^{216}Fr	E_α=7792, 7855 to levels at 409.3, 349.3 keV										Ens075	**
^{221}Po−u	21238	62	21228	21	−0.2	o			GS3	1.0	10Ch19	
	21228	21				2			GS3	1.0	12Ch19	
^{221}At−u	18028	32	18017	15	−0.3	o			GS3	1.0	08ChZI	
	18017	15				2			GS3	1.0	12Ch19	
^{221}Fr−^{133}Cs$_{1.662}$	171390.27	0.77	171390.3	0.7	0.0	1	85	85 ^{221}Fr	SH2	1.0	22Ka.A	*
^{221}Fr−^{226}Ra$_{.978}$	−10590	34	−10596.4	1.8	−0.2	U			MA3	1.0	92Bo28	
^{211}Fr−^{221}Fr$_{.159}$ ^{209}Fr$_{.841}$	−3080	60	−3070	15	0.1	U			P24	2.5	82Au01	
^{221}Rn(α)^{217}Po	6161.8	5.8	6163	3	0.2	3			Dba		77Vy02	*
	6163.5	5.4			−0.1	3			Orm		97Li23	*
	6163.5	5.4			−0.1	3			Orm		04Li28	*
^{221}Fr(α)^{217}At	6457.3	2.0	6457.8	0.9	0.2	–			Orm		62Wa28	*
	6458.5	2.0			−0.4	–			Orm		68Le07	*
ave.	6457.9	1.4			−0.1	1	45	31 ^{217}At			average	
^{221}Ra(α)^{217}Rn	6883.7	5.	6880.4	2.0	−0.7	5					61Ru06	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{221}Ra(α)^{217}Rn	6881.3	3.	6880.4	2.0	−0.3	5					95Ch74	*
	6878.3	3.			0.7	5					97Li12	*
	6884.8	10.2			−0.4	U			GSt		18Sa45	
	6876.7	12.2			0.3	U			Lza		25Gu12	
^{221}Ac(α)^{217}Fr	7786.2	10.	7780	50	−0.0	4					70Bo13	
	7782.1	5.			0.1	4			Lvn		88Hu08	
	7791.3	15.			−0.1	4			Dbb		92An.A	
	7811.4	30.6			−0.5	4			GSa		18Mi11	
	7783.3	3.1			0.5	o			Jya		22Pa09	*
	7784.8	4.0			0.0	4					23Re08	*
	7781.1	11.2			0.1	U			GRa		25To01	*
^{221}Th(α)^{217}Ra	8628.5	5.	8626.1	2.4	−0.5	5					70To07	Z
	8626.0	10.			0.0	5					70Va13	Z
	8626.4	10.			−0.0	5			Dbb		90An19	
	8614.2	10.			1.2	5			GSa		00He17	
	8596.9	66.			0.4	U					05Li17	
	8667.2	30.6			−1.3	U			GSa		18Mi11	
	8381.9	41.			6.0	B			GSa		19Mi08	
	8617.2	19.			0.5	U			Lza		19Ya04	*
	8625.9	3.1			0.1	5			Jya		20Pa44	*
	8637.7	15.3			−0.8	5			Lza		21Hu19	
^{221}Pa(α)^{217}Ac	9247.7	30.				2					89Mi17	
	8250	200	9250	60	4.8	B			GSa		19Mi08	
^{221}U(α)^{217}Th	9889.3	50.9				3			GSa		15Kh09	
*^{221}Fr−^{133}Cs$_{1.662}$	From frequency ratio $^{221}Fr^{++}/(^{133}Cs)^{+}$ =0.8314710318(29)										HWJ262	**
*^{221}Rn(α)^{217}Po	E_α=6035.3(3,Z) to a tentative (11/2$^+$) level at 15(5)keV above (9/2$^+$) ground state										04Li28	**
*^{221}Rn(α)^{217}Po	E_α=6037(2) to a tentative (11/2$^+$) level at 15(5) keV above (9/2$^+$) ground state										04Li28	**
*^{221}Rn(α)^{217}Po	E_α=6037(2) to a tentative (11/2$^+$) level at 15(5) keV above (9/2$^+$) ground state										04Li28	**
*^{221}Fr(α)^{217}At	E_α=6341.1(2,Z), 6125.1(3,Z) to ground state, 5/2$^-$ level at 218.12 keV										Ens039	**
*^{221}Fr(α)^{217}At	E_α=6341.3(2,Z), 6127.2(3,Z) to ground state, 5/2$^-$ level at 218.12 keV										Ens039	**
*^{221}Ra(α)^{217}Rn	E_α=6761.2 6668.2 6613.2 6591.2(5,Z) to ground state, levels at 88.9 149.18 174.3keV										Ens039	**
*^{221}Ra(α)^{217}Rn	E_α=6610(3,Z) to 149.2 level										97Li12	**
*^{221}Ra(α)^{217}Rn	E_α=6754, 6662, 6607, to ground state, 93.02, 149.2 level										97Li12	**
*^{221}Ac(α)^{217}Fr	E_α=7642(2),7440(3),7364(5) to ground state, levels at 209 keV, 276 keV										22Pa09	**
*^{221}Ac(α)^{217}Fr	E_α=7642(4),7547(4),7438(4),7418(4),7373(4) to levels at										23Re08	**
*^{221}Ac(α)^{217}Fr	0, 98.8, 209.4, 231.4 and 275.7 keV										23Re08	**
*^{221}Ac(α)^{217}Fr	also E_α=7432(13) keV to the 209 keV level										25To01	**
*^{221}Th(α)^{217}Ra	E_α=8247(3) keV to level at 226.7(0.2) keV; also E_α=8471(no unc.),										20Pa44	**
*^{221}Th(α)^{217}Ra	7951(8) to ground state and level at 539.8(0.3) keV										20Pa44	**
*^{221}Th(α)^{217}Ra	E_α=8409(16), 8249(19), 8164(15), 7731(17)										21Hu19	*W
^{222}Po−u	24133	72	24140	40	0.1	o			GS3	1.0	10Ch19	
	24140	43				2			GS3	1.0	12Ch19	
^{222}At−u	22459	32	22494	17	1.1	o			GS3	1.0	08ChZI	
	22494	17				2			GS3	1.0	12Ch19	
^{222}Fr−^{133}Cs$_{1.669}$	175383.3	8.0				2			MA8	1.0	14Kr09	
^{222}Fr−^{226}Ra$_{.982}$	−7410	25	−7367	8	1.7	U			MA3	1.0	92Bo28	
^{213}Fr−^{222}Fr$_{.240}$ ^{210}Fr$_{.761}$	−4810	60	−4941	11	−0.9	U			P24	2.5	82Au01	
^{213}Fr−^{222}Fr$_{.096}$ ^{212}Fr$_{.904}$	−1940	60	−1948	9	−0.1	U			P24	2.5	82Au01	
^{221}Fr−^{222}Fr$_{.498}$ ^{220}Fr$_{.502}$	−610	90	−643	4	−0.1	U			P34	2.5	86Au02	
^{222}Rn(α)^{218}Po	5590.39	0.3	5590.4	0.3	−0.0	1	100	100 ^{218}Po			71Gr17	Z
^{222}Ra(α)^{218}Rn	6680.0	5.	6678	4	−0.3	–					56As38	Z
	6681.6	10.2			−0.3	–			Lza		25Gu12	
ave.	6680	5			−0.4	1	73	67 ^{222}Ra			average	
^{222}Ac(α)^{218}Fr	7137.5	2.				3			Bka		82Bo04	Z

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$^{222}Ac^m(\alpha)^{218}Fr^m$	7128.6	22.4				2					72Es03	*
$^{222}Th(\alpha)^{218}Ra$	8127.7	10.	8130.5	2.8	0.3	U					70To07	
	8130.7	8.			−0.0	4					70Va13	
	8126.7	15.			0.3	U			Dbb		92An.A	
	8120.6	10.			1.0	U			GSa		00He17	
	8125.7	10.2			0.5	4			GSa		02CaZU	
	8116.4	48.			0.3	U					05Li17	
	8130.9	3.1			−0.1	4			Jya		16Pa28	*
	8157.3	30.6			−0.9	U			GSa		18Mi11	
$^{222}Pa(\alpha)^{218}Ac$	8788.6	41.	8790	60	0.1	6			GSa		19Mi08	*
	8794.8	15.3			−0.0	6			Lza		21Hu18	*
$^{222}Pa(\alpha)^{218}Ac^m$	8697	30	8736	13	1.3	7					70Bo13	
	8745.5	15.			−0.6	7			GSa		95Ho.C	
$^{222}U(\alpha)^{218}Th$	9481.1	50.9	9416	8	−1.3	U			GSa		15Kh09	
	9416.0	8.1				6			Lza		23Lu04	
$^{222}Np(\alpha)^{218}Pa$	10200.2	33.7				3			Lza		20Ma27	
$*^{222}Ac^m(\alpha)^{218}Fr^m$	E_α=7000(20) keV from 72Es03 to $^{218}Fr^m$ at 86(8) keV										FGK20c	**
$*^{222}Th(\alpha)^{218}Ra$	E_α=7986(3),7603(3),7205(4),7143(4) to ground state,389,793,853 level										16Pa28	**
$*^{222}Pa(\alpha)^{218}Ac$	E_α=8525(16),8437(16),8338(16),8250(16),8058(18)										21Hu18	**
$^{223}At-u$	25172	32	25151	15	−0.7	o			GS3	1.0	08ChZI	
	25151	15				2			GS3	1.0	12Ch19	
$^{223}Rn-^{133}Cs_{1.677}$	180453	11	180446	8	−0.6	1	58	58 ^{223}Rn	MA8	1.0	09Ne03	
$^{223}Rn-u$	21899	32	21889	8	−0.3	o			GS3	1.0	08ChZI	
	21880	13			0.7	1	42	42 ^{223}Rn	GS3	1.0	12Ch19	
$^{213}Fr-^{223}Fr_{.087}\ ^{212}Fr_{.913}$	−1900	60	−1944	9	−0.3	U			P24	2.5	82Au01	
$^{222}Fr-^{223}Fr_{.498}\ ^{221}Fr_{.502}$	790	100	560	7	−0.9	U			P34	2.5	86Au02	
$^{223}Fr(\alpha)^{219}At$	5431.6	80.	5561.4	2.8	1.6	U					55Ad10	
	5562	3			−0.2	1	84	83 ^{219}At			01Li44	
$^{223}Ra(\alpha)^{219}Rn$	5978.9	0.3	5978.98	0.21	0.3	–			Orm		62Wa18	*
	5979.1	0.3			−0.4	–			BIP		71Gr17	*
ave.	5979.00	0.21			−0.1	1	99	90 ^{223}Ra			average	
$^{223}Ac(\alpha)^{219}Fr$	6783.2	1.0				6			Orm		69Le.A	*
$^{223}Th(\alpha)^{219}Ra$	7602	23	7567	4	−1.5	U			ORa		69Ha32	*
	7589	14			−1.6	U					70Va13	*
	7570	25			−0.1	U					84Mi.A	
	7568	10			−0.1	5					87El02	*
	7567.4	10.			−0.0	5			Dbb		90An19	*
	7566.1	5.			0.2	5					92Li09	*
	7570.5	11.2			−0.3	5			GRa		25To01	*
$^{223}Pa(\alpha)^{219}Ac$	8345.0	10.	8340	60	−0.0	5					70Bo13	
	8339.9	10.2			0.1	o			Dbb		89An.A	
	8350.0	15.			−0.1	5			Dbb		90An19	
	8339.9	15.			0.1	5			GSa		95Ho.C	
	8321.6	5.			0.4	5			Jya		99Ho28	
	8370.3	41.			−0.4	5			GSa		19Mi08	*
$^{223}U(\alpha)^{219}Th$	8940.7	40.7	9163	15	3.4	F			Dbb		91An10	*
	9157.5	17.3			0.3	5			Lza		20Su02	*
	9170	39			−0.2	U			Lza		24Ya27	*
	9175	27			−0.5	5			Dbc		25Ku.A	*
$^{223}Np(\alpha)^{219}Pa$	9650.4	44.8				4			Lza		17Su18	*
$^{223}Fr(\beta^-)^{223}Ra$	1170	10	1149.0	0.8	−2.1	U					75We23	*
$*^{223}Ra(\alpha)^{219}Rn$	E_α=5747.0(0.4,Z), 5715.7(0.3,Z), 5606.7(0.3,Z) keV										62Wa18	**
*	to 11/2$^+$ level at 126.77, 7/2$^+$ at 158.64, 3/2$^+$ at 269.48 keV										Ens01a	**
$*^{223}Ra(\alpha)^{219}Rn$	E_α=5747.0(0.40,Z), 5716.23(0.29,Z), 5606.73(0.30,Z) keV										71Gr17	**
*	to 11/2$^+$ level at 126.77, 7/2$^+$ at 158.64, 3/2$^+$ at 269.48 keV										Ens01a	**
$*^{223}Ac(\alpha)^{219}Fr$	E_α=6661.6, 6646.7, 6563.7(1.0,Z) to ground state, 5/2$^+$ at 15.0, 7/2$^-$ at 98.58										Ens01a	**
$*^{223}Th(\alpha)^{219}Ra$	E_α=7330(20) to mixture of excited states at 138(10) keV										GAu	**
$*^{223}Th(\alpha)^{219}Ra$	E_α=7317(10) to mixture of excited states at 138(10) keV										GAu	**
$*^{223}Th(\alpha)^{219}Ra$	E_α=7324(10) to 113.8, 7285(10) 55% to 140.0, 26% to 152.0 level										92Li09	**
$*^{223}Th(\alpha)^{219}Ra$	E_α=7290(10) 55% to 140.0, 26% to 152.0 level										92Li09	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{223}$Th(α)^{219}Ra	E_α=7318(5), 7293(5), 7281(5) to 113.8, 140.0, 152.0 levels										92Li09	**
$*^{223}$Th(α)^{219}Ra	E_α=7297(11) to 140.0-keV level										25To01	**
$*^{223}$U(α)^{219}Th	Might be 8753 to 224 keV level										20Su02	**
$*^{223}$U(α)^{219}Th	Also E_α=8753 to 224 keV level										20Su02	**
$*^{223}$U(α)^{219}Th	Also E_α=8754(47) to 224 keV level										24Ya27	**
$*^{223}$U(α)^{219}Th	E_α=8790(27) to 224 keV level; also E_α=8952(38)										25Ku.A	**
$*^{223}$Np(α)^{219}Pa	Extracted as difference of sum energy 19453(23) and Ea(^{219}Pa)=9976(37)										17Su18	**
$*^{223}$Fr(β^-)^{223}Ra	E_{β^-}=1120(10) to 3/2$^-$ level at 50.128 keV										Ens01a	**
^{224}At−u	29744	63	29749	24	0.1	o			GS3	1.0	10Ch19	
	29749	24				2			GS3	1.0	12Ch19	
^{224}Rn−^{133}Cs$_{1.684}$	183304	16	183315	11	0.7	1	43	43 ^{224}Rn	MA8	1.0	09Ne03	
^{224}Rn−u	24073	32	24096	11	0.7	o			GS3	1.0	08ChZI	
	24104	14			−0.6	1	57	57 ^{224}Rn	GS3	1.0	12Ch19	
^{224}Fr−u	23399	32	23348	12	−1.6	o			GS3	1.0	08ChZI	
	23398	14			−3.6	B			GS3	1.0	12Ch19	
^{224}Fr−^{133}Cs$_{1.684}$	182567	12				2			MA8	1.0	14Kr09	
^{224}Ra−^{133}Cs$_{1.684}$	179430	30	179424.6	0.3	−0.2	U			MA8	1.0	14Bo26	
	179423.73	0.29			3.0	B			CP2	1.0	23Ya.A	
^{224}Ra−^{220}Rn$_{1.018}$	8613.78	0.26	8612.98	0.16	−3.1	B			CP2	1.0	23Ya.A	
^{223}Fr−^{224}Fr$_{.747}$ ^{220}Fr$_{.253}$	−620	70	−768	8	−0.8	U			P34	2.5	86Au02	
^{222}Fr−^{224}Fr$^x_{.496}$ ^{220}Fr$_{.505}$	10	70	−260#	50#	−1.5	U			P24	2.5	82Au01	
^{223}Fr−^{224}Fr$^x_{.747}$ ^{220}Fr$_{.253}$	−410	70	−840#	80#	−2.5	B			P24	2.5	82Au01	
^{223}Fr−^{224}Fr$_{.664}$ ^{221}Fr$_{.336}$	780	110	−519	7	−4.7	F			P34	2.5	86Au02	*
^{223}Fr−^{224}Fr$^x_{.664}$ ^{221}Fr$_{.336}$	−110	70	−590#	70#	−2.7	B			P24	2.5	82Au01	
^{224}Ra(α)^{220}Rn	5788.93	0.15	5788.96	0.15	0.2	1	99	98 ^{224}Ra			71Gr17	Z
^{224}Ac(α)^{220}Fr	6326.9	0.7				6			Orm		69Le.A	*
^{224}Th(α)^{220}Ra	7304.7	10.	7298	5	−0.6	5					61Ru06	
	7304.7	10.			−0.6	5					70Va13	
	7300.7	20.			−0.1	U			Dbb		89An13	
	7286.4	10.			1.2	5			GSa		00He17	
	7294.5	13.2			0.3	5			GRa		25To01	*
	7300.6	17.3			−0.1	5			Lza		25Gu12	
^{224}Pa(α)^{220}Ac	7695.2	10.	7694	4	−0.2	6					70Bo13	*
	7692.6	10.			0.1	F			Dbb		90An19	*
	7680	15			0.9	U			GSa		95Ho.C	
	7693.3	5.			0.1	6					96Li05	*
^{224}U(α)^{220}Th	8624.3	15.	8628	7	0.3	6			Dbb		91An10	
	8612.1	20.			0.8	6			ORa		92To02	
	8631.9	8.1			−0.5	6			Orm		14Lo10	*
^{224}Np(α)^{220}Pa^m	9303.5	20.4				4			Lza		18Hu13	
^{224}Fr(β^-)^{224}Ra	2830	50	2927	11	1.9	U					75We23	*
$*^{223}$Fr−^{224}Fr$_{.664}$ ^{221}Fr.	F : rejection based on line-shape analysis										86Au02	**
$*^{224}$Ac(α)^{220}Fr	E_α=6213.8, 6207.0, 6141.7, 6059.8(0.7,Z) keV										69Le.A	**
*	to ground state, 3$^+$ at 6.92, 3$^+$ at 72.99, 2$^-$ at 156.82 keV										Ens114	**
$*^{224}$Th(α)^{220}Ra	also E_α=6996(20) keV to the 178.4 keV level										25To01	**
$*^{224}$Pa(α)^{220}Ac	E_α=7490(10) to 5$^-$ level at 68.71 keV										Ens114	**
$*^{224}$Pa(α)^{220}Ac	F : intensities in contradiction with reference										96Li05	**
$*^{224}$Pa(α)^{220}Ac	E_α=7488(5), 7375(5) to (5$^-$) level at 68.71 keV and 184.21 level										Ens114	**
$*^{224}$U(α)^{220}Th	E_α=8479(8), 8095(11) to ground state, 386.7(1.8) 2$^+$ level										14Lo10	**
$*^{224}$Fr(β^-)^{224}Ra	E_{β^-}=1780(50) to 1$^-$ level at 1053.041 keV, and other E_{β^-}										Ens15c	**
^{225}Rn−^{133}Cs$_{1.692}$	188484	23	188461	12	−1.0	1	27	27 ^{225}Rn	MA8	1.0	09Ne03	
^{225}Rn−u	28498	32	28486	12	−0.4	o			GS3	1.0	08ChZI	
	28477	14			0.6	1	73	73 ^{225}Rn	GS3	1.0	12Ch19	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{225}Fr−u		25574	14	25573	13	−0.1	1	84	84 ^{225}Fr	GS3	1.0	12Ch19	
^{221}Fr−^{225}Fr$_{.655}$ ^{213}Fr$_{.346}$		−1110	60	−1097	8	0.1	U			P24	2.5	82Au01	
^{224}Fr^{x}−^{225}Fr$_{.747}$ ^{221}Fr$_{.253}$		50	80	700#	100#	3.2	B			P24	2.5	82Au01	
^{224}Fr^{x}−^{225}Fr$_{.498}$ ^{223}Fr$_{.502}$		190	80	760#	100#	2.8	B			P24	2.5	82Au01	
^{225}Ra(α)^{221}Rn		5096.7	5.1				2					00Li37	
^{225}Ac(α)^{221}Fr		5936.1	2.	5935.3	1.4	−0.4	–			Orm		67Ba51	Z
		5934.5	2.			0.4	–					67Dz02	Z
	ave.	5935.3	1.4			0.0	1	94	93 ^{225}Ac			average	
^{225}Th(α)^{221}Ra		6920.7	3.	6921.4	2.1	0.2	6					61Ru06	*
		6922.1	3.			−0.2	6					87Li.A	*
^{225}Pa(α)^{221}Ac		7381.5	20.	7390.6	1.7	0.2	U			ORa		68Ha14	
		7376.4	10.			0.3	F					70Bo13	*
		7392.7	5.1			−0.0	5			Lvn		88Hu08	
		7392.7	5.1			−0.0	5			Lvn		87De.A	
		7383.5	19.			0.1	5					00Sa52	
		7432.1	30.5			−0.7	5			GSa		18Mi11	
		7395.7	3.1			−1.7	5			Jya		22Pa09	*
		7388.3	2.0			1.1	5			Jya		23Re08	
^{225}U(α)^{221}Th		8012.7	20.	8005	6	−0.4	o			Dbb		89An13	
		8022.9	20.			−0.9	o			GSa		89He13	
		8021.9	15.			−1.1	6			ORa		92To02	
		8012.7	20.4			−0.4	6			Dbb		94Ye08	
		8010	10			−0.5	6			GSa		00He17	*
		7992.4	30.5			0.4	U			GSa		18Mi11	
		7992.3	10.2			1.2	6			GSa		19Mi08	
		7988.3	24.4			0.7	6			Dbc		25Ku.A	
^{225}Np(α)^{221}Pa		8786.5	20.	8820	70	0.6	3			Dbb		94Ye08	
		8959.5	102.			−1.2	3			GSa		19Mi08	
^{225}Fr(β^-)^{225}Ra		1820	30	1827	12	0.2	1	16	16 ^{225}Fr			75We23	*
^{225}Ra(β^-)^{225}Ac		360	10	358.2	2.3	−0.2	U					55Ma.A	*
		360	30			−0.1	U					55Pe24	*
*^{225}Th(α)^{221}Ra	E_α=6800.2, 6746.2, 6503.2, 6480.2, 6443.2(3,Z) keV											61Ru06	**
*	to ground state, 7/2$^+$ 53.14, 7/2$^+$ 299.16, 3/2$^+$ 321.39, 5/2$^+$ 359.02 levels											Ens075	**
*^{225}Th(α)^{221}Ra	E_α=6799.3, 6745.3, 6504.3, 6483.3, 6447.3(3,Z) keV											87Li.A	**
*	to ground state, 7/2$^+$ 53.14, 7/2$^+$ 299.16, 3/2$^+$ 321.39, 5/2$^+$ 359.02 levels											Ens075	**
*^{225}Pa(α)^{221}Ac	Also E_α=7182(8),7135(8),7112(8),7084(8) to levels at											22Pa09	**
*	88, 125, 152, 180 keV.											22Pa09	**
*^{225}Pa(α)^{221}Ac	E_α=7258(2),7229(2),7168(2),7078(2) to levels at											23Re08	**
*	0, 27.4, 91.5, 180.0 keV											23Re08	**
*^{225}U(α)^{221}Th	E_α=7868(15), 7621(15) to ground state, 250.9 level											00He17	**
*^{225}Fr(β^-)^{225}Ra	E_{β^-}=1640(10). 28% to 3/2$^-$ level at 225.2											Ens095	**
*^{225}Ra(β^-)^{225}Ac	E_{β^-}=320(10) 320(30) respectively, to 3/2$^+$ level at 40.09 keV											Ens095	**
^{226}Rn−^{133}Cs$_{1.699}$		191490	17	191499	11	0.5	1	44	44 ^{226}Rn	MA8	1.0	09Ne03	
^{226}Rn−u		30864	32	30861	11	−0.1	o			GS3	1.0	08ChZI	
		30868	15			−0.4	1	56	56 ^{226}Rn	GS3	1.0	12Ch19	
^{226}Fr−u		29565	32	29545	7	−0.6	o			GS3	1.0	08ChZI	
		29566	13			−1.7	1	26	26 ^{226}Fr	GS3	1.0	12Ch19	
^{226}Fr−^{133}Cs$_{1.699}$		190173.9	7.8	190182	7	1.0	1	74	74 ^{226}Fr	MA8	1.0	14Kr09	
^{133}Cs−^{226}Ra$_{.588}$		−109487	9	−109487.3	1.0	−0.0	U			MA3	1.0	92Bo28	
		−109499	13			0.9	U			MA4	1.0	99Am05	
^{223}Fr−^{226}Fr$_{.493}$ ^{220}Fr$_{.507}$		−800	80	−1004	4	−1.0	U			P24	2.5	82Au01	
^{225}Fr−^{226}Fr$_{.796}$ ^{221}Fr$_{.204}$		−570	100	−794	13	−0.9	U			P24	2.5	82Au01	
^{225}Fr−^{226}Fr$_{.498}$ ^{224}Fr$^{x}_{.502}$		−260	90	−850#	50#	−2.6	B			P24	2.5	82Au01	
^{226}Ra(α)^{222}Rn		4870.70	0.25	4870.70	0.25	−0.0	1	100	100 ^{222}Rn			71Gr17	Z

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{226}Ac(α)^{222}Fr	5496.1	5.	5505	8	0.2	U			Dba		75Va.A	Z
^{226}Th(α)^{222}Ra	6448.5	3.0	6452.5	1.0	1.3	U					56As38	*
	6454.8	3.6			−0.6	U			Dba		75Va.A	*
	6452.6	1.0			−0.1	1	99	66 ^{226}Th	Gea		12Ma30	
^{226}Pa(α)^{222}Ac	6986.9	10.				4					64Mc21	
^{226}U(α)^{222}Th	7747.4	30.	7701	4	−1.6	U					73Vi10	*
	7706.6	15.			−0.4	5			Dbb		90An22	
	7701.6	5.			−0.1	5			Jya		99Gr28	
	7691.4	10.			0.9	o			GSa		00He17	
	7696.5	10.			0.4	o			GSa		01Ca.B	
	7696.5	10.			0.4	5			GSa		02CaZU	
	7696.4	20.4			0.2	5			RIa		16Ka13	
	7706.6	30.5			−0.2	U			GSa		18Mi11	
^{226}Np(α)^{222}Pa	8189.2	20.4	8240	60	1.0	7			GSa		90Ni05	
	8205.5	20.			0.7	7			Dbb		94Ye08	
	8327.6	20.4			−1.6	7			JSa		19Mi08	
^{226}Pu(α)^{222}U	8912	24				7			Dbc		25Ku.A	
^{226}Ra(p,t)^{224}Ra	−2816	15	−2816.0	1.6	−0.0	U			ANL		74Fr01	
^{226}Ra(d,t)^{225}Ra	−146	10	−141.4	2.3	0.5	U					83Ny01	
^{226}Fr(β^-)^{226}Ra	3804	330	3854	6	0.2	U					75We23	*
	3704	100			1.5	U					87Ve.A	*
^{226}Ac(β^-)^{226}Th	1115	7	1111	4	−0.6	–					68Va17	*
ave.	1115	6			−0.6	1	47	34 ^{226}Th			average	
*^{226}Th(α)^{222}Ra	E_α=6334.6(3,Z), 6224.6(3,Z) to ground state, 2$^+$ level at 111.12 keV										Ens11b	**
*^{226}Th(α)^{222}Ra	E_α=6337.1(1.0,Z), 6233.6(1.0,Z) to ground state, 2$^+$ level at 111.12 keV										Ens11b	**
*^{226}U(α)^{222}Th	E_α=7430(30) to 2$^+$ level at 183.3(0.3) keV										94Ye08	**
*^{226}Fr(β^-)^{226}Ra	E_{β^-}=3550(330) 3450(100) respectively, to 1$^-$ level at 253.73 keV										Ens964	**
*^{226}Ac(β^-)^{226}Th	E_{β^-}=885(7) to 1$^-$ level at 230.37 keV										Ens964	**
^{227}Rn−^{133}Cs$_{1.707}$	196686	19	196698	15	0.6	1	63	63 ^{227}Rn	MA8	1.0	09Ne03	
^{227}Rn−u	35288	33	35304	15	0.5	o			GS3	1.0	08ChZI	
	35325	25			−0.8	1	37	37 ^{227}Rn	GS3	1.0	12Ch19	
^{227}Fr−u	31868	32	31865	6	−0.1	o			GS3	1.0	08ChZI	
	31869	14			−0.3	1	20	20 ^{227}Fr	GS3	1.0	12Ch19	
^{227}Fr−^{133}Cs$_{1.707}$	193258.0	7.1	193259	6	0.1	1	80	80 ^{227}Fr	MA8	1.0	14Kr09	
^{225}Fr−^{227}Fr$_{.708}$ ^{220}Fr$_{.292}$	−410	130	−546	13	−0.4	U			P24	2.5	82Au01	
^{224}Fr^x−^{227}Fr$_{.493}$ ^{221}Fr$_{.507}$	−220	80	490#	100#	3.5	B			P24	2.5	82Au01	
^{227}Ac(α)^{223}Fr	5043.0	2.0	5042.27	0.14	−0.4	U					66Ba19	Z
	5042.27	0.14			−0.0	1	100	99 ^{223}Fr			86Ry04	Z
^{227}Th(α)^{223}Ra	6146.60	0.10	6146.60	0.10	−0.0	1	100	90 ^{227}Th	BIP		71Gr17	*
^{227}Pa(α)^{223}Ac	6581.5	3.	6580.4	2.1	−0.4	7					63Su.A	*
	6579.3	3.			0.4	7					90Sh15	*
^{227}U(α)^{223}Th	7230	30	7235	3	0.2	U			ORa		69Ha32	*
	7206	16			1.8	U					91Ho05	
	7234.7	3.1				6			GSa		15Ka24	
^{227}Np(α)^{223}Pa	7818.0	10.	7819	12	0.0	o			Dbb		90An19	
	7815.0	20.			0.2	6			GSa		90Ni05	
	7818.0	20.			0.1	6			Dbb		94Ye08	
	7825.1	21.4			−0.3	6			Dbc		25Ku.A	*
^{227}Pu(α)^{223}U	8338.2	28.5	8319	19	−0.7	6			Lza		24Ya27	
	8302.6	26.5			0.6	6			Dbc		25Ku.A	
^{226}Ra(n,γ)^{227}Ra	4561.43	0.27				2			ILn		81Vo03	Z
^{227}Fr(β^-)^{227}Ra	2476	100	2506	6	0.3	U					75We23	*
^{227}Ra(β^-)^{227}Ac	1345	20	1326.9	1.8	−0.9	U					53Bu63	*
	1335	15			−0.5	U					71Lo15	*
^{227}Ac(β^-)^{227}Th	45.5	1.0	44.7	0.8	−0.8	–					55Be20	
	43.5	1.5			0.8	–					59No41	
ave.	44.9	0.8			−0.3	1	82	72 ^{227}Ac			average	
*^{227}Th(α)^{223}Ra	E_α=6038.01(0.15,Z), 5977.72(0.10,Z), 5756.89(0.15,Z) keV										71Gr17	**
*	to ground state, 7/2$^+$ at 61.424, 1/2$^+$ at 286.182 keV										Ens01a	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
$*^{227}$Pa(α)^{223}Ac	E_α=6465.8(3,Z), 6423.8(3,Z), 6415.8(3,Z), 6401.7(3,Z), 6356.7(3,Z)											63Su.A	**
*	to ground state, 7/2$^-$ at 42.4, 5/2$^-$ at 50.7, 5/2$^+$ at 64.62, 7/2$^+$ at 110.06											Ens01a	**
$*^{227}$Pa(α)^{223}Ac	E_α=6463, 6421, 6355 keV (all errors 3 keV, estimated by evaluator)											90Sh15	**
*	to ground state, 7/2$^-$ at 42.4, 5/2$^-$ at 50.7, 7/2$^+$ at 110.06 keV											Ens01a	**
$*^{227}$U(α)^{223}Th	E_α=6860(30) to 3/2$^+$ level at 247(1) keV											Ens01a	**
$*^{227}$Np(α)^{223}Pa	also E_α=7626(24)											25Ku.A	**
$*^{227}$Fr(β^-)^{227}Ra	E_{β^-}=1800(100) to 1/2$^-$ level at 675.863 keV											Ens162	**
$*^{227}$Ra(β^-)^{227}Ac	E_{β^-}=1310(20) 1300(15) respectively, to 3/2$^+$ level at 27.369 and 5/2$^+$ at 46.354											Ens162	**
^{228}Rn−^{133}Cs$_{1.714}$	199897	24	199891	19	−0.3	1	63	63	^{228}Rn	MA8	1.0	09Ne03	
^{228}Rn−u	37856	33	37835	19	−0.6	o				GS3	1.0	08ChZI	
	37825	31			0.3	1	37	37	^{228}Rn	GS3	1.0	12Ch19	
^{228}Fr−u	35833	34	35839	7	0.2	o				MA8	1.0	11Kr.A	
	35852	32			−0.4	o				GS3	1.0	08ChZI	
	35821	16			1.2	1	20	20	^{228}Fr	GS3	1.0	12Ch19	
^{228}Fr−^{133}Cs$_{1.714}$	197899.5	8.1	197895	7	−0.6	1	80	80	^{228}Fr	MA8	1.0	14Kr09	
^{224}Fr^x−^{228}Fr$_{.491}$ ^{220}Fr$_{.509}$	−540	320	−390#	100#	0.2	U				P24	2.5	82Au01	
^{228}Th(α)^{224}Ra	5520.17	0.22	5520.22	0.22	0.2	1	99	96	^{228}Th			71Gr17	Z
^{228}Pa(α)^{224}Ac	6266.7	3.	6264.5	1.5	−0.7	7						58Hi.A	*
	6264.7	3.			−0.1	7						93Sh07	*
	6263.5	2.			0.5	7						94Ah03	*
^{228}U(α)^{224}Th	6803.6	10.	6800	9	−0.4	6						61Ru06	
	6774.1	25.4			1.0	6				GSa		18Mi11	
^{228}Np(α)^{224}Pa	7308.5	36.	7540#	100#	3.7	D				JAa		03Ni10	*
^{228}Pu(α)^{224}U	7949.7	20.	7940	18	−0.5	7				Dbb		94An02	
	7911.0	35.			0.8	7				JAa		03Ni10	
^{228}Ra(β^-)^{228}Ac	46.7	2.	45.5	0.6	−0.6	3						61To10	*
	45.5	0.9			0.0	3						72He.A	*
	45.3	1.0			0.2	3						95So11	*
^{228}Ac(β^-)^{228}Th	2240	20	2129.4	1.8	−5.5	B						53Ky19	*
	2158	20			−1.4	U						57Bj56	*
^{228}Pa(ε)^{228}Th	2109	15	2152	4	2.8	B						73Ku09	*
$*^{228}$Pa(α)^{224}Ac	E_α=6119.2(3,Z), 6106.2(3,Z), 6079.2(3,Z) to 37.2, 51.9, 78.4 levels											93Sh07	**
$*^{228}$Pa(α)^{224}Ac	E_α=6118(3) to 37.2 level											93Sh07	**
$*^{228}$Pa(α)^{224}Ac	E_α=6117(2) to 37.1 level											94Ah03	**
$*^{228}$Np(α)^{224}Pa	Trends in Mass Surface TMS suggest ^{228}Np 230 keV less bound											GAu	**
$*^{228}$Ra(β^-)^{228}Ac	E_{β^-}=40(2) 39(1) respectively, to 1$^-$ level at 6.28 keV, and other E_{β^-}											Ens143	**
$*^{228}$Ra(β^-)^{228}Ac	E_{β^-}=39.0(1.0) to 1$^+$ level at 6.28 keV											Ens143	**
$*^{228}$Ac(β^-)^{228}Th	E_{β^-}=2180(20) to 2$^+$ level at 57.773 keV, and other E_{β^-}											Ens143	**
$*^{228}$Ac(β^-)^{228}Th	E_{β^-}=2100(20), 1760, 1180 to 2$^+$ at 57.773, 3$^-$ at 396.094, 2$^+$ at 968.984											Ens143	**
$*^{228}$Pa(ε)^{228}Th	pK=0.33(0.08) to 3$^+$ level at 1944.904 keV, recalculated											Ens143	**
^{229}Rn−^{133}Cs$_{1.722}$	205069	14				2				MA8	1.0	09Ne03	
^{229}Fr−^{133}Cs$_{1.722}$	201262	40	201103	5	−4.0	B				MA8	1.0	08We02	*
	201104.5	6.4			−0.2	1	70	70	^{229}Fr	MA8	1.0	14Kr09	
^{229}Fr−u	38343	32	38292	5	−1.6	o				GS3	1.0	08ChZI	
	38298	15			−0.4	1	13	13	^{229}Fr	GS3	1.0	12Ch19	
^{229}Fr−^{238}U$_{.962}$	−10576	13	−10566	5	0.8	1	17	17	^{229}Fr	MA8	1.0	14Kr09	
^{229}Ra−^{133}Cs$_{1.722}$	197782	21	197768	17	−0.6	2				MA8	1.0	08We02	
	197746	27			0.8	2				MA8	1.0	05He26	
^{229}Ac−u	32947	13				2				GS3	1.0	12Ch19	
^{229}Th(α)^{225}Ra	5167.4	1.2	5167.6	1.0	0.2	–				Kum		71Bb10	*
	5168.2	2.			−0.3	–						87He28	Z
ave.	5167.6	1.0			−0.0	1	100	100	^{225}Ra			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{229}Pa(α)^{225}Ac	5835.6	5.	5835.8	2.9	0.0	1	33	27 ^{229}Pa			63Su.A	*
^{229}U(α)^{225}Th	6475.5	3.				7					61Ru06	Z
^{229}Np(α)^{225}Pa	7012.7	20.	7020	60	0.1	6			ORa		68Ha14	
	7015.8	23.			0.1	6					00Sa52	
	7032.8	30.5			−0.2	6			GSa		18Mi11	
^{229}Pu(α)^{225}U	7592.9	30.	7600	60	0.1	7			Dbb		94An02	
	7598.0	10.			0.0	o			GSa		01Ca.B	
	7589.8	20.			0.2	7			GSa		10Kh06	
	7613	31			−0.3	7			GSa		18Mi11	
^{229}Am(α)^{225}Np	8137.4	20.4				4			GSa		15De22	*
^{229}Ra(β^-)^{229}Ac	1760	40	1872	20	2.8	B					75We23	
^{229}Ac(β^-)^{229}Th	1140	150	1104	12	−0.2	U					73Ch24	
	1090	50			0.3	U					75We23	
*^{229}Fr−^{133}Cs$_{1.722}$	Could be influenced by ^{229}Rn contaminant										08We02	**
*^{229}Th(α)^{225}Ra	E_α=4978.3(1.2,Z), 4967.3(1.2,Z), 4845.1(1.2,Z) keV										71Gr17	**
*	to 100.60, 111.60, 236.25 levels										71Gr17	**
*^{229}Th(α)^{225}Ra	E_α=4979.3(2,Z), 4968.3(2,Z), 4845.1(2,Z) keV										87He28	**
*	to 9/2$^+$ level at 100.50, 7/2$^+$ at 111.60, 5/2$^+$ at 236.25 keV										Ens095	**
*	calibrated with 71BaB2 value for 4845 level										AHW	**
*^{229}Pa(α)^{225}Ac	E_α=5670.2, 5630.2, 5615.2, 5580.2, 5536.2 (all 3,Z) keV to										63Su.A	**
*	5/2$^+$ 64.70, 7/2$^+$ 105.06, 5/2$^-$ 120.80, 5/2$^+$ 155.65, 7/2$^+$ 199.85										Ens095	**
*^{229}Am(α)^{225}Np	E_α=7990(20) and E_α=8000(20)										15De22	**
^{230}Fr−^{133}Cs$_{1.729}$	205878	32	205864	7	−0.4	o			MA8	1.0	05He26	
	205860.1	7.5			0.6	1	88	88 ^{230}Fr	MA8	1.0	14Kr09	
^{230}Fr−u	42401	34	42391	7	−0.3	o			GS3	1.0	08ChZI	
	42421	20			−1.5	1	12	12 ^{230}Fr	GS3	1.0	12Ch19	
^{230}Ra−^{133}Cs$_{1.729}$	200530	13	200528	11	−0.1	2			MA8	1.0	08We02	
	200524	21			0.2	2			MA8	1.0	05He26	
^{230}Ac−u	36328	32	36327	17	−0.0	o			GS3	1.0	08ChZI	
	36327	17				2			GS3	1.0	12Ch19	
^{230}Ra−^{226}Ra$_{1.018}$	11225	35	11191	11	−1.0	U			MA3	1.0	92Bo28	
^{230}Th(α)^{226}Ra	4770.1	1.5	4770.0	1.5	−0.1	1	98	98 ^{226}Ra	Orm		66Ba14	Z
^{230}Pa(α)^{226}Ac	5439.5	0.7	5439.4	0.7	−0.1	1	99	87 ^{226}Ac	Orm		66Ba14	Z
^{230}U(α)^{226}Th	5992.8	0.7	5992.5	0.5	−0.5	2			Orm		66Ba14	Z
	5992.1	0.7			0.5	2			Gea		12Ma30	
^{230}Np(α)^{226}Pa	6778.1	20.				5			ORa		68Ha14	
^{230}Pu(α)^{226}U	7175.0	15.	7178	9	0.2	6			Dbb		90An22	
	7180.1	17.			−0.1	6			Jya		99Gr28	
	7182.2	10.			−0.4	o			GSa		01Ca.B	
	7185.1	20.4			−0.3	6			RIa		16Ka13	
	7174.0	25.4			0.2	6			GSa		18Mi11	
^{230}Th(p,t)^{228}Th	−3550	15	−3566.2	0.5	−1.1	U			ANL		74Fr01	
^{230}Th(p,t)^{228}Th−^{232}Th()^{230}Th	−493.5	1.0	−492.2	0.5	1.3	o					91Gr13	
	−492.5	0.5			0.5	1	93	69 ^{230}Th			94Le22	
^{230}Th(p,t)^{228}Th−^{184}W()^{182}W	1564.0	1.6	1553.9	0.6	−6.3	B					09Le03	
	1564.0	1.8			−5.6	C					09Le.A	
^{230}Th(d,t)^{229}Th	−541	6	−539.0	1.5	0.3	–					90Bu17	
	−525	6			−2.3	–			ANL		67Er02	*
ave.	−533	4			−1.4	1	12	11 ^{229}Th			average	
^{230}Ra(β^-)^{230}Ac	710	300	678	19	−0.1	U					80Gi04	*
^{230}Ac(β^-)^{230}Th	2700	100	2977	16	2.8	U					80Gi04	*
^{230}Pa(ε)^{230}Th	1310.3	3.	1311.2	2.8	0.3	1	88	88 ^{230}Pa			70Lo02	*
^{230}Pa(β^-)^{230}U	561	15	558	4	−0.2	R					70Lo02	
*^{230}Th(d,t)^{229}Th	Q =−525(6) to ^{229}Th^m at 0.0035(0.0010) keV										94He08	**
*^{230}Ra(β^-)^{230}Ac	E_{β^-}=500(200) to level at 211.78 keV										Ens129	**
*^{230}Ac(β^-)^{230}Th	E_{β^-}=1400(100) to 0$^+$ level at 1297.14 keV										Ens129	**
*^{230}Pa(ε)^{230}Th	pK=0.42(0.01) to 3$^-$ level at 1127.789, recalculated										Ens129	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{231}Fr−u	45191	39	45175	8	−0.4	o				GS3	1.0	08ChZI	
	45158	27			0.6	U				GS3	1.0	12Ch19	
^{231}Fr−^{133}Cs$_{1.737}$	209405.3	8.3				2				MA8	1.0	14Kr09	
^{231}Ra−^{133}Cs$_{1.737}$	205267	21	205257	12	−0.5	1	34	34	^{231}Ra	MA8	1.0	05He26	
^{231}Ra−u	41052	32	41027	12	−0.8	o				GS3	1.0	08ChZI	
	41022	15			0.3	1	66	66	^{231}Ra	GS3	1.0	12Ch19	
^{231}Ac−u	38404	32	38393	14	−0.3	o				GS3	1.0	08ChZI	
	38393	14				2				GS3	1.0	12Ch19	
^{231}Pa(α)^{227}Ac	5150.2	1.5	5149.8	0.8	−0.3	o				Orm		66Ba14	
	5146.9	1.0			2.9	B				Kum		68Ba25	*
	5150.7	1.5			−0.6	–				Orm		69Le.A	*
	5149.8	1.0			0.0	–				Kum		76Ba68	*
ave.	5150.1	0.8			−0.3	1	83	56	^{231}Pa			average	
^{231}U(α)^{227}Th	5551.3	50.	5576.3	1.7	0.4	U						53Cr.A	
	5576.9	3.			−0.2	2						94Li12	*
	5576	2			0.1	2						97Mu08	
^{231}Np(α)^{227}Pa	6368.4	8.				8						73Ja06	
^{231}Pu(α)^{227}U	6838.6	20.				7						99La14	
^{231}Am(α)^{227}Np^{p}	7234.5	18.3				8				Lza		26Wa.1	
^{231}Pa(p,t)^{229}Pa	−4133	3	−4132.9	2.6	0.0	1	76	73	^{229}Pa	Mun		91Gr13	*
^{230}Th(n,γ)^{231}Th	5118.00	0.20	5117.98	0.20	−0.1	1	97	85	^{231}Th	ILn		87Wh01	Z
^{230}Th(d,p)^{231}Th	2907	7	2893.41	0.20	−1.9	U				ANL		67Er02	
^{231}Ac(β^-)^{231}Th	2100	100	1948	13	−1.5	U						60Ta19	
^{231}Th(β^-)^{231}Pa	389.2	2.	390.6	1.0	0.7	1	27	22	^{231}Pa			75Ho14	*
*^{231}Pa(α)^{227}Ac	E_α=5057.6(1.0,Z), 4985.9(1.0,Z), 4950.4(1.0,Z) to ground state, 7/2$^-$ at											Ens129	**
*	74.149 keV, and 9/2$^+$ at 109.992 keV											Ens129	**
*^{231}Pa(α)^{227}Ac	E_α=5015.9(1.5,Z),4735.9(1.5,Z) to 5/2$^+$ at 46.354, 3/2$^-$ at 330.040 keV											Ens129	**
*^{231}Pa(α)^{227}Ac	E_α=4736.2(1.0,Z) to 330.040 level											Ens129	**
*^{231}U(α)^{227}Th	E_α=5471(3), 5456(3), 5404(3) to 9.3, 24.4, 77.7 levels											94Li12	**
*^{231}Pa(p,t)^{229}Pa	Q =–4145(3) to 11.6 level											98Le15	**
*^{231}Th(β^-)^{231}Pa	E_{β^-}=305(2) to 5/2$^+$ level at 84.2148 keV											Ens136	**
^{232}Fr−^{133}Cs$_{1.744}$	214353	15				2				MA8	1.0	14Kr09	
^{232}Ra−^{133}Cs$_{1.744}$	208368	13	208367	10	−0.1	1	57	57	^{232}Ra	MA8	1.0	05He26	
^{232}Ra−u	43518	32	43475	10	−1.3	o				GS3	1.0	08ChZI	
	43474	15			0.1	1	43	43	^{232}Ra	GS3	1.0	12Ch19	
^{232}Ac−u	42052	32	42034	14	−0.6	o				GS3	1.0	08ChZI	
	42034	14				2				GS3	1.0	12Ch19	
C_{18} H_{16}−^{232}Th	87142.4	2.	87145.4	1.0	0.6	U				M20	2.5	73Br06	
C_{24} H_{16}−^{232}Th ^{37}Cl ^{35}Cl	152393.4	1.8	152390.2	1.0	−0.7	1	5	5	^{232}Th	M20	2.5	73Br06	
^{232}Th(α)^{228}Ra	4082.5	5.	4081.6	1.4	−0.2	U						57Ha08	Z
	4084.6	5.			−0.6	U						61Ko11	Z
	4083.5	5.			−0.4	U						62Ko12	Z
	4081.6	1.4				2						89Sa01	*
^{232}U(α)^{228}Th	5413.63	0.09	5413.63	0.09	−0.0	1	100	100	^{232}U	BIP		72Go33	*
^{232}Pu(α)^{228}U	6716.0	10.				7						73Ja06	
^{232}Th(p,t)^{230}Th	−3070	15	−3074.0	0.6	−0.3	U				ANL		74Fr01	
^{232}Th(p,t)^{230}Th−^{184}W()^{182}W	2056.4	1.6	2046.2	0.6	−6.4	B						09Le03	
	2056.5	1.8			−5.7	B						09Le.A	
^{232}Th(d,t)^{231}Th	−174	6	−180.6	0.6	−1.1	U				ANL		67Er02	
	−187	10			0.6	U				MIT		72Gr19	
^{232}Ac(β^-)^{232}Th	3700	100	3706	13	0.1	U						90Be.B	
^{232}Pa(β^-)^{232}U	1344	20	1337	7	−0.3	2						63Bj01	*
	1336	8			0.1	2						71Ka42	*
*^{232}Th(α)^{228}Ra	E_α=4012.3(1.4), 3947.2(2.0) to ground state, 2$^+$ level at 63.823 keV											Ens143	**
*^{232}U(α)^{228}Th	E_α=5320.12(0.14,Z), 5263.36(0.09,Z) to ground state, 2$^+$ level at 57.773 level											Ens143	**
*^{232}Pa(β^-)^{232}U	E_{β^-}=1295(20) to 2$^+$ level at 47.573 keV, and other E_{β^-}											Ens06a	**
*^{232}Pa(β^-)^{232}U	E_{β^-}=314(8) to 2$^-$ level at 1016.85 keV, and other E_{β^-}											Ens06a	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{233}Fr−^{133}Cs$_{1.752}$		218166	21				2			MA8	1.0	14Kr09	
^{233}Ra−u		47602	32	47595	9	−0.2	o			GS3	1.0	08ChZI	
		47582	17			0.7	1	30	30 ^{233}Ra	GS3	1.0	12Ch19	
^{233}Ra−^{133}Cs$_{1.752}$		213248	11	213243	9	−0.5	1	70	70 ^{233}Ra	MA8	1.0	14Kr09	
^{233}Ac−u		44363	32	44346	14	−0.5	o			GS3	1.0	08ChZI	
		44346	14				2			GS3	1.0	12Ch19	
^{233}U O$_2$−^{208}Pb$_{1.274}$		59211.9	1.0	59211.7	0.9	−0.2	1	84	84 ^{233}U	TG3	1.0	22La.A	
^{233}U(α)^{229}Th		4908.4	1.2	4908.9	1.2	0.4	1	93	89 ^{229}Th	Kum		68Ba25	Z
^{233}Np(α)^{229}Pa		5626.7	50.9				2					50Ma14	
^{233}Pu(α)^{229}U		6416.3	20.				8					57Th10	
^{233}Am(α)^{229}Np^{p}		6898.6	17.3				8					00Sa52	
^{233}Cm(α)^{229}Pu		7468.5	10.	7474	20	0.5	o			GSa		01Ca.B	
		7468.5	10.			0.5	o			GSa		02CaZU	
		7473.5	20.				8			GSa		10Kh06	
^{233}Bk(α)^{229}Am^{p}		7905.9	20.3				6			GSa		15De22	
^{232}Th(n,γ)^{233}Th		4786.69	0.25	4786.39	0.09	−1.2	–					74Ke13	Z
		4786.34	0.10			0.5	–			Bdn		06Fi.A	
^{232}Th(d,p)^{233}Th		2555	10	2561.82	0.09	0.7	U			MIT		72Gr19	
		2567	7			−0.7	U			ANL		72Vo08	
^{232}Th(n,γ)^{233}Th	ave.	4786.39	0.09	4786.39	0.09	−0.0	1	100	93 ^{233}Th			average	
^{233}Th(β^-)^{233}Pa		1245	3	1244.5	1.1	−0.2	1	12	7 ^{233}Th			57Fr.A	*
^{233}Pa(β^-)^{233}U		568	4	568.6	1.1	0.2	–					54Br37	*
		568	5			0.1	–					55On05	*
		566	5			0.5	–					63Bi03	*
	ave.	567.4	2.6			0.4	1	18	9 ^{233}U			average	
*^{233}Th(β^-)^{233}Pa	PrvCom to reference											58St50	**
*^{233}Pa(β^-)^{233}U	E_{β^-}=568(5), 256(4) to ground state, 3/2$^+$ level at 311.904 keV											Ens057	**
*^{233}Pa(β^-)^{233}U	E_{β^-}=568(5), 257(5) to ground state, 3/2$^+$ level at 311.904 keV											Ens057	**
*^{233}Pa(β^-)^{233}U	E_{β^-}=254(5) to 3/2$^+$ level at 311.904 keV											Ens057	**
^{234}Ra−u		50358	33	50382	9	0.7	o			GS3	1.0	08ChZI	
		50342	33			1.2	U			GS3	1.0	12Ch19	
^{234}Ra−^{133}Cs$_{1.759}$		216692.1	9.0				2			MA8	1.0	14Kr09	
^{234}Ac−u		48137	32	48139	15	0.1	o			GS3	1.0	08ChZI	
		48139	15				2			GS3	1.0	12Ch19	
^{234}U(α)^{230}Th		4857.4	1.0	4858.9	0.5	1.5	–					55Go.A	Z
		4860.4	2.			−0.7	–			Kum		67Ba43	Z
	ave.	4858.0	0.9			1.0	1	30	17 ^{230}Th			average	
^{234}Pu(α)^{230}U		6310.1	5.				3					60Ho.A	*
^{234}Am(α)^{230}Np		6572.6	20.	6740#	150#	8.4	F					90Ha02	*
^{234}Cm(α)^{230}Pu		7365.2	10.	7365	9	0.0	o			GSa		01Ca.B	
		7365.2	10.			0.0	7			GSa		02CaZU	
		7366.1	20.3			−0.0	7			RIa		16Ka13	
^{234}Bk(α)^{230}Am		8087	50	8100	50	0.2	o			RIa		02Mo.B	*
		8098.6	20.3				9			RIa		16Ka13	
^{232}Th(t,p)^{234}Th		2487	20	2495.3	2.3	0.4	U			LAl		69Br11	
^{234}U(p,t)^{232}U		−4099	15	−4120.9	0.6	−1.5	U			ANL		74Fr01	
^{234}U(p,t)^{232}U−^{184}W()^{182}W		1007.6	1.6	999.2	0.6	−5.2	B					09Le03	
		1007.6	1.8			−4.6	C					09Le.A	
^{233}U(d,p)^{234}U		4656	15	4621.6	1.0	−2.3	U			Kop		68Bj05	
^{234}U(d,t)^{233}U		−579	6	−588.9	1.0	−1.7	U			ANL		67Er02	
^{234}Th(β^-)^{234}Pa^{m}		192	2	194.9	1.0	1.5	3					55De40	*
		193	2			1.0	3					63Bj02	*
		197.7	1.5			−1.8	3					73Go40	*
^{234}Pa^{m}(IT)^{234}Pa		79	3	76.5	0.5	−0.8	U					73Go40	
		76.5	0.5				4					16Ri06	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{234}Pa(β^-)^{234}U		2230	40	2197.0	2.5	−0.8	U					62Bj01	
^{234}Pa^m(β^-)^{234}U		2290	20	2273.5	2.4	−0.8	U					63Bj02	
^{234}Np(β^+)^{234}U		1812	10	1810	8	−0.2	2					67Ha04	*
		1805	15			0.3	2					67Wa09	*
*^{234}Pu(α)^{230}U	With correction similar to reference											91Ry01	**
*^{234}Am(α)^{230}Np	F : not believed to be measured in this work, replaced by estimate											GAu	**
*^{234}Bk(α)^{230}Am	E_α=7850(50) to 100 keV excited state											16Ka13	**
*^{234}Th(β^-)^{234}Pa^m	E_{β^-}=191,100(2) reported											55De40	**
*^{234}Th(β^-)^{234}Pa^m	E_{β^-} 194, 100 reported without uncertainties											63Bj02	**
*^{234}Th(β^-)^{234}Pa^m	E_{β^-}=198.5(1.5), 104(2) to ^{234}Pa^m, 1$^-$ at 92.38(0.01) above ^{234}Pa^m											Nub255	**
*^{234}Pa^m(IT)^{234}Pa	E(γ)=106.0(0.5) to ground state in 16Ri06 in parallel with E(γ)=29.49(0.02) to 234											Nub25a	**
*^{234}Np(β^+)^{234}U	E_{β^+}=790(10) pK=0.48(0.03) respectively, to 1$^+$ at 1570.69 and 1$^+$ at 1601.8 keV											Ens074	**
^{235}Ac−u		50872	32	50840	15	−1.0	o			GS3	1.0	08ChZI	
		50840	15				2			GS3	1.0	12Ch19	
^{235}Th−u		47252	32	47255	14	0.1	o			GS3	1.0	08ChZI	
		47255	14				2			GS3	1.0	12Ch19	
^{235}Pa−u		45421	32	45402	14	−0.6	o			GS3	1.0	08ChZI	
		45399	15			0.2	1	91	91 ^{235}Pa	GS3	1.0	12Ch19	
^{235}Pa−^{133}Cs$_{1.767}$		212503	47	212469	14	−0.7	1	9	9 ^{235}Pa	RKM	1.0	23Ni04	
^{235}U−^{206}Pb C$_2$ H$_5$		30341.0	10.	30339.5	0.5	−0.1	U			C4	2.5	71Ke02	
^{235}U−^{133}Cs$_{1.767}$		211024	47	210994.7	0.5	−0.6	U			RKM	1.0	23Ni04	
^{235}U O$_2$−^{208}Pb$_{1.284}$		63742.5	1.5	63738.4	0.5	−2.8	U			TG3	1.0	22La.A	
^{235}U−C$_{18}$ H$_{18}$		−96932.8	3.8	−96922.2	0.5	1.1	U			M20	2.5	73Br06	
C$_{18}$ H$_{20}$−^{235}U		112584.2	4.8	112572.3	0.5	−1.0	U			M20	2.5	73Br06	
^{235}U(α)^{231}Th		4678	2	4679.5	0.5	0.8	–			Kum		60Ba44	*
		4681	3			−0.5	–					60Vo07	*
		4675.5	3.0			1.3	–					64Sc27	*
		4677	3			0.8	–					66Ga03	*
	ave.	4677.9	1.3			1.2	1	15	10 ^{231}Th			average	
^{235}Np(α)^{231}Pa		5197.2	2.0	5194.3	1.2	−1.5	1	35	19 ^{231}Pa	Bka		73Br12	*
^{235}Pu(α)^{231}U		5951.5	20.				3					57Th10	
^{235}Am(α)^{231}Np		6559	100	6576	13	0.2	o			JAa		99Sa.D	*
		6576	15			−0.0	o			JAa		04Sa05	*
		6576	13				9			JAa		04As12	*
^{235}Cm(α)^{231}Pu		7131.6	20.0				8			GSa		20Kh10	
^{235}Bk(α)^{231}Am^p		7764.4	17.3				10			Lza		26Wa.1	
^{233}U(t,p)^{235}U		3668	10	3661.8	0.9	−0.6	U					67Ri.A	*
^{234}U(n,γ)^{235}U		5297.1	0.5	5297.41	0.23	0.6	–					72Ri08	Z
		5297.4	0.3			0.0	–					77Ko15	Z
^{234}U(d,p)^{235}U		3075	7	3072.85	0.23	−0.3	U			ANL		70Br01	
^{235}U(d,t)^{234}U		935	15	959.82	0.23	1.7	U			Kop		68Bj05	
^{234}U(n,γ)^{235}U	ave.	5297.32	0.26	5297.41	0.23	0.4	1	80	65 ^{234}U			average	
^{235}Th(β^-)^{235}Pa		1470	80	1726	19	3.2	B					89Yu01	
^{235}Pa(β^-)^{235}U		1410	50	1373	13	−0.7	U					68Tr07	
^{235}Np(ε)^{235}U		123.5	2.	124.2	0.8	0.3	–					58Gi05	
		123.6	1.			0.6	–					72Mc25	
	ave.	123.6	0.9			0.7	1	87	84 ^{235}Np			average	
*^{235}U(α)^{231}Th	E_α=4596 4398 4372(all 2,Z) to ground state, (7/2$^-$) at 205.310, 9/2$^-$ at 236.906											Ens136	**
*^{235}U(α)^{231}Th	E_α=4598.6(3,Z) , 4402.6(3,Z) to ground state, (7/2$^-$) at 205.310 keV											Ens136	**
*^{235}U(α)^{231}Th	E_α=4595 4394 4364(all 3,Z) to ground state, (7/2$^-$) at 205.310, 9/2$^-$ at 236.906											Ens136	**
*^{235}U(α)^{231}Th	E_α=4595.3, 4397.3, 4365.3(all 3,Z) to ground state, (7/2$^-$) level at 205.310 keV											Ens136	**
*	and 9/2$^-$ at 236.906 keV												
*^{235}Np(α)^{231}Pa	E_α=5105.2(3), 5097.2(3), 5050.8(2,Z), 5024.8(2,Z), 4924.8(2,Z) to ground state,											AHW	**
*	1/2$^-$ at 9.206, 7/2$^-$ at 58.5699, 5/2$^+$ at 84.2148, 5/2$^+$ at 183.4962											Ens136	**
*^{235}Am(α)^{231}Np	E_α=6440(100) to level below 15 keV											04As12	**
*^{235}Am(α)^{231}Np	E_α=6457(14) to level below 15 keV											04As12	**
*^{235}Am(α)^{231}Np	E_α=6457(12) to level below 15 keV											04As12	**
*^{233}U(t,p)^{235}U	Q =3335(10,Ri) to 5/2$^+$ level at 332.845 keV											Ens14b	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{236}Ac−u		55037	73	54990	40	−0.7	o				GS3	1.0	10Ch19	
		54988	41				2				GS3	1.0	12Ch19	
^{236}Th−u		49665	32	49657	15	−0.2	o				GS3	1.0	08ChZI	
		49657	15				2				GS3	1.0	12Ch19	
^{236}Pa−u		48666	32	48668	15	0.1	o				GS3	1.0	08ChZI	
		48668	15				2				GS3	1.0	12Ch19	
^{236}Pa−^{133}Cs$_{1.774}$		216461	79	216396	15	−0.8	U				RKM	1.0	23Ni04	
^{236}U−^{133}Cs$_{1.774}$		213317	79	213294.7	0.5	−0.3	U				RKM	1.0	23Ni04	
^{236}U(α)^{232}Th		4572.2	3.	4571.9	0.8	−0.1	o						60Ko04	Z
		4570.0	3.1			0.6	–						61Ko11	Z
		4573.1	1.0			−1.2	–				Kum		78Ba.C	
	ave.	4572.8	1.0			−1.0	1	67	69	^{232}Th			average	
^{236}Pu(α)^{232}U		5867.15	0.08	5867.15	0.08	−0.0	1	100	100	^{236}Pu			84Ry02	Z
^{236}Am(α)^{232}Np		6256.2	40.				3				JAa		04Sa05	
^{236}Cm(α)^{232}Pu		7074.1	20.	7067	5	−0.4	U				GSa		10Kh06	
		7066.9	5.				8				JAa		10As.A	
^{236}Bk(α)^{232}Am		7447	14	7700#	200#	4.9	D				Bka		20Po07	*
^{236}U(p,t)^{234}U		−3330	15	−3361.0	0.3	−2.1	U				ANL		74Fr01	
^{235}U(n,γ)^{236}U		6545	2	6545.40	0.25	0.2	U						70Ka22	
		6545.1	0.5			0.6	–						74Ju.B	Z
		6545.4	0.5			−0.0	–						75We.A	Z
^{236}U(d,t)^{235}U		−281	6	−288.17	0.25	−1.2	U				ANL		70Br01	
^{235}U(n,γ)^{236}U	ave.	6545.3	0.4	6545.40	0.25	0.4	1	51	33	^{235}U			average	
^{236}Pa(β^-)^{236}U		3350	100	2889	14	−4.6	B						63Wo04	
		2900	200			−0.1	U						68Tr07	*
^{236}Np^m(IT)^{236}Np		60	50				3						Ens06a	
^{236}Np^m(β^-)^{236}Pu		525	10	537	6	1.2	2						56Gr11	
		544	8			−0.9	2						69Le05	*
*^{236}Bk(α)^{232}Am	Trends in Mass Surface TMS suggest ^{236}Bk 250 keV less bound												GAu	**
*^{236}Pa(β^-)^{236}U	E_{β^-}=2000(200) to 1^- level at 687.59 keV, and other E_{β^-}, reinterpreted												Ens06a	**
*^{236}Np^m(β^-)^{236}Pu	E_{β^-}=518(10) 537(8) respectively, to ground state and 2^+ level at 44.63 keV												Ens06a	**
^{237}Th−u		53690	32	53629	17	−1.9	o				GS3	1.0	08ChZI	
		53629	17				2				GS3	1.0	12Ch19	
^{237}Pa−u		51038	32	51032	13	−0.2	o				GS3	1.0	08ChZI	
		51023	14			0.6	1	91	91	^{237}Pa	GS3	1.0	12Ch19	
^{237}Pa−^{133}Cs$_{1.782}$		219605	45	219516	13	−2.0	1	9	9	^{237}Pa	RKM	1.0	23Ni04	
^{237}U−^{133}Cs$_{1.782}$		217225	45	217213.3	0.7	−0.3	U				RKM	1.0	23Ni04	
^{237}Np(α)^{233}Pa		4959.9	3.	4957.0	0.7	−1.0	U						61Ba44	*
		4956.7	1.5			0.2	–				Kum		68Ba25	*
		4959.9	3.			−1.0	U						69Va06	*
		4956.9	0.9			0.1	–				Gea		00Si02	*
	ave.	4956.8	0.8			0.2	1	88	86	^{233}Pa			average	
^{237}Pu(α)^{233}U		5753.3	20.	5745.8	1.5	−0.4	U						57Th10	
		5747	5			−0.2	1	10	7	^{237}Pu	Dba		93Dm02	
^{237}Am(α)^{233}Np^p		6146.2	5.				4						75Ah05	Z
^{237}Cm(α)^{233}Pu		6774.5	10.	6770	50	−0.1	o				JAa		02As08	
		6770.4	10.				9				JAa		06As03	
^{237}Cf(α)^{233}Cm		8220	20				9				GSa		10Kh06	
^{235}U(t,p)^{237}U		3206	20	3189.4	0.5	−0.8	U				Ald		64Mi.A	*
		3178	20			0.6	U				LAl		69Br11	
^{237}Np(p,t)^{235}Np		−3816	15	−3833.6	0.9	−1.2	U				ANL		74Fr01	
^{236}U(n,γ)^{237}U		5125.9	0.5	5125.8	0.5	−0.3	1	85	84	^{237}U	BNn		79Vo05	Z
^{236}U(d,p)^{237}U		2898	8	2901.2	0.5	0.4	U				ANL		67Er02	
^{237}Pa(β^-)^{237}U		2250	100	2145	12	−1.0	U						74Ka05	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{237}U(β^-)^{237}Np	520	5	520.0	0.5	0.0	U					53Wa05	*
	524	5			−0.8	U					56Ba39	*
	523	5			−0.6	U					57Ra04	*
^{237}Pu(ε)^{237}Np	222	8	220.2	1.3	−0.2	U					58Ho02	*
	207	18			0.7	U					59Gi54	*
^{237}Cm^{p}(IT)^{237}Cm	50	1				10					20Kh10	
*^{237}Np(α)^{233}Pa	E_α=4876.7 4774.2 4769.1(3,Z) to ground state, 7/2$^+$ at 103.635, 9/2$^+$ 109.07 keV										Ens057	**
*^{237}Np(α)^{233}Pa	E_α=4787.9(1.5,Z) to 5/2$^+$ level at 86.48 keV										Ens057	**
*^{237}Np(α)^{233}Pa	E_α=4791.0(3,Z), 4774.0(3,Z), 4770.0(3,Z) keV										69Va06	**
*	to 5/2$^+$ at 86.468, 7/2$^+$ at 103.635, 9/2$^+$ at 109.07 keV										Ens057	**
*^{237}Np(α)^{233}Pa	E_α=4788.0(0.9) keV to 5/2$^+$ at 86.468 keV										00Si02	**
*^{235}U(t,p)^{237}U	Q =2980(20) to 7/2$^+$ level at 426.15 keV										Ens068	**
*^{237}U(β^-)^{237}Np	E_{β^-}=245(5), 249(5), 248(5) respectively, to 53% 3/2$^-$ level at 267.556, and										Ens068	**
*	43% 1/2$^-$ level at 281.356 keV										Ens068	**
*^{237}Pu(ε)^{237}Np	LK=2.8(0.8) capture to 5/2$^-$ level at 59.541 keV, recalculated										Ens068	**
*^{237}Pu(ε)^{237}Np	pK=0.38(0.06) to ground state, 7/2$^+$ level at 33.196, 5/2$^-$ at 59.541 keV										Ens068	**
^{238}Pa−u	54648	32	54637	17	−0.3	o			GS3	1.0	08ChZI	
	54637	17				2			GS3	1.0	12Ch19	
^{238}U−^{206}Pb ^{32}S	104253.9	10.	104252.74	0.12	−0.0	U			C4	2.5	71Ke02	
C_{18} H_{22}−^{238}U	121366.0	2.4	121363.081	0.016	−0.5	U			M20	2.5	73Br06	
C_{24} H_{20}−^{238}U $^{35}Cl_2$	168010.8	1.4	168007.62	0.08	−0.9	U			M20	2.5	73Br06	
^{238}U−^{132}Xe$_{1.803}$	223596.003	0.012	223596.003	0.012	0.0	1	100	100 ^{238}U	Hep	1.0	24Kr02	
^{238}U−^{133}Cs$_{1.789}$	219971	21	219934.072	0.018	−1.8	U			RKM	1.0	23Ni04	
^{238}U O_2−^{208}Pb$_{1.298}$	70930.31	0.83	70924.515	0.019	−7.0	B			TG3	1.0	22La.A	
^{238}U−^{235}U	6858.6	10.	6859.3	0.5	0.0	U			C4	2.5	71Ke02	
^{238}U(α)^{234}Th	4271.5	5.	4269.9	2.1	−0.3	2					57Ha08	Z
	4265.1	5.			0.9	2					60Vo07	Z
	4272.9	5.			−0.6	2					61Ko11	Z
	4269.9	3.			−0.0	2			Gea		14Po02	
^{238}Pu(α)^{234}U	5593.20	0.2	5593.30	0.19	0.5	1	90	68 ^{238}Pu			71Gr17	Z
^{238}Am(α)^{234}Np	6041.7	30.				3					72Ah04	
^{238}Cm(α)^{234}Pu	6611.5	50.	6670	10	1.2	U					48St.A	*
	6632.0	50.			0.8	U					52Hi.A	
	6672.3	10.			−0.2	o			JAa		02As08	
	6670.3	10.				4			JAa		06As03	
^{238}U(n,α)^{235}Th	8700	50	8937	13	4.7	B					81Wa11	
^{236}U(t,p)^{238}U	2900	20	2797.4	0.4	−5.1	F			Ald		64Mi.A	*
	2782	10			1.5	U			ANL		67Er02	
	2780	20			0.9	U			LAl		69Br11	
^{238}U(p,t)^{236}U	−2765	15	−2797.4	0.4	−2.2	U			ANL		74Fr01	
^{238}U(p,d)^{237}U	−3951	20	−3928.9	0.6	1.1	U			Ald		64Mi.A	
^{238}U(d,t)^{237}U	116	6	103.8	0.6	−2.0	U			ANL		67Er02	
^{237}Np(n,γ)^{238}Np	5488.32	0.20				2			BNn		79Io01	Z
^{238}Pu(d,t)^{237}Pu	−746	10	−741.4	1.3	0.5	U			Kop		73Gr26	
^{238}Pa(β^-)^{238}U	3600	300	3586	16	−0.0	U					68Tr07	*
	3460	60			2.1	U					85Ba57	*
^{238}Np(β^-)^{238}Pu	1295	10	1290.1	0.4	−0.5	U					55Ra27	*
	1300	15			−0.7	U					56Ba95	*
*^{238}Cm(α)^{234}Pu	PrvCom to reference										58St50	**
*^{236}U(t,p)^{238}U	F : authors not satisfied with target material										AHW	**
*^{238}Pa(β^-)^{238}U	E_{β^-}=1700(300) to (3$^-$) level at 1992.2 keV, and other E_{β^-}, reinterpreted										Ens157	**
*^{238}Pa(β^-)^{238}U	Reports result from thesis										82Gi.A	**
*^{238}Np(β^-)^{238}Pu	E_{β^-}=270(10) 280(10) respectively, to 2$^+$ level at 1028.537 keV, and other E_{β^-}										Ens157	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{239}U−^{133}Cs$_{1.797}$		224266	21	224195.51	0.19	−3.4	B				RKM	1.0	23Ni04	
^{239}Np−^{133}Cs$_{1.797}$		222910	20	222841.5	0.8	−3.4	B				RKM	1.0	23Ni04	
^{239}Pu−^{133}Cs$_{1.797}$		222143	43	222064.8	0.5	−1.8	U				RKM	1.0	23Ni04	
^{239}Pu O−^{208}Pb$_{1.226}$		75704.1	1.4	75703.0	0.5	−0.8	1	11	11	^{239}Pu	TG3	1.0	22La.A	
^{239}Pu(α)^{235}U		5244.60	0.25	5244.60	0.21	0.0	1	67	43	^{235}U			79Ry.A	*
		5242.0	4.3			0.6	U						25Po09	*
^{239}Am(α)^{235}Np		5924.6	2.0	5921.8	1.5	−1.4	o				Bka		71Go01	*
		5920.2	2.0			0.8	o						75Ah05	*
		5921.8	1.5				2						91Ry01	*
^{239}Cm(α)^{235}Pu		6539.7	140.				4				JAa		02Sh.C	*
^{239}Cf(α)^{235}Cm		7760.1	25.	7760	60	0.1	9				GSa		81Mu12	
		7766.2	8.			−0.1	9				GSa		20Kh10	
^{238}U(n,γ)^{239}U		4806.55	0.30	4806.38	0.17	−0.6	2				ANL		72Bo46	Z
		4806.30	0.21			0.4	2				ILn		79Br25	Z
^{238}U(d,p)^{239}U		2588	20	2581.82	0.17	−0.3	U				Ald		64Mi.A	
		2579	7			0.4	U				Tal		66Sh16	
		2585	6			−0.5	U				ANL		67Er02	
^{238}Pu(n,γ)^{239}Pu		5646.7	0.5	5646.1	0.3	−1.2	1	37	32	^{238}Pu			75Ma.A	Z
^{238}Pu(d,p)^{239}Pu		3432	10	3421.5	0.3	−1.0	U				Kop		73Gr26	
^{239}Pu(d,t)^{238}Pu		604	10	611.1	0.3	0.7	U				ANL		73Fr01	
^{239}U(β^-)^{239}Np		1290	20	1261.3	0.8	−1.4	U						64Bl11	*
^{239}Np(β^-)^{239}Pu		722.5	1.0	723.5	0.8	1.0	1	58	49	^{239}Np			59Co63	*
*^{239}Pu(α)^{235}U	E_α=5156.59(0.25,Z) to 1/2$^+$ level at 0.0760 keV												Ens14b	**
*^{239}Pu(α)^{235}U	E_α=5154.01(4.25) to 1/2$^+$ level at 0.0760 keV												Ens14b	**
*^{239}Am(α)^{235}Np	E_α=5824.6(4,Z) 5775.6(2,Z) 5733.6(2,Z); ground state, (5/2)$^-$ 49.10, (7/2)$^-$ 91.6												Ens14b	**
*^{239}Am(α)^{235}Np	E_α=5772.7(2,Z) to (5/2)$^-$ level at 49.10 keV												Ens14b	**
*^{239}Am(α)^{235}Np	E_α=5774.2(1.5,Z) to 5/2$^-$ level at 49.10(0.10) keV												91Ry01	**
*^{239}Cm(α)^{235}Pu	Private communication to reference												08Qi03	**
*^{239}U(β^-)^{239}Np	E_{β^-}=1211(20) to 5/2$^-$ level at 74.6640 keV, and other E_{β^-}												Ens14b	**
*^{239}Np(β^-)^{239}Pu	E_{β^-}=437(1) to 5/2$^+$ level at 285.460 keV, and other E_{β^-}												Ens14b	**
^{240}U−^{133}Cs$_{1.805}$		227329	17	227239.2	1.2	−5.3	B				RKM	1.0	23Ni04	
^{240}Np−^{133}Cs$_{1.805}$		226887	30	226819	18	−2.3	U				RKM	1.0	23Ni04	
^{240}Pu−^{133}Cs$_{1.805}$		224494	31	224471.3	0.4	−0.7	U				RKM	1.0	23Ni04	
^{240}Pu(α)^{236}U		5255.88	0.15	5255.82	0.14	−0.4	1	90	83	^{236}U			72Go33	Z
^{240}Am(α)^{236}Np^p		5468.9	1.0				3						70Go42	Z
^{240}Cm(α)^{236}Pu		6397.8	0.6	6397.8	0.6	−0.0	1	100	100	^{240}Cm	Kum		71Bb10	*
^{240}Cf(α)^{236}Cm		7718.9	10.	7711	4	−0.8	9						70Si19	
		7713.8	20.			−0.1	U				GSa		10Kh06	
		7709.7	4.1			0.3	9				JAa		10As.A	
^{240}Es(α)^{236}Bk		8330	30	8310	70	−0.4	10						17Ko02	*
		8258	77			0.6	10				Bka		20Po07	
		8186.6	31.0			2.1	B				GSt		20Kh10	
^{238}U(t,p)^{240}U		2242	20	2265.3	1.1	1.2	U				Ald		64Mi.A	
		2253	20			0.6	U				LAl		69Br11	
^{240}Pu(p,t)^{238}Pu		−3692	15	−3698.5	0.3	−0.4	U				ANL		74Fr01	
^{239}Pu(n,γ)^{240}Pu		6534.1	1.0	6534.18	0.22	0.1	–						70Ch.A	
		6534.3	0.4			−0.3	–						74Ju.B	Z
		6534.2	0.4			−0.0	–						75We.A	Z
^{239}Pu(d,p)^{240}Pu		4300	10	4309.62	0.22	1.0	U				ANL		73Fr01	
^{239}Pu(n,γ)^{240}Pu	ave.	6534.24	0.27	6534.18	0.22	−0.2	1	68	51	^{239}Pu			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{240}U(β^-)^{240}Np^m	386	20	374	13	−0.6	1	43	42 ^{240}Np^m			53Kn23	*
^{240}Np^m(IT)^{240}Np	20	15	17	14	−0.2	1	83	68 ^{240}Np			81Hs02	*
^{240}Np(β^-)^{240}Pu	2199	30	2187	17	−0.4	1	32	32 ^{240}Np			51Or.A	*
^{240}Np^m(β^-)^{240}Pu	2210	20	2204	13	−0.3	1	43	43 ^{240}Np^m			59Bu20	*
^{240}Am(ε)^{240}Pu	1395	35	1383	14	−0.3	R					72Ah07	*
^{240}Cm(α)^{236}Pu	E_α=6290.5, 6247.7(0.6,Z) to ground state, 2$^+$ level at 44.63 keV										Ens06a	**
^{240}Es(α)^{236}Bk	Highest decay energy is assigned to ground state to ground state transition										HWJ202	**
^{240}U(β^-)^{240}Np^m	E_{β^-}=360(20) to ^{240}Np^m, and 1$^+$ level at 44.17 keV above										Ens08a	**
^{240}Np^m(IT)^{240}Np	From fraction IT =0.0012(0.0001)										AHW	**
^{240}Np(β^-)^{240}Pu	E_{β^-}=890(30) to 5$^-$ level at 1308.74 keV										Ens08a	**
^{240}Np^m(β^-)^{240}Pu	E_{β^-}=2180(20) to ground state and 2$^+$ level at 42.824 keV, and other E_{β^-}										Ens08a	**
^{240}Am(ε)^{240}Pu	pK=0.635(0.020) to 3$^+$ level at 1030.55 keV, recalculated										Ens08a	**
^{241}U−^{133}Cs$_{1.812}$	231693	48				2			RKM	1.0	23Ni04	
^{241}Np−^{133}Cs$_{1.812}$	229729	33	229720	30	−0.3	1	91	91 ^{241}Np	RKM	1.0	23Ni04	
^{241}Pu−^{133}Cs$_{1.812}$	228286	65	228171.1	0.4	−1.8	U			RKM	1.0	23Ni04	
^{241}Am O−^{208}Pb$_{1.236}$	80599.7	2.3	80600.7	0.4	0.4	o			TG3	1.0	22La.A	
	80601.47	0.96			−0.8	1	20	20 ^{241}Am	TG3	1.0	24Ch58	
^{241}Am O−^{238}U O$_2$	11123.12	1.0	11123.9	0.4	0.7	1	19	19 ^{241}Am	TG3	1.0	24Ch58	*
^{241}Am O−C$_{22}$	51744.8	1.9	51740.7	0.4	−1.4	U			TG1	1.5	14Ei01	
^{241}Pu(α)^{237}U	5139.6	3.	5140.1	0.5	0.2	U					68Ah01	*
	5139.3	1.2			0.6	1	16	16 ^{237}U	Kum		68Ba25	*
^{241}Am(α)^{237}Np	5637.81	0.12	5637.81	0.12	0.0	1	100	98 ^{237}Np			71Gr17	*
	5638.6	4.3			−0.2	U					25Po09	*
^{241}Cm(α)^{237}Pu	6182.8	2.0	6185.2	0.6	1.2	U			Kum		67Ba42	*
	6185.2	0.6			−0.1	–			Kum		71Bb10	*
	6185.0	2.0			0.1	–					75Ah05	*
ave.	6185.2	0.6			−0.0	1	99	93 ^{237}Pu			average	
^{241}Cf(α)^{237}Cm^p	7459.0	5.	7455	3	−0.9	11					70Si19	
	7451.9	4.1			0.7	11			JAa		10As.A	
	7451.9	15.3			0.2	U			GSa		20Kh10	
	7484.5	27.5			−1.1	U			Dbg		22Te01	
^{241}Es(α)^{237}Bk	8064.1	30.	8258	15	6.5	C			GSa		85Hi.A	*
	8250.2	20.			0.4	10			GSa		96Ni09	
	8277.7	30.5			−0.6	10			GSt		20Kh08	
	8257.3	30.5			0.0	10			Ara		26Ku.1	
^{239}Pu(t,p)^{241}Pu	3242	20	3293.91	0.23	2.6	U			LAl		69Br11	
^{240}Pu(n,γ)^{241}Pu	5241.3	0.7	5241.522	0.030	0.3	U					75Ma.A	
	5241.52	0.03			0.1	1	100	76 ^{240}Pu	ILn		98Wh01	Z
^{240}Pu(d,p)^{241}Pu	3018	6	3016.955	0.030	−0.2	U			ANL		67Er02	
^{241}Am(d,t)^{240}Am	−388	15	−390	14	−0.1	2			Kop		76Gr19	
^{241}Np(β^-)^{241}Pu	1360	100	1443	29	0.8	1	9	9 ^{241}Np			59Va32	
	1250	100			1.9	B					66Qa02	
^{241}Pu(β^-)^{241}Am	20.8	0.2	22.27	0.03	7.4	B					56Sh31	
	20.7	0.3			5.2	B					99Dr13	
	20.78	0.20			7.5	o					99Ya.A	
	21.6	0.5			1.3	U					10Lo14	*
	22.273	0.033			0.0	1	100	59 ^{241}Am			26Le08	
^{241}Cm(ε)^{241}Am	767.5	1.2	767.5	1.2	0.0	1	94	94 ^{241}Cm			89Su.A	*
^{241}Am O−^{238}U O$_2$	For original doublet ^{241}Am O−(^{238}U O$_2$).952, D_M=13072.7(1.0) μu										HWJ237	**
^{241}Pu(α)^{237}U	E_α=4896.6(3,Z), 4853.6(3,Z) to 5/2$^+$ at 159.962, 11/2$^+$ at 204.06 keV										Ens068	**
^{241}Pu(α)^{237}U	E_α=4896.3(1.2,Z), 4853.3(1.2,Z) to 5/2$^+$ at 159.962, 11/2$^+$ 204.06 keV										Ens068	**
^{241}Am(α)^{237}Np	E_α=5485.56(0.12,Z) 5442.80(0.13,Z) to 5/2$^-$ at 59.54, 7/2$^-$ at 102.96										Ens068	**
^{241}Am(α)^{237}Np	E_α=5486.14(4.25) to 5/2$^-$ at 59.54 keV										Ens068	**
^{241}Cm(α)^{237}Pu	E_α=6080.6(2,Z), 5926.6(2,Z) to ground state, 3/2$^+$ level at 155.456 keV										Ens068	**
^{241}Cm(α)^{237}Pu	E_α=5939.0(0.6,Z), 5884.7(0.6,Z) to 1/2$^+$ at 145.543, 5/2$^+$ 201.179 keV										Ens068	**
^{241}Cm(α)^{237}Pu	E_α=5938.7(2,Z), 5884.7(2,Z) to 1/2$^+$ at 145.543, 5/2$^+$ at 201.179 keV										Ens068	**
^{241}Es(α)^{237}Bk	C : new data from same group (next item) is more reliable										96Ni09	**
^{241}Pu(β^-)^{241}Am	No quoted uncertainty, estimated by evaluator										GAu	**
^{241}Cm(ε)^{241}Am	$Q(\varepsilon)$=5.5(1.2) to 3/2$^-$ level at 636.861 keV										Ens152	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{242}U−^{133}Cs$_{1.820}$		235090	97				2				RKM	1.0	23Ni04	
^{242}Np−^{133}Cs$_{1.820}$		233875	87	233850	80	−0.3	1	86	86	^{242}Np	RKM	1.0	23Ni04	
^{242}Pu−^{133}Cs$_{1.820}$		230800	71	230818.9	0.7	0.3	U				RKM	1.0	23Ni04	
^{242}Pu O$_2$−^{208}Pb$_{1.317}$		79327.8	2.3	79322.0	0.7	−2.5	U				TG3	1.0	22La.A	
^{242}Pu(α)^{238}U		4987.3	2.0	4984.1	0.6	−1.6	–						53As.A	*
		4989.5	3.0			−1.8	U						56Ko67	*
		4982.9	1.2			1.0	–				Kum		68Ba25	*
	ave.	4984.1	1.0			−0.0	1	36	36	^{242}Pu			average	
^{242}Am(α)^{238}Np		5587.5	0.8	5588.49	0.25	1.2	U				Kum		79Ba67	*
		5589.9	0.8			−1.8	U						90Ho02	*
^{242}Cm(α)^{238}Pu		6215.63	0.08	6215.64	0.08	0.0	1	100	100	^{242}Cm			71Gr17	Z
^{242}Cf(α)^{238}Cm		7516.9	4.				5						70Si19	Z
^{242}Es(α)^{238}Bk		8160.2	20.				8				GSa		10An08	
^{240}Pu(t,p)^{242}Pu		3043	20	3069.2	0.6	1.3	U				LAl		69Br11	
^{242}Pu(p,t)^{240}Pu		−3045	15	−3069.2	0.6	−1.6	U				ANL		74Fr01	
^{241}Pu(n,γ)^{242}Pu		6309.5	0.7	6309.4	0.6	−0.1	1	68	56	^{242}Pu			72Ma.A	
^{242}Pu(d,t)^{241}Pu		−49	7	−52.2	0.6	−0.5	U				ANL		67Er02	
^{241}Am(n,γ)^{242}Am		5541.5	1.5	5537.64	0.10	−2.6	U						75Ij.A	
		5537.64	0.1				2				ILn		88Sa18	Z
^{241}Am(d,p)^{242}Am		3308	15	3313.07	0.10	0.3	U				Kop		76Gr19	
^{242}Np(β^-)^{242}Pu		2700	200	2830	80	0.6	1	14	14	^{242}Np			79Ha26	
^{242}Am(β^-)^{242}Cm		651	5	663.0	0.4	2.4	U						50Ok52	*
		667	5			−0.8	U						55Ba.A	
*^{242}Pu(α)^{238}U	E_α=4904.6, 4860.6(2,Z) to ground state, 2$^+$ level at 44.916 keV												Ens157	**
*^{242}Pu(α)^{238}U	E_α=4905.2(3,Z), 4863.2(3,Z) to ground state, 2$^+$ level at 44.916 keV												Ens157	**
*^{242}Pu(α)^{238}U	E_α=4900.4(1.2,Z), 4856.1(1.2,Z) to ground state, 2$^+$ level at 44.916 keV												Ens157	**
*^{242}Am(α)^{238}Np	E_α=5206.6(0.5,Z), 5141.4(0.5,Z) from ^{242}Am^m at 48.60 to 5$^-$ at 342.439,												Ens157	**
*	and 6$^-$ at 407.59 keV; error increased due to conflict with next item												GAu	**
*^{242}Am(α)^{238}Np	E_α=5208.3(0.8,Z), 5144.3(0.9,Z) from ^{242}Am^m to 5$^-$ 6$^-$ levels (see above)												Ens157	**
*^{242}Am(β^-)^{242}Cm	E_{β^-}=628(5) to ground state and 2$^+$ level at 42.13 keV												Ens026	**
^{243}Am O−^{208}Pb$_{1.245}$		85369.89	0.85	85366.2	0.6	−4.3	B				TG3	1.0	22La.A	
		85366.31	1.00			−0.1	1	37	37	^{243}Am	TG3	1.0	24Ch58	
^{243}Am O−^{238}U O$_2$		15679.97	0.89	15679.2	0.6	−0.8	1	47	47	^{243}Am	TG3	1.0	24Ch58	*
^{243}Am O−C$_{22}$		56295.8	1.5	56296.1	0.6	0.1	U				TG1	1.5	14Ei01	
^{243}Am(α)^{239}Np		5438.8	1.0	5439.5	0.7	0.7	–				Kum		68Ba25	*
		5438.5	2.0			0.5	–				Gea		25Po09	*
	ave.	5438.7	0.9			0.9	1	66	51	^{239}Np			average	
^{243}Cm(α)^{239}Pu		6165.4	3.0	6168.8	1.0	1.1	U						57As.A	*
		6165.7	3.0			1.0	U						63Dz07	*
		6165.4	3.0			1.1	o				Kum		66Ba07	*
		6168.8	1.0				2						69Ba57	*
^{243}Bk(α)^{239}Am		6874.4	4.				3				Bka		66Ah.A	Z
^{243}Cf(α)^{239}Cm^p		7178	10				4						67Fi04	*
^{243}Es(α)^{239}Bk		8072.1	10.				11				RIa		89Ha27	
^{243}Es(α)^{239}Bk^m		8022.3	20.	8030.9	2.9	0.4	U						73Es02	
		8031.4	3.			−0.2	12				RIa		89Ha27	
		8027.3	20.			0.2	o				GSa		93Ho.A	
		8025.4	10.			0.6	12				GSa		10An08	
^{243}Fm(α)^{239}Cf		8689.3	25.4	8690	50	−0.0	o				GSa		81Mu12	
		8693.4	20.3			−0.1	o				GSa		08Kh10	
		8689.2	8.1				10				GSa		20Kh10	
^{243}Am(p,t)^{241}Am		−3407	15	−3417.5	0.7	−0.7	U				ANL		74Fr01	
^{242}Pu(n,γ)^{243}Pu		5034.2	3.	5036.7	2.3	0.8	1	59	57	^{243}Pu			76Ca25	
^{242}Pu(d,p)^{243}Pu		2807	8	2812.1	2.3	0.6	U				ANL		67Er02	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{243}Am(d,t)^{242}Am		−111	15	−104.5	0.7	0.4	U				Kop		76Gr19	
^{243}Pu(β^-)^{243}Am		578	10	575.5	2.4	−0.3	–						69Ho10	
		580	10			−0.5	–						77Dr07	
	ave.	579	7			−0.5	1	11	11	^{243}Pu			average	
$*^{243}$Am O−^{238}U O_2	For original doublet ^{243}Am O−(^{238}U O_2).959, D_M=17345.23(0.89) μu												HWJ237	**
$*^{243}$Am(α)^{239}Np	E_α=5275.2(1.0,Z) 5233.3(1.0,Z) to 5/2$^-$ at 74.6640, 7/2$^-$ at 117.715												Ens14b	**
$*^{243}$Am(α)^{239}Np	E_α=5349.0 5275.3 5232.6 to 5/2$^+$ at 0.0,												25Po09	**
$*^{243}$Am(α)^{239}Np	5/2$^-$ at 74.6640, 7/2$^-$ at 117.715; unc=2 keV estimated by FKo												25Po09	**
$*^{243}$Cm(α)^{239}Pu	E_α=6063.7, 5989.7, 5782.7, 5738.7(3,Z) to ground state, 7/2$^+$ level at 75.705,												57As.A	**
*	5/2$^+$ at 285.460, and 7/2$^+$ at 330.124 keV												Ens14b	**
$*^{243}$Cm(α)^{239}Pu	E_α=5990.5, 5783.5, 5738.5(3,Z) to 7/2$^+$ level at 75.705, 5/2$^+$ at 285.46,												Ens14b	**
*	and 7/2$^+$ at 330.124 keV												Ens14b	**
$*^{243}$Cm(α)^{239}Pu	E_α=6067.4, 5992.4(2,Z) to ground state, 7/2$^+$ at 75.705 keV												Ens14b	**
$*^{243}$Cm(α)^{239}Pu	E_α=5785.7(1.0,Z), 5742.8(1.0,Z) to 5/2$^+$ at 285.46, 7/2$^+$ at 330.124keV												Ens14b	**
$*^{243}$Cf(α)^{239}Cm^p	Unhindered E_α=7060(10); there is a weaker E_α=7170(10) keV												AHW	**
^{244}Pu O−^{208}$Pb_{1.250}$		88293.9	1.6	88293.5	0.5	−0.3	–				TG3	1.0	22La.A	
^{244}Pu O_2−^{208}$Pb_{1.327}$		85011.0	1.8	85006.0	0.5	−2.8	o				TG3	1.0	22La.A	
		85006.7	1.1			−0.6	–				TG3	1.0	24Ch58	
^{244}Pu O−^{208}$Pb_{1.250}$	ave.	88294.1	0.9	88293.5	0.5	−0.7	1	33	33	^{244}Pu			average	
^{244}Pu O−C_{22}		59119.3	1.9	59106.6	0.5	−4.5	B				TG1	1.5	14Ei01	
^{244}Pu O_2−^{238}U O_2		13404.7	1.1	13404.3	0.5	−0.3	1	23	23	^{244}Pu	TG3	1.0	24Ch58	*
^{244}Pu(α)^{240}U		4665.6	1.0	4665.6	1.0	−0.0	1	100	100	^{240}U			69Be06	Z
^{244}Cm(α)^{240}Pu		5901.60	0.03	5901.60	0.03	0.0	1	100	100	^{244}Cm	BIP		71Gr17	*
^{244}Bk(α)^{240}Am		6778.8	4.				3						66Ah.B	*
^{244}Cf(α)^{240}Cm		7327.1	2.	7331.0	1.4	1.9	–						67Fi04	Z
		7336.4	4.			−1.3	–						67Si08	Z
		7330.4	20.			0.0	U				GSa		08Kh10	
		7333.5	2.			−1.2	–				Jya		10KeZY	
		7400	92			−0.8	U				Bka		24Po18	*
	ave.	7331.0	1.4			−0.0	1	100	99	^{244}Cf			average	
^{244}Es(α)^{240}Bk^p		7696.4	20.	7703	18	0.3	6						73Es02	
		7728.9	40.7			−0.6	6				Bka		24Po18	*
^{244}Md(α)^{240}Es^p		8446.9	19.3	8950	80	9.3	B				Bka		20Po07	
		8947	61				12				GSt		20Kh08	*
^{242}Pu(t,p)^{244}Pu		2576	20	2583.7	0.8	0.4	U				LAl		69Br11	
^{244}Pu(p,t)^{242}Pu		−2560	15	−2583.7	0.8	−1.6	U				ANL		72Ma15	
^{244}Pu(t,α)^{243}Np		12405	10				2						79Fl02	*
^{244}Pu(d,t)^{243}Pu		234	5	228.4	2.3	−1.1	1	22	21	^{243}Pu	ANL		76Ca25	
^{243}Am(n,γ)^{244}Am^m		5277.90	0.07				2				ILn		84Vo07	Z
^{244}Cm(d,t)^{243}Cm		−530	7	−544.2	1.0	−2.0	U				ANL		67Er02	
^{244}Am^m(IT)^{244}Am		85.0	1.0	90.4	1.2	5.4	F						84Ho02	*
^{244}Am(β^-)^{244}Cm		1427.3	1.0				2						62Va08	*
$*^{244}$Pu O_2−^{238}U O_2	For original doublet ^{244}Pu O_2-(^{238}U O_2)$_{1.022}$, D_M=12511.1(1.1) μu												HWJ237	**
$*^{244}$Cm(α)^{240}Pu	E_α=5804.77(0.05,Z), 5762.65(0.03,Z) to ground state, 2$^+$ level at 42.824 keV												Ens08a	**
$*^{244}$Bk(α)^{240}Am	E_α=6667.5(4,Z), 6625.5(3,Z) to ground state, 2$^+$ level at 42.824 keV												Ens08a	**
$*^{244}$Cf(α)^{240}Cm	average four events 7200, 7225, 7349, 7382												24Po18	**
$*^{244}$Es(α)^{240}Bk^p	average three events 7556, 7622, 7628												24Po18	**
$*^{244}$Md(α)^{240}Es^p	Unweighted average of E_α=8860(30) at E=231.5 MeV and E_α=8740(30)												20Kh08	**
$*^{244}$Md(α)^{240}Es^p	at E=239.8 MeV												20Kh08	**
$*^{244}$Md(α)^{240}Es^p	Trends in Mass Surface TMS favors the higher $Q-\alpha$												GAu	**
$*^{244}$Pu(t,α)^{243}Np	Not necessarily to ground state												Ens928	**
$*^{244}$Am^m(IT)^{244}Am	F : value in Fig. 1 only, no source no error												AHW	**
$*^{244}$Am(β^-)^{244}Cm	E_{β^-}=387(1) to 6$^+$ level at 1040.188 keV; also E_{β^-}=1498(10) from												Ens036	**
*	^{244}Am^m at 90.4(1.2) to ground state and 2$^+$ at 42.965 keV, not used												Nub269	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{245}Cm(α)^{241}Pu		5621.9	0.5	5624.4	0.4	5.1	B			Kum		75Ba65	*
		5624.8	0.5			−0.7	1	73	51 ^{245}Cm	Ara		16Ko.B	*
^{245}Bk(α)^{241}Am		6454.7	4.	6454.5	1.4	−0.0	2					74Po08	*
		6454.5	1.5			0.0	2			Kum		75Ba25	*
^{245}Cf(α)^{241}Cm		7257.5	2.0	7258.5	1.8	0.5	–					67Fi04	*
		7265	5			−1.3	–					96Ma72	*
		7260.8	11.			−0.2	U			GSa		04He28	
	ave.	7258.5	1.9			−0.0	1	98	98 ^{245}Cf			average	
^{245}Es(α)^{241}Bk		7858.5	20.	7909	3	2.5	U					73Es01	
		7884.0	20.			1.3	U			GSa		85He22	
		7909.4	3.				4			RIa		89Ha27	
^{245}Es(α)^{241}Bk^{m}		7827.9	30.	7858.5	1.0	1.0	U					67Mi06	
		7858.5	1.				4			RIa		89Ha27	
		7856.5	20.3			0.1	U			Dbg		22Te01	
^{245}Fm(α)^{241}Cf^{m}		8285.5	20.	8291	6	0.3	13					67Nu01	
		8290.6	7.			0.1	13			GSa		20Kh10	
		8275.3	30.5			0.5	13			GSa		21Kh07	
		8306.8	20.3			−0.8	13			Dbg		22Te01	
^{245}Md(α)^{241}Es^{p}		8783.4	20.	8800	60	0.3	12			GSa		96Ni09	*
		8773.4	30.			0.4	12			GSt		20Kh10	
		8806.8	23.4			−0.1	12			Bka		20Po07	*
		8834.5	30.5			−0.6	12			Ara		26Ku.1	
^{244}Pu(d,p)^{245}Pu		2469	15	2475	13	0.4	2			ANL		75Er.A	*
^{244}Cm(d,p)^{245}Cm		3297	7	3294.1	0.4	−0.4	U			ANL		67Er02	
^{245}Pu(β^-)^{245}Am		1257	30	1266	14	0.3	R					68Da02	*
^{245}Am(β^-)^{245}Cm		905	5	895.8	1.5	−1.8	U					55Br02	
^{245}Es^{p}(IT)^{245}Es^{m}		253.2	0.3				3					05He27	
*^{245}Cm(α)^{241}Pu	E_α=5529.0, 5488.5, 5436.1(0.5,Z), 5303.6, 5234.4 (1.2,Z) keV											75Ba65	**
*^{245}Cm(α)^{241}Pu	to ground state, 7/2$^+$ 41.9722, 9/2$^+$ 95.7795, 9/2$^+$ 231.935, 11/2$^+$ 301.172 levels											Ens15c	**
*	E_α=5361.8(1.2,Z) to 7/2$^+$ 175.05 level in error in 75Ba65; excluded											FGK169	**
*^{245}Cm(α)^{241}Pu	E_α=5532.7, 5491.3(0.5), 5436.6(2.0), 5360.6, 5305.2, 5624.5(0.5) keV											16Ko.B	**
*	to ground state, 41.97, 95.78, 175.05, 231.94, 301.17 levels											Ens15c	**
*^{245}Bk(α)^{241}Am	E_α=6349.0, 6309.0, 6146.0, 5886.0 (all 4,Z)											91Ry01	**
*	to ground state, 7/2$^-$ at 41.176, 5/2$^+$ at 205.883, 3/2$^-$ at 471.810 keV											Ens15c	**
*^{245}Bk(α)^{241}Am	E_α=6347.8, 6307.8, 6146.8, 5885.8 recalibrated as in reference											91Ry01	**
*	to ground state, 7/2$^-$ at 41.176, 5/2$^+$ at 205.883, 3/2$^-$ at 471.810 keV											Ens15c	**
*^{245}Cf(α)^{241}Cm	E_α=7136.8(2.0,Z),7083.8(2.0,Z) to gs+5.6 and 56.1 level											96Ma72	**
*^{245}Cf(α)^{241}Cm	E_α=7145(5),7090(5) to gs+5.6 and 56.1 level											96Ma72	**
*^{245}Md(α)^{241}Es^{p}	Interpreted as e$^-$ summing with E_α=8680(20) keV in reference											21He15	**
*^{245}Md(α)^{241}Es^{p}	Originally assigned to ^{244}Md, re-assigned in reference											21He15	**
*^{244}Pu(d,p)^{245}Pu	Q =2252(15) to 217 level (estimated energy for 15/2$^-$ level)											06Ma.A	**
*^{245}Pu(β^-)^{245}Am	E_{β^-}=1210(40), 930(30) to (9/2$^+$) level at 47.07, 7/2$^+$ at 327.428 keV											Ens112	**
^{246}Cm O−^{208}Pb$_{1.260}$		91560.7	2.5	91557.9	1.1	−1.1	1	18	18 ^{246}Cm	TG3	1.0	22La.A	
^{246}Es−u		72799	122	72904	22	0.9	1	3	3 ^{246}Es	RI1	1.0	18It04	
^{246}Cm(α)^{242}Pu		5475.2	4.	5475.4	0.8	0.1	U					63Dz07	Z
		5474.9	2.			0.3	–			Kum		66Ba07	*
		5475.2	1.			0.2	–					84Sh31	*
	ave.	5475.1	0.9			0.3	1	87	81 ^{246}Cm			average	
^{246}Cf(α)^{242}Cm		6871.0	1.0	6861.8	1.0	−9.2	B					63Fr04	*
		6861.6	1.			0.2	1	99	99 ^{246}Cf	Kum		77Ba69	*
^{246}Es(α)^{242}Bk^{p}		7451.2	30.	7492	4	1.4	U					67Mi06	
		7481.9	30.			0.3	U					73Es01	
		7492.0	4.				2			RIa		89Ha27	
^{246}Fm(α)^{242}Cf		8371.4	20.	8379	5	0.4	U					66Ak01	
		8376.5	20.			0.1	U			Bka		67Nu01	
		8386.7	20.			−0.4	o			GSa		96Ni09	
		8378.4	10.			0.1	6			GSa		10An08	
		8379.5	5.			−0.0	6			GSa		11Ve03	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{246}Md(α)^{242}Es	8884.7	20.				9			GSa		96Ni09	*
	8888.8	40.	8885	20	−0.1	U			GSa		10An08	
^{246}Md^{m}(α)^{242}Es^{p}	8313.5	10.2				11			GSa		10An08	*
^{244}Pu(t,p)^{246}Pu	2085	20	2067	15	−0.9	1	56	56 ^{246}Pu	LAl		79Br19	
^{246}Cm(d,t)^{245}Cm	−196	6	−201.2	1.0	−0.9	U			ANL		67Er02	
^{246}Pu(β^-)^{246}Am^{m}	374	10	369	9	−0.5	1	89	44 ^{246}Am^{m}			56Ho23	*
^{246}Am^{m}(β^-)^{246}Cm	2300	100	2402	15	1.0	U					55En16	*
	2420	20			−0.9	1	56	56 ^{246}Am^{m}			56Sm85	*
^{246}Bk(ε)^{246}Cm	1350	60				2					89Sc26	
^{246}Es^{p}(IT)^{246}Es	151.9	2.0	151.9	2.0	0.0	1	100	97 ^{246}Es			Fgk25a	
^{246}Es^{q}(IT)^{246}Es^{m}	379.5	2.0				3			GSa		19Vo03	
*^{246}Cm(α)^{242}Pu	E_α=5385.3(2,Z), 5342.3(2,Z) to ground state, 2^+ level at 44.54 keV										Ens026	**
*^{246}Cm(α)^{242}Pu	E_α=5385.6(1,Z), 5342.6(1,Z) to ground state, 2^+ level at 44.54 keV										Ens026	**
*^{246}Cf(α)^{242}Cm	E_α=6757.4(1.0,Z), 6718.4(0.7,Z) to ground state, 2^+ level at 42.13 keV										Ens026	**
*^{246}Cf(α)^{242}Cm	E_α=6750.0(1.0,Z), 6708.2(1.0,Z) to ground state, 2^+ level at 42.13 keV										Ens026	**
*^{246}Md(α)^{242}Es	Also a lower E_α=8530(30) keV										96Ni09	**
*^{246}Md^{m}(α)^{242}Es^{p}	E_α=8178(10) to level at 531+x; x estimated to be 100#50 keV										10An08	**
*^{246}Pu(β^-)^{246}Am^{m}	E_{β^-}=150(10) to 1^+ level at 223.74 keV above ^{246}Am^{m}										Ens118	**
*^{246}Am^{m}(β^-)^{246}Cm	E_{β^-}=1222(100) 1350(20) respectively, to 1^- level at 1078.845, 2^- level at 1104.854										Ens118	**
^{247}Cm(α)^{243}Pu	5354.6	4.	5353	3	−0.5	1	70	60 ^{247}Cm			71Fi01	*
^{247}Bk(α)^{243}Am	5889.6	5.				2					69Fr01	*
^{247}Cf(α)^{243}Cm^{p}	6399.6	5.				3					84Ah02	Z
^{247}Es(α)^{243}Bk^{m}	7450.7	30.	7443.8	1.0	−0.2	U					67Mi06	
	7430.5	30.			0.4	U					73Es01	
	7443.8	1.				4			RIa		89Ha27	
^{247}Fm(α)^{243}Cf	8060.8	50.	8255	21	3.9	B			Dba		67Fl15	
	8213	18			2.4	o			GSa		89He03	*
	8287.3	20.			−1.6	o			GSa		04He28	*
	8268.1	18.3			−0.7	o			GSa		06He27	*
^{247}Fm^{m}(α)^{243}Cf	8314.9	30.	8307	16	−0.3	U			Dba		67Fl15	*
	8260.0	30.			1.5	o			GSa		97He29	*
	8304.8	11.			0.2	o			GSa		04He28	
	8306.8	16.1				3			GSa		06He27	*
^{247}Md(α)^{243}Es	8776.6	25.	8780	13	0.1	o			GSa		81Mu12	*
	8772.5	20.			0.4	o			GSa		93Ho.A	*
	8770.5	10.			0.9	o			GSa		05He27	*
	8764.4	10.			1.5	o			GSa		10An08	*
	8779.6	13.2				12			GSa		22He04	*
	8840.2	11.2			−5.4	F			Lzc		25Xu07	*
^{247}Md^{m}(α)^{243}Es	9027.9	40.	8933	8	−2.4	o			GSa		10An08	*
	8932.4	11.2			0.1	12			GSa		22He04	*
	8934.1	11.2			−0.1	12			Lzc		25Xu07	*
^{246}Cm(d,p)^{247}Cm	2931	8	2935	4	0.5	1	21	20 ^{247}Cm	ANL		67Er02	
^{247}Cf(ε)^{247}Bk	646	6	618	15	−4.6	C					56Ch.A	*
^{247}Es^{p}(IT)^{247}Es	293.7	0.1				3					05He27	
*^{247}Cm(α)^{243}Pu	E_α=5267.3(4,Z) 5212.3(4,Z) 4870.3(4,Z) to ground state, $9/2^+$ 58.13, $9/2^-$ 402.6										Ens148	**
*^{247}Bk(α)^{243}Am	E_α=5794, 5710, 5688(5,Z) to ground state, $5/2^+$ level at 84.00, $7/2^+$ at 109.22 keV										Ens148	**
*^{247}Fm(α)^{243}Cf	E_α=8060(15) summed with e^-										AHW	**
*^{247}Fm(α)^{243}Cf	E_α=7840(20) to 318 level										04He28	**
*^{247}Fm(α)^{243}Cf	E_α=7824(10 stat 15 syst) to 315 level										06He27	**
*^{247}Fm^{m}(α)^{243}Cf	Only one event										97He29	**
*^{247}Fm^{m}(α)^{243}Cf	Not observed in later work on ^{251}No decay										01He35	**
*^{247}Fm^{m}(α)^{243}Cf	E_α=8172(5 stat,15 syst)										06He27	**
*^{247}Md(α)^{243}Es	E_α=8428(25) to 209.6 level										10An08	**
*^{247}Md(α)^{243}Es	E_α=8424(20) to 209.6 level										10An08	**
*^{247}Md(α)^{243}Es	E_α=8422(10) to 209.6 level										10An08	**
*^{247}Md(α)^{243}Es	E_α=8616(20), 8416(10) to ground state, 209.6 level										10An08	**
*^{247}Md(α)^{243}Es	E_α=8421(13),8371(13) keV to levels at 219.0, 271.4 keV										22He04	**
*^{247}Md(α)^{243}Es	E_α=8404(11) to level at 303.1 keV, but E_α=8459(11) to										25Xu07	**
*^{247}Md(α)^{243}Es	level at 138.4 keV gives Q_α=8731.4(11.2) - see Table I .										25Xu07	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{247}Md^m(\alpha)^{243}Es$		E_α=8783(40) to 1/2$^-$ level at 100 keV										10An08	**
$*^{247}Md^m(\alpha)^{243}Es$		E_α=8451(11) to 342.1 level										22He04	**
$*^{247}Md^m(\alpha)^{243}Es$		E_α=8458(11) to 342.1 keV level										25Xu07	**
$*^{247}Cf(\varepsilon)^{247}Bk$		LMK=10(3) assuming first-forbidden to (5/2$^-$) level at 447.80 keV, yields 646.8										Ens153	**
*		contradicts LMK=75 allowed to (5/2)$^+$ level at 334.92 keV, yields 550;										WgM10a	**
*		both conflict with 613 from Shiptrap data to ^{255}No plus α chain										WgM10a	**
$^{248}Cm\ O-^{208}Pb_{1.269}$		96883.43	0.83	96881.7	0.5	−2.1	o			TG3	1.0	22La.A	
		96881.11	0.89			0.6	1	35	35 ^{248}Cm	TG3	1.0	24Ch58	
$^{248}Cm\ O-^{238}U\ O_2$		26632.8	1.4	26634.3	0.5	1.1	1	14	14 ^{248}Cm	TG3	1.0	24Ch58	*
$^{248}Cm(\alpha)^{244}Pu$		5161.81	0.25	5161.71	0.25	−0.4	1	93	50 ^{248}Cm	Kum		77Ba69	Z
$^{248}Cf(\alpha)^{244}Cm$		6364.8	5.1	6362	5	−0.6	o					78Gr10	
		6361.2	5.			0.1	1	99	99 ^{248}Cf			84Ah02	*
$^{248}Es(\alpha)^{244}Bk$		7165.8	20.	7160#	50#	−0.3	F					84Li.A	*
$^{248}Es(\alpha)^{244}Bk^p$		6982.8	15.	7020	5	2.5	U					56Ch67	
		6982.8	10.			3.8	B					70Ah01	
		7020.4	5.				5			RIa		89Ha27	
$^{248}Fm(\alpha)^{244}Cf$		8009.4	30.	7996	8	−0.4	–					66Ak01	
		7999.3	20.			−0.2	–					67Nu01	
		8002.3	15.			−0.4	o			GSa		85He.A	
		7992.2	11.			0.3	–			GSa		04He28	
		8019.6	47.8			−0.5	–			Bka		24Po18	*
	ave.	7996	9			−0.0	1	79	78 ^{248}Fm			average	
$^{248}Md(\alpha)^{244}Es^p$		8497.3	30.	8500	30	0.1	8					73Es01	
		8538	116			−0.3	8			Bka		24Po18	
$^{248}Cm(p,t)^{246}Cm$		−2894	15	−2897.5	1.1	−0.2	U			ANL		74Fr01	
$^{248}Cm(d,t)^{247}Cm$		49	8	37	4	−1.4	1	21	20 ^{247}Cm	ANL		67Er02	
$^{248}Bk^m(IT)^{248}Bk$		50	30	36	26	−0.5	1	76	65 ^{248}Bk			S-k25c	
$^{248}Bk^m(\beta^-)^{248}Cf$		870	20	876	19	0.3	1	90	89 ^{248}Bk^m			78Gr10	
$*^{248}Cm\ O-^{238}U\ O_2$		For original doublet ^{248}Cm O−(^{238}U O$_2$).978, D_M=27526.4(1.4) μu										HWJ237	**
$*^{248}Cf(\alpha)^{244}Cm$		E_α=6257.8(5,Z), 6216.8(5,Z) to ground state, 2$^+$ level at 42.965 keV										Ens036	**
$*^{248}Es(\alpha)^{244}Bk$		F : this line is not observed in more recent works										AHW	**
$*^{248}Fm(\alpha)^{244}Cf$		average of 7830(47),7950(47)										24Po18	**
$^{249}Cf-^{208}Pb_{1.197}$		102801.38	0.81	102800.1	0.4	−1.6	o			TG3	1.0	22La.A	
		102799.45	0.79			0.8	–			TG3	1.0	24Ch58	
$^{249}Cf\ O-^{208}Pb_{1.274}$		99514.0	1.9	99512.6	0.4	−0.7	–			TG3	1.0	24Ch58	
$^{249}Cf-^{208}Pb_{1.197}$	ave.	102799.7	0.7	102800.1	0.4	0.4	1	37	37 ^{249}Cf			average	
$^{249}Cf\ O-^{238}U\ O_2$		29150.7	1.6	29148.5	0.4	−1.4	–			TG3	1.0	24Ch58	*
$^{249}Cf-^{238}U\ O_2$		34232.85	0.78	34233.8	0.4	1.3	–			TG3	1.0	24Ch58	*
$^{249}Cf\ O-^{238}U\ O_2$	ave.	29148.1	0.7	29148.5	0.4	0.5	1	40	40 ^{249}Cf			average	
$^{249}Md-u$		82940	239	82810	170	−0.5	1	52	52 ^{249}Md	RI1	1.0	18It04	
$^{249}Cf\ O-C_{22}$		69760.0	1.4	69765.3	0.4	2.5	U			TG1	1.5	14Ei01	
$^{249}Bk(\alpha)^{245}Am$		5520.4	2.0	5521.0	1.4	0.3	3			Bka		66Ah.A	*
		5526.1	1.0			−5.1	B			Kum		71Bb10	*
		5521.6	2.0			−0.3	3			Ara		13Ah03	*
$^{249}Cf(\alpha)^{245}Cm$		6296.0	0.7	6293.2	0.4	−3.9	B			Kum		71Bb10	*
		6293.6	0.5			−0.7	1	73	49 ^{245}Cm	Ara		15Ah03	*
$^{249}Es(\alpha)^{245}Bk^m$		6881.3	5.	6886.0	1.9	0.9	4					70Ah01	Z
		6886.8	2.			−0.4	4			RIa		89Ha27	
$^{249}Fm(\alpha)^{245}Cf$		7718.3	20.	7709	6	−0.5	–					73Es01	*
		7705.1	23.			0.2	o			GSa		85He06	*
		7705.0	14.			0.3	–			GSa		04He28	*
		7710.2	8.1			−0.1	–			Orm		11Lo06	*
	ave.	7710	7			−0.1	1	80	77 ^{249}Fm			average	
$^{249}Md(\alpha)^{245}Es^p$		8161.3	20.	8158	9	−0.2	2					73Es01	
		8157.3	20.			0.0	o			GSa		85He22	
		8165	20			−0.4	o			GSa		01He35	*
		8157.3	10.			0.1	o			GSa		05He27	
		8157.3	10.			0.1	2			GSa		09He20	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{249}Md(α)^{245}Es^{p}	8155.2	31.5	8158	9	0.1	2			Dba		25Br09	
^{249}Md^{m}(α)^{245}Es	8506.9	50.8				3					19Br06	*
	8461.1	31.5	8510	50	1.5	B			Dba		25Br09	
^{249}Md^{m}(α)^{245}Es^{q}	8212.2	10.2				3			GSa		01He35	*
^{249}No(α)^{245}Fm	9248.2	32.1	9318	25	2.2	o			GSa		21Kh07	*
	9318.4	24.5				14			Dbg		22Te01	*
^{248}Cm(n,γ)^{249}Cm	4713.37	0.25				2			ILn		82Ho07	Z
^{248}Cm(d,p)^{249}Cm	2488	6	2488.80	0.25	0.1	U			ANL		67Er02	
^{249}Cm(β^-)^{249}Bk	870	100	892.4	0.8	0.2	U					58Ea06	*
	885	15			0.5	U			ANB		05Ah03	*
^{249}Bk(β^-)^{249}Cf	125	2	123.6	0.4	−0.7	U					59Va02	
	123	2			0.3	U					74Gl10	
	123.6	0.4				2					14Ch47	
^{249}Es^{p}(IT)^{249}Es	353.2	0.4				5					12He09	
*^{249}Cf O−^{238}U O$_2$	For original doublet ^{249}Cf O−(^{238}U O$_2$).981, D_M=29922.4(1.6) μu										HWJ237	**
*^{249}Cf−^{238}U O$_2$	For original doublet ^{249}Cf-(^{238}U O$_2$).922, D_M=37400.91(0.78) μu										HWJ237	**
*^{249}Md−u	Trends in Mass Surface TMS suggest ^{249}Md 100 keV less bound										HWJ262	*G
*^{249}Bk(α)^{245}Am	E_α=5431.8, 5412.8, 5384.8(all 2,Z) to ground state, 7/2$^+$ 19.20, 9/2$^+$ 47.07 keV										Ens112	**
*^{249}Bk(α)^{245}Am	E_α=5437.1(1.0,Z) to gs. Energies of higher branches										71Bb10	**
*	rather different from reference, calibrated with same ground state α										75Ba27	**
*^{249}Bk(α)^{245}Am	E_α=5433(2), 5414(2), 5386(2) to ground state, 7/2$^+$ 19.20, 9/2$^+$ 47.07 keV										Ens112	**
*^{249}Cf(α)^{245}Cm	E_α=6193.8(0.7,Z), 5813.3(1.0,Z) to ground state, 9/2$^-$ level at 388.181 keV										Ens112	**
*^{249}Cf(α)^{245}Cm	E_α=6192.4(0.5), 5810.5(0.5) to ground state, 9/2$^-$ level at 388.181 keV										Ens112	**
*^{249}Fm(α)^{245}Cf	E_α=7540(20) to corresponding 7/2$^+$ [624] level at 55(10) keV										04He28	**
*^{249}Fm(α)^{245}Cf	E_α=7527(23) to corresponding 7/2$^+$ [624] level at 55(10) keV										04He28	**
*^{249}Fm(α)^{245}Cf	Also E_α=7530(10) keV to 7/2$^+$ [624] level at 57(4) keV										11Lo06	**
*^{249}Fm(α)^{245}Cf	E_α=7530(7) to 7/2$^+$ [624] level at 57(4) keV										11Lo06	**
*^{249}Md(α)^{245}Es^{p}	E_α=8022(20) partly summed with conversion electrons										01He35	**
*^{249}Md^{m}(α)^{245}Es	E_α=8370(50) estimated from Figs. 3(α) and 4										FKo256	**
*^{249}Md^{m}(α)^{245}Es^{q}	E_α>8060 shows time component of ^{253}Lr^{m} (1.9s)										01He35	**
*^{249}No(α)^{245}Fm	E_α=9060(30) keV to level estimated at 40(10) keV										21Kh07	**
*^{249}No(α)^{245}Fm	E_α=9129(22)keV to level estimated at 40(10) keV										22Te01	**
*^{249}Cm(β^-)^{249}Bk	E_{β^-}=860(100) 876(15) respectively, to 3/2$^-$ level at 8.777 keV										Ens118	**
^{250}Md−u	84177	155	84257	11	0.5	U			RI1	1.0	18It04	*
^{250}Cf(α)^{246}Cm	6129.1	0.6	6128.51	0.19	−1.0	–			Kum		71Bb10	*
	6128.44	0.2			0.3	–					86Ry04	Z
ave.	6128.51	0.19			−0.0	1	100	100 ^{250}Cf			average	
^{250}Fm(α)^{246}Cf	7550.9	50.	7564	5	0.3	U					57Am47	
	7540.9	30.5			0.8	–					66Ak01	
	7561.1	30.			0.1	–					73Es01	
	7560.1	15.			0.3	–			ORb		77Be36	
	7556.1	35.6			0.2	o			GSa		81Mu06	
	7555.0	12.			0.7	o			GSa		04He28	
	7544.8	35.			0.5	U			Bka		06Fo02	
	7550.0	10.2			1.4	–			GSa		22Br08	
ave.	7553	8			1.4	1	44	43 ^{250}Fm			average	
^{250}Md(α)^{246}Es^{p}	7947.4	30.	7999	18	1.7	F					73Es01	*
	7964.7	20.			1.7	o			GSa		85He22	
	7967.7	40.			0.8	–			GSa		08An16	
	8003.3	20.3			−0.2	–			GSa		19Vo03	*
ave.	7996	18			0.1	1	98	97 ^{246}Es^{p}			average	
^{250}Md^{m}(α)^{246}Es^{q}	7899.7	20.3				4			GSa		19Vo03	*
^{248}Cm(t,p)^{250}Cm	2064	10				2					73Ba72	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{250}Bk^{n}(IT)^{250}Bk	83.88	0.18	83.90	0.18	0.1	1	99	99 ^{250}Bk^{n}			08Ah02	
^{250}Bk(β^-)^{250}Cf	1760	15	1781.7	2.5	1.4	U					59Va02	*
^{250}Md^{q}(IT)^{250}Md	306.8	1.5	306.7	1.5	−0.1	1	100	81 ^{250}Md			19Vo03	
*^{250}Md−u	$M-A$=78472(140) keV for mixture gs+m at 123(28) keV										WgM209	**
*^{250}Cf(α)^{246}Cm	E_α=6030.6(0.6,Z), 5988.9(0.6,Z) to ground state, 2$^+$ level at 42.852 keV										Ens118	**
*^{250}Md(α)^{246}Es^{p}	Lacking spectroscopy information										19Vo03	**
*^{250}Md^{m}(α)^{246}Es^{q}	also E_α=7741(20) to 411.4(2.0) keV above ^{246}Es^{m}										FGK257	**
*^{250}Bk(β^-)^{250}Cf	E_{β^-}=725(15) to 2$^+$ level at 1031.852 keV and 3$^+$ level at 1071.37 keV										Ens01c	**
^{251}Fm−u	81585	45	81545	15	−0.9	1	12	12 ^{251}Fm	RI1	1.0	18It04	
^{251}Md−u	84837	69	84753	4	−1.2	U			RI1	1.0	18It04	*
^{251}No−^{133}Cs$_{1.887}$	267356.7	2.4	267355.5	2.1	−0.5	1	79	79 ^{251}No	SH2	1.0	22Ka45	
^{251}No^{m}−^{133}Cs$_{1.887}$	267468.6	2.4	267469.8	2.1	0.5	1	79	79 ^{251}No^{m}	SH2	1.0	20Ka.A	
^{251}No^{m}−^{251}No	117.5	4.0	114.2	2.6	−0.8	1	42	21 ^{251}No	SH2	1.0	20Ka.A	
^{251}Cf(α)^{247}Cm	6177.2	1.0	6177.0	0.9	−0.2	2			Kum		71Bb10	*
	6176	2			0.5	2					03Ah07	*
^{251}Es(α)^{247}Bk	6593.5	5.	6597.1	1.0	0.7	o					70Ah01	*
	6597.8	3.			−0.2	o					79Ah03	*
	6597.1	1.0				3			Ara		19Ah04	*
^{251}Fm(α)^{247}Cf	7425.1	2.0	7424.5	1.0	−0.3	o					73Ah02	*
	7424.5	1.0				2			Ara		19Ah04	*
^{251}Md(α)^{247}Es^{p}	7672.5	20.3	7670	5	−0.1	U					73Es01	*
	7662.4	10.2			0.8	2			GSa		05He27	*
	7672.5	5.1			−0.4	2			Jya		06Ch52	*
^{251}No(α)^{247}Fm	8739.5	20.	8754	10	0.7	U			Bka		67Gh01	
	8732.4	15.			1.4	o			GSa		89He03	
	8762.9	20.			−0.4	o			GSa		97He29	
	8760.9	20.			−0.7	o			GSa		01He35	
	8747.7	11.			0.6	o			GSa		04He28	
	8751.8	16.3			0.2	2			GSa		06He27	
	8755.8	13.			−0.1	2			Dbg		22Te01	
^{251}No^{m}(α)^{247}Fm^{m}	8619.6	30.	8809	9	6.3	F			GSa		97He29	*
	8805.5	13.			0.3	o			GSa		04He28	
	8808.6	16.3			0.0	2			GSa		06He27	
	8809.7	11.2			−0.0	2			Dbg		22Te01	
^{251}Lr(α)^{247}Md	9396.0	19.3	9388	18	−0.4	13			Ara		22Hu21	
	9342.1	47.8			1.0	13			Bka		24Po17	
^{251}Lr^{m}(α)^{247}Md^{m}	9359.4	19.3	9349	18	−0.6	13			Ara		22Hu21	
	9282.2	47.8			1.4	13			Bka		24Po17	
^{251}Cm(β^-)^{251}Bk	1420	20				4					78Lo13	
^{251}Bk(β^-)^{251}Cf	1093	10				3					84Li05	*
^{251}Es^{p}(IT)^{251}Es	461.40	0.22				4					Ens229	
*^{251}Md−u	possible mixture of gs+p, not corrected										WgM209	**
*^{251}Cf(α)^{247}Cm	E_α=5680.1(1.0,Z) to 1/2$^+$ level at 404.90 keV										Ens153	**
*^{251}Cf(α)^{247}Cm	E_α=6078(2), 5679(2) to ground state, 1/2$^+$ level at 404.90 keV, and others										Ens153	**
*^{251}Es(α)^{247}Bk	E_α=6488.5(5,Z), 6458.5(5,Z) to ground state, (5/2$^-$) level at 29.88 keV										Ens153	**
*^{251}Es(α)^{247}Bk	E_α=6492.8(3,Z), 6462.8(3,Z) to ground state, (5/2$^-$) level at 29.88 keV										Ens153	**
*^{251}Es(α)^{247}Bk	re-evaluated data of ^{1979}Ah$_{03}$; E_α=6492.8(3,Z)										19Ah04	**
*^{251}Fm(α)^{247}Cf	E_α=7305.7(3,Z), 6833.7(2,Z) to ground state and (9/2$^-$) level at 480.40 keV										Ens153	**
*^{251}Fm(α)^{247}Cf	re-evaluated data from 73Ah02; E_α=7306.0(1.0), 6833.4(1.0) keV										19Ah04	**
*^{251}Fm(α)^{247}Cf	to ground state and (9/2$^-$) level at 480.40 keV										19Ah04	**
*^{251}Md(α)^{247}Es^{p}	E_α=7550(20) to 293.7(0.1) level										73Es01	**
*^{251}Md(α)^{247}Es^{p}	E_α=7540(10) to 293.7(0.1) level										05He27	**
*^{251}Md(α)^{247}Es^{p}	E_α=7550(1) to 293.7(0.1) level										06Ch52	**
*^{251}Md(α)^{247}Es^{p}	Original error 1 keV in 06Ch52 increased for calibration										GAu	**
*^{251}No^{m}(α)^{247}Fm^{m}	F : not observed in later work on ^{251}No decay										01He35	**
*^{251}Bk(β^-)^{251}Cf	E_{β^-}=915(10) to 3/2$^+$ level at 177.602 keV										Ens139	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{252}Md−u	86385	98				2				RI1	1.0	18It04	
^{252}No−^{133}Cs$_{1.895}$	268111	34	268134	10	0.7	o				SH1	1.0	10Dw01	
	268133	18			0.0	1	29	29	^{252}No	SH1	1.0	12Mi27	
^{252}Cf(α)^{248}Cm	6216.9	0.5	6216.95	0.04	0.1	U				Kum		71Bb10	Z
	6216.95	0.04			−0.0	1	100	100	^{252}Cf			86Ry04	Z
^{252}Es(α)^{248}Bk	6739.5	3.	6738.6	0.5	−0.3	o						73Fi06	*
	6738.6	0.5			0.0	1	100	65	^{252}Es	Ara		19Ah04	*
^{252}Fm(α)^{248}Cf	7152.7	2.	7153.7	1.0	0.5	o						84Ah02	*
	7153.7	1.0				2				Ara		19Ah04	*
^{252}No(α)^{248}Fm	8545.9	20.	8549	5	0.1	U				Bka		67Gh01	
	8545.9	30.			0.1	U				Dba		67Mi03	
	8551.0	6.			−0.4	–						77Be09	
	8542.8	15.			0.4	o				GSa		85He.A	
	8538.7	13.			0.8	–				GSa		04He28	
	8733	47			−3.9	B				Bka		24Po18	*
ave.	8549	6			−0.0	1	92	71	^{252}No			average	
^{252}Lr(α)^{248}Md	9163.8	20.	9164	17	0.0	9				GSa		01He35	
	9165.8	30.			−0.0	9				Bka		08Ne01	*
	9077	48			1.8	U				Bka		24Po18	*
^{252}Es(ε)^{252}Cf	1260	50	1220	29	−0.8	1	35	35	^{252}Es			73Fi06	*
^{252}Es^{p}(IT)^{252}Es	477	1				2						00Ah02	
*^{252}Es(α)^{248}Bk	E_α=6632.1(3,Z), 6522.1(3,Z) to 0, 7$^+$ level at 70.65 keV above ^{248}Bk											Ens14b	**
*^{252}Es(α)^{248}Bk	re-evaluated data of 73Fi06; E_α=6631.5(0.5), 6050.8(0.5) to ground state and											19Ah04	**
*^{252}Es(α)^{248}Bk	(4+) level at 590 keV											19Ah04	**
*^{252}Fm(α)^{248}Cf	E_α=7038.9(2,Z), 6998.1(2,Z) to ground state, 2$^+$ level at 41.53 keV											Ens14b	**
*^{252}Fm(α)^{248}Cf	re-evaluated data from 84Ah02											19Ah04	**
*^{252}No(α)^{248}Fm	A single observed event could represent a decay from an isomeric											24Po18	**
*^{252}No(α)^{248}Fm	state populated via the EC of ^{252}Lr											24Po18	**
*^{252}Lr(α)^{248}Md	Other E_α=9610(20) unexplained, and 8990, 8820 keV											08Ne01	**
*^{252}Lr(α)^{248}Md	average of 8884(47), 8929(47), 8987(47)											24Po18	**
*^{252}Lr(α)^{248}Md	also broad distribution 8200−8400 and an event at 8643											24Po18	**
*^{252}Es(ε)^{252}Cf	pK=0.45(0.10) to 3$^+$ level at 969.8 keV, recalculated for non-unique first											Ens061	**
*	forbidden or allowed transition; unique first forbidden would give 1440(100)											AHW	**
^{253}No−^{133}Cs$_{1.902}$	270390	13	270392	7	0.2	1	32	32	^{253}No	SH1	1.0	10Dw01	
^{253}Cf(α)^{249}Cm	6127.3	5.	6126	4	−0.3	3						66Rg01	*
	6124.6	5.			0.3	3						68Be21	*
^{253}Es(α)^{249}Bk	6739.24	0.05				3						71Gr17	Z
	6739.7	2.0	6739.24	0.05	−0.2	U				Ara		19Ah04	
^{253}Fm(α)^{249}Cf	7199	3	7197.9	1.0	−0.4	o						67Ah02	*
	7194.2	6.			0.6	U				Orm		11Lo06	*
	7197.9	1.0				2				Ara		19Ah04	*
^{253}Md(α)^{249}Es^{p}	7214.3	15.2	7219	8	0.3	o				GSa		05He27	*
	7219.4	10.1			−0.1	6				Orm		11Lo06	*
	7217.3	15.2			0.1	6				GSa		12He09	*
^{253}No(α)^{249}Fm	8419	20	8415	4	−0.2	U				Bka		67Gh01	*
	8419	30			−0.1	U				Dba		67Mi03	*
	8430	20			−0.8	o				GSa		85He.A	*
	8420	10			−0.5	o				GSa		01He.A	*
	8412.5	11.			0.2	o				GSa		04He28	*
	8411.5	5.			0.6	o				Orm		06Lo12	*
	8415.6	5.0			−0.2	–				Orm		11Lo06	*
	8412.4	11.			0.2	–				GSa		12He09	*
ave.	8415	5			−0.1	1	90	68	^{253}No			average	
^{253}Lr(α)^{249}Md	8941.6	20.	8930	8	−0.6	o				GSa		85He22	
	8935.6	10.			−0.6	o				GSa		01He35	
	8927.4	15.			0.2	o				GSa		09He20	
	8918.3	20.			0.6	–				GSa		10He11	
	8926.3	14.2			0.2	–				Ara		22Hu21	*
	8938.5	13.2			−0.7	–				Lza		24Zh19	
	8925.3	31.5			0.1	–				Dba		25Br09	
ave.	8930	8	8930	8	−0.0	1	100	52	^{253}Lr			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
$^{253}Lr^{m}(\alpha)^{249}Md^{m}$		8862.4	20.	8858	8	−0.2	o				GSa		85He22	
		8862.4	10.			−0.4	o				GSa		01He35	
		8859.4	15.			−0.1	o				GSa		09He20	
		8850.2	20.			0.4	3				GSa		10He11	
		8855.2	14.2			0.2	3				Ara		22Hu21	
		8865.4	13.2			−0.5	3				Lza		24Zh19	
		8853.2	31.5			0.2	3				Dba		25Br09	
$^{253}Rf(\alpha)^{249}No$		9459.9	20.3	9452	15	−0.4	15				Dbg		22Lo03	*
		9418	36			0.9	15				GSa		21Kh07	
		9458	28			−0.2	15				Dbg		22Lo03	
$^{253}Cf(\beta^{-})^{253}Es$		270	50	279	4	0.2	U						59Gh.A	
$*^{253}Cf(\alpha)^{249}Cm$	E_α=5981(5,Z) to 7/2$^+$ level at 48.76 keV												Ens118	**
$*^{253}Cf(\alpha)^{249}Cm$	E_α=5978.4(5,Z), 5920.4(5,Z) to 7/2$^+$ at 48.76, 7/2$^+$ at 110.173 keV												Ens118	**
$*^{253}Fm(\alpha)^{249}Cf$	E_α=7083.2(4,Z), 6943.2(3,Z), 6846.2(3,Z), 6673.2(3,Z) keV												67Ah02	**
*	to ground state and levels 5/2$^+$ at 144.98, 9/2$^+$ at 243.13, 1/2$^+$ at 416.8 keV												Ens118	**
$*^{253}Fm(\alpha)^{249}Cf$	E_α=6670(6) to 416.8 level												11Lo06	**
$*^{253}Fm(\alpha)^{249}Cf$	re-evaluated data from 67Ah02; E_α=7083.9(1.0), 6673.7(1.0) to ground state,												19Ah04	**
$*^{253}Fm(\alpha)^{249}Cf$	1/2$^+$ at 416.8 keV												19Ah04	**
$*^{253}Md(\alpha)^{249}Es^{p}$	E_α=7100(15) to 353.2(0.4) level												05He27	**
$*^{253}Md(\alpha)^{249}Es^{p}$	E_α=7105(10) to 354.6(0.6) level												11Lo06	**
$*^{253}Md(\alpha)^{249}Es^{p}$	E_α=7103(15) to 353.2(0.4) level												12He09	**
$*^{253}No(\alpha)^{249}Fm$	E_α=8010(20) to 279.7 level												04He28	**
$*^{253}No(\alpha)^{249}Fm$	E_α=8010(30) to 279.7 level												04He28	**
$*^{253}No(\alpha)^{249}Fm$	E_α=8021(20) to 279.7 level												04He28	**
$*^{253}No(\alpha)^{249}Fm$	E_α=8011(10) to 279.7 level												04He28	**
$*^{253}No(\alpha)^{249}Fm$	E_α=8004(11) to 279.7(5) level												04He28	**
$*^{253}No(\alpha)^{249}Fm$	E_α=8003(5) to 279.7(5) level; and 8280(10) to ground state												04He28	**
$*^{253}No(\alpha)^{249}Fm$	E_α=8007(4) to 279.8(0.2) level; also E_α=7615(30) to 669(3) and												11Lo06	**
*	E_α=8080(10) to 209.5(0.5) levels												11Lo06	**
$*^{253}No(\alpha)^{249}Fm$	E_α=8004(10) to 279.5(5) level												12He09	**
$*^{253}Lr(\alpha)^{249}Md$	Also E_α=8660(20) to the (11/2$^-$) level												22Hu21	**
$*^{253}Rf(\alpha)^{249}No$	Another observed E_α=9210(20) keV indicates population of another state												22Lo03	**
$^{254}No-^{133}Cs_{1.910}$		271552	15	271551.5	1.6	−0.0	o				SH1	1.0	10Dw01	
		271544	16			0.5	U				SH1	1.0	12Mi27	
$^{254}No-u$		90902	49	90964.7	1.6	1.3	U				RI1	1.0	18It04	
$^{254}No-^{133}Cs_{1.910}$		271551.9	1.6	271551.5	1.6	−0.3	1	98	98	^{254}No	SH2	1.0	22Ka45	
$^{254}Lr-^{133}Cs_{1.910}$		276920.2	7.2	276920	5	0.0	1	49	49	^{254}Lr	SH2	1.0	22Ka45	
$^{254}Lr^{m}-^{133}Cs_{1.910}$		277039.3	4.0	277039	4	−0.0	1	84	84	$^{254}Lr^{m}$	SH2	1.0	22Gi.A	
$^{254}Lr^{m}-^{254}Lr$		118.2	5.8	119	5	0.1	1	67	51	^{254}Lr	SH2	1.0	20Ka.A	
$^{254}Cf(\alpha)^{250}Cm$		5926.9	5.				3						68Be21	Z
$^{254}Es(\alpha)^{250}Bk$		6615.7	1.5	6617.1	0.5	0.9	–				Kum		72Bb24	*
		6617.4	0.5			−0.6	–				Ara		19Ah04	*
	ave.	6617.2	0.5			−0.3	1	91	91	^{254}Es			average	
$^{254}Es(\alpha)^{250}Bk^{n}$		6531.6	1.5	6533.2	0.5	1.0	1	10	9	^{254}Es	Kum		72Bb24	
$^{254}Es^{m}(\alpha)^{250}Bk$		6699.9	2.0	6697.6	1.0	−1.1	o						73Ah04	*
		6697.6	1.0			−0.0	1	100	100	^{250}Bk	Ara		19Ah04	*
$^{254}Fm(\alpha)^{250}Cf$		7306.8	5.	7307.3	1.0	0.1	U				Bka		64As01	Z
		7307.6	2.			−0.2	o						84Ah02	*
		7307.2	1.0			−0.0	1	100	100	^{254}Fm	Ara		19Ah04	*
$^{254}No(\alpha)^{250}Fm$		8229.8	20.	8229	5	−0.1	o				Bka		67Gh01	
		8240.0	30.			−0.4	U				Dba		67Mi03	
		8215.6	20.			0.6	o				GSa		85He22	
		8177.0	30.			1.7	U				Bka		06Fo02	
		8225.7	10.			0.3	o				GSa		10He10	
		8222.7	20.3			0.3	–				RIa		15KaZX	
		8217.6	10.2			1.1	–				GSa		22Br08	
		8222.8	10.2			0.6	–				Dba		23Ku17	
	ave.	8220	7			1.2	1	59	57	^{250}Fm			average	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.		Lab	F	Reference	
^{254}Lr(α)^{250}Md		8804.7	20.	8824	9	1.0	o				GSa		85He22	*
		8804.7	20.3			1.0	1	19	19	^{250}Md	Lza		01Ga20	*
^{254}Lr(α)^{250}Md^{q}		8528.6	30.5	8518	9	−0.4	–				Bka		06Fo02	*
		8519.4	20.3			−0.1	o				GSa		08An16	*
		8520.5	10.2			−0.3	–				GSa		19Vo03	*
	ave.	8521	10			−0.4	1	81	81	^{250}Md^{q}			average	
^{254}Lr^{m}(α)^{250}Md^{p}		8626.1	40.6				2				GSa		19Vo03	*
^{254}Es^{m}(β^-)^{254}Fm		1172	2	1172.0	2.0	−0.0	1	100	100	^{254}Es^{m}			62Un01	*
^{254}Lr^{p}(IT)^{254}Lr		199.1	1.5	199.1	1.5	0.0	1	100	100	^{254}Lr^{p}			19Vo03	
^{254}Lr^{q}(IT)^{254}Lr^{m}		220.7	1.5				2						19Vo03	
*^{254}Es(α)^{250}Bk	re-evaluated 08Ah02; E_α=6430.5(0.5) to 7$^+$ level at 83.88(18) keV												19Ah04	**
*^{254}Es^{m}(α)^{250}Bk	E_α=6558.9(2,Z), 6383.9(2,Z) to 4$^+$ at 35.59, 2$^+$ at 211.82 keV												Ens01c	**
*^{254}Es^{m}(α)^{250}Bk	re-evaluated from 73Ah04; E_α=6383.5(1.0) to 2$^+$ at 211.8(0.2) keV												19Ah04	**
*^{254}Fm(α)^{250}Cf	E_α=7192.3(2,Z), 7150.3(2,Z) to ground state, 5$^+$ level at 42.721 keV												Ens01c	**
*^{254}Fm(α)^{250}Cf	re-evaluated data from 84Ah02												19Ah04	**
*^{254}No(α)^{250}Fm	Also E_α=8050(10) to 44 keV												23Ku17	*W
*^{254}Lr(α)^{250}Md	E_α=8460(20) to 209.1 level												10Vo03	**
*^{254}Lr(α)^{250}Md	E_α=8460(20) to 209.1 level												19Vo03	**
*^{254}Lr(α)^{250}Md^{q}	also E_α=8437(50) to 209.1(1.5) keV												19Vo03	**
*^{254}Lr(α)^{250}Md^{q}	also E_α=8480(20) to 209.1(1.5) keV level												19Vo03	**
*^{254}Lr(α)^{250}Md^{q}	also E_α=8480(15) to 209.1(1.5) keV level												19Vo03	**
*^{254}Lr^{m}(α)^{250}Md^{p}	E_α=8490 keV from fig. 3b in 19Vo03, unc. estimated												FGK257	**
*^{254}Es^{m}(β^-)^{254}Fm	E_{β^-}=1127(2) to 2$^+$ level at 45.000 keV												Ens05b	**
^{255}No−^{133}Cs$_{1.917}$		274440	16	274445	15	0.3	1	89	89	^{255}No	SH1	1.0	12Mi27	
^{255}Lr−^{133}Cs$_{1.917}$		277811	19	277795.2	1.9	−0.8	U				SH1	1.0	12Mi27	*
		277795.6	1.9			−0.2	1	95	95	^{255}Lr	SH2	1.0	22Ka45	
^{255}Lr^{m}−^{133}Cs$_{1.917}$		277831.8	2.0	277832.2	1.9	0.2	1	95	95	^{255}Lr^{m}	SH2	1.0	22Gi.A	
^{255}Lr^{m}−^{255}Lr		35.0	21.0	37.0	2.6	0.1	U				SH2	1.0	20Ka.A	
^{255}Es(α)^{251}Bk		6439.3	3.0	6436.3	1.3	−1.0	4						66Rg01	*
		6435.6	1.5			0.5	4				Kum		71Bb10	*
^{255}Fm(α)^{251}Cf		7237.0	4.	7240.6	0.5	0.9	U						64As01	*
		7240.4	2.			0.1	o						75Ah01	*
		7240.6	0.5				3				Ara		19Ah04	*
^{255}Md(α)^{251}Es		7891.2	15.	7906.3	1.8	1.0	U				GSa		05He27	*
^{255}Md(α)^{251}Es^{p}		7438.9	5.1	7444.9	1.8	1.2	5						70Fi12	
		7457.2	5.1			−2.4	5						71Ho16	
		7444.0	4.1			0.2	o				Ara		00Ah02	
		7444.0	2.0			0.5	5				Ara		19Ah04	*
^{255}No(α)^{251}Fm		8451.1	6.	8428.2	3.0	−3.8	B				ORb		71Di03	*
		8426.4	20.			0.1	o				GSa		98Ho13	*
		8391.9	35.			1.0	U				RIa		04Mo40	*
		8449.3	20.			−1.1	U				Bka		04Fo08	*
		8426.4	10.			0.2	U				GSa		06He20	*
		8403.8	60.			0.4	U				Bka		06Gr24	*
		8428.4	3.			−0.1	1	100	88	^{251}Fm	JAa		11As03	*
^{255}Lr(α)^{251}Md		8563.6	18.	8561	4	−0.2	–				ORb		76BeYM	*
		8534.1	30.			0.9	U				Lza		01Ga20	*
		8554.4	10.			0.6	–				Jya		06Ch52	
		8554.4	10.			0.6	–				Orm		08Ha31	*
	ave.	8556	7			0.7	1	33	28	^{251}Md			average	
^{255}Lr(α)^{251}Md^{m}		8503.6	18.	8503	4	−0.0	2				ORb		76BeYM	*
		8442.7	50.			1.2	F				Bka		95Gh04	*
		8493.5	30.			0.3	U				Lza		01Ga20	*
		8498.6	5.			0.8	2				Jya		06Ch52	
		8506.7	11.			−0.3	2				GSa		08An16	*
		8509.7	10.2			−0.7	2				Orm		08Ha31	
		8509.7	10.2			−0.7	2				RIa		11As.A	*
^{255}Lr^{m}(α)^{251}Md		8592.0	5.	8595	3	0.6	–				Jya		06Ch52	*
		8602.2	11.			−0.6	–				GSa		08An16	
		8603.1	10.2			−0.8	–				Orm		08Ha31	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item		Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{255}Lr$^{m}(\alpha)^{251}$Md		8604.2	10.2	8595	3	−0.9	–			RIa		11As.A	*
	ave.	8597	4			−0.4	1	77	72 ^{251}Md			average	
^{255}Rf$(\alpha)^{251}$No		9042	20	9051	12	0.4	–			Bka		69Gh01	*
		9053	15			−0.2	o			GSa		85He06	*
		9064	20			−0.7	o			GSa		97He29	*
		9062	10			−1.1	o			GSa		01He35	*
		9056	16			−0.3	–			GSa		06He27	*
	ave.	9051	12			−0.0	1	100	100 ^{255}Rf			average	
^{255}Db$(\alpha)^{251}$Lr		9717	27	9390	50	−12.3	B			Jya		05Le.A	
		9385.5	48.				14			Bka		24Po17	*
^{255}No^{m}(IT)^{255}No		225	5				2					24Ke05	
^{255}Rf^{m}(IT)^{255}Rf		135	20	150.0	1.0	0.8	R					15An05	*
		150	1			0.0	1	100	100 ^{255}Rf^{m}			23Ch05	
*^{255}Lr−^{133}Cs$_{1.917}$	D_M=277822(17) μu for mixture 71% ground state + 29% ^{255}Lr^{m} at 41(8); $M-A$=89958(16) keV											Nub269	**
*^{255}Es$(\alpha)^{251}$Bk	E_α=6303(3,Z) 6299.3(1.5,Z) respectively, to (7/2$^+$) level at 35.5 keV											Ens139	**
*^{255}Fm$(\alpha)^{251}$Cf	E_α=7121.5, 7018.5(4,Z) to ground state, 7/2$^+$ level at 106.309 keV											Ens139	**
*^{255}Fm$(\alpha)^{251}$Cf	E_α=7126.8, 7021.8(2,Z) to ground state, 7/2$^+$ level at 106.309 keV											Ens139	**
*^{255}Fm$(\alpha)^{251}$Cf	re-evaluated data from 75Ah01; Also E_α=7022.0(0.5), 6591.3(0.5) to											19Ah04	**
*	7/2$^+$ level at 106.309 keV, 5/2$^+$ level at 543.99 keV											19Ah04	**
*^{255}Md$(\alpha)^{251}$Es	E_α=7323.5(5,Z) 7332.3(5,Z) 7327(4) 7313(15) respectively, to 7/2$^-$ at 461.5 keV											Ens139	**
*^{255}Md$(\alpha)^{251}$Es^{p}	re-evaluated data from 00Ah02; E_α=7327.0(2.0) to 7/2$^-$ at 461.5 keV											19Ah04	**
*^{255}No$(\alpha)^{251}$Fm	E_α=8312(9), 8121(6) to ground state and 199.9 level											06He20	**
*^{255}No$(\alpha)^{251}$Fm	E_α=8296(20), 8092(20) to ground state and 199.9 level											06He20	**
*^{255}No$(\alpha)^{251}$Fm	E_α=8060 to 199.9 level; also E_α=7800 keV											04Mo40	**
*^{255}No$(\alpha)^{251}$Fm	E_α=8341, 8092 to ground state and 199.9 level; also E_α=7873 keV											06He20	**
*^{255}No$(\alpha)^{251}$Fm	E_α=8290(20), 8095(10) to ground state and 199.9 level											06He20	**
*^{255}No$(\alpha)^{251}$Fm	E_α=8150, 8000 to 199.9 level											06He20	**
*^{255}No$(\alpha)^{251}$Fm	E_α=8100(3) to ^{251}Fm^{m} at 200.09(0.11) keV											11As03	**
*^{255}Lr$(\alpha)^{251}$Md	This is the faint α from long-lived isomer to the 7/2$^-$ ground state											06Ch52	**
*^{255}Lr$(\alpha)^{251}$Md	Line is mixed with ^{254}Lr's α											01Ga20	**
*^{255}Lr$(\alpha)^{251}$Md	As interpreted from Fig. 1											08Ha31	**
*^{255}Lr$(\alpha)^{251}$Md^{m}	This is the most intense α from long-lived isomer to 1/2$^-$											06Ch52	**
*^{255}Lr$(\alpha)^{251}$Md^{m}	F : one event in a questionable ^{267}Ds decay chain											AHW	**
*^{255}Lr$(\alpha)^{251}$Md^{m}	No γ observed in coincidence											08An16	**
*^{255}Lr$(\alpha)^{251}$Md^{m}	Original E_α=8371(10) corrected for recoil to E_α=8476(10)											16Hu.A	**
*^{255}Lr$(\alpha)^{251}$Md^{m}	Uncertainty estimated by evaluator											GAu	**
*^{255}Lr$^{m}(\alpha)^{251}$Md	Original error 2 keV increased for calibration											GAu	**
*^{255}Lr$^{m}(\alpha)^{251}$Md	Original E_α=8463(10) corrected for recoil to E_α=8468(10)											16Hu.A	**
*^{255}Lr$^{m}(\alpha)^{251}$Md	Uncertainty estimated by evaluator											GAu	**
*^{255}Rf$(\alpha)^{251}$No	E_α=8700(20) to 203 level											01He35	**
*^{255}Rf$(\alpha)^{251}$No	E_α=8766(15), 8715(15) to 142, 203 levels,											01He35	**
*^{255}Rf$(\alpha)^{251}$No	E_α=8905(20), 8739(20) to ground state, 203 level											01He35	**
*^{255}Rf$(\alpha)^{251}$No	E_α=8722(10) to 203(3) level											01He35	**
*^{255}Rf$(\alpha)^{251}$No	E_α=8716(4 stat, 15 syst) to 203.6(0.2) level											06He27	**
*^{255}Db$(\alpha)^{251}$Lr	average E_α=9245(47),9232(47)											24Po17	**
*^{255}Rf^{m}(IT)^{255}Rf	From 105 keV EC from K shell, error estim. by eval.											WgM164	**
^{256}Lr−^{133}Cs$_{1.925}$		280499	89	280489	6	−0.1	U			SH1	1.0	12Mi27	
		280488.9	6.1				2			SH2	1.0	22Ka45	
^{256}Fm$(\alpha)^{252}$Cf		7027.3	5.	7025.3	1.9	−0.4	2					68Ho13	Z
		7024.9	2.0			0.2	2			Ara		19Ah04	
^{256}Md$(\alpha)^{252}$Es^{p}		7316.6	5.1	7321.6	2.0	1.0	o					70Fi12	*
		7345.0	10.2			−2.3	B					71Ho16	*
		7335.9	3.1			−4.6	B					93Mo18	*
		7320.6	4.1			0.2	o					00Ah02	*
		7321.6	2.0				3					19Ah04	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{256}No(α)^{252}Fm	8553.9	30.	8581	5	0.9	U					67Fl05	*
	8553.9	20.			1.3	U			Bka		67Gh01	*
	8578.3	12.			0.2	3					81Be03	
	8582.3	6.			−0.3	3					90Ho03	
	8578.3	10.2			0.2	3					21Ke10	
^{256}Lr(α)^{252}Md^{p}	8777.4	20.3	8771	8	−0.3	3					71Es01	
	8761.1	25.			0.4	o			ORb		76BeZY	
	8777.4	20.			−0.3	3			ORb		76Di.A	
	8767.2	35.			0.1	F			RIa		04Mo26	K
	8749.9	20.			1.0	F			Bka		04Fo08	K
	8767.3	10.2			0.3	3			RIa		14Sa21	
^{256}Rf(α)^{252}No	8952.1	23.	8926	15	−1.1	o			GSa		85He06	
	8929.8	20.			−0.2	o			GSa		97He29	
	8925.7	15.2				2			GSa		10St14	
^{256}Db(α)^{252}Lr	9336.2	20.				10			Bka		08Ne01	
^{256}Db(α)^{252}Lr^{p}	9157.4	20.	9169	14	0.6	11			GSa		01He35	
	9179.7	20.			−0.6	11			Bka		08Ne01	*
	9075	48			1.9	U			Bka		24Po18	*
*^{256}Md(α)^{252}Es^{p}	Also weak E_α=7670(10) keV presumably to ground state										70Fi12	**
*^{256}Md(α)^{252}Es^{p}	Also weak E_α=7720(20) keV presumably to ground state										71Ho16	**
*^{256}Md(α)^{252}Es^{p}	Also weak E_α=7773(16) keV presumably to ground state										93Mo18	**
*^{256}Md(α)^{252}Es^{p}	Also weak E_α=7676(8) keV presumably to ground state										00Ah02	**
*	re-evaluated data of 00Ah02										19Ah04	**
*^{256}No(α)^{252}Fm	Probably mixture of two branches										AHW	**
*^{256}Lr(α)^{252}Md^{p}	limited statistics following production of ^{272}Rg										FGK256	**
*^{256}Lr(α)^{252}Md^{p}	limited statistics following production of ^{272}Rg										FGK256	**
*^{256}Db(α)^{252}Lr^{p}	5 events E_α=9030 9060 9020 9040 9030 keV										08Ne01	**
*^{256}Db(α)^{252}Lr^{p}	average of E_α=8890(47) 8930(47) 8980(47),										24Po18	**
*^{256}Db(α)^{252}Lr^{p}	also a group in energy 9020−9200 keV										24Po18	**
^{257}Rf−^{133}Cs$_{1.932}$	285680	180	285587	9	−0.5	U			SH2	1.0	22Ka45	
^{257}Db−^{85}Rb$_{3.024}$	374106	250	374250	170	0.6	1	48	48 ^{257}Db	RI1	1.0	21Sc14	*
^{257}Fm(α)^{253}Cf	6862.7	2.	6863.7	0.9	0.5	4			Bka		67As02	*
	6864.4	2.			−0.4	o					82Ah01	*
	6863.9	1.0			−0.2	4			Ara		19Ah04	*
^{257}Md(α)^{253}Es	7549.3	5.	7557.1	0.9	1.6	U					70Fi12	*
	7557.6	1.			−0.5	4					93Mo18	*
	7552.3	3.0			1.6	4			Ara		19Ah04	*
^{257}No(α)^{253}Fm	8474.1	30.	8477	6	0.1	U					70Es02	*
	8480	30			−0.1	U			GSa		96Ho13	*
	8476.6	6.				3			JAa		05As05	*
^{257}Lr(α)^{253}Md	9010.5	20.3	9010	12	−0.0	7					71Es01	
	9001.4	12.2			0.8	o			ORb		77Be36	*
	9015.5	15.2			−0.3	o			GSa		97He29	
	9030.8	50.			−0.4	U			Lza		04Ga29	
	9010.4	15.			−0.0	7			GSa		10St14	
^{257}Rf(α)^{253}No	9079.8	15.	9087	4	0.5	2			ORb		73Be33	*
	9083.7	15.			0.2	o			GSa		85He06	
	9044.0	15.			2.8	B			GSa		97He29	
	9084.1	20.			0.1	o			GSa		07St12	*
	9084.1	10.			0.3	2			GSa		10St14	*
	9106.2	100.			−0.2	U			Ara		09Qi04	*
	9091.2	5.1			−0.9	2			Dbg		22Ha04	*
	9075.0	10.			1.2	2			GSa		16He08	*
^{257}Rf^{m}(α)^{253}No	9142.5	20.	9157	3	0.7	U			Bka		69Gh01	
	9158.8	15.			−0.1	o			ORb		73Be33	
	9155.8	8.			0.1	2			ORb		90Be.A	
	9163.9	15.			−0.5	o			GSa		97He29	
	9144.0	100.			0.1	U			Ara		09Qi04	*
	9148.8	4.3			1.8	2			Gsa		16He08	*
	9167.8	5.1			−2.2	2			Dbg		22Ha04	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{257}Db(α)^{253}Lr	9112.1	20.	9215	10	5.1	B			GSa		85He22	
	9218	10			−0.3	o			GSa		01He35	*
	9209.2	20.			0.3	o			GSa		09He20	*
	9206.6	20.			0.4	–			GSa		10He11	
	9218.7	13.2			−0.3	–			Lza		24Zh19	
	9212	32			0.1	–			Dba		25Br09	*
ave.	9215	10			−0.0	1	100	52 ^{257}Db			average	
^{257}Db^{m}(α)^{253}Lr^{m}	9305.1	20.	9306	10	0.1	o			GSa		85He22	
	9308.2	10.			−0.2	o			GSa		01He35	
	9300.0	20.3			0.3	o			GSa		09He20	
	9313.2	20.3			−0.3	4			GSa		10He11	
	9306.1	13.2			0.0	4			Lza		24Zh19	
	9289.9	31.4			0.5	4			Dba		25Br09	
^{257}Sg(α)^{253}Rf^{p}	9447.3	71.1				17			GSa		25Mo17	
^{257}Rf^{n}(IT)^{257}Rf^{m}	1082	2				3					10Be16	
*^{257}Db−^{85}Rb$_{3.024}$	D_M=374170(250) μu for mixture gs+m at 120#50 keV; $M-A$=100070(230) keV										Nub269	**
*^{257}Fm(α)^{253}Cf	E_α=6518.5(2,Z) to (9/2$^+$) level at 241.01 keV										Ens139	**
*^{257}Fm(α)^{253}Cf	E_α=6756.5(3,Z), 6520.5(2,Z) to ground state, (9/2$^+$) level at 241.01 keV										Ens139	**
*^{257}Fm(α)^{253}Cf	re-evaluated data from 82Ah01; E_α=6519.7(1.0) to (9/2$^+$) level at										19Ah04	**
*^{257}Fm(α)^{253}Cf	241.01 keV										19Ah04	**
*^{257}Md(α)^{253}Es	E_α=7066(5,Z) to 371.4 level										93Mo18	**
*^{257}Md(α)^{253}Es	E_α=7440(2), 7074(1) to ground state, 371.4 level										93Mo18	**
*^{257}Md(α)^{253}Es	re-evaluated data from 70Fi12; E_α=7069.0(3.0) to 371.4 level										19Ah04	**
*^{257}No(α)^{253}Fm	E_α=8320(30) to 22.3 keV										05As05	**
*^{257}No(α)^{253}Fm	E_α=8340(20); one event only; may be summing with e$^-$										AHW	**
*^{257}No(α)^{253}Fm	E_α=8222(6) to 124.1(1) level; also E_α=8323(7) to 22.3 keV										05As05	**
*^{257}Lr(α)^{253}Md	See also 76BeYM E_α=8861(12) keV										77Be36	**
*^{257}Rf(α)^{253}No	E_α=8778(15) to 166.7 level										07St12	**
*^{257}Rf(α)^{253}No	E_α=8778(20) and 8495(20) to 166.7 and 455 level										07St12	**
*^{257}Rf(α)^{253}No	E_α=8778(10) to 166.7; and 8950(15) to ground state is sum with conversion e$^-$										10St14	**
*^{257}Rf(α)^{253}No	E_α=8800(100) to 166.7 level										09Qi04	**
*^{257}Rf(α)^{253}No	E_α=8785(5) to 167.1 level										22Ha04	**
*^{257}Rf(α)^{253}No	E_α=8769 to 167.1 level, assumed 1% error										16He08	**
*^{257}Rf^{m}(α)^{253}No	E_α=9000(100) and 8950(100) to ground state and 54(14) level										09Qi04	**
*^{257}Rf^{m}(α)^{253}No	E_α=8942, 8296 to 62.6, 724 keV										16He08	**
*^{257}Rf^{m}(α)^{253}No	E_α=9023(5),8965(5),8286(5) to gs., 63.4, 750.0 keV										22Ha04	**
*^{257}Db(α)^{253}Lr	E_α=9074(10) partly summed with conversion e$^-$										01He35	**
*^{257}Db(α)^{253}Lr	E_α=8965(20) coinc. E(γ)=102.2; E_α=9066(20) summed with conversion e$^-$										09He20	**
*^{257}Db(α)^{253}Lr	E_α=8925(31) keV and 9009(31) keV to levels at 145 keV and 60 keV										25Br09	**
^{258}Md(α)^{254}Es	7266.8	5.	7271.3	1.9	0.9	2					70Fi12	*
	7272	2			−0.4	2					93Mo18	*
^{258}Lr(α)^{254}Md	8870	50	8904	19	0.7	3			ORb		76BeYM	*
	8910	20			−0.3	3					88Gr30	*
^{258}Rf(α)^{254}No	9192.8	30.5	9196	13	0.1	2					08Ga08	
	9196.8	14.2			−0.1	2			GSa		16He15	
	9197.8	15.2			−0.1	o			GSa		19Vo03	
^{258}Db(α)^{254}Lr	9531.0	50.	9488	10	−0.6	o			GSa		97Ho14	
	9500.6	15.			−0.2	1	4	4 ^{258}Db	GSa		09He20	
^{258}Db(α)^{254}Lr^{m}	9341.1	10.	9377	11	0.7	o			GSa		09He20	
	9330	21			2.3	F			GSa		19Vo03	*
^{258}Db(α)^{254}Lr^{p}	9288.3	10.2	9289	10	0.0	1	96	96 ^{258}Db	GSa		19Vo03	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{258}Db$^{m}(\alpha)^{254}$Lr	9445.7	15.	9548	11	6.8	F			GSa		85He22	
	9446.8	20.			5.1	F			Lza		01Ga20	
^{258}Db$^{m}(\alpha)^{254}$Lr^{q}	9217.2	10.2				3			GSa		19Vo03	*
^{258}Db^{q}(IT)^{258}Db	156.5	0.5				2					09He20	
*^{258}Md$(\alpha)^{254}$Es	E_α=6713(5) to 447.9 level										93Mo18	**
*^{258}Md$(\alpha)^{254}$Es	E_α=6763(4), 6718(2) to 403.8, 447.9 levels										93Mo18	**
*^{258}Lr$(\alpha)^{254}$Md	E_α=8648(10) in coincidence with X(L) not X(K) $\rightarrow$ E(γ)=90(50) keV										AHW	**
*^{258}Lr$(\alpha)^{254}$Md	E_α=8752 observed as sum of α's and conversion electrons										AHW	**
*^{258}Lr$(\alpha)^{254}$Md	Mass assignment confirmed										92Gr02	**
*^{258}Db$(\alpha)^{254}$Lr^{m}	E_α=9185(15), unc. as given in reference										19Vo03	**
*^{258}Db$(\alpha)^{254}$Lr^{m}	not trusted, lack of spectroscopy information										FGK25a	**
*^{258}Db$(\alpha)^{254}$Lr^{p}	other E_α=9165(15) to 134.5(1.5)+42.9(1.5) keV level										19Vo03	**
*^{258}Db$^{m}(\alpha)^{254}$Lr^{q}	other E_α=9185(15) to unobserved 112(14) keV level										19Vo03	**
^{259}No$(\alpha)^{255}$Fm	7849.2	15.	7854	5	0.3	U					73Si40	*
	7869.6	15.			−1.0	U					93Mo18	*
	7854	5				4			JAa		13As02	*
^{259}Lr$(\alpha)^{255}$Md^{p}	8582.8	20.	8574	9	−0.4	7					71Es01	
	8571.6	10.			0.2	7					92Ha22	
	8577.7	29.			−0.1	U			Bka		92Kr01	
^{259}Rf$(\alpha)^{255}$No	9054.9	20.3	9050	12	−0.2	2			Bka		69Gh01	*
	9054.9	20.3			−0.2	2					81Be03	*
	9040.6	20.3			0.5	2			GSa		98Ho13	*
	9026.6	50.			0.5	C			RIa		04Mo40	
	9026.6	50.			0.5	o			Bka		04Fo08	
	9017	50			0.7	C			Bka		06Gr24	
	8940.4	11.			10.0	F			GSa		10Ni14	*
	8968.8	50.0			1.6	U			Lza		12Zh04	
^{259}Db$(\alpha)^{255}$Lr^{q}	9618.8	20.				3			Lza		01Ga20	
^{259}Sg$(\alpha)^{255}$Rf	9771.2	30.5	9765	8	−0.1	o			GSa		85Mu11	
	9807.7	23.			−0.8	U			Bka		09Fo02	*
	9784	50			−0.4	o			GSa		09He20	*
	9765.0	8.1				2			GSa		15An05	
^{259}Sg$^{m}(\alpha)^{255}$Rf	9852.4	20.3	9850	8	−0.1	1	14	14 ^{259}Sg^{m}	GSa		15An05	
^{259}Sg$^{m}(\alpha)^{255}$Rf^{m}	9700.0	8.1	9700	8	0.0	1	86	86 ^{259}Sg^{m}	GSa		15An05	
*^{259}No$(\alpha)^{255}$Fm	suffer summing effect; highest E_α seen 7685(10); extra unc. added										WgM147	**
*^{259}No$(\alpha)^{255}$Fm	suffer summing effect; highest E_α seen 7689(4); extra unc. added										WgM147	**
*^{259}No$(\alpha)^{255}$Fm	E_α=7505(5) to 231.1 level										13As02	**
*^{259}Rf$(\alpha)^{255}$No	E_α=8770(20) to 147 keV level suggested by 09As.A; also E_α=8860(20)										FGK261	**
*^{259}Rf$(\alpha)^{255}$No	E_α=8770(20) to 147 keV level suggested by 09As.A; also E_α=8870(20)										FGK261	**
*^{259}Rf$(\alpha)^{255}$No	E_α=8756(20) to 147 keV level suggested by 09As.A; also E_α=8861(20)										FGK261	**
*^{259}Rf$(\alpha)^{255}$No	F : lifetime 107 ms is much shorter than T=2.63 s in Nubase										Nub269	**
*^{259}Sg$(\alpha)^{255}$Rf	One event only, resolution 23 keV; also a wide group at lower 9593(46)keV										09Fo02	**
*^{259}Sg$(\alpha)^{255}$Rf	One event with E_α 9050 in coincidence with 593 γ; also groups at										09He20	**
*	E_α=9607(10), 9550(10) keV										09He20	**
^{260}Lr$(\alpha)^{256}$Md^{p}	8155.6	20.3				6					71Es01	
^{260}Db$(\alpha)^{256}$Lr	9191.5	30.	9500#	40#	10.3	F			RIa		04Mo26	*
	9516.5	30.			−0.5	F			RIa		04Mo26	*
	9563.2	20.			−3.1	F			Bka		04Fo08	*
^{260}Db$(\alpha)^{256}$Lr^{p}	9283.1	20.	9271	13	−0.6	4			Bka		70Gh02	
	9262.8	17.			0.5	4			ORb		77Be36	
	9316.5	60.			−0.8	U			GSa		95Ho04	*
	9285.1	60.			−0.2	U			GSa		02Ho11	*
	9181.3	60.			1.5	U			Lza		04Ga29	*
	9310.4	60.			−0.7	U			RIa		04Mo26	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{260}Sg(α)^{256}Rf	9923.0	30.	9907	5	−0.5	o			GSa		85Mu11	
	9900.6	10.			0.7	3			GSa		09He20	
	9909.8	6.1			−0.4	3			Dba		26Is.A	
^{260}Bh(α)^{256}Db	10400.3	30.5				11					08Ne01	*
*^{260}Db(α)^{256}Lr	Highest energy event; other two E_α=8810 and 8500 keV										04Mo26	**
*^{260}Db(α)^{256}Lr	F : not observed in experiments with greater statistics										77Be36	**
*^{260}Db(α)^{256}Lr^{p}	Two events E_α=9200 and 9146; error estimated by evaluator										FGK126	**
*^{260}Db(α)^{256}Lr^{p}	Two events E_α=9156 and 9129; error estimated by evaluator										FGK126	**
*^{260}Db(α)^{256}Lr^{p}	Eight events out of 14, E_α=9170 9050 9340 9400 9010 9100 9140 9130 keV										04Mo26	**
*^{260}Db(α)^{256}Lr^{p}	Two longer-lived α-escapes are assigned to the daughter										FGK126	**
*^{260}Bh(α)^{256}Db	Other events E_α=10170, 10170 and 10190; 10080 and 10030										08Ne01	**
^{261}Rf(α)^{257}No^{p}	8409.1	20.	8410	13	0.0	5			Bka		70Gh01	
	8388.8	30.			0.7	o			GSa		98Tu01	*
	8429.4	30.5			−0.6	o			Dba		00La34	
	8470.0	50.			−1.2	U					08Ga08	*
	8419.3	50.			−0.2	5			GSa		08Dv02	
	8409.1	50.			0.0	o			RIm		11Ha13	
	8409.1	40.6			0.0	5			RIm		13Mu08	
	8409	20			0.0	5			Dba		24Og02	
^{261}Rf^{m}(α)^{257}No	8652.9	20.	8640	30	−0.5	o			GSa		96Ho13	
	8652.9	20.			−0.5	o			GSa		02Ho11	
	8659	52			−0.3	o			GSa		03Tu05	*
	8642.6	50.			−0.0	4			GSa		08Dv02	
	8652.9	50.			−0.2	o			RIm		11Ha13	*
	8642.6	60.			−0.0	4			RIm		12Ha05	*
	8642.7	60.9			−0.0	4			Dba		24Og02	
^{261}Db(α)^{257}Lr^{p}	9069.2	20.	9068	14	−0.1	9			Bka		71Gh01	
	9069.2	40.			−0.0	9			Lza		04Ga29	
	9066.2	20.			0.1	9			GSa		10St14	
^{261}Sg(α)^{257}Rf	9709.0	30.	9714	15	0.2	o			GSa		85Mu11	
	9713.1	20.			0.0	F			GSa		95Ho03	*
	9770.0	20.3			−2.8	o			GSa		07St12	
	9713.7	15.				3			GSa		10St14	*
^{261}Bh(α)^{257}Db	10562.1	25.	10500	70	−1.1	o			GSa		89Mu09	
	10507.3	75.			−0.1	o			Bka		06Fo02	*
	10492.1	75.			0.1	2			Bka		08Ne08	*
	10504.3	40.			−0.1	2			GSa		10He11	*
	10218	13			5.5	B			Lza		24Zh19	*
*^{261}Rf(α)^{257}No^{p}	In addition 60% E_α=8380(30) keV										98Tu01	**
*^{261}Rf(α)^{257}No^{p}	Single event, decay time 103.2 s										08Ga08	**
*^{261}Rf^{m}(α)^{257}No	Two events with E_α=8500(+70–30) keV										GAu	**
*^{261}Rf^{m}(α)^{257}No	From direct production (fusion-evaporation)										11Ha13	**
*^{261}Rf^{m}(α)^{257}No	Decay chain of ^{265}Sg, observation is independent of previous item										12Ha05	**
*^{261}Sg(α)^{257}Rf	F : α's to 157 level summed with conversion electron										FGK10a	**
*^{261}Sg(α)^{257}Rf	E_α=9410(15) to 157(1) level										10St14	**
*^{261}Bh(α)^{257}Db	E_α=10346(75) one event; error estimated by evaluator										06Fo02	**
*^{261}Bh(α)^{257}Db	Highest E_α; error estimated; others 10054, 10285, 10113, 10165, 9989										08Ne08	**
*^{261}Bh(α)^{257}Db	Average of 2 highest 10331, 10355 as read from graph; error estimated										10He11	**
*^{261}Bh(α)^{257}Db	broad distribution at higher energies										HWJ25a	**
^{262}Db(α)^{258}Lr^{p}	8794.5	20.	8806	11	0.6	5			Bka		71Gh01	
	8815.8	20.			−0.5	5					88Gr30	
	8804.7	20.			0.1	5			GSa		99Dr09	
	8875.8	20.			−1.4	o			RIa		09Mo12	
	8814.8	30.5			−0.3	5			RIa		14Ha04	
^{262}Sg(α)^{258}Rf	9599.8	15.2				3			GSa		10Ac.A	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{262}Bh(α)^{258}Db^{p}	9978.7	15.2				3			GSa		09He20	
^{262}Bh^{m}(α)^{258}Db^{q}	10216.2	25.	10163	15	−2.1	F			GSa		89Mu09	*
	10300.5	25.4			−5.4	B			GSa		97Ho14	
	10231.4	25.4			−2.7	B			Bka		06Fo02	
	10239.2	30.			−2.5	B			Bka		08Ne08	*
	10163.5	15.2				3			GSa		09He20	
^{262}Bh^{n}(α)^{258}Db	10531.1	25.4	10530	50	0.1	o			GSa		89Mu09	
	10605.2	25.4			−1.3	o			GSa		97Ho14	
	10508.7	76.2			0.3	o			Bka		06Fo02	*
	10544.3	76.2			−0.1	U			Bka		08Ne08	
	10534.1	15.				2			GSa		09He20	
*^{262}Bh^{m}(α)^{258}Db^{q}	F : not highest line, see reference										97Ho14	**
*^{262}Bh^{m}(α)^{258}Db^{q}	average E_α=10096, 10025, 10125 keV to the 156.5(0.5) keV level										08Ne08	**
*^{262}Bh^{n}(α)^{258}Db	Single event, error estimated by evaluator										GAu	**
^{263}Rf(α)^{259}No^{p}	8022.2	40.6	8022	29	−0.0	6					93Gr.C	
	8022.2	40.6			−0.0	6					99Ga.A	
^{263}Db(α)^{259}Lr^{p}	8484.3	27.				9			Bka		92Kr01	
^{263}Sg(α)^{259}Rf	9393.1	40.	9400	60	0.3	o			Bka		74Gh04	
	9403.3	60.				3			Bka		06Gr24	*
^{263}Sg(α)^{259}Rf^{q}	9200.2	40.	9200	60	−0.1	o			Bka		74Gh04	
	9149.4	60.9			0.8	o			Bka		94Gr08	
	9198.1	60.9				4			Bka		06Gr24	*
^{263}Sg^{m}(α)^{259}Rf^{p}	9391.1	20.	9390	13	−0.1	6			GSa		98Ho13	
	9382.9	50.8			0.1	o			Bka		03Gi05	
	9393.1	35.			−0.1	6			RIa		04Mo40	*
	9388.0	20.			0.1	6			Bka		04Fo08	
	9198.1	11.			17.5	F			GSa		10Ni14	*
^{263}Hs(α)^{259}Sg	10733.5	60.				3			Bka		09Dr02	
^{263}Hs^{m}(α)^{259}Sg	11058.5	60.				3			Bka		09Dr02	*
*^{263}Sg(α)^{259}Rf	Two events E_α=9290 and 9230 keV										06Gr24	**
*^{263}Sg(α)^{259}Rf^{q}	Four events E_α=9010, 9100, 9060 and 9060 keV										06Gr24	**
*^{263}Sg^{m}(α)^{259}Rf^{p}	Also lower E_α=9130, 9040, 9150 keV										04Mo40	**
*^{263}Sg^{m}(α)^{259}Rf^{p}	F : the α chain originating from ^{267}Hs is in conflict with other data										10Ni14	**
*^{263}Hs^{m}(α)^{259}Sg	Assignment assumed by evaluator										GAu	**
^{264}Bh(α)^{260}Db^{p}	9767.3	20.	9760	18	−0.4	6			GSa		95Ho04	*
	9636.0	60.			2.1	U			Lza		04Ga29	*
	9737.8	35.5			0.6	6			RIa		04Mo26	*
^{264}Hs(α)^{260}Sg	10870	210	10591	20	−1.3	o			GSa		87Mu15	*
	10590.8	20.				4			GSa		95Ho.B	
	10966.4	80.			−4.7	B			RIa		11Sa41	*
*^{264}Bh(α)^{260}Db^{p}	Three more events in reference E_α=9365, 9514 and 9113 keV										02Ho11	**
*^{264}Bh(α)^{260}Db^{p}	Three more events E_α=9501, 9481,9440 keV										04Ga29	**
*^{264}Bh(α)^{260}Db^{p}	Six events; also two E_α=9830 keV										04Mo26	**
*^{264}Hs(α)^{260}Sg	Q_α=11000(+100–300) derived from T(1/2), one event only										87Mu15	**
*^{264}Hs(α)^{260}Sg	Also E_α=10610(40) keV										11Sa41	**
^{265}Sg(α)^{261}Rf	8945.3	60.	8979	29	0.6	F			Dba		94La22	*
	8904.7	30.			2.5	F			GSa		96Ho13	*
	8975.7	30.			0.1	o			GSa		98Tu01	
	9077.3	30.			−3.3	F			GSa		98Tu01	*
	9036.6	50.8			−1.1	o			GSa		03Tu05	
	8985.9	50.			−0.1	6			GSa		08Du02	
	8975.7	50.8			0.1	6			RIm		12Ha05	
	8975.7	50.8			0.1	6			Dba		24Og02	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$^{265}Sg^{m}(\alpha)^{261}Rf^{p}$	8823.5	50.	8832	27	0.2	o			GSa		03Tu05	
	8813.3	40.			0.5	o			GSa		06Dv01	
	8823.5	50.			0.2	7			GSa		08Dv02	
	8843.8	40.			−0.3	7			RIa		08Mo09	
	8823.5	50.			0.2	o			RIa		12Ha05	
	8792.9	81.2			0.5	o			RIa		13Su04	
	8823.5	50.8			0.2	7			Dba		24Og02	
$^{265}Bh(\alpha)^{261}Db^{p}$	9381.9	50.				11			Lza		04Ga29	
$^{265}Hs(\alpha)^{261}Sg$	10524.2	25.	10470	15	−2.1	o			GSa		87Mu15	
	10468.3	20.			0.1	o			GSa		95Ho03	
	10459.2	15.			0.7	o			GSa		99He11	*
	10470.3	15.2				4			GSa		09He20	*
$^{265}Hs^{m}(\alpha)^{261}Sg$	10712.0	20.	10700	15	−0.6	o			GSa		95Ho03	
	10734.3	15.2			−2.3	o			GSa		99He11	*
	10699.8	15.				4			GSa		09He20	
$*^{265}Sg(\alpha)^{261}Rf$	F : average but probably due to several groups, see reference										98Tu01	**
$*^{265}Sg(\alpha)^{261}Rf$	F : this event is not trusted, see reference from same group										02Ho11	**
$*^{265}Sg(\alpha)^{261}Rf$	F : most probably not from ^{265}Sg										GAu	**
$*^{265}Hs(\alpha)^{261}Sg$	Also E_α=10426(15) keV										99He11	**
$*^{265}Hs(\alpha)^{261}Sg$	Most intense line, also E_α=10282(15), 10428(15), 10573(15) keV										09He20	**
$*^{265}Hs^{m}(\alpha)^{261}Sg$	Also E_α=10726(15) keV										99He11	**
$^{266}Sg(\alpha)^{262}Rf$	8762.0	50.	8800#	100#	0.8	F			Dba		94La22	*
	8904.1	40.			−2.6	F			GSa		98Tu01	*
	8853.4	50.			−1.1	F			GSa		02Tu05	*
$^{266}Bh(\alpha)^{262}Db^{p}$	9432	50	9380	30	−1.1	7			Bka		00Wi15	
	9245.3	81.2			1.6	7			Lza		06Qi03	
	9371.2	50.			0.1	7			RIa		09Mo12	*
$^{266}Hs(\alpha)^{262}Sg$	10335.7	20.3	10346	16	0.5	4			GSa		01Ho06	
	10360	24			−0.6	4			GSa		11Ac.A	
$^{266}Hs^{m}(\alpha)^{262}Sg^{p}$	10592.6	20.				7			GSa		11Ac.A	
$^{266}Mt(\alpha)^{262}Bh$	10995.7	25.				4			GSa		97Ho14	
$^{266}Mt^{m}(\alpha)^{262}Bh^{m}$	11269.7	50.	11460	40	3.7	B			GSa		84Mu07	*
	11168.1	30.			9.6	B			GSa		89Mu16	
	11479.0	50.8			−0.5	4			GSa		97Ho14	*
	11432.3	50.8			0.5	4			Bka		09Ne02	*
$*^{266}Sg(\alpha)^{262}Rf$	Average of two groups										02Tu05	**
$*^{266}Sg(\alpha)^{262}Rf$	F : no α decay from ^{266}Sg, all re-assigned to ^{265}Sg										08Dv02	**
$*^{266}Bh(\alpha)^{262}Db^{p}$	Also E_α=9770(40), 9080(40) keV from ^{278}Nh decay chain										08Mo09	**
$*^{266}Mt^{m}(\alpha)^{262}Bh^{m}$	One E_α only; may be ground state										AHW	**
$*^{266}Mt^{m}(\alpha)^{262}Bh^{m}$	One E_α=11739(50), one 11306(50); several smaller										AHW	**
$*^{266}Mt^{m}(\alpha)^{262}Bh^{m}$	One E_α=11260, one 10670; several escape events										FGK257	**
$^{267}Sg(\alpha)^{263}Rf^{p}$	8325.0	50.	8386	19	1.2	8			GSa		08Dv02	
	8396	20			−0.5	8			Dba		24Og02	
$^{267}Bh(\alpha)^{263}Db^{p}$	8964.5	30.5	8970	26	0.2	11			Bka		00Wi15	
	8984.8	50.			−0.3	11			GSa		02Tu05	
$^{267}Hs(\alpha)^{263}Sg$	10015.3	60.	10038	13	0.4	4			Dba		95La20	*
	10032.6	20.			0.2	4			GSa		98Ho13	
	10035.7	50.			0.0	o			Bka		03Gi05	
	10069.1	35.			−0.9	4			RIa		04Mo40	*
	10034.6	20.			0.1	4			Bka		04Fo08	*
	10145.2	50.			−2.2	U			GSa		10Ni14	*
$^{267}Hs(\alpha)^{263}Sg^{m}$	9978.8	20.	9987	13	0.4	5			GSa		98Ho13	*
	9979.8	35.			0.2	5			RIa		04Mo40	*
	9997.0	20.			−0.5	5			Bka		04Fo08	*

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$^{267}Hs^{m}(\alpha)^{263}Sg^{p}$	9979.8	20.				7			Bka		04Fo08	
$^{267}Ds(\alpha)^{263}Hs$	11776.8	50.8				4			Bka		95Gh04	*
$*^{267}Hs(\alpha)^{263}Sg$	Selecting two events at E_α=9860,9870; and one E_α=9740(60) keV										95La20	**
$*^{267}Hs(\alpha)^{263}Sg$	Selecting four events at E_α=9970, 9890, 9900, 9910 keV										04Mo40	**
$*^{267}Hs(\alpha)^{263}Sg$	Selecting two events at E_α=9880, 9888 keV										04Fo08	**
$*^{267}Hs(\alpha)^{263}Sg$	Directly produced ^{267}Hs; daughter and grand-daughter also conflicting										10Ni14	**
$*^{267}Hs(\alpha)^{263}Sg^{m}$	And one E_α=9749(20) at 13 ms										98Ho13	**
$*^{267}Hs(\alpha)^{263}Sg^{m}$	Selecting 7 events at E_α=9830, 9820, 9820, 9860, 9820, 9830, 9830 keV										04Mo40	**
$*^{267}Hs(\alpha)^{263}Sg^{m}$	Selecting 2 events at E_α=9864, 9830 keV										04Fo08	**
$*^{267}Ds(\alpha)^{263}Hs$	Maybe the upper isomer at about 250 keV excitation energy										AHW	**
$^{268}Hs(\alpha)^{264}Sg$	9622.9	16.	9760#	100#	8.6	D			GSa		10Ni14	*
$^{268}Mt(\alpha)^{264}Bh^{p}$	10395.5	20.	10438	19	2.1	o			GSa		95Ho04	*
	10431.9	20.			0.3	8			GSa		02Ho11	*
	10476.7	50.			−0.8	8			RIa		04Mo26	*
	10268	20			8.5	B			Bka		04Fo08	*
$*^{268}Hs(\alpha)^{264}Sg$	Trends in Mass Surface TMS suggest ^{268}Hs 140 keV less bound										GAu	**
$*^{268}Mt(\alpha)^{264}Bh^{p}$	Two events E_α=10221 coinc. E(γ)=93 and 10259; event #3 E_α=10097 keV										95Ho04	**
*	could be decay of an isomer with lifetime=171 ms										02Ho11	**
$*^{268}Mt(\alpha)^{264}Bh^{p}$	Average of event 95Ho04 E_α=10259 and present 10294 keV										02Ho11	**
$*^{268}Mt(\alpha)^{264}Bh^{p}$	Also E_α=10340 keV										04Mo26	**
$*^{268}Mt(\alpha)^{264}Bh^{p}$	One event only										04Fo08	**
$^{269}Sg(\alpha)^{265}Rf$	8699.6	100.	8550	50	−1.3	U					10El06	
	8628.5	60.9			−1.0	o			Dba		15Ut02	
	8536.9	40.6			0.2	o			Dba		18Ut02	
	8550	15				7			DJa		25Og03	
$^{269}Hs(\alpha)^{265}Sg^{p}$	9369.6	30.	9303	25	−2.2	B			GSa		96Ho13	*
	9349.3	30.			−1.5	o			GSa		02Ho11	*
	9278.2	50.8			0.5	o			GSa		03Tu05	*
	9268.0	50.8			0.7	8			GSa		08Dv02	
	9349.2	71.1			−0.7	o			RIa		08Mo09	
	9288.3	40.6			0.4	8			RIa		13Su04	
	9339.1	40.6			−0.9	8			Dba		24Og02	
$^{269}Hs^{m}(\alpha)^{265}Sg^{q}$	9207.1	30.5	9215	14	0.3	8			GSa		06Dv01	
	9217.3	15.2			−0.1	8			Dba		24Og02	
$^{269}Ds(\alpha)^{265}Hs^{m}$	11280.1	20.				5			GSa		95Ho03	
$*^{269}Hs(\alpha)^{265}Sg^{p}$	Event number 2 only; first event rejected, see reference										02Ho11	**
$*^{269}Hs(\alpha)^{265}Sg^{p}$	Two events E_α=9230, 9180 both following 300 μs ^{273}Ds										02Ho11	**
$*^{269}Hs(\alpha)^{265}Sg^{p}$	Three events E_α=9180, 9100, 8880; latter probably due to energy loss										03Tu05	**
$^{270}Db(\alpha)^{266}Lr$	8019.0	30.5	8310#	200#	5.0	D			GSa		14Kh04	*
$^{270}Bh(\alpha)^{266}Db$	9064.5	81.2	9110	60	0.5	o			Dba		07Og02	
	9115	31				5			Dba		22Og08	
$^{270}Hs(\alpha)^{266}Sg$	9324.3	52.8	9070	40	−4.8	F			GSa		03Tu05	*
	9024	52			1.5	o			GSa		06Dv01	*
	9013.8	50.			1.1	5			GSa		08Dv02	*
	9123.4	77.			−0.7	5			GSa		10Gr04	*
	9155.8	80.			−1.1	5			Dba		13Og03	
$^{270}Mt(\alpha)^{266}Bh$	10181.1	70.	10180#	100#	−0.0	o			RIa		08Mo09	
	10414.5	71.			−2.7	D			RIa		12Mo25	*
$^{270}Ds(\alpha)^{266}Hs$	11196.2	50.8	11117	28	−1.6	o			GSa		01Ho06	
	11117.0	28.4				5			GSa		11Ac.A	
$^{270}Ds^{m}(\alpha)^{266}Hs$	12333	50	12510	50	2.5	B			GSa		01Ho06	
	12508.6	20.				5			GSa		11Ac.A	
$^{270}Ds^{m}(\alpha)^{266}Hs^{m}$	11318	50	11410	70	1.2	o			GSa		01Ho06	
	11405.2	52.				6			GSa		11Ac.A	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{270}$Db(α)^{266}Lr	Trends in Mass Surface TMS suggest ^{270}Db 290 keV less bound										GAu	**
$*^{270}$Hs(α)^{266}Sg	F : re-assigned to ^{269}Hs										06Dv01	**
$*^{270}$Hs(α)^{266}Sg	4 events at 8850(1.62s), 8900(0.05s), 8920(0.5s), 8880(0.4s)										GAu	**
$*^{270}$Hs(α)^{266}Sg	2 events at 8760(0.27s), 8810(0.27s); E_α=8880(50) for both experiments										GAu	**
$*^{270}$Hs(α)^{266}Sg	Symmetrized from E_α=9020(+50–100); independent from previous item										GAu	**
$*^{270}$Mt(α)^{266}Bh	Trends in Mass Surface TMS suggest ^{270}Mt 230 keV more bound										GAu	**
^{271}Sg(α)^{267}Rf^{p}	8658	80	8629	16	−0.3	o			Dba		04Og12	
	8668.2	81.2			−0.4	o			Dba		06Og05	
	8628.6	16.2				10			Dba		22Og05	*
^{271}Bh(α)^{267}Db	9490.3	162.4	9370	50	−0.7	o			Dba		12St.A	
	9409.1	71.0			−0.5	o			GSa		13Ru11	
	9420	70			−0.6	o			SRv		17Og01	
	9367	21				8			Dba		22Og08	
^{271}Hs(α)^{267}Sg^{p}	9439.6	50.7	9475	19	0.7	10			GSa		08Dv02	
	9480.2	20.3			−0.3	10			Dba		24Og02	
^{271}Hs^{m}(α)^{267}Sg^{q}	9185.8	20.3				12			Dba		24Og02	
^{271}Ds(α)^{267}Hs	10869.8	20.	10870	18	0.0	5			GSa		98Ho13	*
	10870.8	35.			−0.0	5			RIa		04Mo40	*
^{271}Ds^{m}(α)^{267}Hs	10937.8	20.				5			Bka		04Fo08	*
	10803.8	50.	10938	20	2.7	F					12Zh04	*
^{271}Ds^{m}(α)^{267}Hs^{m}	10899.2	20.	10899	13	−0.0	6			GSa		98Ho13	
	10880.8	50.			0.3	o			Bka		03Gi05	
	10883.0	35.			0.4	6			RIa		04Mo40	*
	10903.3	20.			−0.2	6			Bka		04Fo08	*
$*^{271}$Sg(α)^{267}Rf^{p}	includes data from previous experiments										22Og05	**
$*^{271}$Ds(α)^{267}Hs	Decay chain number 6 for the long-lived isomer, GAu interpretation										04Mo40	**
$*^{271}$Ds(α)^{267}Hs	Trends in Mass Surface TMS suggest ^{291}Ds 110 keV less bound										HWJ266	**
$*^{271}$Ds^{m}(α)^{267}Hs	Decay chain number 6, GAu interpretation										04Fo08	**
$*^{271}$Ds^{m}(α)^{267}Hs	F : α escaped ?										GAu	**
$*^{271}$Ds^{m}(α)^{267}Hs^{m}	GAu : average of decay chains number 2, 5, 10, 13 for short-lived isomer										04Mo40	**
$*^{271}$Ds^{m}(α)^{267}Hs^{m}	Decay chains number 5 and 7, GAu interpretation										04Fo08	**
^{272}Bh(α)^{268}Db	9303.0	20.				7			Bka		15Ga24	*
	9210	10	9300	50	9.3	o			SRv		17Og01	*
	9327.4	60.9			−0.4	o			Bka		18Ga34	
^{272}Bh(α)^{268}Db^{p}	9154.9	60.	9161	16	0.1	o			Dba		04Og03	
	9144.7	60.9			0.2	o			Dba		12Og02	
	9126.4	30.4			0.6	o			Dba		13Og01	*
	9187.3	30.4			−0.5	8			GSa		13Ru11	*
	9159	17			0.1	8			Dba		22Og08	*
	9154.9	60.9			0.1	8			Bka		18Ga34	
^{272}Hs(α)^{268}Sg	9772.0	21.3				6			Dba		23Og03	
^{272}Rg(α)^{268}Mt	10981.9	20.	11197	13	10.8	B			GSa		95Ho04	*
	11191.9	20.			0.3	9			GSa		02Ho11	*
	11184.7	35.			0.4	9			RIa		04Mo26	*
	11207.1	20.3			−0.5	9			Bka		04Fo08	*
$*^{272}$Bh(α)^{268}Db	Average 9.16(2) 9.20(5) 9.18(5) 9.16(5)										15Ga24	**
$*^{272}$Bh(α)^{268}Db^{p}	Average E_α of 8 events; uncertainty increased										13Ru11	**
$*^{272}$Bh(α)^{268}Db^{p}	Unweighted average E_α of 23 events; unc=smallest experimental unc										22Og08	**
$*^{272}$Rg(α)^{268}Mt	B : one event only; E(K) in coincidence may explain disagreement										GAu	**
$*^{272}$Rg(α)^{268}Mt	Two events E_α=11008 and 11046 keV										02Ho11	**
$*^{272}$Rg(α)^{268}Mt	Also others up to E_α=11560 keV										04Mo26	**
$*^{272}$Rg(α)^{268}Mt	One event only										04Fo08	**
^{273}Hs(α)^{269}Sg	9732.9	40.	9650	50	−1.3	o					10El06	
	9671.9	40.6			−0.3	o			Dba		15Ut02	
	9670	40			−0.3	o			SRv		16Ho09	
	9670	40			−0.3	o			SRv		17Og01	
	9650	40			0.0	o			Dba		18Ut02	
	9650.7	17.2				8			DJa		25Og03	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{273}Ds(α)^{269}Hs^{p}	9875.0	20.	11093	20	22.6	F			GSa		96Ho13	*
	11194.3	81.2			−1.1	U			RIa		13Su04	
	11092.8	20.3				10			Dba		24Og02	
^{273}Ds^{m}(α)^{269}Hs^{m}	11248.1	30.	11330	40	1.4	F			GSa		96Ho13	*
	11519.1	60.			−2.4	F			Dba		96La12	*
	11366.9	20.3			−0.7	9			GSa		02Ho11	*
	11311.0	70.			0.3	U			RIa		08Mo09	
	11265.4	71.0			0.9	9			Dba		24Og02	
*^{273}Ds(α)^{269}Hs^{p}	F : this event is distrusted, see reference										02Ho11	**
*^{273}Ds^{m}(α)^{269}Hs^{m}	F : event number 2, probably to excited state in ^{269}Hs										GAu	**
*^{273}Ds^{m}(α)^{269}Hs^{m}	F : this event is distrusted, see reference;										02Ho11	**
*^{273}Ds^{m}(α)^{269}Hs^{m}	And one E_α=11080 keV										02Ho11	**
^{274}Bh(α)^{270}Db	8930.6	101.5	8940	60	0.1	o			Dba		11Og04	*
	8890.0	50.7			0.7	o			Dba		13Og04	
	8971.2	30.5			−0.5	o			GSa		14Kh04	
	8940	60				6			SRv		17Og01	
^{274}Mt(α)^{270}Bh^{p}	9904.8	101.5				7			Dba		07Og02	
^{274}Rg(α)^{270}Mt	11477.9	70.				9			RIa		08Mo09	*
*^{274}Bh(α)^{270}Db	All results from this work were first published in reference										10Og01	**
*^{274}Bh(α)^{270}Db	E_α=8.73−8.84 MeV, unc. increased from 30 to 60 keV for "O" state										17Og01	**
*^{274}Rg(α)^{270}Mt	Also one E_α=11150(70) keV										08Mo09	**
^{275}Hs(α)^{271}Sg	9437.5	71.0	9460	50	0.3	o			Dba		04Og12	
	9437.5	60.9			0.3	o			Dba		06Og05	
	9450	15			0.2	o			SRv		16Ho09	
	9450	20			0.2	o			SRv		17Og01	
	9461	15				11			Dba		22Og05	*
^{275}Mt(α)^{271}Bh	10482.8	90.	10490	60	0.1	o			Dba		04Og03	
	10503.0	60.			−0.2	o			Dba		12St.A	
	10482.7	10.1			0.1	9			GSa		13Ru11	
	10498	21			−0.1	9			Dba		22Og08	
^{275}Ds(α)^{271}Hs^{m}	11366	20	11200#	300#	−3.1	D			Dba		24Og02	*
*^{275}Hs(α)^{271}Sg	includes data from previous experiments										22Og05	**
*^{275}Ds(α)^{271}Hs^{m}	Trends in Mass Surface TMS suggest ^{275}Ds 166 keV less bound										FGK267	**
^{276}Mt(α)^{272}Bh	10253	65	10155	17	−1.5	o			Dba		04Og03	*
	10212	84			−0.7	o			Dba		12Og02	*
	10160	16			−0.3	o			Dba		13Og01	*
	10102.3	10.1			5.2	o			GSa		13Ru11	*
	10093.0	20.			3.1	o			Bka		15Ga24	*
	10100	10			5.5	o			SRv		17Og01	
	10155	17				8			Dba		22Og08	*
^{276}Ds(α)^{272}Hs	10904.3	28.4				7			Dba		23Og03	
*^{276}Mt(α)^{272}Bh	E_α=9710(60) to 434 and 363 keV levels										13Ru11	**
*^{276}Mt(α)^{272}Bh	E_α=9670(80) to 434 and 363 keV levels, and one 9260(64) keV										12Og02	**
*^{276}Mt(α)^{272}Bh	Average for 18 events E_α=9622(16) to 434 and 363 keV levels										13Ru11	**
*^{276}Mt(α)^{272}Bh	E_α=9530(10) to 434 keV level; also E_α=9600(10) to 362 keV level										13Ru11	**
*^{276}Mt(α)^{272}Bh	Average of 9 events E_α=9570−9630 =9590 to 362 keV level										15Ga24	**
*	also one event E_α=9890(20) to 60 keV level										15Ga24	**
*^{276}Mt(α)^{272}Bh	E_α=8.52−10.01 MeV										17Og01	**
*^{276}Mt(α)^{272}Bh	Unweighted average for 21 events E_α=9580(17) to 434-keV level										13Og01	**
^{277}Ds(α)^{273}Hs^{p}	10725.2	40.	10693	17	−0.5	o					10El06	
	10704.8	40.6			−0.2	o			Dba		15Ut02	
	10720	40			−0.4	o			SRv		16Ho09	
	10710	40			−0.3	o			SRv		17Og01	
	10700	40			−0.1	o			Dba		18Ut02	
	10692.7	17.3				10			DJa		25Og03	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
^{277}Cn(α)^{273}Ds	11622.2	30.				11			GSa		96Ho13	*
	11821.0	30.	11620	60	−3.4	F			GSa		96Ho13	*
	11486.2	40.6			2.1	C			RIa		04MoZU	
	11486.2	40.			2.1	B			RIa		08Mo09	*
	11232.4	81.2			4.1	B			RIa		13Su04	
*^{277}Cn(α)^{273}Ds	And one E_α=11170(20) Q_α=11334.(20.) keV										02Ho11	**
*^{277}Cn(α)^{273}Ds	F : this event is distrusted, see reference										02Ho11	**
*^{277}Cn(α)^{273}Ds	And one E_α=11090(70) keV										08Mo09	**
^{278}Mt(α)^{274}Bh^{p}	9689.7	190.	9580	30	−0.6	U			Dba		11Og04	
	9527.3	71.			0.6	o			Dba		12Og06	
	9689.6	30.4			−1.9	o			Dba		13Og04	
	9588.2	30.			−0.1	o			GSa		14Kh04	
	9580	30				8			SRv		17Og01	*
^{278}Rg(α)^{274}Mt	10846.3	81.2	10840	50	−0.1	o			Dba		07Og02	
	10840	22				8			Dba		22Og08	
^{278}Nh(α)^{274}Rg	11850.8	40.	11990	80	2.2	o			RIa		08Mo09	*
	11992.8	61.				10			RIa		12Mo25	
*^{278}Mt(α)^{274}Bh^{p}	E_α=9.38−9.55 MeV										17Og01	**
*^{278}Nh(α)^{274}Rg	Also one E_α=11520(40) keV										08Mo09	**
^{279}Ds(α)^{275}Hs^{p}	9841.3	60.9	9830	60	−0.2	o			Dba		04Og12	
	9841.3	60.9			−0.2	o			Dba		06Og05	
	9847	15			−0.3	o			Dba		06Og05	
	9850	20			−0.4	o			Dba		06Og05	
	9827	15			0.0	13			Dba		22Og05	*
	9831	10			−0.0	13			Bka		26Ga05	*
^{279}Rg(α)^{275}Mt^{p}	10521.1	162.3	10440	50	−0.5	o			Dba		04Og03	
	10530	160			−0.5	U			SRv		17Og01	
	10442	21				11			Dba		22Og08	
*^{279}Ds(α)^{275}Hs^{p}	includes data from 06Og05 and 04Og12										22Og05	**
*^{279}Ds(α)^{275}Hs^{p}	includes data from previous experiments										26Ga05	**
^{280}Rg(α)^{276}Mt^{p}	10128.6	60.	10145	10	0.3	o			Dba		04Og03	
	10128.6	60.			0.3	U			Dba		12Og02	
	10148.8	10.2			−0.4	o			GSa		13Ru11	
	10088.0	15.2			3.8	o			Dba		13Og01	
	10148.8	10.2			−0.4	10			Bka		15Ga24	
	10080	60			1.1	o			Bka		18Ga34	
	10119	27			1.0	10			Dba		22Og08	
^{281}Ds(α)^{277}Hs^{p}	8885.5	20.3				5			GSt		23Co04	*
	8957.8	180.	8885	20	−0.4	F			Dba		99Og10	*
	8825.9	100.			0.5	F			Dba		04Mo15	*
	8856.3	30.4			0.5	o			GSt		10Du06	
	8853.3	25.			0.6	o			GSt		11Ga19	
	8853.3	15.			0.6	o			GSa		12Ho12	
	8853	30			0.6	U			SRv		16Ho09	
	8850	30			0.6	o			SRv		17Og01	
^{281}Ds^{m}(α)^{277}Hs^{m}	9449.8	15.				5			GSa		12Ho12	*
^{281}Rg(α)^{277}Mt	9454.8	40.6	10020#	400#	8.8	o			Dba		13Og04	
	9410	50			8.6	D			SRv		17Og01	*
^{281}Cn(α)^{277}Ds	10459.2	40.	10420	50	−0.6	o					10El06	
	10448.9	40.6			−0.5	o			Dba		15Ut02	
	10450	40			−0.5	o			SRv		16Ho09	
	10450	40			−0.5	o			SRv		17Og01	
	10430	40			−0.2	o			Dba		18Ut02	
	10418	17			−0.0	o			DJa		25Og03	
	10418	17				11			Dba		26Og02	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{281}$Ds(α)^{277}Hs^{p}	E_α=8670(20) keV to level at 90 keV										23Co04	**
$*^{281}$Ds(α)^{277}Hs^{p}	F : wrong α chain, see ^{285}Cn and ^{289}Fl										GAu	**
$*^{281}$Ds(α)^{277}Hs^{p}	F : non traceable information										GAu	**
$*^{281}$Ds^{m}(α)^{277}Hs^{m}	Assignment of ^{293}Lvm-^{289}Flm-^{285}Cnm-^{281}Dsm-^{277}Hsm chain is tentative										12Ho12	**
$*^{281}$Rg(α)^{277}Mt	Trends in Mass Surface TMS suggest ^{281}Rg 610 keV less bound										HWJ266	**
^{282}Mt(α)^{278}Bh	8960	180	8660#	200#	−1.6	D			SRv		16Ho09	*
^{282}Rg(α)^{278}Mt^{p}	9129.7	101.4	9160	30	0.3	o			Dba		11Og04	
	9139.8	50.7			0.3	o			Dba		13Og04	
	9180.4	30.4			−0.3	o			GSa		14Kh04	*
	9160	30				10			SRv		17Og01	*
^{282}Cn(α)^{278}Ds	9667.4	100.	10150#	200#	4.3	F			Dba		04Mo15	*
^{282}Nh(α)^{278}Rg	10783.2	81.2	10780	50	−0.0	o			Dba		07Og02	
	10782	22				9			Dba		22Og08	
$*^{282}$Mt(α)^{278}Bh	re-interpreted results from ^{99}Og$_{10}$										16Ho09	**
$*^{282}$Mt(α)^{278}Bh	Trends in Mass Surface TMS suggest ^{282}Mt 300 keV more bound										GAu	**
$*^{282}$Rg(α)^{278}Mt^{p}	also observed E_α=8860(30) keV										14Kh04	**
$*^{282}$Rg(α)^{278}Mt^{p}	E_α=8.86−9.05 MeV										17Og01	**
$*^{282}$Cn(α)^{278}Ds	F : non traceable information										GAu	**
^{283}Cn(α)^{279}Ds^{p}	9677	70	9670	50	−0.1	o			Dba		04Og07	
	9677.0	60.9			−0.1	o			Dba		04Og12	
	9677.0	60.9			−0.1	o			Dba		06Og05	
	9606.0	60.9			0.8	o					07Ei02	
	9656.7	15.2			0.2	o			GSa		07Ho18	
	9788.6	100.			−1.1	o			Bka		09St21	
	9658	15			0.2	o			SRv		16Ho09	
	9660	20			0.1	o			SRv		17Og01	
	9585.5	51.			1.1	o			RIa		17Ka31	
	9668	15			−0.0	o			Dba		22Og05	*
	9667	10				15			Bka		26Ga05	*
^{283}Nh(α)^{279}Rg^{p}	10265.4	90.	10360	60	0.9	o			Dba		04Og03	
	10376.9	10.1			−0.3	13			GSa		13Ru11	
	10346	25			0.3	13			Dba		22Og08	
$*^{283}$Cn(α)^{279}Ds^{p}	includes data from previous experiments										22Og05	**
$*^{283}$Cn(α)^{279}Ds^{p}	includes data from 22Og05, 17Ka31, 17Og01, 16Ho09, 09St21,										26Ga05	**
$*^{283}$Cn(α)^{279}Ds^{p}	07Ho18, 07Ei02, 06Og05, 04Og12 and 04Og07										26Ga05	**
^{284}Cn(α)^{280}Ds	9301.3	50.	9670#	150#	7.4	F			Dba		01Og01	*
	9269.0	100.			4.0	F			Dba		04Mo15	*
	9869	193			−1.0	D			RIa		17Ka66	*
	9463.5	10.1			20.4	D			GSt		21Sa01	*
^{284}Nh(α)^{280}Rg	10254.6	20.3				11			GSa		13Ru11	*
	10305.3	20.	10250	50	−0.9	o			Bka		15Ga24	*
	10120	50			2.7	B			SRv		17Og01	*
^{284}Nh(α)^{280}Rg^{p}	10143.1	60.9	10086	16	−0.7	o			Dba		04Og03	
	10112.6	50.			−0.4	o			Dba		12Og02	
	10097.4	15.			−0.2	o			Dba		13Og01	
	10072.1	60.9			0.2	12			Bka		18Ga34	
	10087.3	16.2			−0.1	12			Dba		22Og08	*
^{284}Fl(α)^{280}Cn	10721.3	17.2				9			DJa		25Og03	
$*^{284}$Cn(α)^{280}Ds	F : no α observed in later work by same group; re-assigned to ^{285}Cn										04Og07	**
$*^{284}$Cn(α)^{280}Ds	F : non traceable information										GAu	**
$*^{284}$Cn(α)^{280}Ds	Alternatively ^{284}Cn could decay by SF										17Ka66	*W
$*^{284}$Cn(α)^{280}Ds	Trends in Mass Surface TMS suggest ^{284}Cn 200 keV more bound										GAu	**
$*^{284}$Cn(α)^{280}Ds	from 1 decay chain ^{288}Fl(α)^{284}Cn(α)^{280}Ds(sf) event										21Sa01	**
$*^{284}$Cn(α)^{280}Ds	Trends in Mass Surface TMS suggest ^{284}Cn 200 keV less bound										GAu	**
$*^{284}$Nh(α)^{280}Rg	Highest E_α=10110(10), error increased										13Ru11	**
$*^{284}$Nh(α)^{280}Rg	Highest E_α=10160(20)										15Ga24	**
$*^{284}$Nh(α)^{280}Rg	E_α=9.10−10.11 MeV; unc. increased from 10 to 50 keV for "O" state										17Og01	**
$*^{284}$Nh(α)^{280}Rg	Trends in Mass Surface TMS disagrees with the Superheavy Review SRv										GAu	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{284}$Nh(α)^{280}Rg^{p}	Also E_α=9886(62) keV										12Og02	*G
$*^{284}$Nh(α)^{280}Rg^{p}	Average E_α of 21 events										13Og01	*G
$*^{284}$Nh(α)^{280}Rg^{p}	Unweighted average E_α of 27 focal-plane events										22Og08	**
^{285}Cn(α)^{281}Ds^{p}	8793.7	50.	9396	19	8.5	F			Dba		99Og10	*
	8793.7	100.			5.4	F			Dba		04Mo15	*
	9290.6	60.9			1.3	o			Dba		04Og07	
	9280.5	50.			1.6	o			Dba		07Og01	
	9341.3	30.4			0.9	o			GSt		10Du06	
	9341.3	30.4			0.9	o			GSt		11Ga19	
	9314.9	15.			1.5	o			GSa		12Ho12	
	9320	15			1.4	o			SRv		16Ho09	
	9320	20			1.4	o			SRv		17Og01	
	9320.8	30.4			1.3	7			RIa		17Ka66	
	9394.4	20.3			0.1	o			GSt		23Co04	*
	9404.5	20.3			−0.4	7			Bka		26Ga05	*
^{285}Cn^{m}(α)^{281}Ds^{m}	9845.4	15.2				6			GSa		12Ho12	*
^{285}Nh(α)^{281}Rg	9878.9	81.1	10010	60	1.4	B			Dba		11Og04	*
	10026.9	62.			−0.2	o			Dba		12Og02	*
	9767.3	183.			1.3	B			Dba		12Og06	
	10031.0	162.			−0.1	o			Dba		13Og01	
	9990.4	41.			0.3	o			Dba		13Og04	
	10008.7	20.3			0.0	o			Dba		16Fo16	*
	10010	40			−0.0	o			SRv		17Og01	*
	10010	33				11			Dba		22Og08	
^{285}Fl(α)^{281}Cn	10558.4	50.7	10600	50	0.6	o			Dba		15Ut02	
	10560	50			0.6	o			SRv		16Ho09	
	10560	50			0.6	o			SRv		17Og01	
	10560	50			0.6	o			Dba		18Ut02	
	10599.1	17.2			−0.0	o			DJa		25Og03	
	10599.1	17.2				12			Dba		26Og02	
$*^{285}$Cn(α)^{281}Ds^{p}	F : one event at 15.4 m, later work yields much shorter half-lives										GAu	**
$*^{285}$Cn(α)^{281}Ds^{p}	F : non traceable information										GAu	**
$*^{285}$Cn(α)^{281}Ds^{p}	E_α=9140(20) keV to level at 124 keV										23Co04	**
$*^{285}$Cn(α)^{281}Ds^{p}	combined result, E_α=9150(20) keV to level at 124 keV; analysis includes										26Ga05	**
$*^{285}$Cn(α)^{281}Ds^{p}	29 events from previous experiments (see 23Co04)										26Ga05	**
$*^{285}$Cn(α)^{281}Ds^{p}	and 2 events from 26Ga05										26Ga05	**
$*^{285}$Cn^{m}(α)^{281}Ds^{m}	Assignment of ^{293}Lvm-^{289}Flm-^{285}Cnm-^{281}Dsm-^{277}Hsm chain is tentative										12Ho12	**
$*^{285}$Nh(α)^{281}Rg	And E_α=9480(110) keV										11Og04	**
$*^{285}$Nh(α)^{281}Rg	Also E_α=9740(80) and 9480(110) keV										12Og02	**
$*^{285}$Nh(α)^{281}Rg	reanalyzed data of 11Og04, 12Og02, 12Og06, 13Og01 and 13Og04										16Fo16	**
*	unweighted average of the 4 highest energy events as selected by										GAu	**
*	the evaluator: s06=9857 s11=9902 s12=9845 s13=9867 in Table 2										GAu	**
$*^{285}$Nh(α)^{281}Rg	E_α = 9.47−10.18 MeV										17Og01	**
^{286}Rg(α)^{282}Mt	8793	45				4			SRv		16Ho09	
^{286}Cn(α)^{282}Ds	9432.1	40.6	9280#	100#	−3.7	D			GSt		23Co04	*
^{286}Nh(α)^{282}Rg	9766.9	100.	9790	50	0.2	o			Dba		11Og04	
	9817.5	111.6			−0.2	o			Dba		12Og06	
	9432.1	304.3			1.2	F			GSa		14Kh04	*
	9790	50				11			SRv		17Og01	*
^{286}Fl(α)^{282}Cn	10142.1	100.	10325	20	1.8	F			Dba		04Mo15	*
	10172.5	314.4			0.5	o			Dba		04Og07	*
	10345	60			−0.3	o			Dba		04Og12	*
	10334.7	60.9			−0.2	o			Dba		06Og05	*
	10375.4	40.			−1.3	o			Bka		09St21	
	10456.5	100.			−1.3	o			Bka		10El06	
	10355.1	40.6			−0.8	o			SRv		16Ho09	
	10355.1	40.6			−0.8	o			SRv		17Og01	*
	10334.7	30.2			−0.3	o			GSt		21Sa01	*
	10335.8	10.1			−1.1	o			Dba		22Og05	*
	10324.6	20.3				11			Bka		24Ga31	*
^{286}Mc(α)^{282}Nh	10862.2	20.3				10			Dba		22Og08	

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{286}$Cn(α)^{282}Ds	Assignment to ^{286}Cn(α) made in 23Sa03										23Co04	**
$*^{286}$Cn(α)^{282}Ds	Trends in Mass Surface TMS suggest ^{286}Cn 150 keV more bound										FKo265	**
$*^{286}$Nh(α)^{282}Rg	Only partial energy										19Kh04	**
$*^{286}$Nh(α)^{282}Rg	E_α = 9.61−9.75 MeV										17Og01	**
$*^{286}$Fl(α)^{282}Cn	F : non traceable information										GAu	**
$*^{286}$Fl(α)^{282}Cn	including previous data from Dubna										04Og07	**
$*^{286}$Fl(α)^{282}Cn	including previous data from Dubna										04Og12	**
$*^{286}$Fl(α)^{282}Cn	E_α=10.16(9) MeV plus data from 04Og12 and 04Og07										06Og05	**
$*^{286}$Fl(α)^{282}Cn	from the compilation of all the preceding published data										16Ho09	*W
$*^{286}$Fl(α)^{282}Cn	from the compilation of all the preceding published data										17Og01	*W
$*^{286}$Fl(α)^{282}Cn	E_α=10.16(1) MeV; final result includes data from 12Og06, 10El06										21Sa01	**
$*^{286}$Fl(α)^{282}Cn	includes data from 21Sa01, 18Br13, 12Og06, 10El06,09St21,										22Og05	**
$*^{286}$Fl(α)^{282}Cn	06Og05, 04Og07 and 04Og12										22Og05	**
$*^{286}$Fl(α)^{282}Cn	for 56 events, including all the events from previous publications										24Ga31	**
^{287}Cn(α)^{283}Ds	9168.1	60.9				4			GSt		23Co04	*
^{287}Fl(α)^{283}Cn	10435.7	20.3	10160	50	−5.1	F			Dba		99Og07	*
	10182.2	70.			−0.2	o			Dba		04Og07	
	10161.9	60.8			0.0	o			Dba		04Og12	
	10161.9	60.8			0.0	o			Dba		06Og05	
	10166.8	15.2			−0.1	o			SRv		16Ho09	
	10170	20			−0.1	o			SRv		17Og01	
	10158	15			0.1	o			Dba		22Og05	*
	10162.0	10.1				16			Bka		26Ga05	*
^{287}Mc(α)^{283}Nh	10740	90	10710	50	−0.3	o			Dba		04Og03	
	10740	60			−0.4	o			Dba		12St.A	
	10780.5	50.7			−1.0	o			GSa		13Ru11	
	10760	50			−0.7	o			SRv		17Og01	
	10707	21				14			Dba		22Og08	
$*^{287}$Cn(α)^{283}Ds	E_α observed in 23Co04; assignment to ^{287}Cn(α) is tentative										23Co04	**
$*^{287}$Fl(α)^{283}Cn	F : 2 evts at 1.32 s, 14.4 s, later work yields T=1.7 s										GAu	**
$*^{287}$Fl(α)^{283}Cn	includes data from 16Ho09, 15Og05, 12Ho12, 10Ei01,										22Og05	**
$*^{287}$Fl(α)^{283}Cn	09St21, 08Ei03, 07Ho18, 06Og05, 04Og07 and 04Og12										22Og05	**
$*^{287}$Fl(α)^{283}Cn	includes data from 22Og05, 17Og01, 16Ho09, 06Og05,										26Ga05	**
$*^{287}$Fl(α)^{283}Cn	04Og12, 04Og07 and 99Og07										26Ga05	**
^{288}Fl(α)^{284}Cn	9968.8	50.	10064	9	1.9	F			Dba		01Og01	*
	9958.8	100.			1.1	F			Dba		04Mo15	*
	10090.3	80.			−0.3	o			Dba		04Og07	
	10090.3	70.			−0.4	o			Dba		04Og12	
	10080.2	60.8			−0.3	U			Dba		07Og01	
	10090.3	30.			−0.9	o			GSt		10Du06	
	10090.3	30.			−0.9	o			GSt		11Ga19	
	10067.0	15.			−0.2	o			GSa		12Ho12	
	10074.1	15.2			−0.7	o			SRv		16Ho09	
	10070	30			−0.2	o			SRv		17Og01	
	10080.2	20.3			−0.8	11			RIa		17Ka66	
	10059.9	10.1			0.4	o			GSt		21Sa01	*
	10060.0	10.1			0.4	11			Bka		26Ga05	*
^{288}Mc(α)^{284}Nh	10607.6	60.8	10580	50	−0.3	o			Dba		04Og03	
	10627.8	60.			−0.6	o			Dba		12Og02	
	10727.2	82.1			−1.5	o			Dba		13Og01	*
	10698.8	10.1			−2.3	o			GSa		13Ru11	*
	10754.6	20.3			−3.2	o			Bka		15Ga24	*
	10650	50			−1.3	o			SRv		17Og01	*
	10440	60			1.8	o			Bka		18Ga34	
	10583	16				12			Dba		22Og08	*
^{288}Lv(α)^{284}Fl	11240.3	15.2				10			DJa		25Og03	
$*^{288}$Fl(α)^{284}Cn	F : T=1800(+2100–600) ms, later work yields shorter half-lives										GAu	**
*	re-assigned to ^{289}Fl										04Og07	**
$*^{288}$Fl(α)^{284}Cn	F : non traceable information										GAu	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{288}$Fl(α)^{284}Cn	E_α=9.92(1) MeV; analysis includes 32 events from 22Ya.1, 17Ka66,										21Sa01	**
$*^{288}$Fl(α)^{284}Cn	14Ya33, 12Ho12, 11Ga19, 10Du06, 04Og.A, 04Og12 , 04Og07										21Sa01	**
$*^{288}$Fl(α)^{284}Cn	and 11 events from 21Sa01										21Sa01	**
$*^{288}$Fl(α)^{284}Cn	E_α=9.92(1) MeV; analysis includes 43 events from previous										26Ga05	**
$*^{288}$Fl(α)^{284}Cn	experiments (see 21Sa01) and 6 events from 26Ga05										26Ga05	**
$*^{288}$Mc(α)^{284}Nh	Highest E_α=10578(81); average 24 events would yield E_α=10480 Q_α=10627.8										13Og01	**
$*^{288}$Mc(α)^{284}Nh	Average of 2 highest E_α=10560(10) 10540(10)										13Ru11	**
$*^{288}$Mc(α)^{284}Nh	Average of 2 highest E_α=10610(20) 10600(20)										15Ga24	**
$*^{288}$Mc(α)^{284}Nh	E_α=10.21−10.67 MeV, Q_α=10.65(0.01) MeV in reference; unc. increased										17Og01	**
$*^{288}$Mc(α)^{284}Nh	Unweighted average E_α of 31 focal-plane events										22Og08	**
^{289}Fl(α)^{285}Cn	9846.5	50.7	9940	70	1.3	F			Dba		99Og10	*
	9846.5	101.4			0.8	F			Dba		04Mo15	*
	9958.1	60.			−0.3	o			Dba		04Og07	
	9958.1	50.			−0.3	o			Dba		07Og01	
	10008.8	30.			−1.3	o			GSt		10Du06	
	10008.8	30.			−1.3	o			GSt		11Ga19	
	9956.0	15.2			−0.4	o			GSa		12Ho12	
	9974	15			−0.7	o			SRv		16Ho09	
	9980	20			−0.8	o			SRv		17Og01	
	9917.3	50.7			0.3	8			RIa		17Ka66	
	9947.9	30.2			−0.2	8			GSt		23Co04	*
^{289}Fl^m(α)^{285}Cn^m	10169.9	15.				7			GSa		12Ho12	*
^{289}Mc(α)^{285}Nh	10455.0	90.	10500	50	0.5	o			Dba		11Og04	
	10522.8	62.			−0.3	o			Dba		12Og02	*
	10404.2	71.			1.1	o			Dba		12Og06	
	10546.2	81.			−0.5	o			Dba		13Og01	
	10607.0	152.			−0.7	o			Dba		13Og04	
	10510.7	20.3			−0.2	o			Dba		16Fo16	*
	10490	50			0.2	o			SRv		17Og01	*
	10502	18				12			Dba		22Og08	*
^{289}Lv(α)^{285}Fl	11057.3	15.2	11060	50	0.0	o			DJa		25Og03	
	11059.4	13.2				13			Dba		26Og02	
$*^{289}$Fl(α)^{285}Cn	F : one event at 30.4 s, later work yields much shorter half-lives										GAu	**
$*^{289}$Fl(α)^{285}Cn	F : non traceable information										GAu	**
$*^{289}$Fl(α)^{285}Cn	E_α=9800(20) keV to level at 30 keV										23Co04	**
$*^{289}$Fl^m(α)^{285}Cn^m	Assignment of ^{293}Lvm-^{289}Flm-^{285}Cnm-^{281}Dsm-^{277}Hsm chain is tentative										12Ho12	**
$*^{289}$Mc(α)^{285}Nh	Also E_α=10310(90) keV										12Og02	**
$*^{289}$Mc(α)^{285}Nh	reanalyzed data of 11Og04, 12Og02, 12Og06, 13Og01 and 13Og04										16Fo16	**
*	unweighted average of the 3 highest energy events selected by										GAu	**
*	the evaluator: s06=10370 s12=10364 s14=10362 in Table 2										GAu	**
$*^{289}$Mc(α)^{285}Nh	E_α=10.15−10.54 MeV										17Og01	**
$*^{289}$Mc(α)^{285}Nh	average of 8 events from 22Og07 and 22Og08										FGK239	**
^{290}Nh(α)^{286}Rg	9846	45	9250#	300#	−8.9	D			SRv		16Ho09	*
^{290}Fl(α)^{286}Cn	9856.2	30.4				5			RIa		17Ka66	*
	10130	41	9860	30	−6.7	B			GSt		23Co04	*
^{290}Mc(α)^{286}Nh	10454.4	40.6	10410	40	−0.7	o			GSa		14Kh04	
	10410	40				12			SRv		17Og01	*
^{290}Lv(α)^{286}Fl	10920.9	100.	10990	70	0.6	F			Dba		04Mo15	*
	11002.0	81.1			−0.1	o			Dba		04Og07	
	10991.8	81.1			−0.0	o			Dba		06Og05	
	10999.9	65.9			−0.1	o			Dba		12Og06	
	11002.0	60.8			−0.2	o			SRv		16Ho09	
	11002.0	71.0			−0.1	o			SRv		17Og01	
	11013.2	142.0			−0.2	o			Dba		18Br13	
	10952.3	178.5			0.2	o			Dba		20Vo07	*
	10991.9	71.0				12			Bka		24Ga31	*
$*^{290}$Nh(α)^{286}Rg	Authors say assignment is tentative										16Ho09	**
$*^{290}$Nh(α)^{286}Rg	Trends in Mass Surface TMS suggest ^{290}Nh 580 keV more bound										GAu	**
$*^{290}$Fl(α)^{286}Cn	Alternatively it could be ^{289}Fl(α)^{285}Cn										17Ka66	**
$*^{290}$Fl(α)^{286}Cn	Assignment to ^{290}Fl(α) made in 23Sa03										23Co04	**

Table I. Comparison of input data and adjusted values (continued, Explanation of Table on page 120002-32)

Item	Input value		Adjusted value		v_i	Dg	Signf.	Main infl.	Lab	F	Reference	
$*^{290}$Mc(α)^{286}Nh	E_α=9.78−10.31 MeV										17Og01	**
$*^{290}$Lv(α)^{286}Fl	F : non traceable information										GAu	**
$*^{290}$Lv(α)^{286}Fl	Same event as in 18Br13										20Vo07	**
$*^{290}$Lv(α)^{286}Fl	for 18 events, including all the events from previous publications										24Ga31	**
^{291}Fl(α)^{287}Cn	9703.7	30.4				5			GSt		23Co04	*
^{291}Lv(α)^{287}Fl	10890	70	10890	90	−0.0	o			Dba		04Og07	
	10890	70			−0.0	o			Dba		06Og05	
	10890	70			−0.0	o			SRv		16Ho09	
	10890	70				17			SRv		17Og01	
$*^{291}$Fl(α)^{287}Cn	E_α observed in 23Co04; assignment to ^{291}Fl(α) is tentative										23Co04	**
^{292}Lv(α)^{288}Fl	10707.0	50.	10791	12	1.7	F			Dba		01Og01	*
	10676.5	100.			1.1	F			Dba		04Mo15	*
	10808.3	71.0			−0.2	U			Dba		04Og12	
	10772.8	15.2			1.2	o			GSa		12Ho12	
	10775.9	15.2			1.0	12			SRv		16Ho09	
	10777.9	20.3			0.7	o			SRv		17Og01	
	10818.4	20.3			−1.3	12			RIa		17Ka66	
$*^{292}$Lv(α)^{288}Fl	F : daughter and grand-daughter re-assigned to ^{289}Fl and ^{285}Cn										GAu	**
$*^{292}$Lv(α)^{288}Fl	F : non traceable information										GAu	**
^{293}Lv(α)^{289}Fl	10676	60	10680	60	0.0	o			Dba		04Og07	
	10686.1	60.8			−0.1	o			Dba		07Og01	
	10679.0	15.2			−0.0	o			GSa		12Ho12	
	10705	15			−0.5	9			SRv		16Ho09	
	10710	20			−0.6	o			SRv		17Og01	
	10635.2	40.6			0.7	9			RIa		17Ka66	
^{293}Lv^m(α)^{289}Fl^m	10647.6	15.2				8			GSa		12Ho12	*
^{293}Ts(α)^{289}Mc	11182.9	81.1	11310	40	1.3	o			Dba		11Og04	*
	10949.7	71.			4.1	B			Dba		12Og06	*
	11253.9	91.2			0.5	o			Dba		13Og04	*
	11293.4	20.3			0.3	13			Dba		16Fo16	*
	11320	50			−0.2	13			SRv		17Og01	*
$*^{293}$Lv^m(α)^{289}Fl^m	Assignment of ^{293}Lvm-^{289}Flm-^{285}Cnm-^{281}Dsm-^{277}Hsm chain is tentative										12Ho12	**
$*^{293}$Ts(α)^{289}Mc	reanalyzed data of 11Og04, 12Og02, 12Og06, 13Og01 and 13Og04										16Fo16	**
*	unweighted average of the 8 highest energy events selected by										GAu	**
*	the evaluator: s02=11140 s03=11080 s07=11142 s08=11114 s12=11183										GAu	**
*	s13=11203 s14=11059 s16=11190 in Table 2										GAu	**
$*^{293}$Ts(α)^{289}Mc	E_α=10.60−11.20 MeV										17Og01	**
^{294}Ts(α)^{290}Mc	11202.6	40.6	11180	40	−0.4	o			GSa		14Kh04	
	11180	40				13			SRv		17Og01	*
^{294}Og(α)^{290}Lv	11800.9	100.	11870	30	0.7	F			Dba		04Mo15	*
	11810.9	60.			0.9	o			Dba		04Og12	
	11810.9	60.			0.9	o			Dba		06Og05	
	11840.3	65.9			0.4	o			Dba		12Og06	
	11821.1	60.8			0.8	o			SRv		16Ho09	
	11821.1	60.8			0.8	13			SRv		17Og01	
	11878.9	30.4			−0.4	o			Dba		18Br13	
	11883.9	36.5			−0.5	13			Dba		20Vo07	*
$*^{294}$Ts(α)^{290}Mc	E_α=10.81−11.07 MeV										17Og01	**
$*^{294}$Og(α)^{290}Lv	F : non traceable information										GAu	**
$*^{294}$Og(α)^{290}Lv	Same event as in 18Br13										20Vo07	**
^{295}Og(α)^{291}Lv	11810.4	71.0	11700#	200#	−1.3	F			Dba		04Og05	*
$*^{295}$Og(α)^{291}Lv	F: Reassigned to ^{294}Og in later work										04Og12	**